# **BMJ Open** Retrospective study of COVID-19 outcomes among healthcare workers in **Rivers State, Nigeria**

Chidinma Eze-Emiri <sup>1,2</sup> Foster Patrick,<sup>1,2</sup> Ezinne Igwe,<sup>3</sup> Golden Owhonda<sup>2</sup>

To cite: Eze-Emiri C, Patrick F, Igwe E, et al. Retrospective study of COVID-19 outcomes among healthcare workers in Rivers State, Nigeria. BMJ Open 2022;12:e061826. doi:10.1136/ bmjopen-2022-061826

Prepublication history for this paper is available online. To view these files, please visit the journal online (http://dx.doi. org/10.1136/bmjopen-2022-061826).

Received 15 February 2022 Accepted 21 October 2022

#### Check for updates

C Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Epidemiology, School of Public Health, University of Port Harcourt, Choba, Rivers State, Nigeria <sup>2</sup>Department of Public Health & Disease Control, Rivers State Ministry of Health, Port Harcourt, Nigeria

<sup>3</sup>University of Wollongong Faculty of Science Medicine and Health, Wollongong, New South Wales, Australia

#### **Correspondence to**

Dr Golden Owhonda: goldenowhonda@yahoo.com

#### ABSTRACT

**Objective** To determine the illness severity and mortality among COVID-19-infected healthcare workers (HCWs). Design A retrospective cohort study using populationlevel data. Secondary analysis was conducted on collated data from the Public Health Emergency Operations Centre (PHEOC) at the State Ministry of Health, Rivers State, Nigeria, Data were gathered from the COVID-19 patient database of the PHEOC on demographics, place of work, illness severity and outcome.

Participants The cohort included all documented HCWs with confirmed COVID-19 infection (diagnosed by PCR). Primary and secondary outcome measures Illness severity defined as 'hospitalisation required' and treatment outcome labelled as 'alive' or 'dead' were the outcomes of interest.

**Results** The mean age was 43 years and 50.5% of the cohort were female. Of the 301 HCWs infected. 187 patients were symptomatic with 32 requiring hospitalisation. Seven infected HCWs died of their COVID-19 infection, resulting in a case fatality ratio (CFR) of 2.3%. Population proportions for age groups, case presentation and mortality, would be significantly greater than those seen in the study population. Health professionals made up 79.7% (240) of the study cohort, with 68.8% (165) of them working at the teaching hospitals; the association between HCWs and health facilities they worked in, was significant. Symptomatic cases were more inclined to progress to severe illness ( $\chi^2_{(1)}$  = 15.219,  $\alpha = < 0.0001$ ; adjusted OR (aOR) 10.658, 95% CI 2.494 to 45.552); patients also had greater odds of dying from COVID-19  $(\chi^2_{(5)} = 13.7, \ \alpha = 0.003; \text{ aOR } 1.079, 95\% \text{ Cl } 1.02$ 

to 1.141) per year increase in age adjusted for sex, case class and illness severity.

Conclusions Frontl-ine HCWs are at an increased risk of exposure to COVID-19 infections. In Nigeria, there is a higher risk of experiencing severe illness if symptomatic while infected with COVID-19. Preventive strategies, proper education and awareness must be put in place to protect HCWs.

Objective To determine the illness severity and mortality among COVID-19-infected HCWs.

# **INTRODUCTION**

Healthcare workers (HCWs) have a higher risk of encountering infectious agents due to

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- $\Rightarrow$  The use of population registry data enabled the representation of the population, giving a snapshot of the burden of COVID-19 on health workers in the study region.
- $\Rightarrow$  It also limited bias due to selection and recall.
- $\Rightarrow$  The use of secondary data also implied that some variables that would have better informed the study were not available.
- $\Rightarrow$  The reliance on reported infections and deaths made it impossible to estimate how many cases were missed by non-reporting.

for uses related their work environments. With the COVID-19 pandemic, front-line HCWs face a higher đ risk of infection and mortality as well as ē being the drivers of community-level infection. Recent evidence shows that compared with individuals in the general community, front-line HCWs have a 12-fold risk of testing positive for COVID-19, with a higher risk observed for workers with inadequate access to personal protective equipment (PPE).<sup>1</sup> In addition to increased exposure to COVID-19 ≥ in the pandemic, Wang *et al*<sup>2</sup> found that poor can increase the risk of nosocomial SARSpossible to extrapolate these results to the Nigerian healthcare setting; the density of the health workforce (1.95-1000 persons) in the country,<sup>3</sup> in an addition to an estimated doctor-to-patient ratio of 1:2753<sup>4</sup> is reportedly 'still very low' to effectively deliver essential health services.<sup>3</sup> the country,<sup>3</sup> in an addition to an estimated

SARS-CoV-2 appears to have tropism for 8 diverse tissues, this underscores the difficulty in predicting the severity of COVID-19. Nevertheless, factors, such as age, comorbidities, immune response, radiographic findings, laboratory markers and indicators of organ dysfunction, might predict worse outcomes independently or collectively.<sup>5</sup> It was suggested that age, gender and the number of comorbidities showed a good

Protected by copyright, including

Symptomatic

Table 1Demogamong healthcare	raphics and characte e workers (n=301)	eristics of COVID-19
Variable	N (%)	z-test (95% CI) p value
Sex		
Male	149 (49.5)	-0.17 (-0.12 to 0.10)
Female	152 (50.5)	H <sub>a</sub> ≠ 0: 0.862
Age	43±11.7*	
20–29	25 (8.3)	†7.61 (0.38 to 0.60)
30–39	118 (39.2)	H <sub>a</sub> ≠ 0: 0.000; H <sub>a</sub> >0: 0.000
40–49	81 (26.9)	
50–59	44 (14.6)	
60–69	23 (7.6)	
>70	10 (3.3)	
‡Case presentation		

187 (62.1)

4.08 (0.13 to 0.35)

Asymptomatic	114 (37.9)	H <sub>a</sub> ≠ 0: 0.000; H <sub>a</sub> >0: 0.000
‡Illness severity (requiring hospitalis	sation)	
Yes	32 (10.6)	-10.75 (-0.90 to 0.68)
No	269 (89.4)	$H_a \neq 0: 0.000; H_a < 0: 0.000$
Contact with probable case		
Yes	108 (35.9)	
No	156 (51.8)	
Non-response/incomplete data	37 (12.3)	
Knowledge of suspected exposure		
Yes	101 (33.6)	
No	157 (52.3)	
Non-response/incomplete data	47 (15.7)	
Exposure		
Church	12 (4)	
Home	12 (4)	
Social event	26 (8.7)	
Workplace	51 (17)	
Ootcome		
Recovered	294 (97.7)	9.94 (0.43 to 1.06)
Dead	7 (2.3)	$H_a \neq 0: 0.000; H_a > 0: 0.000$

\*Mean±SD, case fatality ratio=2.33%.

†Difference in proportion between 20–49 years and ≥50 years

‡At the time of testing.

predictive ability of whether confirmed patients would develop severe disease,<sup>6</sup> and evidence showed that advanced age, male sex, current smoking status, preexisting comorbidities (especially chronic kidney, respiratory and cardiocerebrovascular diseases) were important predictors associated with mortality.<sup>78</sup>

The estimated mortality rate among HCWs attributed to COVID-19 has increased progressively.<sup>9</sup> In May 2020, the total number of reported HCW deaths from 67 countries was 1413. Consequently, it suggests that for every 100 HCWs that got infected, one died—the deaths were also 0.5% of the total number of 270 426 COVID-19 deaths worldwide.<sup>10</sup> In addition, a survey of 37 countries estimated median death of 0.05 HCWs per 100 000 population. A report from the Pan American Health

BMJ Open: first published as 10.1136/bmjopen-2022-061826 on 11 November 2022. Downloaded from Enseignement Superieur (ABE ŝ copyright, including ₫ sesn. ſe a data m http://bmjopen.bmj.com/ on June 12, 2025 at Agence Bibliographique de <u>م</u> Dd <u>0</u>

Organization in collaboration with the WHO stated that approximately 570 000 HCWs got infected with COVID-19 with above 2500 dying from the virus across the American region alone.<sup>11</sup> The WHO also estimated that between 80 000 and 180 000 HCWs died of COVID-19 in the period between January 2020 and May 2021, converging to a medium scenario of 115 500 deaths; although HCWs mortality in the African region was estimated at 2003, it was also acknowledged that several uncertainties and limitations surrounded the measurement of the death toll of HCWs due to COVID-19.<sup>12</sup>

A subnational study highlighting the burden of COVID-19 among HCWs is paramount to understanding the effect of the pandemic on the healthcare workforce of in Nigeria. The study's aim was to determine the illness severity and mortality among COVID-19 infected HCWs in Rivers State, Nigeria.

# METHOD

## **Study location**

The study was conducted in Rivers State, located in the South-South geopolitical zone of Nigeria.

#### Study design and population

The study was a retrospective cohort study using existing population-level records data. The cohort included only HCWs with confirmed COVID-19 infection that were reported to the Public Health Emergency Operations Centre (PHEOC). The HCWs were categorised using the WHO and International Labour Organization International Standard Classification of Occupations.<sup>13</sup> There were five categories based on their roles in patient management and healthcare services:

- Health professionals—medical doctors, nurses, dentists, pharmacists and health safety professionals.
- Health associate professional—all technologists and assistants in health professions, CHWs.
- Personal care workers—healthcare and home-based care workers.
- Health management and support personnel—administrative and management staff, trade workers, social workers and life science professionals.
- Other health service providers—armed forces staff, interns and hospital volunteers.

Furthermore, the health facilities were classified based on services rendered. Hospital classification by services: Teaching hospitals—offering tertiary health services; General hospitals—offering secondary health services; Community hospitals—offering primary and community-based care; specialised outpatient clinics rendering specialty outpatient services like dentistry, radiology and diagnostic services; Corporate/occupational health clinics—ffering general and occupational health services, restricted to employees only and Health-allied organisations.

### Data source

Secondary data were collated from the COVID-19 case investigation form dataset at the PHEOC data centre,

Rivers State Ministry of Health. The data sources included reports from public and private-owned health facilities, containment centres, offshore platforms and other health-allied facilities. The duration of data extraction was from 24 March 2020 to 30 November 2021. The dataset characterised demographics, pre-existing comorbidities, symptoms, facility managed, patient status, treatment outcome and dates of related events, without personal identifiers. Hence, this secondary analysis waived the required individual informed consent. Patient information was retrieved from the dataset based on the occupation of interest-HCW and their respective designation-alongside demographic data on age, sex, place of work defined as 'health facility', hospitalisation required', and case presentation, knowledge of exposure, place of exposure and treatment outcome labelled as 'recovered' or 'dead'.

Based on the National Interim Guidelines for Clinical Management of COVID-19 (Nigeria Centre for Disease Control, 2020), all patients requiring hospitalisation at the time of testing were classified as severely ill, and it was the definition for illness severity; case presentation at the time of testing was also categorised as 'symptomatic' or 'asymptomatic'.

#### **Data analysis**

Data were analysed using Stata Statistical Software: Release V.12.<sup>14</sup> Descriptive statistics were used to report on the cohort characteristics. Means and SDs were reported for continuous variables and proportions for categorical variables, qualitative responses and subgroup analysis of variables. A test of proportions was conducted for both risk factors and outcome proportions. Univariate analysis of categorical variables was conducted by using  $(\chi^2)$  and Fischer's exact test where appropriate. A two-tailed p<0.05 was statistically significant. Multivariate logistic regression was used to evaluate risk factors of illness severity and mortality among HCWss with COVID-19. The aOR with a 95% CI was used to report the measure of association between the following: illness severity and risk factorsage, sex and case presentation; mortality and risk factors -age, sex, illness severity and case presentation.

### Patient and public involvement

It was not possible to involve patients or the public in the design, conduct, reporting or dissemination plans of our research as the study used secondary data without personal identifiers.

#### RESULTS Patient characteristics

Patient characteristics Data on 301 HCWs infected with COVID-19 were identi-fied and extracted into a spreadsheet. The demographic and clinical characteristics of the patients are found in ŝ table 1. The mean age was 43 years and 50.5% of the cohort were female. Of the 301 HCWs, 187 patients were 8 symptomatic with 10% (32) of the study cohort requiring hospitalisation-a measure of illness severity. A total of 108 HCWs were in contact with known probable cases, and 101 persons knew the place of exposure. From the available data, 7 (26.4%) infected HCWs died of their COVID-19 infection, resulting in a CFR of 2.3%. A test of proportion revealed a significant difference in proportions for age groups, case presentation, illness severity and use outcome. Statistics showed that population proportions for illness severity would be significantly lower than seen in the study population, while they would be significantly greater for age groups, case presentation and mortality.

The distribution of HCWs by the WHO classification<sup>13</sup> and health facilities were cross-tabulated in table 2 to determine association. Health professionals made up 79.7% (240) of the study cohort, with 68.8% (165) of them working at the teaching hospitals. The teaching hospitals are also the two major government own tertiary facilities in the state; HCWs in these facilities made up 66.1% (199) of the study population. A significant association was portrayed between HCWs and the health facilities they worked in.

training, Table 3 is a cross-tabulation of HCWs by risk factors and outcomes. Health professionals aged 30-49 years were the most affected subgroup, making up 53.2% of the study population; they also made up 80.7% (151/187) of all symptomatic cases. All mortality cases were also health similar technologies

	Health facili	ty classification, n (	%)				
Healthcare workers	Teaching hospitals	General service hospitals	Health allied organisations	Corporate health clinics	Community hospitals	Outpatient clinics	Total
Health professionals	165 (54.8)	32 (10.6)	9 (3.0)	17 (5.7)	15 (5.0)	2 (0.7)	240 (79.7)
Health associate professionals	11 (3.7)	0 (0.0)	3 (1.0)	3 (1.0)	1 (0.3)	1 (0.3)	19 (6.3)
Personal care workers	12 (4.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.7)	0 (0.0)	14 (4.7)
Health management and support personnel	10 (3.3)	4 (1.3)	10 (3.3)	2 (0.7)	0 (0.0)	0 (0.0)	26 (8.7)
Other health service providers	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)
Missing values	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	1 (0.3)
Total	199 (66.1)	36 (12.0)	22 (7.3)	23 (7.6)	18 (6.0)	3 (1.0)	301 (100%)

2

⊳

, and

Table 3 Contingency table of hea	althcare v	workers by	risk facto	ors and or	Itcomes									
	Risk fact	tors, n (%)									Outcomes, I	(%) ו		
	Age						Sex		Symptomat	ic	Illness seve	rity	Mortality	
Healthcare workers	20-29	30–39	40-49	50-59	69-09	>70	Female	Male	No	Yes	No	Yes	Alive	Dead
Health professionals	21 (7.0)	99 (32.9)	61 (20.3)	33 (11.0)	18 (6.0)	8 (2.7)	125 (41.5)	115 (38.2)	89 (29.6)	151 (50.2)	215 (71.4)	25 (8.3)	233 (77.4)	7 (2.3)
Health associate professionals	1 (0.3)	10 (3.3)	5 (1.7)	2 (0.7)	1 (0.3)	0.0) 0	6 (2.0)	13 (4.3)	7 (2.3)	12 (4.0)	17 (5.7)	2 (0.7)	19 (6.3)	0 (0.0)
Personal care workers	1 (0.3)	2 (0.7)	5 (1.7)	2 (0.7)	3 (1.0)	1 (0.3)	9 (3.0)	5 (1.7)	4 (1.3)	10 (3.3)	13 (4.3)	1 (0.3)	14 (4.7)	0 (0.0)
Health management and support personnel	2 (0.7)	7 (2.3)	9 (3.0)	7 (2.3)	0 (0.0)	1 (0.3)	11 (3.7)	15 (5.0)	13 (4.3)	13 (4.3)	23 (7.6)	3 (1.0)	26 (8.7)	0 (0.0)
Other health service providers	0.0) 0	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	0.0) 0	1 (0.3)	0 (0.0)	0 (0.0)	1 (0.3)	0.0) 0	1 (0.3)	1 (0.3)	0 (0.0)
Missing value*	0.0) 0	0 (0.0)	1 (0.3)	0 (0.0)	0 (0.0)	0.0) 0	0 (0.0)	1 (0.3)	1 (0.3)	0 (0.0)	1 (0.3)	0.0) 0	1 (0.3)	0 (0.0)
Total	25 (8.3)	118 (39.2)	81 (26.9)	44 (14.6)	23 (7.6)	10 (3.0)	152 (50.5)	149 (49.5)	114 (37.9)	187 (62.1)	269 (89.4)	32 (10.6)	294 (97.7)	7 (2.3)
$\chi^{2}$	$\chi^2$ (20) =	= 27.8331					$\chi^{2}(4) = 5.70$	15	$\chi^{2}$ (4) = 2.8	223	$\chi^{2}(4) = 8.56$	143	$\chi^{2}(4) = 1.79$	918
α	0.113						0.223		0.5888		0.072		0.774	
Fishers exact							0.204		0.6360		0.264		1.000	
*Not included in analysis.														

professionals. The association between HCWs and either risk factors, or outcomes were not significant.

The cross-tabulation of health facilities by risk factors and outcomes showed that health workers aged 30-49 years were also the most dominant in the teaching hospitals, and they made up 44.2% of the total HCWs. The majority of symptomatic patients-73.2%, persons who required hospitalisation for testing-87.5%, and six out of the seven mortality cases were also staff of the teaching hospitals. Age and case presentation were significantly associated with the health facilities and are shown in table 4.

Table 5 shows a subgroup analysis conducted on the ş health professionals infected-doctors (71.7%), nurses copyright, incl (27.3%), others (1%); and teaching hospitals by ownership: public (78.6%), private (21.4%).

# Predictors of illness severity and mortality among HCWs

The outcome proportions by risk factors were reported in tables 6 and 7. The effects of age, sex and case class on illness severity were evaluated using both univariate (table 6) and multivariate logistic regression (table 7), respectively. Symptomatic cases were more likely to uses rela advance to severe illness ( $\chi^2(1) = 15.219$ ,  $\alpha = < 0.0001$ ; aOR 10.658, 95% CI 2.494 to 45.552). The overall model was statistically significant  $(\chi^2 (8) = 19.112, \alpha < 0.0001)$ ); it explained 12.5% (Nagelkerke  $\mathbb{R}^2$ ) of the variance in illness severity, and correctly classified 89.4% of cases. 6

Predictors of mortality assessed included age, sex, case class and illness severity. The logistic regression model was statistically significant,  $\chi^{2}$ , (9) = 16.965,  $\alpha$  = 0.049. The model explained 27.6% (Nagelkerke  $R^2$ ) of the variance in mortality and correctly classified 97.7% of cases. Age  $(\chi^2(1) = 19.24, \alpha = 0.002; \text{ aOR } 1.079, 95\% \text{ CI } 1.02)$ to 1.141 per year increase) was identified as a risk factor for mortality among HCWs with COVID-19 patients.

# DISCUSSION

ing, Al training Using comprehensive data on COVID-19 infections and in HCWs in Rivers State Nigeria, this study showed a mortality proportion of 2.3% in the study cohort of 301 participants. Health professionals and HCWs in the teaching hospitals made up a majority of the study population, and HCWs between 30 and 49 years were the most affected. It was also noted that the most probable source of infection was the workplace, followed by a social event. Ten per cent of the study cohort experienced severe & illness, the result agrees with available evidence from a meta-analysis that reported a 9.9% incidence of severe disease in HCWs.<sup>15</sup> Age and gender are predictors with established association with COVID-19 mortality. Age is a crucial risk factor in the epidemiology of COVID-19; our study found an association between age and mortality; prior studies revealed patients above 65 years are at a greater risk of both disease severity and mortality from infection with SARS-CoV2.16 17 Consistent with research findings, mortality was higher in male patients

e

an

đ

∃

	Risk facto	rs, n (%)									Outcomes,	(%) u		
	Age						Sex		Symptoma	tic	Illness seve	erity	Mortality	
Health facility	20-29	30-39	40-49	50-59	69-09	>70	Female	Male	No	Yes	No	Yes	Alive	Dead
Teaching hospitals	14 (4.7)	79 (26.3)	54 (17.9)	34 (11.3)	14 (4.7)	4 (1.3)	101 (33.6)	98 (32.6)	62 (20.6)	137 (45.5)	171 (56.8)	28 (9.3)	193 (64.1)	6 (2.0)
General service hospitals	7 (2.3)	9 (3.0)	10 (3.3)	4 (1.3)	4 (1.3)	2 (0.7)	21 (7.0)	15 (5.0)	20 (6.6)	16 (5.3)	32 (10.6)	4 (1.3)	35 (11.6)	1 (0.3)
Health allied organisations	1 (0.3)	8 (2.7)	3 (1.0)	6 (2.0)	2 (0.7)	2 (0.7)	7 (2.3)	15 (5.0)	11 (3.7)	11 (3.7)	22 (7.3)	0.0) 0	22 (7.3)	0.0) 0
Corporate health clinics	2 (0.7)	13 (4.3)	7 (2.3)	0 (0.0)	1 (0.3)	0.0) 0	8 (2.7)	15 (5.0)	14 (4.7)	9 (3.0)	23 (7.6)	0.0) 0	23 (7.6)	0.0) 0
Community hospitals	1 (0.3)	6 (2.0)	7 (2.3)	0 (0.0)	2 (0.7)	2 (0.7)	13 (4.3)	5 (1.7)	6 (2.0)	12 (4.0)	18 (6.0)	0.0) 0	18 (6.0)	0.0) 0
Outpatient clinics	0 (0.0)	3 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.7)	1 (0.3)	1 (0.3)	2 (0.7)	3 (1.0)	0.0) 0	3 (1.0)	0 (0.0)
Total	25 (8.3)	118 (39.2)	81 (26.9)	44 (14.6)	23 (7.6)	10 (3.0)	152 (50.5)	149 (49.5)	114 (37.9)	187 (62.1)	269 (89.4)	32 (10.6)	294 (97.7)	7 (2.3)
$\chi^2$	$\chi^{2}(30) = :$	37.7053					$\chi^{2}(5) = 9.5$	447	$\chi^{2}(5) = 15.$	3283	$\chi^{2}(5) = 10.$	3373	$\chi^{2}(5) = 2.0$	203
α	0.049						0.077		0.009		0.066		0.846	
Fishers exact							0.067		0.007		0.068		1.000	

en access		D
	oper	2222
240 (100)		Þ
172 (71.7)	ž	÷
63 (26.3)		5
276 (100)		2
217 (78.6)	60 0	2
59 (21.4)	2	5
276 (100) 217 (78.6) 59 (21.4) ociation was ction among the of testing, nis might be ts of positive required to atable to the art to conclu- e manifesta- ofessionals. <sup>20</sup> ion between or evidence SARS-CoV-2 ess onset and viral load in a cohort was of 0.5%. <sup>22</sup> atable to the and may be d workforce. mostly been ope and the red with the ing country. ed subgroup stly involved tudy though s as against ne front-line e state, and Other plau- re not in the HCWs were y than in the oth adher- PC) protocol work during Ws as study picts a work- o association younger age our results. ill in service. age ranges	Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.	lished as 10 1132/kmianan 2003 021020 an 11 Navambar 2003 Dawalandad from http://kmianan.hmi.com/ on lung 13 2005 at Agence Bibliograp
epending on		5
5	due de	1.20
-		

in our study<sup>18</sup><sup>19</sup>; although no significant associa deduced in the cohort evaluated. Also, infectio HCWs was typically asymptomatic at the time of with 89.4% not requiring hospitalisation, this due to the active tracing and testing of contacts of cases in the State; however, more research is redetermine whether these findings are attributab healthy worker bias. The results were similar to sions from a study that observed less severe m tions of COVID-19 infection in medical profes Our results also showed a significant association symptomatic cases and illness severity; prior found an association between prolonged SA RNA shedding and the interval between illness of treatment.<sup>21</sup> and may be indicative of a high vira these persons. The proportion of mortality in the study co higher, compared with available evidence of The difference in mortality is perhaps attributat geographical location of studies conducted and suggestive of better working conditions and w Studies on COVID-19-related mortality have mo conducted in developed regions (Asia, Europe USA), and showed lower mortality compared current study conducted in Nigeria developing Health professionals being the most infected s aligns with various evidence,<sup>12</sup> as they are mostly in patient-facing roles. A difference in our stud was most infections were seen in doctors a nurses in other studies: the doctors were the responders for COVID-19 response in the s that may be the reason for this observation. Ot sible explanations are that all exposures were n workplace, and there was an indication that HCWs were infected to a greater extent in the community than in the workplace<sup>23</sup>; thoughts are also to be given to the adherence to infection prevention and control (IPC) protocol **g**. workplace, and there was an indication that HCWs were among the HCWs, and their willingness to work during **g** the COVID-19 pandemic.<sup>24 25</sup> A random selection of studies using HCWs as study participants conducted in the study region depicts a workforce majorly aged under  $50^{26-29}$ ; although no association

 
 Table 5
 Subgroup analysis
Health professionals

Hospitals by ownership

Doctors Nurses

Public

Private

can be inferred, it is suggestive of a mostly younger age distribution of HCWs and was indicated in our results. Also indicated, were persons aged above 70, still in service. Although the constitutional retirement age ranges between 60 and 70 years from civil service depending on

	Risk facto	ors, n (%)										
	Age						Sex		Symptom	atic	Illness Sev	rerity
Outcome	20-29	30–39	40-49	50-59	69-09	>70	Female	Male	No	Yes	No	Yes
Illness severity, n=32	3 (9.4)	13 (40.6)	4 (12.5)	6 (18.8)	5 (15.6)	1 (3.1)	16 (50)	16 (50)	2 (6.3)	30 (93.8)	I	I
$\chi^2$	$\chi^2 (5) = 6.$	2195					$\chi^{2}(1) = 0.00$	36	$\chi^{2}(1) = 15$	.2187	I	
α	0.285						0.952		0.000		I	
Fishers exact	0.199						1.000		0.000		I	
Mortality, n=7	0 (0.0)	1 (14.3)	0 (0.0)	2 (28.6)	3 (42.9)	1 (14.3)	1 (14.3)	6 (85.7)	2 (28.6)	5 (71.4)	6 (85.7)	1 (14.3)
x <sup>2</sup>	$\chi^{2}(1) = 15$	9.240					$\chi^{2}(1) = 3.75.$	96	$\chi^{2}(1) = 0.5$	:636	$\chi^{2}(1) = 0.1$	200
α	0.002						0.053		0.608		0.754	
Fishers exact	0.003						0.065		0.713		0.549	

e

Table 7 Multivariate analysis of risk factors for COVID-19 outcomes

Outcome	Risk factors	aOR	95% CI	P value
Illness severity	Age	0.98	0.455 to 2.111	0.959
	Sex*	1.003	0.971 to 1.036	0.859
	Case class†	10.658	2.494 to 45.552	0.001
Classification tal	ole -89.4% correctly c	lassified, c	onstant=-4.139	
Mortality	Age	1.079	1.02 to 1.141	0.008
	Sex*	4.274	0.486 to 37.582	0.190
	Case class†	1.166	0.198 to 6.869	0.865
	Illness severity‡	1.305	0.130 to 13.123	0.821
*Female				

+Asymptomatic +No

Protected by copyright, inc the profession (70 years for the medical professional), the Civil Service Commission offers contract appointments to pensioners. Due to several challenges, evidence showed that Nigerians generally have a poor attitude towards retirement; with a preference to continue working ₫ privately in some capacity, after retirement from the civil service.<sup>30–32</sup>

The use of routinely collected data is beneficial in this scenario as the data were readily available and could be representative of the study population, following the integrated testing and reporting approach used in data collection. The completeness of the data also minimises the effects of selection bias due to non-response and lost to follow-up. The independent mode of prospective data collection reduces recall bias on exposure.<sup>33</sup> This study gives a snapshot into the impact of COVID-19 on ta the healthcare workforce of Rivers State and serves as a model for a more holistic research.

The existent significant challenge in giving an accurate report on deaths due to COVID-19, let alone those among HCWs for several reasons, was acknowledged by the WHO.<sup>12</sup> As applicable to our setting, reports were for ğ deaths with a confirmed COVID-19 test; hence, untested individuals and persons who died outside a hospital facility would not have been included in the death counts. The reliance on reported infections and deaths implies that there is a probability of missing unreported cases and mild cases of COVID-19. Some reasons for non-reporting may include: the fear of stigmatisation from colleagues, no and poor health-seeking behaviour among this cohortthe practice of self-medication and reluctance to obtain medical care.<sup>35 36</sup> The health-seeking behaviour of the HCWs in Nigeria and poor reporting of COVID-19 infection cases within this cohort is to be considered. There is evidence that the practice of self-medication and reluctance to obtain medical care is high among doctors and nurses in Nigeria<sup>35–39</sup>; therein lies the possibility of nonreporting of mild cases. This behavourial pattern emphasises the need for more awareness and education on these issues within this group of healthcare professionals. These are limitations to this study. The viral load of SARS-CoV-2

and comorbidities are useful markers for assessing disease severity and prognosis<sup>21</sup>; the availability of information on these variables would have better informed the study. Data on the interval between illness onset and treatment onset would have given a more concise inference of disease severity also.

HCWs are the most important human resource for hospitals; the workplace-related mortality in HCWs not only compromises the workforce in healthcare settings but also affects the mental health of colleagues.<sup>40 41</sup> A CFR of 2.33% though comparable with global statistics for HCWs<sup>10</sup> is higher than both the CFR of the study area— Rivers state (0.98%) and Nigeria (1.23%).<sup>42</sup> There is a need for re-evaluation of compliance to the COVID-19 IPC protocol, the adequacy of PPE and working conditions in place for HCWs in Rivers State; because, exposure to numerous infected individuals, may demonstrate that HCWs, if infected, could be characterised by higher viral load, thereby, associated with worse clinical outcomes.<sup>21</sup> The results from the study also further emphasise the need to protect HCWs; ensure they are knowledgeable in both IPC, and that the healthcare space is safe against nosocomial infections. The density of HCWs in the state and country at large is also a point of concern; although it is estimated that 74543 doctors and 301579 nurses are registered in Nigeria, the Medical and Dental Council Of Nigeria stated that only about 59% of the doctors and 35% of nurses are in active service. Nigeria has also suffered a mass exodus of HCWs over the years; in 2020 during the height of COVID-19, it was reported that 7256 Nigerian nurses migrated from Nigeria.<sup>43–45</sup> Better working conditions for health workers need to be advocated, to regulate the export of human capital.

To our knowledge, this research is the foremost study representing a relatively comprehensive analysis of COVID-19-related mortality and disease severity in HCWs from available state records in Rivers State. As an emerging research area in the current pandemic, there are other factors worth considering. For example, the effect of time of hospitalisation on disease severity and mortality, and viral load count. As a secondary analysis, we were unable to analyse this variable. Future studies to investigate this variable is essential. The impact of nosocomial versus community transmission is also a vital area of research. A national study is required to extrapolate the findings from this study to the nation, as surveillance of the impact of COVID-19 by occupation and industry will benefit not only HCWs but all workers in the nation.

### CONCLUSION

In conclusion, front-line HCWs are at an increased risk of exposure to COVID-19 infections. In Nigeria, there is the possibility of a higher risk of experiencing a severe disease if symptomatic while infected. It is imperative that preventive strategies are established and implemented, alongside proper education, and awareness to protect HCWs. Acknowledgements We highly appreciate the editor and anonymous reviewers whose comments and suggestions helped greatly improve and clarify the manuscript. The authors acknowledge the Rivers State Public Health Emergency Operations Centre and the Rivers State Ministry of Health.

**Contributors** CE-E: conception, design of the work; data collection; data analysis and interpretation; first draft of the manuscript; critical revisions of the article. FP: drafting the article; data collection, critical revisions of the article. EI: data analysis and interpretation; drafting the article; critical revisions of the article. GO: conception, design of the work; drafting the article; critical revisions of the article, supervision of work. All authors read and gave final approval of the manuscript version submitted for publication. Guarantor, GO.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. According to the Data Protection Law of the Federal Republic of Nigeria, the authors cannot publicly release the data accessed from the Rivers State Health Records Database. Data are available on reasonable request with permission from the Rivers State Ministry of Health.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

#### ORCID iD

Chidinma Eze-Emiri http://orcid.org/0000-0002-6423-3175

#### REFERENCES

- Nguyen LH, Drew DA, Graham MS, et al. Risk of COVID-19 among front-line health-care workers and the general community: a prospective cohort study. Lancet Public Health 2020;5:e475–83.
- 2 Wang X, Jiang X, Huang Q, et al. Risk factors of SARS-CoV-2 infection in healthcare workers: a retrospective study of a nosocomial outbreak. Sleep Med X 2020;2:100028.
- 3 Adeloye D, David RA, Olaogun AA, et al. Health workforce and governance: the crisis in Nigeria. Hum Resour Health 2017;15:32.
- 4 World Health Organisation. Global health workforce statistics database, 2022. Available: https://www.who.int/data/gho/data/ themes/topics/health-workforce [Accessed 30 June 2022].
- 5 Gallo Marin B, Aghagoli G, Lavine K, *et al.* Predictors of COVID-19 severity: a literature review. *Rev Med Virol* 2021;31:e2146
- 6 Yao Y, Tian J, Meng X, et al. Progression of severity in coronavirus disease 2019 patients before treatment and a self-assessment scale to predict disease severity. *BMC Infect Dis* 2022;22:409.
- 7 Izcovich A, Ragusa MA, Tortosa F, et al. Prognostic factors for severity and mortality in patients infected with COVID-19: a systematic review. PLoS One 2020;15:e0241955.
- 8 Shi C, Wang L, Ye J, *et al.* Predictors of mortality in patients with coronavirus disease 2019: a systematic review and meta-analysis. *BMC Infect Dis* 2021;21:663.
- 9 World Health Organization. The impact of COVID-19 on health and care workers: a closer look at deaths. health workforce department – working paper 1. (WHO/HWF/WorkingPaper/2021.1). Licence: CC BY-NC-SA 3.0 IGO. 2021. Geneva: World Health Organization, 2021.

- 10 Bandyopadhyay S, Baticulon RE, Kadhum M, et al. Infection and mortality of healthcare workers worldwide from COVID-19: a systematic review. BMJ Glob Health 2020;5:e003097.
- 11 Pan American Health Organization-World Health Organisation. COVID-19 has infected some 570,000 health workers and killed 2,500 in the Americas, PAHO Director says - PAHO/WHO | Pan American Health Organization, 2021. Available: https://www.paho. org/en/news/2-9-2020-covid-19-has-infected-some-570000health-workers-and-killed-2500-americas-paho [Accessed 06 Dec 2021].
- 12 World Health Organization. The impact of COVID-19 on health and care workers: a closer look at deaths Geneva: World Health organization;, 2021. Available: https://apps.who.int/iris/handle/ 10665/345300 accessed WHO/HWF/WorkingPaper/2021.1.
- 13 World Health Organisation. Healthcare workers classification, 2020. Available: https://www.who.int/hrh/statistics/Health\_workers\_ classification.pdf [Accessed 09 Jan 2022].
- 14 Stata Statistical Software. Release 12. College Station, TX: StataCorp LP. [program], 2011.
- 15 Sahu AK, Amrithanand VT, Mathew R, et al. COVID-19 in health care workers - A systematic review and meta-analysis. Am J Emerg Med 2020;38:1727–31.
- 16 Chen R, Liang W, Jiang M, et al. Risk Factors of Fatal Outcome in Hospitalized Subjects With Coronavirus Disease 2019 From a Nationwide Analysis in China. Chest 2020;158:97–105.
- 17 CDC. Severe outcomes among patients with coronavirus disease 2019 (COVID-19) United States 2020;69.
- 18 Cao M, Zhang D, Wang Y. Clinical features of patients infected with the 2019 novel coronavirus (COVID-19) in Shanghai, China. *MedRxiv* 2020.
- 19 Jin J-M, Bai P, He W. Higher severity and mortality in male patients with COVID-19 independent of age and susceptibility. *MedRxiv* 2020.
- 20 Li J, Long X, Zhang Q, *et al.* Mild manifestations of COVID-19 in healthcare workers. *PLoS Negl Trop Dis* 2020;14:e0008950.
- 21 Liu Y, Yan L-M, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. Lancet Infect Dis 2020;20:656–7.
- 22 Gómez-Ochoa SA, Franco OH, Rojas LZ, et al. COVID-19 in healthcare workers: a living systematic review and meta-analysis of prevalence, risk factors, clinical characteristics, and outcomes. Am J Epidemiol 2021;190:161–75.
- 23 Jacob JT, Baker JM, Fridkin SK, *et al.* Risk factors associated with SARS-CoV-2 seropositivity among US health care personnel. *JAMA Netw Open* 2021;4:e211283.
- 24 Chiedozie I. How work pressure forces Nigerian doctors, nurses to relocate abroad | the ICIR. *ICIR* 2021.
- 25 Mbaba AN, Ogolodom MP, Abam R. Willingness of health care workers to respond to covid-19 pandemic in Port Harcourt, Nigeria. *Health Science Journal* 2021;15.
- 26 Kadiri-Eneh NP, Azuike EC, Tobin-West C. An assessment of the potentials for retention of primary health care workers in rivers state, Nigeria 2018;6:35–51.

- 27 Akpuh N, Ajayi I, Adebowale A, et al. Occupational exposure to HIV among healthcare workers in PMTCT sites in Port Harcourt, Nigeria. BMC Public Health 2020;20:451.
- 28 Bell-Gam H, Amadi E. Perception and attitude of doctors on ageing, old age and geriatric care in rivers state, Nigeria; 2021.
- 29 Ogba A. Occupational stress and its management among health care workers in the University of port Harcourt teaching Hospital, rivers state. *Health Science Journal* 2020;14.
- 30 Onuoha BC, Creation W. Retirement and succession planning of entrepreneurs in south-east, Nigeria. *International Business and Management* 2013;7:99–105.
- 31 Ogunbameru OA, Bamiwuye S. Attitudes toward retirement and PRERETIREMENT education among Nigerian bank workers. *Educ Gerontol* 2004;30:391–401.
- 32 Eremie MD. Challenges and Expectations of Retirees in Rivers State, Nigeria : Implications for Career Counselling. *AJBMR* 2015;3:1–6.
- 33 Thygesen LC, Ersbøll AK. When the entire population is the sample: strengths and limitations in register-based epidemiology. *Eur J Epidemiol* 2014;29:551–8.
- 34 Kwaghe AV, Ilesanmi OS, Amede PO, et al. Stigmatization, psychological and emotional trauma among frontline health care workers treated for COVID-19 in Lagos state, Nigeria: a qualitative study. BMC Health Serv Res 2021;21:855.
- 35 Agaba PA, Ocheke AN, Akanbi MO. Health seeking behavior of physicians at the Jos university teaching hospital. *Nigerian Medical Journal* 2011;52:90–4.
- 36 Fawibe AE, Odeigah LO, Akande TM, et al. Self-Reported medical care seeking behaviour of doctors in Nigeria. *Alexandria Journal of Medicine* 2017;53:117–22.
- 37 Adamu H, Yusuf A, Inalegwu C, et al. Factors influencing healthseeking behavior of health workers in a tertiary health institution in Sokoto, northwest Nigeria. Sahel Medical Journal 2018;21:162–70.
- 38 Adewoye KR, Aremu ŠK, Ipinnimo TM, et al. Awareness and practice of proper health seeking behaviour and determinant of self-medication among physicians and nurses in a tertiary hospital in Southwest Nigeria. Open Journal of Epidemiology 2019;09:36–49.
- 39 Fadare J, Desalu O. Self-prescription practices by Nigerian medical doctors. South African Family Practice 2014;56:253–7.
- 40 Badrfam R, Zandifar A, Arbabi M. Mental health of medical workers in COVID-19 pandemic: restrictions and barriers. J Res Health Sci 2020;20:e00481–e81.
- 41 Javed B, Sarwer A, Soto EB, et al. The coronavirus (COVID-19) pandemic's impact on mental health. Int J Health Plann Manage 2020;35:993–6.
- 42 Nigeria Centre for Disease Control. Ncdc coronavirus COVID-19 microsite, 2021. Available: https://covid19.ncdc.gov.ng [Accessed 24 Dec 2021].
- 43 Newspaper H-P. Alarming exodus of health professionals; 2022.
- 44 Jannamike L. Mass exodus of doctors from Nigeria worrisome Buhari; 2019.
- 45 Newspaper H-P. 7,256 Nigerian nurses left for UK in one year Report - Healthwise; 2022.