To cite: Goldsmith LP.

Prepublication history and

for this paper are available

online. To view these files, please visit the journal online

(http://dx.doi.org/10.1136/

LPG, MR-P and KH are joint first

Received 10 February 2022

Accepted 22 September 2022

Check for updates

C Author(s) (or their

Published by BMJ.

Correspondence to

Dr Sally Hargreaves;

s.hargreaves@sgul.ac.uk

end of article.

employer(s)) 2022. Re-use

For numbered affiliations see

permitted under CC BY.

bmjopen-2022-061896).

authors.

additional supplemental material

BMJ Open Use of social media platforms by migrant and ethnic minority populations during the COVID-19 pandemic: a systematic review

Lucy Pollyanna Goldsmith ⁽¹⁾, ^{1,2} May Rowland-Pomp,¹ Kristin Hanson,³ Anna Deal,^{1,4} Alison F Crawshaw ⁽¹⁾, ¹ Sally E Hayward ⁽¹⁾, ¹ Felicity Knights,^{1,2} Jessica Carter ⁽¹⁾, ¹ Ayesha Ahmad, ⁵ M Razai ⁽¹⁾, ² Tushna Vandrevala,⁶ Sally Hargreaves ¹

ABSTRACT

Rowland-Pomp M. Hanson K. **Objective** Migrants and ethnic minority groups have been et al. Use of social media disproportionately impacted by COVID-19 and have lower platforms by migrant and ethnic levels of vaccine uptake in some contexts. We aimed to minority populations during determine the extent and nature of social media use in the COVID-19 pandemic: a migrant and ethnic minority communities for COVID-19 systematic review. BMJ Open information, and implications for preventative health 2022;12:e061896. doi:10.1136/ measures including vaccination intent and uptake. bmjopen-2022-061896

> **Design** A systematic review of published and grey literature following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. We searched databases including Embase, Web of Science, PubMed NIH, CINAHL, facilitated through the WHO Global Research on COVID-19 database from 31 December 2019 to 9 June 2021.

Eligibility criteria for study selection Research reporting the use of social media by migrants and/or ethnic minority groups in relation to COVID-19.

Data extraction We extracted data on key outcomes, study design, country, population under study and sample size.

Results 1849 unique records were screened, and 21 data sources were included, including populations in the UK, USA, China, Jordan, Qatar and Turkey. We found evidence of consistent use of a range of social media platforms for COVID-19 information in some migrant and ethnic minority populations (including WeChat, Facebook, WhatsApp, Instagram, Twitter, YouTube), which may stem from difficulty in accessing COVID-19 information in their native languages or from trusted sources. Some evidence suggested circulating misinformation and social media use may be associated with lower participation in preventative health measures, including vaccine intent and uptake, findings which are likely relevant to multiple population aroups.

Conclusions Social media platforms are an important source of information about COVID-19 for some migrant and ethnic minority populations. Urgent actions and further research are now needed to better understand effective approaches to tackling circulating misinformation, and to seize on opportunities to better use social media platforms to support public health communication and improve vaccine uptake.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- \Rightarrow Comprehensive systematic review methods were used, following Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.
- \Rightarrow Both published and grev literature were searched and papers from all countries and all regions were included, allowing available evidence to be synthesised.
- \Rightarrow We acknowledge the limited geographical scope of included studies, with no data available from lowincome countries.

Registration This study has been registered with PROSPERO (CRD42021259190).

INTRODUCTION

Protected by copyright, including for uses related to text and data mining, The pandemic has been accompanied by an ≥ infodemic, defined as an excess of information during a disease outbreak—including false or misleading information in digital and physical environments¹—that makes it difficult to distinguish reliable information from misinformation including disinformation <u>0</u> (deliberate misinformation) and conspiracy theories. The WHO highlights that in all communities, infodemics cause 'confusion and risk-taking behaviours that can harm health...it leads to mistrust in health authorities and undermines the public health response, and can intensify or lengthen 8 outbreaks'.1 The rapid expansion of internet and social media use, in particular, in recent years (including platforms such as Twitter, WhatsApp and YouTube; table 1) has meant that both useful and potentially harmful health information can spread rapidly. Although social media can be used to disseminate factual, appropriate and useful information, a large proportion of the most popular

BMJ

 Table 1
 Popular social media platforms statistics from Statistica⁷¹

Platform	Primary feature	Country of origin	Organisation	Users	Notes
YouTube	Online video sharing and social media platform. Free to use.	USA	Google	Approximately >2 billion monthly	Searchable.
WhatsApp	Messaging platform, allows users to send text messages and voice messages, make voice and video calls, and share images, documents, user locations, and other content. Free to use.	USA	Meta	Approximately >2 billion monthly	Not searchable. Groups can have 512 users in them.
Instagram	Photo and video sharing social networking site. Free to use.	USA	Meta	Approximately 1 billion monthly	Searchable. Some content is limited to connections only.
Facebook	Social networking service, allows messaging, image and video sharing, marketplace online shopping, live video sharing. Free to use.	USA	Meta	2.89 billion active monthly	Searchable. Some content is limited to connections only.
WeChat	Instant messaging, social media, mobile payment. Free to use.	China	Tencent Holdings Limited	1.25 billion monthly	Searchable. Some content is limited to connections only.
TikTok (Known in China as Douyin)	Video sharing focused on short form videos (15 s to 3 min). Free to use.	China	ByteDance	837 million monthly active	Searchable. Some content is limited to connections only.
Snapchat	Photo sharing multimedia app with video features. Free to use.	USA	Snap	347.3 million monthly active	Searchable. Some content is limited to connections only.
Twitter	Microblogging focused on short messages known as 'tweets'. Live chat event function Tweetchat.	USA	Twitter	330 million monthly active	Searchable. Some content is limited to followers only.
COVID-19 vide ound to conta ion, reaching considered a n the control of 0 of 'undesirabl	eos on YouTube, for example, ain misinformation, or no factur millions of people worldwide. ²¹ najor platform for information COVID-19, but most COVID-19 le quality' containing few go recommendations according	, have been tal informa- ³ YouTube is concerning videos were overnment/ to a recent	as a source may also di informatior tionately im migrants, a minority gro were at high	of health information. raw on diaspora media a n. ¹⁰ The COVID-19 pand pacted and exacerbated and ethnically diverse c pups (including some mig ner risk of contracting, be	These communities s a source of health emic has dispropor inequalities faced by ommunities—ethnic grant populations)— ing hospitalised with

COVID-19 videos on YouTube, for example, have been found to contain misinformation, or no factual information, reaching millions of people worldwide.²³ YouTube is considered a major platform for information concerning the control of COVID-19, but most COVID-19 videos were of 'undesirable quality' containing few government/ public health recommendations according to a recent study.⁴ A review of YouTube videos on general vaccination found 65% expressed antivaccination sentiment,⁵ with antivaccine posts more likely to be recirculated on Twitter.² The spread of misinformation and disinformation has been highlighted as a major risk to ending the COVID-19 pandemic-including undermining trust in vaccines⁶—with researchers highlighting links between misinformation on social media and public doubts around vaccine safety, self-reported compliance with public health guidelines, and intent to vaccinate.⁷⁸

Although social media platforms are commonly used in the general population, and patterns of use are complex across different population groups;^{8 9} some migrant and ethnic minority groups—who may experience barriers to accessing health information and health systems may be more reliant on social media and the internet

minority groups (including some migrant populations)were at higher risk of contracting, being hospitalised with and dying from COVID-19.^{10–15} They are also more likely to be vaccine hesitant-with lower take-up of preventative health measures, such as vaccines, noted in some groups due to a range of personal, societal and physical barriers.^{12 14 16} Some migrant and ethnic minority communities may be more exposed to social media misinformation because of access barriers to accurate information (eg, from official government sources),^{17 18} due to restricted eligibility and access to services, language barriers and low health literacy. However, little is known about the extent and nature of social media use in these populations, nor the impact that social media use has had on preventative health measures during the pandemic, including COVID-19 vaccine uptake. In addition, there is an opportunity now to explore the extent to which social media platforms could be better used to

support information sharing and promote public health messaging in marginalised communities during the pandemic and beyond.

We therefore did a systematic review to explore and assess the extent and nature of social media use by migrant and ethnic minority groups to access COVID-19 health information, the extent to which misinformation on social media may have influenced views about COVID-19 preventative measures including vaccination intention and uptake, and to explore good practice.

METHODS

Search strategy

The review¹⁹ followed Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.²⁰ The study protocol is available from the International Prospective Register of Systematic Review (PROSPERO) database (CRD42021259190). A Boolean search strategy was developed containing terms relating to migrants, ethnic minorities, COVID-19, social media and misinformation (see online supplemental file 1). We included papers covering any prevention topic, including social distancing, hand washing, mask wearing, testing, isolation, test and trace activities, and vaccination. We searched the following databases: Embase, Web of Science, Oxford Academic Journals, PubMed NIH, Clinical Trials, China CDC MMWR, CDC reports, ProQuest Central (Proquest), CINAHL, Africa Wide Information (Ebsco), Scopus, PsycInfo, CAB Abstracts, Global Health, J Stage, Science Direct, Wiley Online Journals, JAMA Network, British Medical Journal, Mary Ann Liebert, New England Journal of Medicine, Sage Publications, Taylor and Francis Online, Springer Link, BioMed Central, MDPI, ASM, PLOS, The Lancet, Cell Press, and preprint sites chemRxiv, SSRNbioRxiv and medRxiv. This was facilitated through the WHO Global Research on COVID-19 database. We searched records from the date the WHO was first informed of COVID-19, 31 December 2019,²¹ to 9 June 2021(https:// search.bvsalud.org/global-literature-on-novel-coronavirus-2019-ncov/). The WHO COVID-19 Database²² is a daily updated multilingual resource of all literature (peer-reviewed literature, preprints and grey literature) pertaining to COVID-19.

Records were imported to Rayvan OCRI.²³ Both title and abstract screening and full text screening were conducted independently by two reviewers (MR-P and LPG) using Rayyan QCRI.²³ Additional relevant papers and grey literature (eg, from third-sector organisation websites) were identified using hand searching including backwards and forwards citation tracking.

Selection criteria and primary outcomes

Papers reporting the use of social media platforms and implications for preventative health measures and vaccination intent of migrants and/or ethnic minority groups to COVID-19 globally were eligible. All types of COVID-19 preventative health measures, including

social distancing, hand washing, mask wearing, testing, isolating, tracing close contacts of people who have COVID-19, alongside preventative measures based on misinformation were included. To include all available evidence, all types of scientific articles, reports and commentaries, editorials, correspondence letters were eligible for inclusion. Social media platforms were defined as any medium whereby content (including images, videos and messages) is circulated to the general public and may include YouTube, Facebook, Twitter, T TikTok and Snapchat. 'Migrants' were defined as foreignborn, residing outside of their country of birth. An ethnic minority group was defined as a group of people with a shared culture, tradition, language, history, living in a country where most people are from a different ethnic group, and will include migrants/foreign-born populations alongside individuals born in the host country. Where studies reported a general population sample, results about migrant/ethnic minority groups within that sample were eligible for inclusion. No papers were excluded based on language or geographical origin. Studies were excluded if it was not possible to determine ₫ whether individual(s) in the population studied were uses related migrants or from an ethnic minority group.

Data extraction, critical appraisal and synthesis

Data extraction was completed independently by two ō researchers (MR-P and LPG) using a piloted, structured data extraction sheet in Microsoft Excel. Fields in the data extraction sheet included author and year, dates for data collection, location of study, location of population of interest, whether qualitative methods were used, whether quantitative methods were used, study design, whether there was an intervention, the type of intervention, methodology, population of interest, further information about population, sample size, type of social \triangleright media, misinformation type, participant recruitment strategy, and all outcomes. Outcomes were extracted as reported. Risk of bias was assessed independently by two researchers (LPG, MR-P) using the Quality assessment for Survey Studies in Psychology for Surveys²⁴ for quantitative studies. The 20 items on this scale can be rated as 'yes', or 'no', 'not stated clearly', or 'not applicable'. Scores are calculated by dividing the 'yes' answers by the total number of applicable items, with scores over 70% indicating 'acceptable' quality. The Critical Appraisal Skills programme (CASP) checklist was used for qualitative 2 studies.²⁵ The 10 items can be rated 'yes', 'can't tell' or 'no'. We rated the CASP by dividing the 'yes' answers by the total number of applicable items, with a score of over 60% indicating 'acceptable' quality. We did not exclude any papers on the basis of quality. The selection of risk of bias rating instruments was finalised once we had a complete list of the type methods used in the included studies. We used a mixed-methods²⁶ narrative synthesis²⁷ approach, synthesising the qualitative and quantitative data together by theme.

Identification

Screening

Eligibility

ncluded



Patient and public involvement

Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.However, three of the authors are from ethnic minority groups.

RESULTS

Figure 1

Overview of data sources

Following de-duplication, 1849 unique data sources were identified and screened and ultimately 129 were full-text screened. Twenty-one data sources were included in the final analysis (figure 1). Six studies were conducted in the UK,²⁸⁻³³ two were jointly conducted in the UK and USA.³⁴³⁵ An additional eight studies were conducted in the USA,^{36–43} and one each in China,⁴⁴ Jordan,⁴⁵ Qatar⁴⁶ and Turkey.⁴⁷ Eight studies reported on migrants,^{31 33 36 44-48} including migrants in the host countries of China,44 Jordan,⁴⁵ Qatar,⁴⁶ Turkey,⁴⁷ and the USA³⁶ and UK,^{31 33} and one study involved predominantly migrants from Venezuela residing in other countries.⁴⁸ Nine studies reported about a specific ethnic minority or group (Latino individuals,^{36 38 40 41} black American citizens,^{39 43} Jain community members³² and Syrian migrants).^{45 47} Seven studies reported about ethnic minority groups generally.^{28–30 34 35 37 42} A survey design was the most common design, used in half of included studies.

Characteristics of included studies are presented in table 2, including the risk of bias assessment scores. Quality scores ranged from 76% to 90% for included

papers, suggesting acceptable quality of all included data sources where quality assessment was applicable. The most common shortcomings for studies related to reporting about the ethics and participants. None of the included studies were preprints.

Online supplemental material 2 shows the geographical location of data sources, highlighting the absence of published and unpublished data on this topic from most regions of the world.

Use of social media platforms as a source of information about COVID-19

For some migrants and ethnic minority groups, consistent use of social media platforms for sharing and receiving COVID-19-related health information was reported in several included studies.^{33 38-40 44-48} Figure 2 highlights quantitative data sets showing use of social media for information about COVID-19. Social media was reported to no be the preferred source of information about COVID-19 for international migrants in China (WeChat was used by 94.5% of respondents for COVID-19 information).⁴⁴ Among 389 Syrian refugee mothers in Jordan,⁴⁵ Facebook and WhatsApp were the main sources of information for 87% and 69% of respondents, respectively, for COVID-19 information; with 21% indicating that they accessed information from professional databases or government websites, and 53% via television (this survey was circulated via Facebook and WhatsApp). Migrants from Venezuela (residing in numerous countries) reported Facebook and WhatsApp were their two primary sources of information

data

		:					
Table 2 Characteris	tics of included s	tudies					
Included study	Location of study	Study design	Population under study	Type of publication	Main topic of the paper	Sample size	Quality rating*
Alabdulla ⁴⁶	Qatar	Cross-sectional survey	Non-Qatari residents	Peer-reviewed journal	Vaccine hesitancy	7821	76%†
Allington (a) ³⁰	NK	Cross-sectional survey	Non-white ethnic groups	Peer-reviewed journal	Vaccine attitudes, trust and COVID-19 information source as predictors of vaccine hesitancy	4343	82%†
Allington (b) ³⁵	UK	Cross-sectional survey	Non-white ethnic groups	Peer-reviewed journal	Media usage predicts intention to be vaccinated against COVID-19	8988	89%†
Behbahani ³⁶	NSA	Organisational case study	Latino migrants	Peer-reviewed journal	Helping vulnerable migrant populations in the COVID-19 crisis	N/K	NA⁺
Campos-Castillo ³⁷	USA	Cross-sectional survey	Non-white ethnic groups	Peer-reviewed journal	Racial and ethnic digital divides in posting COVID-19 content on social media	10510	88%†
Cervantes ³⁸	USA	Qualitative interviews with thematic analysis	Low-income Latino individuals	Peer-reviewed journal	Experiences of Latinx individuals hospitalised for COVID-19— misinformation and disbelief	60	\$00%
Chandler ³⁹	USA	Qualitative interviews with thematic analysis	Black women (18– 31 years)	Peer-reviewed journal	Evaluating the perspectives and sources of information of black women about COVID-19	15	\$00%
Crawshaw ³¹	ЛК	Evidence synthesis linked to outputs from participatory workshops with migrants	International migrants	Peer-reviewed journal	Vaccine hesitancy and barriers to COVID-19 vaccination in migrants	N/K	N/A§
Despres ⁴⁰	USA	Organisational case study	Latino community living in America	Peer-reviewed journal	A digital content curation model to challenge the inequitable impacts of COVID-19 on US Latinos	N/K	N/A§
Danish Refugee Council (DRC) ⁴⁷	Turkey	Cross-sectional survey	Syrian refugees in South-East Turkey	Research- based needs assessment report	The impact of COVID-19 on refugees in South-East Turkey	774	82%†
Hamadneh ⁴⁵	Jordan	Cross-sectional survey	Syrian refugee mothers	Peer-reviewed journal	Knowledge and attitudes about COVID-19 among Syrian refugee women in Jordan	389	78%†
Lockyer ²⁸	UK	Qualitative interview; reflective thematic analysis	People from different ethnic groups in Bradford	Peer-reviewed journal	COVID-19 misinformation and vaccine hesitancy in context	20	\$%06
							Continued

BMJ Open: first published as 10.1136/bmjopen-2022-061896 on 17 November 2022. Downloaded from http://bmjopen.bmj.com/ on June 9, 2025 at Agence Bibliographique de I Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

Table 2 Continued							
Included study	Location of study	Study design	Population under study	Type of publication	Main topic of the paper	Sample size	Quality rating*
Loomba ³⁴	UK and USA	Randomised controlled experiment	Other ethnic groups than white	Peer-reviewed journal	The impact of COVID-19 vaccine misinformation on vaccination intent	4000 (UK) 4001 (USA)	N/A§
Moyce et al ⁴¹	NSA	Qual interviews narrative synthesis	Latino individuals	Peer-reviewed journal	Perceptions of COVID-19, news about COVID-19 and approaches to protecting health	14	\$%06
Paul ²⁹	ПК	Repeated measures survey; cohort study	Other ethnic groups than white	Peer-reviewed journal	Attitudes towards COVID-19 vaccines, vaccine intent and implications for public health messaging	32361	89%†
Regional interagency coordination platform (R4V) ⁴⁸	Any host county for migrants from Venezuela	Cross-sectional survey	Predominantly migrants from Venezuela	Research-based report	The difficulties encountered by refugees and migrants in the COVID-19 infodemic. Misinformation and vaccine hesitancy	334	\$%06
Vekemans ³²	UK	Organisational case study	Jain community members	Peer-reviewed journal	The relocation of the Jain community into the digital realm during the COVID-19 pandemic	25000 estimate	N/A§
Viswanath ⁴²	NSA	Cross-sectional survey	Non-white ethnic groups	Peer-reviewed journal	Individual and social determinants of COVID-19 vaccine uptake	1012	78%†
Wang ⁴⁴	China	Cross-sectional survey	International migrants	Peer-reviewed journal	COVID-19 knowledge, attitudes and sources of knowledge among international migrants in China	1426	78%†
Woko ⁴³	USA	Cross-sectional survey	Black American citizens	Peer-reviewed journal	The role of beliefs and trust in COVID-19 information sources in low COVID-19 vaccination intention among black Americans	1074	77%†
Deal ³³	лк	Qualitative in-depth interview study	Precarious migrants (asylum seekers, undocumented migrants, refugees)	Peer-reviewed journal	Action points to promote the equitable uptake of COVID-19 vaccinations for precarious migrants	32	\$%06
*Scores were calculated †Quality assessment for ‡Critical Appraisal Skills I §N/A=not applicable due	on both scales by d Survey Studies in P Programme (CASP) to research item de	lividing the 'yes' answers t 'sychology for Surveys (Q- checklist. ssign.	oy the total number of ap SSP) Checklist for survey	plicable items. /s.			

6

Goldsmith LP, et al. BMJ Open 2022;12:e061896. doi:10.1136/bmjopen-2022-061896



Figure 2 Data on use of social media platforms as a source of information about COVID-19.

about COVID-19 in a survey of 334 migrants.⁴⁸ A survey of 774 refugee households in South-East Turkey⁴⁷ found the majority (75%) obtained COVID-19 information from Facebook, YouTube, Twitter and the internet in general, 15% via SMS/WhatsApp messages, followed by radio/TV (64%) and members of their community/ family (34%); only 10% reported getting information from non-governmental organisations (NGOs)/United Nations sources. This study concluded that the heavy reliance on social media for information may expose a sizeable proportion of refugee households to fake or inaccurate information. In a US study of black women aged 18–31 years, 58% of respondents reported using social media (Instagram and Facebook) to obtain COVID-19 information.³⁹

Participants from the US Latino community described relying on social media for information about the pandemic.³⁸ In Qatar, migrants reported they preferred to find out about COVID-19 using their own personal research or searching for information, including using social media as a source.⁴⁶ A study of precarious migrants (asylum seekers, undocumented migrants) in the UK found many relied on social media (WhatsApp groups, Facebook) for information on the pandemic and the ongoing vaccination programme.³³

A key theme emerging in one UK study of ethnic minority groups²⁸ was that the 'avalanche' of information surrounding COVID-19 had led to interviewees feeling overwhelmed and confused: participants reported using a variety of sources of information, including TV, radio, news stations in Pakistan, India, Slovakia and Poland, online newspapers, Facebook, WhatsApp, Twitter, Google, and medical journals. A number of these participants said they dismissed some stories encountered on WhatsApp and Facebook; however, the sheer volume of messages coupled with the fact that people they trusted were sharing them, proved difficult to ignore, with participants raising concerns about how quickly social media

stories were shared. One study exploring the views of US Latinos reported that they consulted national and local news reports for information about COVID-19 and many reported that they got their news from Spanish-language news due to difficulty in understanding news in English; some received their news from social media sources, including Facebook, but expressed caution around messages from social media as there was no way to ensure the accuracy of the reports.⁴¹ Language barriers were also reported among the Syrian refugee population in South-East Turkey who typically prefer information in Arabic,⁴⁷ as well as the Latino population in the USA,^{36 38 40} for people from a range of ethnic groups in Bradford²⁸ and for international migrants in another study.³¹

According to one study, members of ethnic minority \overline{g} groups were also more likely to post COVID-19 content on social media than white individuals,³⁷ with respondents who identified as black (OR 1.29, 95% CI 1.02 to 1.64; p=0.03), Latino (OR 1.66, 95% CI 1.36 to 2.04; p<0.001) or other races/ethnicities (OR 1.33, 95% CI 1.02 to 1.72; p=0.03) having higher odds than respondents who identified as white of reporting posting COVID-19 content on social media.

Drivers of social media reliance

Studies reported that some migrant and ethnic minority groups turned to social media as a result of a need for connection and to acquire accessible information from people they considered to be reliable sources.^{32 41} For the Latino community in the USA, faith and community bonds were valued ways of coping with the difficulties of the pandemic which included feelings of social isolation, stress and uncertainty and—according to one study—social media facilitated these connections in a virtual space.⁴¹ The Jain community in London used social media to communicate news and knowledge about COVID-19 and stay connected online, with events moving to a virtual

space; individuals reportedly benefited from and were grateful for this community use of social media.³²

Several studies highlight concerns that some migrant and ethnic minority groups were unable to find official information in their host country in their native language about various aspects of COVID-19, hence their reliance on social media.^{28 31 33 44 47} For example, a UK study of precarious migrants (asylum seekers, undocumented migrants) reported that those feeling most abandoned or scared due to a lack of understandable, clear official information in the early stages of the pandemic were more likely to rely on word of mouth or social media (WhatsApp groups, Facebook) for information, including around the vaccination programme.³³ One study of international migrants in China (94.5% of whom preferred social media for news about COVID-19) had lower rates of correct knowledge about COVID-19 compared with rates reported for Chinese residents.⁴⁵ The authors speculate that this might be due to a lack of available public health information in a range of languages.

Other studies showed positive associations with use of social media and access to information. One study highlighted that social media can support migrants to navigate the complex medicolegal context of their host countries by accessing information about public health measures and how to access medical help.36 Social media use was associated with improved knowledge about COVID-19 and how to stay safe, in studies of Syrian refugee mothers⁴⁵ and US Latinos.⁴⁰ In another study specifically curated, culturally relevant digital content was considered to be an effective health promotion tool to share knowledge about practical actions to be taken to address the inequitable impact of the pandemic on US Latinos.⁴⁰

Misinformation and social media use

A summary of some of the key misinformation narratives identified in studies is provided in table 3. Some studies made links between social media and circulating misinformation in migrant and ethnic minority groups. For example, a UK cohort study found that belonging to an ethnic minority group and socioeconomic disadvantage was associated with exposure to misinformation about vaccines and mistrust in information about COVID-19.²⁹ A study of Syrian refugee mothers in Jordan, who reported receiving most of their COVID-19 information through social media, identified some erroneous beliefs about pregnancy, COVID-19 and breast milk.⁴⁵ A UK study among ethnic minority groups reported that participants encountered a range of misinformation, usually through social media sources and that vaccine hesitancy could be attributed to safety concerns, negative stories and personal knowledge, all of which had been amplified by recent exposure to misinformation via social media.²⁸ Myths identified included the idea that health professionals at the local hospital were injecting people with COVID-19 or killing people with the COVID-19 vaccine; there were wider beliefs reported about vaccines containing a microchip; making people infertile, or that

<page-header><text><text><text><text><text>

Table 3 Examples of circulating misinformation on social media platforms relating tfrom Loomba et al^{34} and Lockyer et al^{28}	o COVID-19,	reproduced and	compiled
Misinformation identified	Source	Engagement*	Reach†
"Scientists have expressed doubts over the effectiveness of a COVID-19 vaccine that has been rushed to human trials, after all the monkeys used in initial testing later contracted COVID-19."	Twitter	1.59 K	1.5m
"The new vaccine for COVID-19 will be the first of its kind EVER. It will be an mRNA vaccine which will literally alter your DNA. It will wrap itself into your system. You will essentially become a genetically modified human being"	Twitter	27	19.6 K
"They said it was just to flatten the curve. Now it's a battle for human survival." The only must-see action thriller for 2020. Starring: Bill Gates, Anthony Fauci, Chris Witty, Matt Hancock. Guest mask appearances: Clintons, Boris Johnson, Nicola Sturgeon, Joe Biden & Tedros. (Graphic featuring Mr. Bill Gates with the following quote.) "If we do a really good job with vaccines, we can reduce population by up to 15%. But if we create a worldwide pandemic first, killing people and making many of the survivors sterile, then create the vaccine, we may achieve the Georgia Guidestones first commandment!"	Twitter	11	1.49 K
Something is very fishy about all this indeed. "A VIRUS WITH A 99.6% SURVIVAL RATE FOR PEOPLE UNDER 70 BUT THE ENTIRE WORLD NEEDS TO TAKE A VACCINE? I'M NO SHERLOCK HOLMES BUT SOMETHING'S FISHY ABOUT ALL THAT"	Twitter	NK	32.5 K
"Big Pharma whistle-blower: '97% of corona vaccine recipients will become infertile"	Twitter	6.95 K	336 K
"I've been in Twitter jail for the last 12 hours for posting a link to a peer reviewed scientific study published in Vaccine showing that in military personnel prior receipt of the influenza shot increased COVID-19 risk by 36%. Censorship is vile & unAmerican."	Twitter	25.1 K	1.41 K
"So we know for a fact that the influenza vaccine worsens COVID-19 symptoms. So what are they mandating now? The influenza vaccine, of course."	Facebook	NK	NK
"PREPARING THE PROPAGANDA BLITZ. Yale University and the U.S. government are running clinical trials to develop propaganda messaging to persuade Americans to take experimental, genetically engineered, unlicensed, "Warp Speed," zero liability, expedited vaccines with limited short duration safety testing. Researchers compared reactions in 12 focus groups using "guilt, embarrassment, bravery, anger, trust" and "fear" to overcome vaccines hesitancy"	Instagram	28.2 K	NK
 COVID-19 is not real, it is an effort to control society COVID-19 has been manufactured by China or other governments for control purposes COVID-19 is caused by 5G COVID-19 has been invented to make people use contactless payments so that the government can track individuals COVID-19 testing gives so many false positives that it is ineffective and you should not self-isolate COVID-19 exists but is not as virulent as the government says it is If children test positive for COVID-19 during school hours, they can be taken away into care and will not be able to see their parents until they test negative The COVID-19 vaccine contains a chip that will track individuals, stop them travelling, and so on The COVID-19 vaccine will make people infertile and is an attempt to reduce the population, particularly targeted at people from BAME communities BAME people are being used as 'guinea pigs' to test out the COVID-19 vaccine The COVID-19 vaccine will negatively disrupt your natural immune system Herbal remedies will be more effective than the COVID-19 vaccine 	Multiple platforms/ unknown	NK	NK
*Engagement measures the number of likes and retweets. †Reach measures the number of followers and thus potential audience size.			

NK, not known.

vaccine hesitancy due to misinformation circulating on social media and word of mouth combined with a lack of accurate, translated and clear guidance.³¹ Similarly, in a US qualitative study of Latino adults, some participants reported encountering a lack of knowledge accompanied by misinformation on social media causing them to dismiss preventative measures.³⁸ Another US study among Latino people reported that social media acted as a potential deterrent for following some public health measures to prevent infection by encouraging rulebreaking behaviour through socially normalising such behaviour by enabling people to observe the negative, guideline-breaking behaviours of others in social media posts.⁴¹

On the other hand, a large (8001 participants) US/UK randomised controlled experiment³⁴ found no significant differences in the response of different ethnic groups to misinformation in relation to vaccine intent. A large US/UK study³⁵ found membership of an ethnic minority group was associated with reduced vaccine intention, a relationship which was significant in three out of four studies (p<0.001, n=3890; p=0.017, n=1663; p<0.001, n=2237). The relationship persisted even when use of legacy (print and broadcast media) and frequency of use of social media was controlled for. High levels of social media use was not associated with vaccine intent in any of the three studies exploring this relationship; however, high information reliance on social media was significantly associated with negative vaccine intent (p=0.028, n=2237), suggesting a reliance on social media for information can make users vulnerable to misinformation. This study did not include interaction terms between ethnicity and information reliance on social media, which could have indicated whether the effect of information reliance on social media on vaccine intent differs by ethnicity.

Good practice in promoting information and countering misinformation

Evidence suggests the important role of strong connections with the local community to identify and counter misinformation and rumours by trusted and valued sources of information. Most studies recommended improving the accessibility of public health information for migrant and ethnic minority communities.^{29 31 36 38 39 41-44 48} For example, providing public health information in the media channel preferred by that group,³⁸ in multiple languages,²⁸ and using local, trusted voices delivering specific and targeted messages to counter fake news.^{28 38} A strong interest in online, personalised information was identified.^{40 41} Where social media was used to share personalised and culturally tailored public health information, it has a positive influence with good health knowledge, health-seeking behaviours and vaccine intent.^{28–31 36 38–40 42–45 48} Studies indicated the need for culturally tailored health messaging to ensure equitable health knowledge for improving vaccine intent and health-seeking behaviours.^{31 38-40}

<page-header><text><text><text><text><text>

about COVID-19 for some migrant and ethnic minority populations during the pandemic. While we know social media is used by many people, and misinformation has been circulating widely in the general population, it may be the case that those excluded from national public health responses and/or who faced specific barriers to accurate public health information and support may have been disproportionately impacted. Urgent actions and further research are now needed to better understand use of social media platforms for health information in different population groups, find effective approaches to tackling misinformation, and to seize on opportunities to make better use of social media platforms to support public health communication and improve vaccine uptake globally. Furthermore, the findings highlight the crucial role of locally trusted sources in identifying and tackling misinformation, and underscores the benefits of disseminating personalised and culturally relevant health messages, including via social media.

This review is the first attempt to synthesise global studies exploring the use and impact of social media on migrant and ethnic minority populations during the COVID-19 pandemic. However, it is limited by the availability and quality of the data sets available. We acknowledge the limited geographical scope of included studies, with 16 of 21 studies focused on migrant and ethnic minority populations residing in the UK and USA and no data at all from low-income countries. It may be that the lower availability of research funding in low-income countries may explain the lack of studies from these countries. We acknowledge that definitions and terms pertaining to migrants and ethnic minorities and social media are used inconsistently in research; this is an ongoing challenge within the field, which has previously been evidenced in similar reviews, and may mean we have missed papers. This was mitigated against by searching the published and grey literature more widely. We also acknowledge that many of the surveys didn't formally report whether the social media feeds their responders were following were from 'official' sources (such as government or NGOs or from 'unofficial' sources), or from friends, relatives, or from accounts simply with many followers. However, the included data sets, and further qualitative work our group is currently doing in the UK, suggest they will be predominantly unofficial sources, with government public health teams in several countries very slow to make effective use of social media as a platform of communication at the start of the pandemic. A further limitation is that there were insufficient studies to reliably compare use of social media across type of migrants (refugees/asylum seeker, undocumented migrants), and future research should explore this. We acknowledge that migrants and ethnic minorities are a highly diverse group with a range of health and socioeconomic situations making it hard to generalise; however, there is evidence in several contexts that these populations may have been disproportionately impacted by the COVID-19 pandemic.^{13 16 49}

The findings of our review have been confirmed by more recent studies. For example, a survey of migrants in Greece found their main source of information about the vaccine was via social media platforms and the internet in general, and that vaccine hesitancy was linked to a lack of adequate information and driven by fear, anxiety, exposure to negative news and misinformation.^{50 51} In Turkey, a 2021 survey and feedback mechanism --information gaps, misconceptions and rumours as--COVID-19 vaccines circulating mainly by word of mouth rescial media, undermining health information.⁵² 70% said they had access to a mobile phone, with the main communication channels being WhatsApp and Facebook, yet half said they felt uninformed.⁵³ We also found that **8** some migrants and ethnic minorities used diaspora media as a source of COVID-19-related information during the pandemic, which merits further consideration in terms of understanding how to better engage these groups in preventative healthcare and vaccination, and has been previously reported in studies as influencing views and beliefs around vaccination.⁵⁴ Misinformation on social media correlated negatively with vaccine intention and our findings align uses rela with other research in this area and will undoubtedly be relevant to many other population groups.^{2 7 33} A recent study among migrants and nationals in Qatar acknowledged 'personal research' via social media as important to them for seeking information about COVID-19 vaccines, underlining the key role social media has in influencing people's attitudes towards vaccine uptake.⁴⁶

The European Centre for Disease Prevention and Control and other public health bodies have raised <u>o</u> concerns around barriers to public health information a among migrant populations and ethnic minority groups \blacksquare residing in Europe and other high-income countries during the pandemic.¹³ ¹⁴ Public health guidance in some countries was not initially tailored to the needs of migrant and ≥ ethnic minority groups.^{18 55–57} A review of the availability of training, government-produced risk communications across Council of Europe member states in June 2021 found only 48% (23/47) of countries translated COVID-19 information into at least one migrant language, with information on testing S or healthcare entitlements in common migrant languages only found in 6% (3/47), suggesting individuals not able to access information in the host country language may have been excluded to some extent from governments' public health messaging.¹⁸ In Denmark, a series of qualitative interviews with migrants found that they felt uncertain & regarding government guidance for COVID-19; although **3** written material was translated into 19 languages, it was not effectively disseminated.⁵⁸ In Montreal, Canada, there were delays to publishing official multilingual fact sheets on COVID-19 guidelines, and information phone lines only operate in French and English; those who had arrived most recently, had lower language (French/English) ability or lower literacy, had more difficulty accessing local COVID-19 information.⁵⁹ Lack of English or French language at the time of immigration to Canada were associated with lower

BMJ Open: first published as 10.1136/bmjopen-2022-061896 on 17 November 2022. Downloaded from http://bmjopen.bmj.com/ on June 9, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES)

rates of testing and higher per cent positivity for COVID-19 in recently arrived adult immigrants and refugees.⁶⁰ A study among refugees and migrants in deprived areas in Greece found that migrants may have difficulties understanding public health messaging due to cultural and language barriers.^{18 56 61} Merely translating public health information is not likely to be sufficient; information needs to be tailored and targeted so it is conveyed in ways that resonate with the target population. A range of key resources and guidelines on risk communication and engagement strategies for COVID-19 public health responses, including vaccination, among marginalised populations globally are available, as well as a social media toolkit for healthcare practitioners (https://www.who.int/publications/m/item/a-socialmedia-toolkit-for-healthcare-practitioners-desktop). However, it will be vitally important that the lessons learnt around communication of public health information to marginalised groups during the pandemic are meaningfully carried forward.

Where social media is used to share personalised and culturally tailored public health information, it has a positive correlation with good health knowledge, health seeking behaviours and vaccine intent.^{40 45} Our research shows the need for culturally tailored health messaging to ensure equitable health knowledge and to improve vaccine uptake, by accurate public health messaging through trusted sources of information.^{31 38–40} We make a number of recommendations for policy and practice, which include the need for systematic monitoring of information and attitudes circulating on social media,⁶⁵ as well as timely rebuttal of misinformation from trusted professionals (see box 1). Several resources are now available to support addressing misinformation about COVID-19 vaccines as well as fostering demand for vaccines.^{66–68}

There is a stark lack of data on social media use from low-income and middle-income countries, which merits greater consideration as COVID-19 vaccination gathers pace in these contexts. Studies from high-income countries are also limited, with the majority of studies focused on the USA and UK. In addition, more evidence is needed to examine the role social media platforms play in positively or negatively influencing health behaviours such as vaccine intent and uptake for COVID-19 in all populations (including other excluded groups, for example, homeless, internally displaced people). Social media is an important source of health information for some migrant and ethnic minority communities and tackling misinformation needs to be done using this medium given the lack of trust in government messaging in some of these communities.⁶⁹ Our findings are consistent with those of others working in this field, which show that social media can have a crucial role in disseminating health information, tackling infodemics and misinformation.⁴ There is an opportunity now to more effectively use social media to make vaccine intent desirable, appealing and normative among migrants and ethnic minority groups. There is an urgent need to address infodemic-related challenges in a rapidly changing information environment,

Box 1 Key messages and recommendations

- ⇒ Social media is an important source of health information for some migrant and ethnic minority communities, who may face barriers to accurate public health information, health and vaccinations systems. More evidence is urgently needed to examine the role social media platforms play in positively or negatively influencing health behaviours such as vaccine intent and uptake for COVID-19 in marginalised populations.
- ⇒ There is a stark lack of data on social media use from low-income and middle-income countries which merits greater consideration as COVID-19 vaccination gathers pace in these contexts.
- ⇒ More emphasis must be placed on exploring opportunities for sharing and transmitting accurate information via social media platforms, for example, to make vaccine intent desirable, appealing and normative.
- ⇒ Use of diaspora media by migrant populations, as a source of COVID-19-related information during the pandemic and for other health information, merits further research and greater consideration when designing and delivering public health interventions.
- ⇒ Proactively monitor social media platforms and other media sources to identify antivaccine sentiment, misinformation, fake news and rumours, and address them in real time.
- ⇒ There is a need to promote targeted and tailored health information to marginalised populations who face access barriers to health and vaccination systems, through preferred and trusted sources and channels of information including social media platforms, and to ensure investment in workforce and infrastructure to support this.
- ⇒ Engage with and involve communities in developing culturally specific messages and approaches, and support community-driven initiatives to identify at-risk groups, map local influencers, and define content for locally meaningful communication campaigns. Facilitate partnership working at the local level through involvement of diverse stakeholders and ensure community partners are recognised and reimbursed for their contributions and expertise.
- ⇒ Social media platforms should exercise more accountability and sign pledges to systematically track and remove harmful content that undermine public health measures, particularly during a public health crisis. The public must be empowered to identify and flag misinformation on social media.
- ⇒ Public health bodies and healthcare professionals should avoid a narrow focus on misinformation and a one-way communication of 'more accurate' information. They should seek to understand the underlying causes of exposure to and belief in misinformation including genuine knowledge void, access barriers and health literacy.
- ⇒ Lessons must be learnt around shortfalls in the communication of public health information to marginalised groups during this pandemic. Importantly, countries should gather and evaluate innovations and models of best practice in this area, which must be meaningfully carried forward to strengthen uptake of routine vaccinations and other public health interventions.

including real-time monitoring of social media messages and misinformation and the development of online tools to fight disinformation, with a focus on collecting stratified population data to enable targeted and tailored responses. Robust interventions relying on behavioural science to tackle misinformation using social media and evaluations are a plausible next step to address immunisation challenges for COVID-19 vaccines and also routine vaccines. Building trust in public health messaging, <u>ð</u>

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

identifying information gaps, finding innovative ways of disseminating health information, and detecting and responding to misinformation as it emerges remain a priority for public health.^{69 70}

Author affiliations

¹The Migrant Health Research Group, Institute for Infection and Immunity, St George's University of London, London, UK

²Population Health Research Institute, St George's University of London, London, UK ³Faculty of Health, Social Care and Education, Kingston University, Kingston-Upon-Thames, UK

⁴Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, London, UK

⁵Institute of Medical and Biomedical Education, St George's University of London, London, UK

⁶Faculty of Health, Social Care and Education, Centre for Applied Health and Social Care Research, Kingston University, Kingston, UK

Twitter Alison F Crawshaw @AlisonCrawshaw, Tushna Vandrevala @psych_tush and Sally Hargreaves @sal_hargreaves

Contributors The study was conceptualised by SH, and the protocol and research question were developed by SH, LPG and MR-P. Searches were developed by MR-P and LPG, with input from SH and SEH. Screening was done by LPG and MR-P. Data extraction and analysis were done by LPG and MR-P, with input from SH and KH. The first draft of the manuscript was produced by LPG, MR-P and SH, and developed with KH and TV, who all contributed to interpretation of the results. All authors, including AD, AFC, SEH, FK, JC, AA and MR commented upon and approved the final manuscript. SH is guarantor of this study.

Funding This study was funded by the National Institute for Health Research (NIHR300072), the Academy of Medical Sciences (SBF005\1111), and the Medical Research Council (MRC/N013638/1). LPG, SH and AFC are funded by the NIHR (NIHR300072); AFC and SH are funded by the Academy of Medical Sciences (SBF005\1111). SH acknowledges funding from the Novo Nordisk Foundation (Mobility–Global Medicine and Health Research) and the WHO. AD and SEH are funded by the Medical Research Council (MRC/N013638/1). MR is funded by an NIHR In-Practice Clinical Fellowship (NIHR 302007). JC is funded by an NIHR In-Practice Clinical Fellowship (NIHR300290).

Disclaimer The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care. The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. This is a systematic review of published data.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: https://creativecommons.org/ licenses/by/4.0/.

ORCID iDs

Lucy Pollyanna Goldsmith http://orcid.org/0000-0002-6934-1925 Alison F Crawshaw http://orcid.org/0000-0003-0450-7258 Sally E Hayward http://orcid.org/0000-0002-4105-0990 Jessica Carter http://orcid.org/0000-0001-9590-3146 M Razai http://orcid.org/0000-0002-6671-5557 Sally Hargreaves http://orcid.org/0000-0003-2974-4348

REFERENCES

- 1 WHO. Health topics/Infodemic. Available: https://www.who.int/ health-topics/infodemic#tab=tab_1
- 2 Puri N, Coomes EA, Haghbayan H, et al. Social media and vaccine hesitancy: new updates for the era of COVID-19 and globalized infectious diseases. *Hum Vaccin Immunother* 2020;16:2586–93.
- 3 Li HO-Y, Bailey A, Huynh D, et al. Youtube as a source of information on COVID-19: a pandemic of misinformation? BMJ Glob Health 2020;5:e002604.
- 4 Tsao S-F, Chen H, Tisseverasinghe T, *et al.* What social media told us in the time of COVID-19: a scoping review. *Lancet Digit Health* 2021;3:e175–94.
- 5 Basch CH, Zybert P, Reeves R, et al. What do popular YouTube[™] videos say about vaccines? Child Care Health Dev 2017;43:499–503.
- 6 Larson HJ. The biggest pandemic risk? viral misinformation. *Nature* 2018;562:309.
- 7 Wilson SL, Wiysonge C. Social media and vaccine hesitancy. BMJ Glob Health 2020;5:e004206.
- 8 Roozenbeek J, Schneider CR, Dryhurst S, et al. Susceptibility to misinformation about COVID-19 around the world. R Soc Open Sci 2020;7:201199.
- 9 American Press Institute. Race, Ethnicity, and the Use of Social Media for News. How Millennials Use Technology to Get News, 2015. Available: https://www.americanpressinstitute.org/publications/ reports/survey-research/race-ethnicity-social-media-news/
- 10 Gorman DR, Bielecki K, Willocks LJ, et al. A qualitative study of vaccination behaviour amongst female Polish migrants in Edinburgh, Scotland. *Vaccine* 2019;37:2741–7.
- 11 World Health Organization. 10 global health issues to track in 2021. Spotlight. Available: https://www.who.int/news-room/spotlight/10global-health-issues-to-track-in-2021
- 12 Činelli M, Quattrociocchi W, Galeazzi A, *et al*. The COVID-19 social media infodemic. *Sci Rep* 2020;10:16598.
- 13 Hayward SE, Deal A, Cheng C, et al. Clinical outcomes and risk factors for COVID-19 among migrant populations in high-income countries: a systematic review. J Migr Health 2021;3:100041.
- 14 ECDC. Reducing COVID-19 transmission and strengthening vaccine uptake among migrant populations in the EU/EEA. technical report. Stockholm: ECDC, 2021.
- 15 Khanijahani A, lezadi S, Gholipour K, et al. A systematic review of racial/ethnic and socioeconomic disparities in COVID-19. Int J Equity Health 2021;20:248.
- 16 Hargreaves S, Hayward SE, Noori T, *et al*. COVID-19: counting migrants in. *Lancet* 2021;398:211–2.
- 17 Turkish Red Crescent Society CCs, Fatma Nur Bakkalbaşı, Onurcan Ceyhan, Tenadi Gölemerz. COVID-19 Rumour tracking report: Turkish red crescent Society (TRCS) in collaboration with international Federation of red cross and red crescent societies (IFRC, 2021.
- 18 Nezafat Maldonado BM, Collins J, Blundell HJ, et al. Engaging the vulnerable: a rapid review of public health communication aimed at migrants during the COVID-19 pandemic in Europe. J Migr Health 2020;1:100004.
- 19 Rowland-Pomp M, Hargreaves S, Goldsmith LP. The impact of social media and misinformation on migrant and ethnic minorities populations and their response to the COVID-19 pandemic: a systematic review. prospero 2021 CRD42021259190 2021.
- 20 Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- Organisation. WH. Archived: WHO Timeline COVID-19, 2020.
 World Health Organization. COVID-19 global literature on coronavirus
- 22 World Health Organization. COVID-19 global interature on coronavirus disease database. Available: https://www.who.int/emergencies/ diseases/novel-coronavirus-2019/global-research-on-novelcoronavirus-2019-ncov
- 23 Ouzzani M, Hammady H, Fedorowicz Z, *et al.* Rayyan-a web and mobile APP for systematic reviews. *Syst Rev* 2016;5:210.
- 24 Protogerou C, Hagger MS. A checklist to assess the quality of survey studies in psychology. *Meth Psychol* 2020;3:100031.
- 25 Critical Appraisal Skills Programme. Casp qualitative checklist, 2018. Available: https://casp-uk.net/casp-tools-checklists/

Open access

- 26 Pluye P, Hong QN. Combining the power of stories and the power of numbers: mixed methods research and mixed studies reviews. *Annu Rev Public Health* 2014;35:29–45.
- 27 Lisy K, Porritt K. Narrative synthesis. *Int J Evid Based Healthc* 2016;14:201.
- 28 Lockyer B, Islam S, Rahman A, et al. Understanding COVID-19 misinformation and vaccine hesitancy in context: findings from a qualitative study involving citizens in Bradford, UK. *Health Expect* 2021;24:1158–67.
- 29 Paul E, Steptoe A, Fancourt D. Attitudes towards vaccines and intention to vaccinate against COVID-19: implications for public health communications. *Lancet Reg Health Eur* 2021;1:100012–12.
- 30 Allington D, McAndrew S, Moxham-Hall V, et al. Coronavirus conspiracy suspicions, general vaccine attitudes, trust and coronavirus information source as predictors of vaccine hesitancy among UK residents during the COVID-19 pandemic. *Psychol Med* 2021:1–12.
- 31 Crawshaw AF, Deal A, Rustage K, et al. What must be done to tackle vaccine hesitancy and barriers to COVID-19 vaccination in migrants? *J Travel Med* 2021;28:taab048.
- 32 Vekemans T. Crisis and continuation: the digital relocation of Jain Socio-Religious praxis during the COVID-19 pandemic. *Religions* 2021;12:342–42.
- 33 Deal A, Hayward SE, Huda M, et al. Strategies and action points to ensure equitable uptake of COVID-19 vaccinations: a national qualitative interview study to explore the views of undocumented migrants, asylum seekers, and refugees. J Migr Health 2021;4:100050.
- 34 Loomba S, de Figueiredo A, Piatek SJ, et al. Measuring the impact of COVID-19 vaccine misinformation on vaccination intent in the UK and USA. Nat Hum Behav 2021;5:337–48.
- 35 Allington D, McAndrew S, Moxham-Hall VL, et al. Media usage predicts intention to be vaccinated against SARS-CoV-2 in the US and the UK. Vaccine 2021;39:2595–603.
- 36 Behbahani S, Smith CA, Carvalho M, et al. Vulnerable immigrant populations in the new York metropolitan area and COVID-19: lessons learned in the epicenter of the crisis. Acad Med 2020;95:1827–30.
- 37 Campos-Castillo C, Laestadius LI. Racial and ethnic digital divides in Posting COVID-19 content on social media among US adults: secondary survey analysis. J Med Internet Res 2020;22:e20472–e72.
- 38 Cervantes L, Martin M, Frank MG, et al. Experiences of Latinx individuals hospitalized for COVID-19: a qualitative study. JAMA Netw Open 2021;4:e210684–e84.
- 39 Chandler R, Guillaume D, Parker AG, et al. The impact of COVID-19 among black women: evaluating perspectives and sources of information. *Ethn Health* 2021;26:80–93.
- 40 Despres C, Aguilar R, McAlister A, et al. Communication for awareness and action on inequitable impacts of COVID-19 on Latinos. *Health Promot Pract* 2020;21:859–61.
- 41 Moyce S, Velazquez M, Claudio D, et al. Exploring a rural Latino community's perception of the COVID-19 pandemic. Ethn Health 2021;26:126–38.
- 42 Viswanath K, Bekalu M, Dhawan D, et al. Individual and social determinants of COVID-19 vaccine uptake. BMC Public Health 2021;21:818–18.
- 43 Woko C, Siegel L, Hornik R. An investigation of low COVID-19 vaccination intentions among black Americans: the role of behavioral beliefs and trust in COVID-19 information sources. *J Health Commun* 2020;25:819–26.
- 44 Wang C, Tian Q, Zhao P, et al. Disease knowledge and attitudes during the COVID-19 epidemic among international migrants in China: a national cross-sectional study. Int J Biol Sci 2020;16:2895–905.
- 45 Hamadneh S, Hamadneh J, Amarin Z, et al. Knowledge and attitudes regarding Covid-19 among Syrian refugee women in Jordan. Int J Clin Pract 2021;75:e14021–e21.
- 46 Alabdulla M, Reagu SM, Al-Khal A, et al. COVID-19 vaccine hesitancy and attitudes in Qatar: a national cross-sectional survey of a migrant-majority population. *Influenza Other Respir Viruses* 2021;15:361–70.
- 47 Danish Refugee Council. COVID-19 impact on refugees in South East turkey, 2020. Available: https://data2.unhcr.org/en/needsassessment/1432
- 48 (R4V) RI-aCP. Information and communication needs assessment -U-Report Uniendo Voces Regional Poll, 2021.
- 49 Buikema AR, Buzinec P, Paudel ML, et al. Racial and ethnic disparity in clinical outcomes among patients with confirmed COVID-19 infection in a large US electronic health record database. EClinicalMedicine 2021;39:101075.

- 50 Hellenic Red Cross. CEA. Survey on the migrant's population informatio needs, regarding health issues (COVID-19). Available: https://communityengagementhub.org/resource/survey-on-themigrants-population-information-needs-regarding-health-issuescovid-19/
- 51 Hellenic Red Cross. Perceptions survey on Covid-19 vaccination, Greece. Available: https://communityengagementhub.org/resource/ perceptions-survey-on-covid-19-vaccination-greece/
- 52 Turkish Red Crescent Society. COVID-19 rumour tracking report. Ankara, 2021. Available: https://communityengagementhub.org/wpcontent/uploads/sites/2/2021/10/Rumour-tracking-report-2021.pdf [Accessed 11 Jan 2022].
- 53 Only half of refugees and migrants from Venezuela feel informed, survey finds. Panama City: coordination platform for refugees and migrants from Venezuela, 2020. Available: https://www.r4v.info/sites/ default/files/2021-06/CWC%20EN.pdf [Accessed 11 Jan 2022].
- 54 Ganczak M, Bielecki K, Drozd-Dąbrowska M, et al. Vaccination concerns, beliefs and practices among Ukrainian migrants in Poland: a qualitative study. BMC Public Health 2021;21:93.
- 55 Patel P, Hiam L, Orcutt M. Policy brief: Including migrants and refugees in the British government's response to COVID-19 2020.
- 56 Doctors of the World. *An unsafe distance: the impact of the COVID-19 pandemic on excluded people in England*, 2020.
- 57 Kondilis E, Papamichail D, McCann S, *et al.* The impact of the COVID-19 pandemic on refugees and asylum seekers in Greece: a retrospective analysis of national surveillance data from 2020. *EClinicalMedicine* 2021;37:100958.
- 58 Institut for Menneskerettigheder. Corona rammer skævt etnicitet og smitte. Copenhagen, Denmark, 2020.
- 59 Cleveland J, Hanley J, Jaimes A. Impacts de la crise de la COVID-19 sur les «communautés culturelles »montréalaises: Enquête sur les facteurs sociocul- turels et structurels affectant les groupes vulnérables. Montréal: Institut universitaire SHERPA, 2020.
- 60 Guttmann A, Gandhi S, Wanigaratne S. COVID-19 in immigrants, refugees and other newcomers in Ontario: characteristics of those tested and those confirmed positive. Toronto, ON: ICES, 2020.
- 61 Cholera R, Falusi OO, Linton JM. Sheltering in place in a xenophobic climate: COVID-19 and children in immigrant families. *Pediatrics* 2020;146:e20201094.
- 62 World Health Organization. Conducting community engagement for COVID-19 vaccines: interim guidance. Geneva: World Health Organization, 2021. https://apps.who.int/iris/handle/10665/339451
- 63 World Health Organization. COVID-19 immunization in refugees and migrants: principles and key considerations: interim guidance, 31 August 2021. Geneva: World Health Organization, 2021. https://www. who.int/publications/i/item/covid-19-immunization-in-refugees-andmigrants-principles-and-key-considerations-interim-guidance-31august-2021
- 64 British Red Cross. Good practice guidance for risk communication and community engagement (RCCE) for refugees, internally displaced persons (IDPs), migrants and host communities particularly vulnerable to COVID-19 pandemic. London: British Red Cross, 2020. https://communityengagementhub.org/wp-content/ uploads/sites/2/2020/06/Practical-Guidance-RCCE-Refugees-IDPs-Migrants.pdf
- 65 LSHTM. The vaccine confidence project. media monitoring report UK COVID-19. London: LSHTM, 2021. https://www.vaccineconfidence. org/research-feed/social-media-conversations-and-attitudes-in-the-uk-towards-covid-19
- 66 United Nations Children's Fund. *Vaccine misinformation management field guide: guidance for addressing a global infodemic and fostering demand for immunization.* New York: United Nations Children's Fund, 2020. https://vaccinemisinformation.guide
- 67 Pan American Health Organization. Understanding the infodemic and misinformation in the fight against COVID-19: digital transformation toolkit. Washington, DC: Pan American Health Organization, 2020. https://iris.paho.org/bitstream/handle/10665.2/52052/Factsheetinfodemic_eng.pdf?sequence=16
- 68 United States Centers for Disease Control and Prevention. *How to address COVID-19 vaccine misinformation*. Atlanta, GA, 2021. https://www.cdc.gov/vaccines/covid-19/health-departments/ addressing-vaccine-misinformation.html
- 69 Vandrevala, Hendy, Hanson, Alidu & Ala. Unpacking COVID-19 and conspiracy theories in the UK black community 2022.
- 70 Vandrevala T, Alidu L, Hendy J, et al. 'It's possibly made us feel a little more alienated': how people from ethnic minority communities conceptualise COVID-19 and its influence on engagement with testing. J Health Serv Res Policy 2022;27:141–50.
- Statista, 2021. Available: https://www.statista.com [Accessed 17 Sep 2021].