



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy: Protocol for a randomised controlled trial.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058683
Article Type:	Protocol
Date Submitted by the Author:	25-Oct-2021
Complete List of Authors:	Contreras-Hernandez, Ignacio; University of Birmingham Falla, Deborah; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences Martinez-Valdes, Eduardo; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences
Keywords:	SPORTS MEDICINE, Adult orthopaedics < ORTHOPAEDIC & TRAUMA SURGERY, Foot & ankle < ORTHOPAEDIC & TRAUMA SURGERY, Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY

SCHOLARONE™  
Manuscripts

Title

Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy: Protocol for a randomised controlled trial.

**Authors:** Ignacio Contreras-Hernandez<sup>1</sup>, Deborah Falla<sup>1</sup>, Eduardo Martinez-Valdes<sup>1</sup>

Corresponding author: Eduardo Martinez-Valdes<sup>1</sup>  
Lecturer (Assistant Professor) in Physiotherapy  
Centre of Precision Rehabilitation for Spinal Pain (CPR Spine)  
School of Sport, Exercise and Rehabilitation Sciences  
College of Life and Environmental Sciences  
University of Birmingham  
Edgbaston B15 2TT, United Kingdom  
T: +44 (0) 121 4158187  
M: +44 (0) 7961 548851  
Email: [e.a.martinezvaldes@bham.ac.uk](mailto:e.a.martinezvaldes@bham.ac.uk)

Author Affiliations

<sup>1</sup>Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, UK.

Email: Ignacio Contreras-Hernandez<sup>1</sup> [iac921@student.bham.ac.uk](mailto:iac921@student.bham.ac.uk) – Deborah Falla<sup>1</sup> [d.falla@bham.ac.uk](mailto:d.falla@bham.ac.uk) – Eduardo Martinez-Valdes<sup>1</sup> [e.a.martinezvaldes@bham.ac.uk](mailto:e.a.martinezvaldes@bham.ac.uk)

**WORD COUNT MAIN TEXT: 5486 (excluding title page, abstract, references, figures, tables and acknowledgments)**

## Abstract

**Introduction.** There is limited evidence on the neural strategies employed by the central nervous system to control muscle force in the presence of non-insertional Achilles tendinopathy. Additionally, the neuromuscular mechanisms by which exercise may help to resolve tendon pain remain unclear.

**Objective.** This study aims to firstly determine the differences in pain and function, mechanical and structural properties of the Achilles tendon, and motor unit firing properties of the calf muscles in individuals with non-insertional Achilles tendinopathy compared with asymptomatic controls. Secondly, we want to establish changes in these properties following a 6-week training protocol based on either eccentric or concentric exercise in individuals with non-insertional Achilles tendinopathy.

**Methods and analysis.** A total of 26 individuals with chronic (> 3 months) non-insertional Achilles tendinopathy and 13 healthy controls will participate in the study. Individuals with non-insertional Achilles tendinopathy will be randomised to perform eccentric or concentric training for 6 weeks. Participants will complete a battery of questionnaires to document their physical activity level, symptoms of Achilles tendinopathy, physical function, pain catastrophizing, and fear of movement. Achilles tendon length, cross-sectional area, thickness, and stiffness will be assessed using B-mode ultrasonography and shear wave elastography, as well as motor unit firing properties of the medial gastrocnemius, lateral gastrocnemius and soleus muscles using high-density surface electromyography.

**Ethics and dissemination.** Ethical approval (ERN\_20-0604) for the study was obtained from the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham. The results of the study will be published in peer-review journals.

**Protocol Registration.** This protocol has been registered on ISRCTN registry (ISRCTN46462385).

**Keywords:** Achilles tendinopathy, morphological properties, mechanical features, HD-sEMG, motor unit.

**WORD COUNT: 261**



## Introduction

Achilles tendinopathy (AT) is a painful overuse injury of the Achilles tendon. This injury is common among athletes, especially those involved in running and jumping sports<sup>1-3</sup>. Among elite track and field athletes, 43% reported having either current or prior symptoms of AT, with the highest prevalence (83%) in middle-distance runners<sup>3</sup>. However, this pathology can also be present in a sedentary population.

AT is clinically diagnosed when the patient presents with a combination of localized pain, swelling of the Achilles tendon, and loss of function<sup>4</sup>. Achilles tendon injuries can be separated into non-insertional tendinopathy (55%-65% of the injuries), insertional tendinopathy (20%-25%), and proximal musculotendinous junction (9%-25%) injuries, according to the location of pain<sup>4</sup>. Nevertheless, individuals may present with symptoms at the insertion and mid-portion concurrently, and approximately 30% have bilateral pain<sup>5</sup>.

The essence of tendinopathy is a failed healing response, with degeneration and proliferation of tenocytes, disruption of collagen fibres, and subsequent increase in non-collagenous matrix<sup>6</sup>. In tendinopathic samples, there is unequal and irregular crimping, loosening and increased waviness of collagen fibres, with an increase in Type III (reparative) collagen<sup>6</sup>. These structural changes in the tendon result in increased cross-sectional area, reduced tendon stiffness, and altered viscoelastic properties in both symptomatic and asymptomatic tendons<sup>7</sup>.

The aetiology of AT remains debated and is likely caused by intrinsic and extrinsic factors. Decreased plantar-flexor strength, deficits in hip neuromuscular control, abnormal ankle dorsiflexion and subtalar-joint range of motion, increased foot pronation, and an increased body weight are intrinsic risk factors that can be addressed during treatment<sup>8</sup>. Systemic disease, genetic variants, and family history of tendinopathy have also been identified as intrinsic risk factors<sup>9</sup>. Instead, the type of footwear, sport participation, surface, training-load adjustment (i.e. sudden increase in training duration, mileage, or intensity) are considered as extrinsic risk factors<sup>8-10</sup>. One of the most accepted theories is that pain perception during early support loading may trigger inhibition of

neuromuscular activity of the calf muscles detected as a reduction in electromyographic (EMG) amplitude<sup>11 12</sup>. Thus, the decrease in the ability to generate force in patients with AT could also be a reflection of the decline in neuromuscular activity observed,<sup>13 14</sup> and it is believed that these motor adaptations may generate greater torsional stress in the tendon<sup>15 16</sup>. These motor impairments not only affect the maximal force exerted but also influence the ability of an individual to produce steady force during submaximal voluntary contractions<sup>17</sup>.

Until now, most studies examining the neuromuscular impairments induced by AT have focused on investigating changes in interference EMG amplitude which is an indirect estimate of neural activity with many factors of influence<sup>18-20</sup>. Clearer information about the neural strategies employed by the central nervous system to control muscle force in the presence of AT can be obtained through motor unit recordings, since motor unit firing properties represent the direct neural output from the spinal cord to muscles<sup>21</sup>. Nevertheless, there are no studies that have measured motor unit firing properties in individuals with non-insertional AT.

Although eccentric exercise has been widely used for the treatment of AT, evidence of histological changes following a program of eccentric exercise is lacking, and the mechanisms by which eccentric exercises may help to resolve tendon pain remain unclear<sup>22</sup>. Regarding eccentric exercise alone, two prospective studies have reported a significant reduction in pain intensity and change on the Victorian Institute of Sports Assessment-Achilles questionnaire (VISA-A) in recreational athletes following a 12-week exercise program<sup>23 24</sup>. In contrast, another study in non-athletic individuals found no significant improvement after a similar 12-week exercise program<sup>25</sup>. Concerning eccentric exercises with an adjunctive treatment (e.g., pulsed ultrasound, ice, sensory motor training), a 4-week intervention study resulted in decreased pain and higher plantarflexion peak torque in individuals with AT compared to controls<sup>26</sup>. However, studies that include eccentric exercises with an adjunctive treatment showed limited evidence of improvement over eccentric exercises alone<sup>27</sup>.

There are few studies where the effectiveness of eccentric versus concentric

exercises has been compared. Mafi et al. showed that patient satisfaction and return to previous activity were significantly superior after participating in a 12-week rehabilitation protocol based on eccentric exercise compared to concentric exercise. Although pain intensity decreased significantly in both groups, the amount of pain reduction was significantly greater for those that performed eccentric exercise<sup>28</sup>. Likewise, Yu et al.<sup>29</sup> demonstrated that eight weeks of eccentric exercise was more effective at reducing pain than concentric in individuals with chronic AT. Additionally, they found that eccentric exercise was more effective than concentric exercise at increasing muscle strength and endurance, and improving function<sup>29</sup>. In these investigations, participants performed the rehabilitation protocols with insufficient control over the load, speed, pain tolerance, or the range of motion in which the exercises were performed. Moreover, it is essential to consider that when participants perform an eccentric plantar flexion exercise without adequate equipment, it is difficult to achieve pure eccentric contractions, which could have influenced the results obtained in these studies.

Based on the above, the aims of this study are to: (1) compare the mechanical and structural properties of the Achilles tendon, and gastrocnemius-soleus motor unit firing properties in individuals with non-insertional AT compared with asymptomatic controls; (2) establish changes in the level of pain and function, mechanical and structural properties of the Achilles tendon, and gastrocnemius-soleus motor unit firing properties after applying a training protocol of 6-week based on either controlled eccentric or concentric contractions in individuals with non-insertional AT.

## Methods and Analysis



1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

**Participants**

Twenty-six individuals with non-insertional AT and thirteen asymptomatic controls will be recruited from the University of Birmingham staff/student population and the local community via leaflets, e-mail, and social media.

Men or women aged 18 to 55 years old will be recruited. This age range was selected based on previous findings showing lower stiffness and Young’s modulus of the Achilles tendon in older than younger population<sup>30</sup>. Inclusion criteria for those with AT are non-insertional AT determined by an experienced physiotherapist based on defined clinical findings, VISA-A and NRS (Numerical Rating Scale) scores, physical examination, and ultrasound assessment, as well as having pain for at least 3 months<sup>31</sup>. VISA-A scores less than 90 will be considered as a reference to identify individuals with AT<sup>32</sup>. Ultrasound evaluation of the tendon’s mid-portion will include identifying local thickening of the tendon and/or irregular tendon structure with hypoechoic areas and/or irregular fiber orientation<sup>33</sup>.

The exclusion criteria for both groups will include: (1) systemic or inflammatory conditions including rheumatic, neuromuscular disorders, and malignancy, (2) current or history of chronic respiratory, neurological, or cardiovascular diseases, (3) history of lower limb surgery. Specific exclusion criteria for the participants with AT are participation in any other treatment or rehabilitation program for AT, corticosteroid injections in the previous 12 months, and insertional AT. Additionally, if any participants present non-insertional and insertional AT concurrently in the same limb, they will be excluded. Specific exclusion criteria for the control group are pain/injury in the lower limbs within the previous 6 months, history of AT or lower limb surgery.

**Study Design**


This two-arm, parallel-group, randomised controlled trial will be conducted from October 2021 to December 2022 at a laboratory within the Centre of Precision Rehabilitation for Spinal Pain, University of Birmingham, UK. The Science, Technology, Engineering and Mathematics Ethical Review Committee, University of Birmingham, United Kingdom, approved the study (ERN\_20-0604). All participants will provide written informed consent prior to participation. The study will be conducted according to the

Declaration of Helsinki. Reporting will follow the CONSORT statement and the CONSORT flow diagram will be used to describe the flow of participants throughout the trial.

Participants with non-insertional AT will visit the laboratory over six consecutive weeks for the experimental sessions (at weeks 1, 3, and 6) and training sessions (2-3 sessions per week) (**Fig.1**). We will randomly allocate these participants into two groups: eccentric (ECC) or concentric (CON) training. Healthy participants will visit the laboratory once to allow baseline comparison with ECC and CON groups. Additionally, we will randomise the assessed leg in the healthy control group, and the most symptomatic leg in the ECC and CON groups will be evaluated. Finally, foot preference in specific daily activities (foot dominance) will be determined using a behavioral foot-preference inventory<sup>34</sup>. Each experimental session will last 2.5 hours, and each training session will last 40 minutes.

### Sample size

According to power calculations (G\*Power software)<sup>35</sup>, a total of 26 individuals with non-insertional AT (ECC group=13, and CON group= 13) and 13 healthy controls will be required for this study. This sample size considers a power=0.80, alpha=0.01, 25% loss of participants and an effect size (d) of 1.7 calculated from the study of Yu et al.<sup>29</sup>, where the authors compared reductions in pain after an 8-week concentric and eccentric training protocol in individuals with AT.



**Figure 1.** Overview of the study design.

**Experimental sessions**


These sessions will involve the completion of questionnaires, ultrasound imaging of the gastrocnemius-soleus muscles and the Achilles tendon, surface electromyography and torque recordings.

Anthropometric data (age, gender, weight, height, leg dominance, and body mass index) will be obtained, and the participants will then be asked to complete a battery of questionnaires. This includes the International Physical Activity Questionnaire short form (IPAQ-SF),<sup>36</sup> VISA-A,<sup>37</sup> Foot and Ankle Ability Measure (FAAM),<sup>38</sup> Pain Catastrophizing Scale (PCS),<sup>39</sup> and Tampa Scale for Kinesiophobia (TSK)<sup>40</sup>. Additionally, participants will be asked to report their current level of pain using the NRS. After that, participants will lie prone on the chair of a Biodex System 3 dynamometer (Biodex Medical Systems), and ultrasonography (LOGIQ S8 GE Healthcare, Milwaukee, USA) will be used to measure the length, thickness, and cross-sectional area of the Achilles tendon, fascicle length, thickness, and pennation angle of the medial gastrocnemius (MG), and thickness of the lateral gastrocnemius (LG), and soleus (SO) muscles during rest. Then, we will prepare the skin and place the electrodes on the MG, LG, and SO muscles, and using high-density surface electromyography (HD-sEMG); we will ensure minimal electrical activity of these

muscles during rest conditions for the measurements of the Achilles tendon stiffness (passive elastography).

Following the ultrasound assessment the maximal voluntary contraction (MVC) will be recorded during three isometric plantarflexion contractions of 5 seconds each<sup>18</sup>. Between each MVC, the volunteers will have two minutes of rest<sup>18</sup> and all MVCs will be performed at 0° of plantarflexion. The highest MVC value will be used as the reference maximal torque. We will use this MVC value as a reference for the isometric and dynamic plantarflexion contractions during the experimental and training sessions, to avoid multiple MVC measurements that may be produce pain and discomfort in individuals with non-insertional AT. Afterwards, we will measure the stiffness of the tendon during two isometric plantarflexion contractions at 10% MVC (1 s ramp-up, 12 s hold, 1 s ramp-down, and 30 s rest) (active elastography). Subsequently, using HD-sEMG, we will record motor unit activity of the MG, LG, and SO muscles during two isometric plantarflexion contractions at 10%, 40%, and 70% MVC (10% MVC/s ramp-up, 10 s hold, 10% MVC/s ramp-down, and 30 s rest), one concentric-eccentric plantarflexion contraction at 25% and 50% MVC, and one eccentric-concentric plantarflexion contraction at 25% and 50% MVC (the order of the different types of contractions will be randomly selected). Volunteers will have five minutes of rest at the end of each type of contraction (isometric, concentric-eccentric, and eccentric-concentric). After 15 minutes of rest, HD-sEMG will be recorded from the MG, LG, and SO muscles during six explosive (fast force development) isometric plantarflexion contractions at 75% MVC (1 s ramp-up, 3 s hold, 1 s ramp-down, and 10 s rest)<sup>41</sup>. Finally, both the rate of perceived exertion and the level of pain will be monitored regularly throughout the session (**Fig. 2**).

During all contractions, visual feedback of the target torque output will be provided via computer monitor positioned 1 m from the participant. Prior to the contractions, participants will be instructed to match the force output as closely as possible to the target force for the full duration of the contraction. For the dynamic contractions, the range of motion will be set at the total of 30° (neutral position 0° to 30° of plantarflexion) and the angular speed will be set at 3°/s.



## Training sessions

Preceding the contractions, participants will be instructed to match the torque output as closely as possible to the target torque for the full duration of the contraction.

## Outcome measures

### Primary outcomes measure

The primary outcomes for this study will be level of pain and function assessed using the NRS and VISA-A questionnaire, Achilles tendon length, thickness, cross-sectional area, and stiffness assessed using ultrasonography, and motor unit discharge rate, recruitment and de-recruitment thresholds, and discharge rate variability of the GM, GL and SO muscles assessed using HD-sEMG and decomposition techniques.

### Secondary outcomes measure

Secondary outcomes will include the level of physical activity, physical function, pain catastrophizing and fear of movement assessed using the IPAQ-SF, FAAM, PCS, and TSK questionnaires, respectively, GM, GL, and SO muscles thickness, and GM muscle fascicle length and pennation angle evaluated using ultrasonography, and participants' characteristics as age, gender, weight, height, BMI, and foot dominance.

### Questionnaires

In each experimental session, participants will be asked to complete the IPAQ-SF, VISA-A, FAAM, PCS, and TSK questionnaires to measure physical activity level, symptoms in individuals with AT, physical function, pain catastrophizing and fear of movement, respectively. The IPAQ has become the most widely used physical activity questionnaire,<sup>42</sup> and it has acceptable measurement properties for monitoring population levels of physical activity among 18 to 65 years old adults in diverse settings<sup>36</sup>. The VISA-A was developed with the aim of evaluating the symptoms of AT and their impact on physical activity. This questionnaire is valid, reliable, easy to use, and ideal for comparing patients' progress in clinical settings<sup>37</sup>. The FAAM was developed to meet the need for a self-reported evaluative instrument that comprehensively assesses the physical function of individuals with musculoskeletal disorders of the leg, foot, and ankle. The FAAM is a reliable, valid, and responsive measure of self-reported physical function<sup>38</sup>. Additionally, we will use the PCS to understand the psychological processes that lead to heightened physical and emotional distress in response to aversive stimulation. This questionnaire is a reliable and valid measure of catastrophizing<sup>39</sup>. Finally, we will apply the TSK to



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

measure the fear of movement/(re)injury. This questionnaire has been validated in patients with chronic back pain <sup>43-45</sup>, acute back pain <sup>46 47</sup>, osteoarthritis <sup>48</sup>, and fibromyalgia <sup>44 45</sup>.

**Measurement set-up**

For the measurements of the Achilles tendon, MG, LG, and SO muscles, participants will lie prone on the dynamometer, with their knees extended, and their tested foot tightly strapped on the footplate. The pelvis will be stabilized with another strap to minimise compensatory movements. The ankle will be positioned in 0° of plantarflexion and the axis of the dynamometer will be aligned with the inferior tip of the lateral malleolus. The setting and position of the set up (i.e., chair and isokinetic device) will be saved, so the participants’ position will be similar in each experimental session.

**Ultrasound measurements**

All ultrasound images will be obtained using an ultrasound imaging device equipped with shear wave elastography (SWE) (LOGIQ S8 GE Healthcare, Milwaukee, USA). For the measurements of the length, thickness, and cross-sectional area of the Achilles tendon, and the measurements of the fascicle length, thickness, and pennation angle of the calf muscles, B-mode will be used with a 16-linear array probe (50 mm, 4-15 MHz). Subsequently, for the measurements of the Achilles tendon’s stiffness during rest conditions and isometric plantarflexion contraction, the elastography mode will be used with a 9-linear array probe (44 mm, 2-8 MHz).

An adaptation of the protocol developed by Arya and Kulig<sup>49</sup> will be used to measure the structural properties of the Achilles tendon. Briefly, to obtain tendon length, the ultrasound transducer will be placed longitudinally over the posterior aspect of the heel, and the distal part of the Achilles tendon will be imaged, and the corresponding point will be marked on the skin with a marker. Then, the ultrasound probe will be moved proximally to locate the musculotendinous junction of the MG, and this point will be marked on the skin. The distance between these two points will be measured with a tape and this distance will represent the resting length of the Achilles tendon. Marks at 2 cm,


4 cm, and 6 cm above the Achilles tendon insertion will then be made on the skin. Later, these marks will be used to place the probe in the transversal plane and determine the cross-sectional area of the Achilles tendon at 2 cm, 4 cm, and 6 cm of its insertion. Additionally, we will use these marks to place the probe in the sagittal plane and determine the thickness of the Achilles tendon at 2 cm, 4 cm, and 6 cm of its insertion.

For muscle ultrasound images, the mid-line of the leg will be marked in the direction of the Achilles tendon. Additionally, a mark will be made on the leg 10 cm above the musculotendinous junction of the MG muscle and 4 cm medial to the mid-line. In this position, we will place the middle point of the HD-sEMG electrode grid and mark the contour of the grid on the skin. We will use these marks to place the probe in the sagittal plane and obtain the images of the MG muscle. Similarly, the leg will be marked 10 cm above the musculotendinous junction of the MG muscle and 4 cm lateral to the mid-line. Then, we will repeat the procedure mentioned above, but now for the LG muscle. Next, the leg will be marked 5 cm below the musculotendinous junction of the MG and 4 cm lateral to the mid-line. Again, we will repeat the procedure mentioned above, but now for the SO muscle (**Fig. 3**). The middle column of the HD-sEMG electrode grid will be used as a reference (see HD-sEMG and torque section below) to place the probe in the same position during all the experimental sessions and the images will be acquired with the probe oriented in the sagittal plane, and perpendicular to the skin, according to the recommendations of Bolsterlee et al<sup>50</sup>. The software Image J (<https://imagej.nih.gov/ij/>) will be used to analyze these images and determine the fascicle length, thickness, and pennation angle.

For tendon stiffness measurements, the probe will be placed in the sagittal plane with the middle part of the probe located at 4 cm above the Achilles tendon insertion. A probe holder will be used to avoid movements that may interfere with the measurements. The passive elastography images will be acquired for 12 seconds (twice) and the active elastography images will be acquired during two isometric plantarflexion contractions at 10% MVC (1 s ramp-up, 12 s hold, 1 s ramp down, 1 min rest). Using elastography ultrasound tools, a shear elastography color map (height x width, 2.5 cm x 1 cm) will be chosen to allow a complete visualization of the Achilles tendon width, and a region of interest (ROI) of 3 mm diameter<sup>51</sup> will be selected in the middle of each image to



determine the shear wave velocity (m/s) and Young’s modulus (kPa). Finally, we will average the mean shear wave velocity and Young’s modulus over the ROIs of consecutives images<sup>52</sup>.



**Figure 3.** A) Anatomical landmarks used for ultrasonography and B) position of the electrodes in the MG, LG and SO muscles.

**HD-sEMG and torque recording**

Prior to electrode placement, the skin will be shaved (if necessary), gently abraded (Nuprep, Skin Prep Gel, Weaver and Company, Aurora, Colorado) to reduce skin impedance, and cleaned with water. Three two-dimensional (2D) adhesive grids (SPES Medica, Salerno, Italy) of 13 x 5 equally spaced electrodes (each of 1 mm diameter, with an inter-electrode distance of 8 mm) will be used to record the HD-EMG signals. Conductive paste (AC-CREAM, SPES Medica, Genova, Italy) will be placed into the

cavities of the grid, and the HD-sEMG electrodes will be placed in the exact position described for the ultrasound measurements of the triceps surae (one electrode grid for each muscle).

All signals will be converted from analog-to-digital by a 16-bit converter (Quattrocento- OTBioelettronica, Torino, Italy). The sampling frequency will be 2048 Hz and the amplifier gain will be set to 150. HD-sEMG signals will be digitally filtered with a bandwidth set up to 10 Hz for high pass cut frequency and to 500 Hz for low pass cut frequency<sup>53</sup>. HD-sEMG will be acquired in monopolar mode with reference electrodes (WhiteSensor WS, Ambu A/S, Ballerup, Denmark) positioned in the head of the fibula and with a strip in the thigh of the evaluated leg. All the electrode grids and reference electrodes will be connected to the same bioelectrical amplifier (Quattrocento-OT-Bioelettronica, Torino, Italy).

The torque exerted by the volunteers will be assessed with the isokinetic dynamometer, which will be synchronized with the HD-sEMG signals. Synchronization will be obtained by recording torque signals generated by the isokinetic dynamometer through the auxiliary input of the EMG amplifier<sup>53</sup>.

## Signal Analysis

The torque signal will be low pass filtered at 15 Hz and then used to quantify torque steadiness (coefficient of variation of torque,  $SD \text{ torque} / \text{mean torque} * 100$ ) from the stable part of the contractions<sup>54</sup>.

The HD-sEMG signals will be decomposed into motor unit spike trains with an algorithm based on blind source separation, which provides automatic identification of motor unit activity<sup>54</sup>, and the accuracy of the decomposition will be tested with the silhouette measure, which will be set to  $\geq 0.90$ <sup>55</sup>. The signals will be decomposed during the entire duration of the contractions, and the discharge times of the motor units will be transformed in binary spike trains<sup>56</sup>. The mean discharge rate and the discharge rate variability (CoV of the interspike interval [ $CoV_{isi}$ ]) will be determined during the stable plateau of torque signal. Additionally, motor unit recruitment and de-recruitment thresholds will be defined as the ankle plantarflexion torques (%MVC) at the times when the motor units began and stopped discharging action potentials, respectively<sup>54</sup>.

Discharges rates at recruitment and de-recruitment will be determined using the first and last six discharges of the motor units.<sup>54</sup> Erroneous discharges will be visually inspected and edited using a custom algorithm<sup>54 57</sup>. Motor unit activity will be monitored longitudinally with a recent method proposed by Martinez-Valdes et al.<sup>58</sup> which allows tracking the same motor units across different experimental sessions.

**Adverse event management**

Participants will be informed that they may experience some pain during or after the experimental and training sessions. In order to minimise this, appropriate rest time will be provided throughout the experimental and training sessions, and extra rest periods will be given to the participants at any time if required. If a participant experiences moderate pain (>6 NRS) during the contractions, they will be given additional time to rest. If the pain intensity is maintained or worsens then, we will terminate and reschedule the session.

**Randomisation and blinding**

Before the first experimental session, individuals with non-insertional AT will be randomised by an independent researcher in a 1:1 allocation ratio to either ECC or CON groups using computer-generated simple scheme randomisation (<https://sigdaan.com/randomization/app/randomization-app>).

In order to achieve double-blinding, one researcher will perform the experimental sessions and another the training sessions. The first researcher will use the blindr (<https://github.com/U8NWXD/blindr>) software to encode the results from different subjects and will be blinded to the allocation of the participants. The second researcher will be blind to the participant's results, but not to their allocation. An independent researcher blind to the training sessions will unmask the results after the data analysis is performed.

## Statistical analysis

IBM SPSS Statistics for Windows, Version 25.0 (Armonk, New York, USA) computer software will be used for statistical analysis of the data. The analysis will be performed on an intention-to-treat basis. Descriptive statistics will be used to interpret the data which will be presented as mean  $\pm$  SD. The Shapiro-Wilk Test will be used to assess data normality. The level of significance for all statistical procedures will be set at  $\alpha=0.05$  and 95% confidence intervals (CI) will be reported. Independent t-test will be used to determine the differences between individuals with non-insertional Achilles tendinopathy and healthy controls at baseline. If the data are normally distributed, repeated measures analysis of variance (ANOVA) will be used. Factors of group (ECC and CON) and time (at week 1, 3, and 6) will be used to analyze each variable. Bonferroni post hoc analysis will be used if ANOVA is significant. The partial eta-squared ( $\eta_p^2$ ) for ANOVA will be used to examine the effect size of changes after the training intervention. An  $\eta_p^2$  less than 0.06 will be classified as “small”, 0.07-0.14 as “moderate”, and greater than 0.14 as “large”<sup>59</sup>. If data is not normally distributed, appropriate nonparametric tests will be used.

## Discussion

To our knowledge this is the first study aiming to determine the relationship between pain, function, mechanical and structural properties of the Achilles tendon, and motor unit firing properties of the gastrocnemius-soleus muscles in individuals with non-insertional AT. Additionally, this study will be the first to establish changes in these variables after applying a training protocol based in controlled eccentric or concentric contractions.

Regarding the different questionnaires applied, we hypothesize that individuals with non-insertional AT will obtain lower scores in the IPAQ-SF, VISA-A, and FAAM, and higher scores in the PCS and TSK questionnaires compared with healthy controls. These differences could be explained by the symptoms duration of the non-insertional AT group since it is assumed that individuals with symptoms for more than 3 months are in a chronic stage<sup>31</sup>. Thus, it is likely that we will observe high levels of pain and reduced functionality

in this population. Additionally, we expect that individuals in the ECC group will obtain higher scores in the VISA-A and FAAM, and lower scores in the PCS and TSK questionnaires compare to the CON group after participating in a program of controlled eccentric or concentric contractions, respectively. The characteristics of the training protocols could explain these differences since eccentric contractions generate higher tendon load, produce greater force variability<sup>60</sup> and are more demanding for the neuromuscular system. Furthermore, we do not expect changes in the IPAQ-SF questionnaire between groups, because this questionnaire only informs about general physical function.

A strength of this study is that we will perform a detailed assessment of the mechanical and structural properties of the Achilles tendon and the calf muscles. One study showed that tendinopathy alters both the mechanical and material properties of the human Achilles tendon<sup>49</sup>. Morphological comparisons of tendinopathic and healthy tendons demonstrated a larger cross-sectional area for the degenerated Achilles tendon. Typically, a larger tendon is considered mechanically stronger due to its ability to dissipate stresses across the tendon and yield lower strain energy. Nonetheless, in the study of Arya and Kulig<sup>49</sup>, they demonstrated that despite having a larger cross-sectional area, the degenerated tendon had lower stiffness and Young's modulus compared with healthy tendons. Additionally, our study includes the use of SWE. This procedure has been used to measure tissue elasticity in tendons and might add to a better understanding of the effects of eccentric exercises in tendons<sup>61</sup>. Furthermore, SWE is able to measure the Young modulus (slope of the stress-strain curve in the linear region<sup>62</sup>) of Achilles tendon with high reliability<sup>63</sup>. Previous studies suggest that SWE might be a useful tool for diagnosing and monitoring AT<sup>64</sup>. For instance, one study demonstrated that symptomatic Achilles tendons had lower Young modulus compared with healthy tendons and that stiffness increases in correlation with VISA-A scores after 6 months<sup>65</sup>.

In line with what has been observed in previous studies<sup>49</sup>, we hypothesize that individuals with non-insertional AT will present a larger cross-sectional area, thickness, and less stiffness (passive and active) of the Achilles tendon compared with healthy

controls. Furthermore, we believe that individuals in the ECC and CON groups will show similar increases in strength and calf-muscle thickness, since similar improvements in strength and muscle thickness have been observed after applying a 10-week concentric or eccentric resistance training protocol<sup>66</sup>, in the quadriceps muscles. Additionally, we hypothesize individuals in the ECC group will have an increase in the medial gastrocnemius fascicle length, passive, and active Achilles tendon stiffness. In contrast, we expect that individuals in the CON group will have a greater increase in the medial gastrocnemius pennation angle and lower increase in the passive and active Achilles tendon stiffness. These expectations are based on the observations of Franchi et al.<sup>66</sup>, where they compared the effects in quadriceps muscle architecture of eccentric and concentric exercises, showing a higher increase in the fascicle length in the eccentric group and a more significant increase in the pennation angle in the concentric group.

The last variables to be evaluated in our study will be those related to the electromyographic activity and motor unit firing properties (i.e., discharge rate, recruitment and discharge rate variability) of the gastrocnemius-soleus muscles. Previous studies have estimated the activation of the triceps surae muscles in people with AT during walking,<sup>67</sup> running,<sup>68</sup> isometric plantarflexion tasks,<sup>69</sup> and dynamic plantarflexion tasks<sup>70 71</sup>. Currently, there is no agreement in the literature in terms of plantarflexion torque measured during maximal contractions in individuals with AT, some authors did not find any difference between groups,<sup>49 72</sup> while others found statistically significant differences<sup>69 73</sup>. Interestingly, an investigation has observed a significant increase in LG activation during isometric plantarflexion tasks in people with AT following a 12-week training program<sup>69</sup>. Despite all of these efforts, currently there are no studies that have evaluated motor unit firing properties of calf muscles in individuals with AT.

We believe that individuals with non-insertional AT will show differences in the motor unit firing properties of the gastrocnemius-soleus muscles compared with healthy controls. Specifically, we expect that the discharge rate will be higher in the AT group compared to the control group. This increase in discharge rate could be potentially explained as part of a neuromuscular strategy aiming to compensate for the decrease in



Achilles tendon stiffness present in individuals with AT. In addition, we hypothesize that individuals in the ECC and CON groups will show differences in the motor unit firing properties of the gastrocnemius-soleus muscles following a period of eccentric or concentric exercises. Particularly, we expect that individuals in the ECC group will show a decrease in motor unit discharge rate variability compared to the CON group. We believe that the differences between the training protocols could explain a decrease in the discharge rate variability because greater force variability has been observed during eccentric contractions<sup>60</sup>. Indeed, eccentric contractions are more challenging to control and therefore might provide a more potent stimulus to reduce discharge rate variability and improve torque steadiness. Furthermore, we hypothesize that individuals in the CON group will have an increase in discharge rate to compensate for the lower increase in Achilles tendon stiffness.

The study of motor units is an area in continuous development, which in recent years has allowed a more profound understanding of the neural mechanisms involved in muscle contractions. However, much of the research in this area has focused on the normal neurophysiology of muscle rather than its relationship with alterations of the musculoskeletal system.

This research will therefore provide new insights regarding the neuromechanical effects of ECC and CON exercises in the management of individuals with non-insertional AT. A more precise understanding of the mechanisms involved in this pathology is essential to improve the rehabilitation programs commonly used in the management of this condition.

Additionally, we expect this study will enable further investigation into the effects of different treatment strategies on the mechanical properties of the Achilles tendon and the motor unit firing properties of the gastrocnemius-soleus muscles.

## Patient and Public Involvement

The research question in this study forms part of a larger discussion about exercise and pain relief within our patient and public involvement meetings. Patients will not be involved in the analysis and data collection but will contribute to data interpretation and production of a lay summary of findings.

## Ethics and Dissemination

The research protocol has been approved by the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham (ERN\_20-0604).

Researchers will inform all participants of the characteristics of the research and will obtain written consent. Participants will be informed that they are free to withdraw from the study at any time without needing to provide a reason. In any unlikely adverse events, this will be immediately reported by the principal investigator to the ethics committee.

The results of this study will be submitted for publication in a peer review journal and presented at conferences.

## Author Contributions

ICH and EMV are responsible for the conception, design and development of the protocol. EMV is the lead supervisor of ICH and DF is the co-supervisor. EMV and DF have provided guidance on methodological decisions and critical revision. All authors have read and subsequently approved the final manuscript.

## Funding Statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

## Competing Interests Statement

The authors have no competing interests to report.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Data Sharing Statement**

No additional data are available.

For peer review only

## References

1. Sobhani S, Dekker R, Postema K, et al. Epidemiology of ankle and foot overuse injuries in sports: A systematic review. *Scand J Med Sci Sports* 2013;23(6):669-86. doi: 10.1111/j.1600-0838.2012.01509.x [published Online First: 2012/08/01]
2. Magnussen RA, Dunn WR, Thomson AB. Nonoperative treatment of midportion Achilles tendinopathy: a systematic review. *Clin J Sport Med* 2009;19(1):54-64. doi: 10.1097/JSM.0b013e31818ef090 [published Online First: 2009/01/07]
3. Janssen I, van der Worp H, Hensing S, et al. Investigating Achilles and patellar tendinopathy prevalence in elite athletics. *Research in Sports Medicine* 2018;26(1):1-12. doi: 10.1080/15438627.2017.1393748
4. Kvist M. Achilles tendon injuries in athletes. *Ann Chir Gynaecol* 1991;80(2):188-201. [published Online First: 1991/01/01]
5. Silbernagel KG, Thomee R, Eriksson BI, et al. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med* 2007;35(6):897-906. doi: 10.1177/0363546506298279 [published Online First: 2007/02/20]
6. Longo UG, Ronga M, Maffulli N. Achilles tendinopathy. *Sports Med Arthrosc Rev* 2009;17(2):112-26. doi: 10.1097/JSA.0b013e3181a3d625 [published Online First: 2009/05/15]
7. Scott A, Backman LJ, Speed C. Tendinopathy: Update on Pathophysiology. *J Orthop Sports Phys Ther* 2015;45(11):833-41. doi: 10.2519/jospt.2015.5884 [published Online First: 2015/09/22]
8. Martin RL, Chimenti R, Cuddeford T, et al. Achilles Pain, Stiffness, and Muscle Power Deficits: Midportion Achilles Tendinopathy Revision 2018. *J Orthop Sports Phys Ther* 2018;48(5):A1-a38. doi: 10.2519/jospt.2018.0302 [published Online First: 2018/05/02]
9. Magnan B, Bondi M, Pierantoni S, et al. The pathogenesis of Achilles tendinopathy: a systematic review. *Foot Ankle Surg* 2014;20(3):154-9. doi: 10.1016/j.fas.2014.02.010 [published Online First: 2014/08/12]
10. van der Vlist AC, Breda SJ, Oei EHG, et al. Clinical risk factors for Achilles tendinopathy: a systematic review. *Br J Sports Med* 2019;53(21):1352-61. doi: 10.1136/bjsports-2018-099991 [published Online First: 2019/02/06]
11. Henriksen M, Alkjaer T, Lund H, et al. Experimental quadriceps muscle pain impairs knee joint control during walking. *J Appl Physiol (1985)* 2007;103(1):132-9. doi: 10.1152/jappphysiol.01105.2006 [published Online First: 2007/04/07]
12. Tucker K, Butler J, Graven-Nielsen T, et al. Motor unit recruitment strategies are altered during deep-tissue pain. *J Neurosci* 2009;29(35):10820-6. doi: 10.1523/jneurosci.5211-08.2009 [published Online First: 2009/09/04]
13. Haglund-Akerlind Y, Eriksson E. Range of motion, muscle torque and training habits in runners with and without Achilles tendon problems. *Knee Surg Sports Traumatol Arthrosc* 1993;1(3-4):195-9. doi: 10.1007/bf01560205 [published Online First: 1993/01/01]
14. Mahieu NN, Witvrouw E, Stevens V, et al. Intrinsic risk factors for the development of achilles tendon overuse injury: a prospective study. *Am J Sports Med* 2006;34(2):226-35. doi: 10.1177/0363546505279918 [published Online First: 2005/11/02]
15. Henriksen M, Aaboe J, Graven-Nielsen T, et al. Motor responses to experimental Achilles tendon pain. *Br J Sports Med* 2011;45(5):393-8. doi: 10.1136/bjsm.2010.072561 [published Online First: 2010/06/15]

16. Wyndow N, Cowan SM, Wrigley TV, et al. Neuromotor control of the lower limb in Achilles tendinopathy: implications for foot orthotic therapy. *Sports Med* 2010;40(9):715-27. doi: 10.2165/11535920-000000000-00000 [published Online First: 2010/08/24]

17. Tracy BL, Enoka RM. Older adults are less steady during submaximal isometric contractions with the knee extensor muscles. *J Appl Physiol (1985)* 2002;92(3):1004-12. doi: 10.1152/japplphysiol.00954.2001 [published Online First: 2002/02/14]

18. Martinez-Valdes E, Negro F, Falla D, et al. Surface electromyographic amplitude does not identify differences in neural drive to synergistic muscles. *J Appl Physiol (1985)* 2018;124(4):1071-79. doi: 10.1152/japplphysiol.01115.2017 [published Online First: 2018/02/09]

19. Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG. *J Appl Physiol (1985)* 2004;96(4):1486-95. doi: 10.1152/japplphysiol.01070.2003 [published Online First: 2004/03/16]

20. Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG: an update. *J Appl Physiol (1985)* 2014;117(11):1215-30. doi: 10.1152/japplphysiol.00162.2014 [published Online First: 2014/10/04]

21. Farina D, Negro F, Muceli S, et al. Principles of Motor Unit Physiology Evolve With Advances in Technology. *Physiology* 2016;31(2):83-94. doi: 10.1152/physiol.00040.2015

22. Maffulli N, Longo UG, Kadakia A, et al. Achilles tendinopathy. *Foot Ankle Surg* 2019 doi: 10.1016/j.fas.2019.03.009 [published Online First: 2019/04/30]

23. Alfredson H, Pietila T, Jonsson P, et al. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med* 1998;26(3):360-6. doi: 10.1177/03635465980260030301 [published Online First: 1998/06/09]

24. Maffulli N, Walley G, Sayana MK, et al. Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil* 2008;30(20-22):1677-84. doi: 10.1080/09638280701786427 [published Online First: 2008/07/09]

25. Sayana MK, Maffulli N. Eccentric calf muscle training in non-athletic patients with Achilles tendinopathy. *J Sci Med Sport* 2007;10(1):52-8. doi: 10.1016/j.jsams.2006.05.008 [published Online First: 2006/07/11]

26. Mayer F, Hirschmüller A, Müller S, et al. Effects of short-term treatment strategies over 4 weeks in Achilles tendinopathy. *Br J Sports Med* 2007;41(7):e6. doi: 10.1136/bjsm.2006.031732 [published Online First: 2007/01/31]

27. Jarin I, Backer HC, Vosseller JT. Noninsertional Achilles Tendinopathy: Topical Review. *Foot Ankle Int* 2020;1071100720914605. doi: 10.1177/1071100720914605 [published Online First: 2020/05/01]

28. Mafi N, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 2001;9(1):42-7. doi: 10.1007/s001670000148 [published Online First: 2001/03/28]

29. Yu J, Park D, Lee G. Effect of eccentric strengthening on pain, muscle strength, endurance, and functional fitness factors in male patients with achilles tendinopathy. *American journal of physical medicine & rehabilitation* 2013;92(1):68-76. doi: 10.1097/phm.0b013e31826eda63

30. Lindemann I, Coombes BK, Tucker K, et al. Age-related differences in gastrocnemii muscles and Achilles tendon mechanical properties in vivo. *J Biomech* 2020;112:110067. doi: 10.1016/j.jbiomech.2020.110067 [published Online First: 2020/10/10]

31. Beyer R, Kongsgaard M, Hougs Kjær B, et al. Heavy Slow Resistance Versus Eccentric Training as Treatment for Achilles Tendinopathy: A Randomized Controlled Trial. *Am J Sports Med* 2015;43(7):1704-11. doi: 10.1177/0363546515584760 [published Online First: 2015/05/29]

32. Iversen JV, Bartels EM, Langberg H. The victorian institute of sports assessment - achilles questionnaire (visa-a) - a reliable tool for measuring achilles tendinopathy. *Int J Sports Phys Ther* 2012;7(1):76-84.
33. Rompe JD, Furia J, Maffulli N. Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. *Am J Sports Med* 2009;37(3):463-70. doi: 10.1177/0363546508326983 [published Online First: 2008/12/18]
34. Chapman JP, Chapman LJ, Allen JJ. The measurement of foot preference. *Neuropsychologia* 1987;25(3):579-84. doi: 10.1016/0028-3932(87)90082-0 [published Online First: 1987/01/01]
35. Faul F, Erdfelder E, Buchner A, et al. Statistical power analyses using G\*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009;41(4):1149-60. doi: 10.3758/brm.41.4.1149 [published Online First: 2009/11/10]
36. Craig CL, Marshall AL, Sjoström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35(8):1381-95. doi: 10.1249/01.MSS.0000078924.61453.Fb [published Online First: 2003/08/06]
37. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med* 2001;35(5):335-41. doi: 10.1136/bjsm.35.5.335 [published Online First: 2001/10/02]
38. Martin RL, Irrgang JJ, Burdett RG, et al. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int* 2005;26(11):968-83. doi: 10.1177/107110070502601113 [published Online First: 2005/11/29]
39. Sullivan MJL, Bishop S, Pivik J. The Pain Catastrophizing Scale: Development and Validation. *Psychological Assessment* 1996;7:524-32. doi: 10.1037//1040-3590.7.4.524
40. Miller RP, Kori SH, Todd DD. The Tampa Scale: a Measure of Kinesiophobia. *The Clinical Journal of Pain* 1991;7(1):51.
41. Del Vecchio A, Negro F, Holobar A, et al. You are as fast as your motor neurons: speed of recruitment and maximal discharge of motor neurons determine the maximal rate of force development in humans. *J Physiol* 2019;597(9):2445-56. doi: 10.1113/jp277396 [published Online First: 2019/02/16]
42. van Poppel MN, Chinapaw MJ, Mokkink LB, et al. Physical activity questionnaires for adults: a systematic review of measurement properties. *Sports Med* 2010;40(7):565-600. doi: 10.2165/11531930-000000000-00000 [published Online First: 2010/06/16]
43. Vlaeyen JW, Kole-Snijders AM, Boeren RG, et al. Fear of movement/(re)injury in chronic low back pain and its relation to behavioral performance. *Pain* 1995;62(3):363-72. doi: 10.1016/0304-3959(94)00279-n [published Online First: 1995/09/01]
44. Goubert L, Crombez G, Van Damme S, et al. Confirmatory factor analysis of the Tampa Scale for Kinesiophobia: invariant two-factor model across low back pain patients and fibromyalgia patients. *Clin J Pain* 2004;20(2):103-10. doi: 10.1097/00002508-200403000-00007 [published Online First: 2004/02/11]
45. Roelofs J, Goubert L, Peters ML, et al. The Tampa Scale for Kinesiophobia: further examination of psychometric properties in patients with chronic low back pain and fibromyalgia. *Eur J Pain* 2004;8(5):495-502. doi: 10.1016/j.ejpain.2003.11.016 [published Online First: 2004/08/25]
46. Swinkels-Meewisse EJ, Swinkels RA, Verbeek AL, et al. Psychometric properties of the Tampa Scale for kinesiophobia and the fear-avoidance beliefs questionnaire in acute low back pain. *Man Ther* 2003;8(1):29-36. doi: 10.1054/math.2002.0484 [published Online First: 2003/02/15]
47. Swinkels-Meewisse IE, Roelofs J, Verbeek AL, et al. Fear of movement/(re)injury, disability and participation in acute low back pain. *Pain* 2003;105(1-2):371-9. doi: 10.1016/s0304-3959(03)00255-0 [published Online First: 2003/09/23]

48. Heuts PH, Vlaeyen JW, Roelofs J, et al. Pain-related fear and daily functioning in patients with osteoarthritis. *Pain* 2004;110(1-2):228-35. doi: 10.1016/j.pain.2004.03.035 [published Online First: 2004/07/28]

49. Arya S, Kulig K. Tendinopathy alters mechanical and material properties of the Achilles tendon. *J Appl Physiol (1985)* 2010;108(3):670-5. doi: 10.1152/japplphysiol.00259.2009 [published Online First: 2009/11/07]

50. Bolsterlee B, Gandevia SC, Herbert RD. Effect of Transducer Orientation on Errors in Ultrasound Image-Based Measurements of Human Medial Gastrocnemius Muscle Fascicle Length and Pennation. *PLoS One* 2016;11(6):e0157273. doi: 10.1371/journal.pone.0157273 [published Online First: 2016/06/15]

51. Siu W-l, Chan C-h, Lam C-h, et al. Sonographic evaluation of the effect of long-term exercise on Achilles tendon stiffness using shear wave elastography. *Journal of Science & Medicine in Sport* 2016;19(11):883-87.

52. Coombes BK, Tucker K, Vicenzino B, et al. Achilles and patellar tendinopathy display opposite changes in elastic properties: A shear wave elastography study. *Scand J Med Sci Sports* 2018;28(3):1201-08. doi: 10.1111/sms.12986 [published Online First: 2017/10/04]

53. Arvanitidis M, Falla D, Martinez-Valdes E. Can visual feedback on upper trapezius high-density surface electromyography increase time to task failure of an endurance task? *J Electromyogr Kinesiol* 2019;49:102361. doi: 10.1016/j.jelekin.2019.102361 [published Online First: 2019/10/13]

54. Martinez-Valdes E, Negro F, Farina D, et al. Divergent response of low- versus high-threshold motor units to experimental muscle pain. *J Physiol* 2020;598(11):2093-108. doi: 10.1113/jp279225 [published Online First: 2020/03/19]

55. Negro F, Muceli S, Castronovo AM, et al. Multi-channel intramuscular and surface EMG decomposition by convolutive blind source separation. *J Neural Eng* 2016;13(2):026027. doi: 10.1088/1741-2560/13/2/026027 [published Online First: 2016/03/01]

56. Martinez-Valdes E, Laine CM, Falla D, et al. High-density surface electromyography provides reliable estimates of motor unit behavior. *Clin Neurophysiol* 2016;127(6):2534-41. doi: 10.1016/j.clinph.2015.10.065 [published Online First: 2016/01/19]

57. Boccia G, Martinez-Valdes E, Negro F, et al. Motor unit discharge rate and the estimated synaptic input to the vasti muscles is higher in open compared with closed kinetic chain exercise. *J Appl Physiol (1985)* 2019;127(4):950-58. doi: 10.1152/japplphysiol.00310.2019 [published Online First: 2019/08/02]

58. Martinez-Valdes E, Negro F, Laine CM, et al. Tracking motor units longitudinally across experimental sessions with high-density surface electromyography. *J Physiol* 2017;595(5):1479-96. doi: 10.1113/jp273662 [published Online First: 2016/12/30]

59. Cohen J. Statistical Power Analysis for the Behavioral Sciences. 2nd ed. Hillsdale (NJ) ed: L. Erlbaum Associates; 1988:p.567.

60. CHRISTOU EA, CARLTON LG. Motor output is more variable during eccentric compared with concentric contractions. *Medicine & Science in Sports & Exercise* 2002;34(11):1773-78.

61. Washburn N, Onishi K, Wang J. Ultrasound elastography and ultrasound tissue characterisation for tendon evaluation. *Journal of Orthopaedic Translation* 2018;15 doi: 10.1016/j.jot.2018.06.003


62. Maganaris CN, Narici MV, Maffulli N. Biomechanics of the Achilles tendon. *Disabil Rehabil* 2008;30(20-22):1542-7. doi: 10.1080/09638280701785494 [published Online First: 2008/08/23]

63. Leung W, Chu K, Lai C. Sonographic evaluation of the immediate effects of eccentric heel drop exercise on Achilles tendon and gastrocnemius muscle stiffness using shear wave elastography. *PeerJ* 2017;5 doi: 10.7717/peerj.3592




64. Dirrichs T, Quack V, Gatz M, et al. Shear Wave Elastography (SWE) for the Evaluation of Patients with Tendinopathies. *Acad Radiol* 2016;23(10):1204-13. doi: 10.1016/j.acra.2016.05.012 [published Online First: 2016/06/20]
65. Dirrichs T, Quack V, Gatz M, et al. Shear Wave Elastography (SWE) for Monitoring of Treatment of Tendinopathies: A Double-blinded, Longitudinal Clinical Study. *Acad Radiol* 2018;25(3):265-72. doi: 10.1016/j.acra.2017.09.011 [published Online First: 2017/11/21]
66. Franchi MV, Atherton PJ, Reeves ND, et al. Architectural, functional and molecular responses to concentric and eccentric loading in human skeletal muscle. *Acta Physiol (Oxf)* 2014;210(3):642-54. doi: 10.1111/apha.12225 [published Online First: 2014/01/07]
67. Baur H, Divert C, Hirschl Müller A, et al. Analysis of gait differences in healthy runners and runners with chronic Achilles tendon complaints. *Isokinetics and Exercise Science* 2004;12:111-16. doi: 10.3233/IES-2004-0161
68. Wyndow N, Cowan SM, Wrigley TV, et al. Triceps surae activation is altered in male runners with Achilles tendinopathy. *J Electromyogr Kinesiol* 2013;23(1):166-72. doi: 10.1016/j.jelekin.2012.08.010 [published Online First: 2012/11/13]
69. Masood T, Kalliokoski K, Bojsen-Møller J, et al. Plantarflexor muscle function in healthy and chronic Achilles tendon pain subjects evaluated by the use of EMG and PET imaging. *Clin Biomech (Bristol, Avon)* 2014;29(5):564-70. doi: 10.1016/j.clinbiomech.2014.03.003 [published Online First: 2014/04/10]
70. Reid D, McNair PJ, Johnson S, et al. Electromyographic analysis of an eccentric calf muscle exercise in persons with and without Achilles tendinopathy. *Phys Ther Sport* 2012;13(3):150-5. doi: 10.1016/j.ptsp.2011.08.003 [published Online First: 2012/07/21]
71. Yu J. Comparison of Lower Limb Muscle Activity during Eccentric and Concentric Exercises in Runners with Achilles Tendinopathy. *J Phys Ther Sci* 2014;26(9):1351-3. doi: 10.1589/jpts.26.1351 [published Online First: 2014/10/03]
72. Child S, Bryant AL, Clark RA, et al. Mechanical properties of the achilles tendon aponeurosis are altered in athletes with achilles tendinopathy. *Am J Sports Med* 2010;38(9):1885-93. doi: 10.1177/0363546510366234 [published Online First: 2010/05/29]
73. O'Neill S, Barry S, Watson P. Plantarflexor strength and endurance deficits associated with mid-portion Achilles tendinopathy: The role of soleus. *Phys Ther Sport* 2019;37:69-76. doi: 10.1016/j.ptsp.2019.03.002 [published Online First: 2019/03/19]


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41



11-058683 on 29 July 2022. Downloaded from <http://bmjopen.bmj.com/> on June 13, 2025 at Agence Bibliographique de l'Enseignement Supérieur (ABES).  
t, including for uses related to text and data mining, AI training, and similar technologies.







## CONSORT 2010 checklist of information to include when reporting a randomised trial\*

Section and topic	Items N°	Checklist item	Page N°
<b>Title and abstract</b>			
	1a	Identification as a randomised trial in the title	1
	1b	Structured summary of trial design, methods, results, and conclusions	2
<b>Introduction</b>			
Background and objectives	2a	Scientific background and explanation rationale	4-6
	2b	Specific objectives or hypotheses	6
<b>Methods</b>			
Trial design	3a	Description of trial design (such as parallel, factorial) including allocation ratio	7
	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	N/A
Participants	4a	Eligibility criteria for participants	7
	4b	Settings and locations where the data were collected	7
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were administered	11
Outcomes	6a	Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed	12
	6b	Any changes to trial outcomes after the trial commenced, with reasons	N/A
Sample size	7a	How sample size was determined	8
	7b	When applicable, explanation of any interim analyses and stopping guidelines	N/A
Randomisation:			
Sequence generation	8a	Method used to generate the random allocation sequence	17
	8b	Type of randomisation; details of any restriction (such as blocking and block size)	17
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	17
Implementation	10	Who generated the random allocation sequence, who enrolled the participants, and who assigned participants to interventions	17
Blinding	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how	17
	11b	If relevant, description of the similarity of interventions	N/A
Statistical methods	12a	Statistical methods used to compare groups for primary and secondary outcomes	18
	12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	N/A

<b>Results</b>			
Participant flow (a diagram is strongly recommended)	13a	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome	N/A
	13b	For each group, losses and exclusions after randomisation, together with reasons	N/A
Recruitment	14a	Dates defining the periods of recruitment and follow-up	N/A
	14b	Why the trial ended or was stopped	N/A
Baseline data	15	A table showing baseline demographic and clinical characteristics for each group	N/A
Numbers analysed	16	For each group, number of participants (denominator) included in each analysis and whether analysis was by original assigned groups	N/A
Outcomes and estimation	17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)	N/A
	17b	For binary outcomes, presentation of both absolute and relative effect size is recommended	N/A
Ancillary analysis	18	Results of any other analyses performed, including subgroup analyses and adjusted analyses distinguishing pre-specified from exploratory	N/A
Harms	19	All important harms or unintended effects in each group	N/A
<b>Discussion</b>			
Limitations	20	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant multiplicity of analyses	18-21
Generalisability	21	Generalisability (external validity, applicability) of the trial findings	18-21
Interpretation	22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	18-21
<b>Other information</b>			
Registration	23	Registration number and name of trial registry	2
Protocol	24	Where the full trial protocol can be accessed, if available	N/A
Funding	25	Sources of funding and other support (such as supply of drugs), role of funders	22

\*We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration<sup>13</sup> for important clarifications on all the items. If relevant, we also recommend reading CONSORT extensions for cluster randomised trials,<sup>11</sup> non-inferiority and equivalence trials,<sup>12</sup> non-pharmacological treatments,<sup>32</sup> herbal interventions,<sup>33</sup> and pragmatic trials.<sup>34</sup> Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see [www.consort-statement.org](http://www.consort-statement.org).

From: Schulz K, Altman D, Moher D. CONSORT 2010 Statement: updated guideline for reporting parallel group randomised trials. BMJ 2010; 340:c 332

# BMJ Open

## Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy: Protocol for a randomised controlled trial.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058683.R1
Article Type:	Protocol
Date Submitted by the Author:	03-Mar-2022
Complete List of Authors:	Contreras-Hernandez, Ignacio; University of Birmingham Falla, Deborah; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences Martinez-Valdes, Eduardo; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences
<b>Primary Subject Heading</b>:	Sports and exercise medicine
Secondary Subject Heading:	Rehabilitation medicine
Keywords:	SPORTS MEDICINE, Adult orthopaedics < ORTHOPAEDIC & TRAUMA SURGERY, Foot & ankle < ORTHOPAEDIC & TRAUMA SURGERY, Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY

SCHOLARONE™  
Manuscripts

Title

Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy: Protocol for a randomised controlled trial.

**Authors:** Ignacio Contreras-Hernandez<sup>1</sup>, Deborah Falla<sup>1</sup>, Eduardo Martinez-Valdes<sup>1</sup>

Corresponding author: Eduardo Martinez-Valdes<sup>1</sup>  
Centre of Precision Rehabilitation for Spinal Pain (CPR Spine)  
School of Sport, Exercise and Rehabilitation Sciences  
College of Life and Environmental Sciences  
University of Birmingham  
Edgbaston B15 2TT, United Kingdom  
T: +44 (0) 121 4158187  
M: +44 (0) 7961 548851  
Email: [e.a.martinezvaldes@bham.ac.uk](mailto:e.a.martinezvaldes@bham.ac.uk)

**Author Affiliations**

<sup>1</sup>Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, UK.

Email: Ignacio Contreras-Hernandez<sup>1</sup> [iac921@student.bham.ac.uk](mailto:iac921@student.bham.ac.uk) – Deborah Falla<sup>1</sup> [d.falla@bham.ac.uk](mailto:d.falla@bham.ac.uk) – Eduardo Martinez-Valdes<sup>1</sup> [e.a.martinezvaldes@bham.ac.uk](mailto:e.a.martinezvaldes@bham.ac.uk)

**WORD COUNT MAIN TEXT: 5867 (excluding title page, abstract, references, figures, tables, and acknowledgments)**

## Abstract

**Introduction.** There is limited evidence on the neural strategies employed by the central nervous system to control muscle force in the presence of non-insertional Achilles tendinopathy (NIAT). Additionally, the neuromuscular mechanisms by which exercise may help to resolve tendon pain remain unclear.

**Objective.** This study aims to firstly establish changes in the gastrocnemius-soleus motor unit firing properties after applying a training protocol of 6-weeks based on either controlled eccentric or concentric contractions in individuals with NIAT. Secondly, we want to determine changes in the level of pain and function and mechanical and structural properties of the Achilles tendon after applying the same training protocol. Additionally, we want to compare these variables at baseline between individuals with NIAT and asymptomatic controls.

**Methods and analysis.** A total of 26 individuals with chronic (> 3 months) NIAT and 13 healthy controls will participate in the study. Individuals with NIAT will be randomised to perform eccentric or concentric training for 6 weeks. Motor unit firing properties of the medial gastrocnemius, lateral gastrocnemius and soleus muscles will be assessed using high-density surface electromyography, as well as Achilles tendon length, cross-sectional area, thickness, and stiffness using B-mode ultrasonography and shear wave elastography. Moreover, participants will complete a battery of questionnaires to document their level of pain and function.

**Ethics and dissemination.** Ethical approval (ERN\_20-0604) for the study was obtained from the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham. The results of the study will be published in peer-review journals.

**Protocol Registration.** This protocol has been registered on ISRCTN registry (ISRCTN46462385).

**Keywords:** Achilles tendinopathy, HD-sEMG, motor unit, morphological properties, mechanical features.

**WORD COUNT:** 271

1

2

3 **Article Summary**

4

5 **Strengths and limitations of this study**

6

7

8

- 9
- 10 • This will be the first study to examine whether gastrocnemius-soleus motor unit firing
  - 11 properties change in response to a training protocol in individuals with non-insertional
  - 12 Achilles tendinopathy (NIAT).
  - 13
  - 14 • This will also be the first study to determine gastrocnemius-soles motor unit firing
  - 15 properties in individuals with NIAT compared with asymptomatic controls.
  - 16
  - 17 • The training protocol will be based on pure eccentric or concentric contractions
  - 18 controlled using an isokinetic dynamometer.
  - 19
  - 20 • We will apply a relatively short training protocol, therefore longer training interventions
  - 21 might be required to assess changes in the mechanical and structural properties of
  - 22 the Achilles tendon in the long term.
  - 23
  - 24 • Due to the nature of the interventions, blinding of the participants is not possible.
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60

## Introduction

Achilles tendinopathy (AT) is a painful overuse injury of the Achilles tendon and it is common among athletes, especially those involved in running and jumping sports<sup>1-3</sup>. AT is clinically diagnosed when the patient presents with a combination of localized pain, swelling of the Achilles tendon, and loss of function<sup>4</sup>. The essence of tendinopathy is a failed healing response, with degeneration and proliferation of tenocytes, disruption of collagen fibres, and subsequent increase in non-collagenous matrix<sup>5</sup>. These structural changes in the tendon result in increased cross-sectional area, reduced tendon stiffness, and altered viscoelastic properties in both symptomatic and asymptomatic tendons<sup>6</sup>.

The aetiology of AT remains debated and is likely caused by intrinsic and extrinsic factors<sup>7</sup>. One of the most accepted theories is that pain perception during early support loading may trigger inhibition of neuromuscular activity of the calf muscles detected as a reduction in electromyographic (EMG) amplitude<sup>8 9</sup>. Thus, the decrease in the ability to generate force in patients with AT could also reflect the decline in neuromuscular activity observed<sup>10 11</sup>. Moreover, it has been observed that this motor inhibition also affects synergist and antagonist muscles<sup>12</sup>. Additionally, individuals with tendinopathy tend to use movement patterns that place an excessive or abnormal load on their tendons<sup>6</sup>, and it is believed that these motor adaptations may generate greater torsional stress in the tendon<sup>12 13</sup>. Finally, studies have shown that AT reduces tendon's stiffness<sup>14</sup>, which impairs the mechanisms responsible for transmitting force to the bone. Therefore, it is very likely that these alterations in tendon properties may produce changes in the neural drive received by the calf muscles.

Until now, most studies examining the neuromuscular impairments induced by AT have focused on investigating changes in interference EMG amplitude which is an indirect estimate of neural activity with many factors of influence<sup>15-17</sup>. Clearer information about the neural strategies employed by the central nervous system to control muscle force in the presence of AT can be obtained through motor unit recordings, since motor unit firing properties represent the direct neural output from the spinal cord to muscles<sup>18</sup>. Nevertheless, there are no studies that have measured motor unit firing properties in



individuals with NIAT.

Although eccentric exercise has been widely used for the treatment of NIAT, the mechanisms by which eccentric exercises may help to resolve tendon pain remain unclear<sup>7</sup>. Regarding eccentric exercise alone, two prospective studies have reported a significant reduction in pain intensity and change on the Victorian Institute of Sports Assessment-Achilles questionnaire (VISA-A) in recreational athletes following a 12-week exercise program<sup>19 20</sup>. In contrast, another study in non-athletic individuals found no significant improvement after a similar 12-week exercise program<sup>21</sup>. Concerning eccentric exercises with an adjunctive treatment (e.g., pulsed ultrasound, ice, sensory motor training), a 4-week intervention study resulted in decreased pain and higher plantarflexion peak torque in individuals with NIAT compared to controls<sup>22</sup>. However, studies that include eccentric exercises with an adjunctive treatment showed limited evidence of improvement over eccentric exercises alone<sup>23</sup>.

There are few studies where the effectiveness of eccentric versus concentric exercises has been compared. Mafi et al. showed that patient satisfaction and return to previous activity were significantly superior after participating in a 12-week rehabilitation protocol based on eccentric exercise compared to concentric exercise. Although pain intensity decreased significantly in both groups, the amount of pain reduction was significantly greater for those that performed eccentric exercise<sup>24</sup>. Likewise, Yu et al.<sup>25</sup> demonstrated that eight weeks of eccentric exercise was more effective at reducing pain than concentric in individuals with chronic NIAT. Additionally, they found that eccentric exercise was more effective than concentric exercise at increasing muscle strength and endurance, and improving function<sup>25</sup>. In these investigations, participants performed the rehabilitation protocols with insufficient control over the load, speed, pain tolerance, or the range of motion in which the exercises were performed. Moreover, it is essential to consider that when participants perform an eccentric plantar flexion exercise without adequate equipment, it is difficult to achieve pure eccentric contractions, which could have influenced the results obtained in these studies.

Based on the above, the aims of this study are to (1) establish changes in the

gastrocnemius-soleus motor unit firing properties after applying a training protocol of 6-weeks based on either controlled eccentric or concentric contractions in individuals with NIAT; (2) determine changes in the level of pain and function and mechanical and structural properties of the Achilles tendon after applying the same training protocol; (3) compare these properties at baseline between individuals with NIAT and asymptomatic controls.

## Methods

### Participants

Twenty-six individuals with NIAT and thirteen asymptomatic controls will be recruited from the University of Birmingham staff/student population and the local community via leaflets, e-mail, and social media.

Men or women aged 18 to 55 years old will be recruited. This age range was selected based on previous findings showing lower stiffness and Young's modulus of the Achilles tendon in older than younger population<sup>26</sup>. Inclusion criteria are NIAT determined by an experienced physiotherapist based on defined clinical findings (VISA-A<sup>27</sup> and NRS (Numerical Rating Scale)<sup>28</sup> scores), physical examination, and ultrasound assessment, as well as having pain for at least 3 months<sup>29</sup>. VISA-A scores less than 90 will be considered as a reference to identify individuals with NIAT<sup>30</sup>. Regarding the NRS scores, previous studies have shown high variability in individuals with NIAT<sup>29 31</sup>, thus we will consider individuals with an NRS score  $\geq 2$ . Physical examination will include palpation of the Achilles tendon along its whole length in a proximal to distal direction, and gentle squeezing the tendon between the thumb and the index finger to identify tenderness over the tendon<sup>20</sup>. Ultrasound evaluation of the tendon's mid-portion will include identifying local thickening of the tendon and/or irregular tendon structure with hypoechoic areas and/or irregular fiber orientation<sup>31</sup>.

The exclusion criteria for both groups will include: (1) systemic or inflammatory conditions including rheumatic, neuromuscular disorders, and malignancy, (2) current or history of chronic respiratory, neurological, or cardiovascular diseases, (3) history of lower limb surgery. Specific exclusion criteria for the participants with NIAT are participation in any other treatment or rehabilitation program for AT, corticosteroid injections in the

previous 12 months, and insertional AT. Additionally, if any participants present non-insertional and insertional AT concurrently in the same limb, they will be excluded. Specific exclusion criteria for the control group are pain/injury in the lower limbs within the previous 6 months, history of AT or lower limb surgery.

Study Design

This two-arm, parallel-group, randomised controlled trial will be conducted from October 2021 to December 2022 at a laboratory within the Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), University of Birmingham, UK. The Science, Technology, Engineering and Mathematics Ethical Review Committee, University of Birmingham, United Kingdom, approved the study (ERN\_20-0604). All participants will provide written informed consent prior to participation. The study will be conducted according to the Declaration of Helsinki. This protocol has been designed following the SPIRIT 2013 statement<sup>32</sup>. Reporting will follow the CONSORT statement and the CONSORT flow diagram will be used to describe the flow of participants throughout the trial (ISRCTN registry number: ISRCTN46462385).

Participants with NIAT will visit the laboratory over six consecutive weeks for the experimental sessions (at weeks 1, 3, and 6) and training sessions (2-3 sessions per week) (**Fig.1**). We will randomly allocate these participants into two groups: eccentric (ECC) or concentric (CON) training. Healthy participants will visit the laboratory once to allow baseline comparison with ECC and CON groups. Additionally, we will randomise the assessed leg in the healthy control group, and the most symptomatic leg in the ECC and CON groups will be evaluated. Finally, foot preference in specific daily activities (foot dominance) will be determined using a behavioral foot-preference inventory<sup>33</sup>. Each experimental session will last 2.5 hours, and each training session will last 40 minutes.

Sample size

According to power calculations (G\*Power software)<sup>34</sup>, a total of 26 individuals with NIAT (ECC group=13, and CON group= 13) and 13 healthy controls will be required for this study. This sample size considers a power=0.80, alpha=0.01, 25% loss of participants and an effect size (d) of 1.7 calculated from the study of Yu et al.<sup>25</sup>, where the authors compared reductions in pain after an 8-week concentric and eccentric training protocol in individuals with NIAT.

## Experimental sessions

These sessions will involve the completion of questionnaires, ultrasound imaging of the gastrocnemius-soleus muscles and the Achilles tendon, surface electromyography and torque recordings. All the procedures during the experimental sessions will be done by Ignacio Contreras-Hernandez (IC-H) and Joeri van Helden (JVH). IC-H is a PhD student at the University of Birmingham, Master in Physiology, Physiotherapist, and member of the CPR Spine group. Joeri is a PhD student at the University of Birmingham, Master in Neuroscience, Psychologist, and member of the CPR Spine group.

Anthropometric data (age, gender, weight, height, leg dominance, and body mass index) will be obtained, and the participants will then be asked to complete a battery of questionnaires. This includes the International Physical Activity Questionnaire short form (IPAQ-SF),<sup>35</sup> VISA-A,<sup>27</sup> Foot and Ankle Ability Measure (FAAM),<sup>36</sup> Pain Catastrophizing Scale (PCS),<sup>37</sup> and Tampa Scale for Kinesiophobia (TSK)<sup>38</sup>. Additionally, participants will be asked to report their current level of pain using the NRS score. After that, participants will lie prone on the chair of a Biodex System 3 dynamometer (Biodex Medical Systems), and ultrasonography (LOGIQ S8 GE Healthcare, Milwaukee, USA) will be used to measure the length, thickness, and cross-sectional area of the Achilles tendon, fascicle length, thickness, and pennation angle of the medial gastrocnemius (MG), and thickness of the lateral gastrocnemius (LG), and soleus (SO) muscles during rest. Then, we will prepare the skin and place the electrodes on the MG, LG, and SO muscles, and using high-density surface electromyography (HD-sEMG); we will ensure minimal electrical activity of these muscles during rest conditions for the measurements of the Achilles tendon stiffness (passive elastography).

Following the ultrasound assessment the maximal voluntary contraction (MVC) will be recorded during three isometric plantarflexion contractions of 5 seconds each<sup>15</sup>. Between each MVC, the volunteers will have two minutes of rest<sup>15</sup> and all MVCs will be performed at 0° of plantarflexion. The highest MVC value will be used as the reference maximal torque. We will use this MVC value as a reference for the isometric and dynamic plantarflexion contractions during the experimental and training sessions, to avoid multiple MVC measurements that may be produce pain and discomfort in individuals with NIAT. Afterwards, we will measure the stiffness of the tendon during two isometric plantarflexion contractions at 10% MVC (1 s ramp-up, 12 s hold, 1 s ramp-down, and 30 s rest) (active elastography). Subsequently, using HD-sEMG, we will record motor unit activity of the MG, LG, and SO muscles during two isometric plantarflexion contractions at 10%, 40%, and 70% MVC (10% MVC/s ramp-up, 10 s hold, 10% MVC/s ramp-down, and 30 s rest), one concentric-eccentric plantarflexion contraction at 25% and 50% MVC, and one eccentric-concentric plantarflexion contraction at 25% and 50% MVC (the order of the different types of contractions will be randomly selected). Volunteers will have five minutes of rest at the end of each type of contraction (isometric, concentric-eccentric, and eccentric-concentric). After 15 minutes of rest, HD-sEMG will be recorded from the MG, LG, and SO muscles during six explosive (fast force development) isometric plantarflexion contractions at 75% MVC (1 s ramp-up, 3 s hold, 1 s ramp-down, and 10 s rest)<sup>39</sup>. Finally, both the rate of perceived exertion and the level of pain will be monitored regularly throughout the session, using the Borg ratings of perceived exertion scale<sup>40</sup> and the NRS. (Fig. 2).

During all contractions, visual feedback of the target torque output will be provided via computer monitor positioned 1 m from the participant. Prior to the contractions, participants will be instructed to match the force output as closely as possible to the target force for the full duration of the contraction. For the dynamic contractions, the range of motion will be set at the total of 30° (neutral position 0° to 30° of plantarflexion) and the angular speed will be set at 3°/s.

## Training sessions

The training sessions will be done by Michalis Arvanitidis (MA). MA is a PhD student at the University of Birmingham, Master in Advanced Manipulative Physiotherapy, and Specialist Musculoskeletal Physiotherapist (Member of the Musculoskeletal Association of Chartered Physiotherapist) and member of the CPR Spine group.

All the training sessions will be done in prone position on the Biodex System 3 dynamometer.

The participants in the ECC group will be asked to perform a warm-up consisting of three eccentric plantarflexion contractions at 25% MVC; this will be followed by the eccentric training protocol. This protocol consists of 4 x 15 eccentric plantarflexion contractions at 50% MVC, range of motion of 30° (neutral position 0° to 30° of plantarflexion), time under tension of 10 seconds, angular speed of 3°/s and 3 minutes of rest between each series. Visual feedback of the exerted torque will be provided. Participants in the CON group will perform a warm-up consisting of three concentric plantarflexion contractions at 25% MVC, and then, the concentric training protocol. This protocol consists of 4 x 15 concentric plantarflexion contractions at 50% MVC, range of motion of 30° (neutral position 0° to 30° of plantarflexion), time under tension of 10 seconds, angular speed of 3°/s, and 3 minutes of rest between each series.

Preceding the contractions, participants will be instructed to match the torque output as closely as possible to the target torque for the full duration of the contraction.

## Follow-up

Participants with NIAT will be asked to report their level of pain and function at 3 and 6 months after completing the training protocol.

## Outcome measures

### Primary outcomes measure

The primary outcomes for this study will be GM, GL, and SO muscles motor unit firing properties assessed using HD-sEMG and decomposition techniques. These properties include motor unit discharge rate, recruitment and de-recruitment thresholds, and discharge rate variability.



**Secondary outcomes measure**

Secondary outcomes will include level of pain and function assessed using the NRS and VISA-A questionnaire, Achilles tendon length, thickness, cross-sectional area, and stiffness using B-mode ultrasonography and shear wave elastography. Additionally, secondary outcomes will include GM, GL, and SO muscles thickness, and GM muscle fascicle length and pennation angle evaluated using B-mode ultrasonography, as well as the level of physical activity, physical function, pain catastrophizing and fear of movement assessed using the IPAQ-SF, FAAM, PCS, and TSK questionnaires, respectively.

**Questionnaires**

In each experimental session, participants will be asked to complete the IPAQ-SF, VISA-A, FAAM, PCS, and TSK questionnaires to measure physical activity level, symptoms in individuals with AT, physical function, pain catastrophizing and fear of movement, respectively. The IPAQ has become the most widely used physical activity questionnaire,<sup>41</sup> and it has acceptable measurement properties for monitoring population levels of physical activity among 18 to 65 years old adults in diverse settings<sup>35</sup>. The VISA-A was developed with the aim of evaluating the symptoms of AT and their impact on physical activity. This questionnaire is valid, reliable, easy to use, and ideal for comparing patients' progress in clinical settings<sup>27</sup>. The FAAM was developed to meet the need for a self-reported evaluative instrument that comprehensively assesses the physical function of individuals with musculoskeletal disorders of the leg, foot, and ankle. The FAAM is a reliable, valid, and responsive measure of self-reported physical function<sup>36</sup>. Additionally, we will use the PCS to understand the psychological processes that lead to heightened physical and emotional distress in response to aversive stimulation. This questionnaire is a reliable and valid measure of catastrophizing<sup>37</sup>. Finally, we will apply the TSK to measure the fear of movement/(re)injury. This questionnaire has been validated in patients with chronic back pain <sup>42-44</sup>, acute back pain <sup>45 46</sup>, osteoarthritis <sup>47</sup>, and fibromyalgia <sup>43 44</sup>.

**Measurement set-up**

For the measurements of the Achilles tendon, MG, LG, and SO muscles, participants will lie prone on the dynamometer, with their knees extended, and their tested foot tightly strapped on the footplate. The pelvis will be stabilized with another strap to minimise compensatory movements. The ankle will be positioned in 0° of plantarflexion and the axis of the dynamometer will be aligned with the inferior tip of the lateral malleolus. The setting and position of the set up (i.e., chair and isokinetic device) will be saved, so the participants' position will be similar in each experimental session.

### Ultrasound measurements

All ultrasound images will be obtained using an ultrasound imaging device equipped with shear wave elastography (SWE) (LOGIQ S8 GE Healthcare, Milwaukee, USA). For the measurements of the length, thickness, and cross-sectional area of the Achilles tendon, and the measurements of the fascicle length, thickness, and pennation angle of the calf muscles, B-mode will be used with a 16-linear array probe (50 mm, 4-15 MHz). Subsequently, for the measurements of the Achilles tendon's stiffness during rest conditions and isometric plantarflexion contraction, the elastography mode will be used with a 9-linear array probe (44 mm, 2-8 MHz).

An adaptation of the protocol developed by Arya and Kulig<sup>14</sup> will be used to measure the structural properties of the Achilles tendon. Briefly, to obtain tendon length, the ultrasound transducer will be placed longitudinally over the posterior aspect of the heel, and the distal part of the Achilles tendon will be imaged, and the corresponding point will be marked on the skin with a marker. Then, the ultrasound probe will be moved proximally to locate the musculotendinous junction of the MG, and this point will be marked on the skin. The distance between these two points will be measured with a tape and this distance will represent the resting length of the Achilles tendon. Marks at 2 cm, 4 cm, and 6 cm above the Achilles tendon insertion will then be made on the skin. Later, these marks will be used to place the probe in the transversal plane and determine the cross-sectional area of the Achilles tendon at 2 cm, 4 cm, and 6 cm of its insertion. Additionally, we will use these marks to place the probe in the sagittal plane and determine the thickness of the Achilles tendon at 2 cm, 4 cm, and 6 cm of its insertion.

For muscle ultrasound images, the mid-line of the leg will be marked in the direction of the Achilles tendon. Additionally, a mark will be made on the leg 10 cm above the musculotendinous junction of the MG muscle and 4 cm medial to the mid-line. In this position, we will place the middle point of the HD-sEMG electrode grid and mark the contour of the grid on the skin. We will use these marks to place the probe in the sagittal plane and obtain the images of the MG muscle. Similarly, the leg will be marked 10 cm above the musculotendinous junction of the MG muscle and 4 cm lateral to the mid-line. Then, we will repeat the procedure mentioned above, but now for the LG muscle. Next, the leg will be marked 5 cm below the musculotendinous junction of the MG and 4 cm lateral to the mid-line. Again, we will repeat the procedure mentioned above, but now for the SO muscle (**Fig. 3**). The middle column of the HD-sEMG electrode grid will be used as a reference (see HD-sEMG and torque section below) to place the probe in the same position during all the experimental sessions and the images will be acquired with the probe oriented in the sagittal plane, and perpendicular to the skin, according to the recommendations of Bolsterlee et al<sup>48</sup>. To ensure that we are measuring the exact location of interest, we will use the procedure described above in each experimental session, and we also will mark the middle point of the ultrasound probe. Then, during the acquisition of the ultrasound images, we will align the mark in the ultrasound probe with the marks on the skin at 2 cm, 4 cm, and 6 cm from the Achilles tendon insertion and with the mark of the middle point of the HD-sEMG electrode grid of each muscle. This procedure will allow us to identify the location of interest during the analysis of the ultrasound images since we know that the middle point of the image represents the location of interest. The software Image J (<https://imagej.nih.gov/ij/>) will be used to analyze these images and determine the fascicle length, thickness, and pennation angle.

For tendon stiffness measurements, the probe will be placed in the sagittal plane, with the middle part of the probe located at 4 cm above the Achilles tendon insertion. A probe holder will be used to avoid applying pressure over the tendon and movements that may interfere with the measurements. Then, we will perform a trial SWE to check for possible voids. If voids are detected at this stage, we will remove the probe holder and place the ultrasound probe again. The passive elastography images will be acquired for 12 seconds (twice), and the active elastography images will be acquired during two

isometric plantarflexion contractions at 10% MVC (1 s ramp-up, 12 s hold, 1 s ramp down, 1 min rest). We will check the elastography images following each measurement to determine possible voids that may affect our results. If voids in the middle part of the tendon (at 4 cm of the insertion) are detected in this stage, we will repeat the procedure. A shear elastography colour map (height x width, 2.5 cm x 1 cm) will be chosen using elastography ultrasound tools to allow complete visualization of the Achilles tendon width, and a region of interest (ROI) of 3 mm diameter<sup>49</sup> will be selected in the middle of each image to determine the shear wave velocity (m/s) and Young's modulus (kPa). Finally, we will average the mean shear wave velocity and Young's modulus over the ROIs of consecutive images<sup>50</sup>.

### HD-sEMG and torque recording

Prior to electrode placement, the skin will be shaved (if necessary), gently abraded (Nuprep, Skin Prep Gel, Weaver and Company, Aurora, Colorado) to reduce skin impedance, and cleaned with water. Three two-dimensional (2D) adhesive grids (SPES Medica, Salerno, Italy) of 13 x 5 equally spaced electrodes (each of 1 mm diameter, with an inter-electrode distance of 8 mm) will be used to record the HD-EMG signals. Conductive paste (AC-CREAM, SPES Medica, Genova, Italy) will be placed into the cavities of the grid, and the HD-sEMG electrodes will be placed in the exact position described for the ultrasound measurements of the triceps surae (one electrode grid for each muscle).

All signals will be converted from analog-to-digital by a 16-bit converter (Quattrocento- OTBioelettronica, Torino, Italy). The sampling frequency will be 2048 Hz and the amplifier gain will be set to 150. HD-sEMG signals will be digitally filtered with a bandwidth set up to 10 Hz for high pass cut frequency and to 500 Hz for low pass cut frequency<sup>51</sup>. HD-sEMG will be acquired in monopolar mode with reference electrodes (WhiteSensor WS, Ambu A/S, Ballerup, Denmark) positioned in the head of the fibula and with a strip in the thigh of the evaluated leg. All the electrode grids and reference electrodes will be connected to the same bioelectrical amplifier (Quattrocento-OT-Bioelettronica, Torino, Italy).

The torque exerted by the volunteers will be assessed with the isokinetic dynamometer, which will be synchronized with the HD-sEMG signals. Synchronization will be obtained by recording torque signals generated by the isokinetic dynamometer through the auxiliary input of the EMG amplifier<sup>51</sup>.

**Signal Analysis**

The torque signal will be low pass filtered at 15 Hz and then used to quantify torque steadiness (coefficient of variation of torque, SD torque/mean torque \* 100) from the stable part of the contractions<sup>52</sup>.

The HD-sEMG signals will be decomposed into motor unit spike trains with an algorithm based on blind source separation, which provides automatic identification of motor unit activity <sup>52</sup>, and the accuracy of the decomposition will be tested with the silhouette measure, which will be set to  $\geq 0.90$ <sup>53</sup>. The signals will be decomposed during the entire duration of the contractions, and the discharge times of the motor units will be transformed in binary spike trains <sup>54</sup>. The mean discharge rate and the discharge rate variability (CoV of the interspike interval [CoV<sub>isi</sub>]) will be determined during the stable plateau of torque signal. Additionally, motor unit recruitment and de-recruitment thresholds will be defined as the ankle plantarflexion torques (%MVC) at the times when the motor units began and stopped discharging action potentials, respectively<sup>52</sup>. Discharges rates at recruitment and de-recruitment will be determined using the first and last six discharges of the motor units.<sup>52</sup> Erroneous discharges will be visually inspected and edited using a custom algorithm<sup>52 55</sup>. Motor unit activity will be monitored longitudinally with a recent method proposed by Martinez-Valdes et al.<sup>56</sup> which allows tracking the same motor units across different experimental sessions.

**Adverse event management**

Participants will be informed that they may experience some pain during or after the experimental and training sessions. Monitoring of participants' pain will be done in each experimental and training sessions using the NRS. Appropriate rest time will be provided throughout the experimental and training sessions, and extra rest periods will be given to the participants at any time if required. If a participant experiences moderate

pain (>6 NRS) during the contractions, they will be given additional time to rest. If the pain intensity is maintained or worsens, we will terminate and reschedule the session. The session will be rescheduled in the upcoming three days. If pain intensity is maintained or worsened during these days and rescheduling the session is not possible, the participant will be removed from the study. This will be considered as an adverse effect, and it will be reported to the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham.

## Randomisation and blinding

Individuals with NIAT will be randomised by an independent researcher (Dr. Eduardo Martinez-Valdes (EM-V)) in a 1:1 allocation ratio to either ECC or CON groups (parallel-groups) using computer-generated simple scheme randomisation (<https://sigdaan.com/randomization/app/randomization-app>). Allocation concealment will be ensured, as EM-V will secure the randomisation code using password-protected files. EM-V will give access to MA to the randomisation code once each participant has completed the first experimental session.

In order to achieve double-blinding, IC-H and JVH will perform the experimental sessions and MA will perform the training sessions. IC-H will use the blindr (<https://github.com/U8NWXD/blindr>) software to encode the results from different participants and will be blinded to the training protocol applied to the participants. MA will be blind to the participant's results, but not to the training protocol applied. EM-V will unmask the results after the data analysis is performed. Due to the nature of the interventions, participants' blinding is not possible.

## Statistical analysis

IBM SPSS Statistics for Windows, Version 25.0 (Armonk, New York, USA) computer software will be used for statistical analysis of the data. Intention-to-treat and per protocol analysis will be performed. Descriptive statistics will be used to interpret the data which will be presented as mean  $\pm$  SD. The Shapiro-Wilk Test will be used to assess



data normality. The level of significance for all statistical procedures will be set at  $\alpha=0.05$  and 95% confidence intervals (CI) will be reported. Independent t-test will be used to determine the differences between individuals with NIAT and healthy controls at baseline. If the data are normally distributed, repeated measures analysis of variance (ANOVA) will be used. Factors of group (ECC and CON) and time (at week 1, 3, and 6) will be used to analyze each variable. Bonferroni post hoc analysis will be used if ANOVA is significant. The partial eta-squared ( $\eta_p^2$ ) for ANOVA will be used to examine the effect size of changes after the training intervention. An  $\eta_p^2$  less than 0.06 will be classified as “small”, 0.07-0.14 as “moderate”, and greater than 0.14 as “large”<sup>57</sup>. If data is not normally distributed, appropriate nonparametric tests will be used.

## Discussion

To our knowledge this is the first study aiming to establish changes in motor unit firing properties of the gastrocnemius-soleus muscles after applying a training protocol based on either controlled eccentric or concentric contractions in individuals with NIAT. Additionally, this study will be the first to determine motor unit firing properties of the gastrocnemius-soleus muscles in individuals with NIAT compared with asymptomatic controls.

Regarding the variables related to the electromyographic activity and motor unit firing properties (i.e., discharge rate, recruitment and discharge rate variability) of the gastrocnemius-soleus muscles; previous studies have estimated the activation of the triceps surae muscles in people with AT during walking,<sup>58</sup> running,<sup>59</sup> isometric plantarflexion tasks,<sup>60</sup> and dynamic plantarflexion tasks<sup>61 62</sup>. Currently, there is no agreement in the literature in terms of plantarflexion torque measured during maximal contractions in individuals with NIAT; some authors did not find any difference between groups,<sup>14 63</sup> while others found statistically significant differences<sup>60 64</sup>. Interestingly, an investigation has observed a significant increase in LG activation during isometric plantarflexion tasks in people with AT following a 12-week training program<sup>60</sup>. Despite all of these efforts, currently there are no studies that have evaluated motor unit firing properties of calf muscles in individuals with NIAT.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

A strength of this study is that we will perform a detailed assessment of the mechanical and structural properties of the Achilles tendon and the calf muscles. One study showed that tendinopathy alters both the mechanical and material properties of the human Achilles tendon<sup>14</sup>. Morphological comparisons of tendinopathic and healthy tendons demonstrated a larger cross-sectional area for the degenerated Achilles tendon. Typically, a larger tendon is considered mechanically stronger due to its ability to dissipate stresses across the tendon and yield lower strain energy. Nonetheless, in the study of Arya and Kulig<sup>14</sup>, they demonstrated that despite having a larger cross-sectional area, the degenerated tendon had lower stiffness and Young's modulus compared with healthy tendons. Additionally, our study includes the use of SWE. This procedure has been used to measure tissue elasticity in tendons and might add to a better understanding of the effects of different types of exercises in tendons<sup>65</sup>. Furthermore, SWE is able to measure the Young modulus (slope of the stress-strain curve in the linear region<sup>66</sup>) of Achilles tendon with high reliability<sup>67</sup>. Previous studies suggest that SWE might be a useful tool for diagnosing and monitoring AT<sup>68</sup>. For instance, one study demonstrated that symptomatic Achilles tendons had lower Young modulus compared with healthy tendons and that stiffness increases in correlation with VISA-A scores after 6 months of treatment<sup>69</sup>.

Another strength of our study is using an isokinetic dynamometer to perform the training sessions. This device will allow us to control the intensity (50% MVC), range of motion (0° to 30° of plantarflexion), and angular speed (3°/s) of the contractions, enabling us to have close control over the time under tension (10 seconds) of the Achilles tendon. Although the use of isokinetic dynamometers is common to measure peak torque in musculoskeletal research, its use during a training protocol is limited. To our knowledge, there are no studies using isokinetic dynamometers to train individuals with NIAT. This represents an essential aspect of our RCT because previous studies investigating the effects of exercise in individuals with NIAT usually apply training protocols with insufficient control over the load, speed, pain tolerance, or the range of motion, and this could have influenced their results.

Regarding the study's limitations, the relatively short training protocol (6 weeks) might influence the changes expected in the mechanical and structural properties of the Achilles tendon; therefore, longer training interventions might be required to assess changes in these parameters in the long term. Moreover, due to the nature of the training protocol applied (eccentric and concentric exercises), blinding of participants is not possible because we need to explain how to do the different types of exercises on the isokinetic dynamometer. We are aware this introduces bias into the RCT, but unfortunately, it is not possible to achieve participants' blinding. Another study limitation is the inclusion of participants with bilateral symptoms, which could potentially affect the results. Since morphological changes to the asymptomatic tendon are common in this condition<sup>70</sup> and 45% of thickened Achilles tendons progress to develop clinical symptoms within 12 months<sup>71</sup>, we decided to also include these patients in the study. Additionally, participants' age range is another limitation of our study, as it might affect the reproducibility of our findings in older populations; however, we have decided to recruit participants in this age range based on previous studies showing age-related differences in Achilles tendon's stiffness and Young's modulus, which could confound the results of the intervention.

The study of motor units is an area in continuous development, which in recent years has allowed a more profound understanding of the neural mechanisms involved in muscle contractions. However, much of the research in this area has focused on the normal neurophysiology of muscle rather than its relationship with alterations of the musculoskeletal system.

This research will therefore provide new insights regarding the neuromechanical effects of ECC and CON exercises in the management of individuals with NIAT. A more precise understanding of the mechanisms involved in this pathology is essential to improve the rehabilitation programs commonly used in the management of this condition.

## Patient and Public Involvement

The research question in this study forms part of a larger discussion about exercise and pain relief within our patient and public involvement meetings. Patients will not be

involved in the analysis and data collection but will contribute to data interpretation and production of a lay summary of findings.

## **Ethics and Dissemination**

### **Ethical approval and trial registration**

The research protocol has been approved by the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham (ERN\_20-0604). Additionally, this protocol has been registered on ISRCTN registry (ISRCTN46462385).

Researchers will inform all participants of the characteristics of the research and will obtain written consent. Participants will be informed that they are free to withdraw from the study at any time without needing to provide a reason. In any unlikely adverse events, this will be immediately reported by the principal investigator to the ethics committee.

The results of this study will be submitted for publication in a peer review journal and presented at conferences.

### **Author Contributions**

ICH and EMV are responsible for the conception, design, and development of the protocol. EMV is the lead supervisor of ICH and DF is the co-supervisor. EMV and DF have provided guidance on methodological decisions and critical revision. All authors have read and subsequently approved the final manuscript.

### **Funding Statement**

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

### **Competing Interests Statement**

The authors have no competing interests to report.

**Data Sharing Statement**

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

**Figure captions**

**Figure 1.** Overview of the study design.

**Figure 2.** Experimental session design.

**Figure 3.** A) Anatomical landmarks used for ultrasonography and B) position of the electrodes in the MG, LG and SO muscles.

**References**

1. Sobhani S, Dekker R, Postema K, et al. Epidemiology of ankle and foot overuse injuries in sports: A systematic review. *Scand J Med Sci Sports* 2013;23(6):669-86. doi: 10.1111/j.1600-0838.2012.01509.x [published Online First: 2012/08/01]
2. Magnussen RA, Dunn WR, Thomson AB. Nonoperative treatment of midportion Achilles tendinopathy: a systematic review. *Clin J Sport Med* 2009;19(1):54-64. doi: 10.1097/JSM.0b013e31818ef090 [published Online First: 2009/01/07]
3. Janssen I, van der Worp H, Hensing S, et al. Investigating Achilles and patellar tendinopathy prevalence in elite athletics. *Research in Sports Medicine* 2018;26(1):1-12. doi: 10.1080/15438627.2017.1393748
4. Kvist M. Achilles tendon injuries in athletes. *Ann Chir Gynaecol* 1991;80(2):188-201. [published Online First: 1991/01/01]
5. Longo UG, Ronga M, Maffulli N. Achilles tendinopathy. *Sports Med Arthrosc Rev* 2009;17(2):112-26. doi: 10.1097/JSA.0b013e3181a3d625 [published Online First: 2009/05/15]
6. Scott A, Backman LJ, Speed C. Tendinopathy: Update on Pathophysiology. *J Orthop Sports Phys Ther* 2015;45(11):833-41. doi: 10.2519/jospt.2015.5884 [published Online First: 2015/09/22]
7. Maffulli N, Longo UG, Kadakia A, et al. Achilles tendinopathy. *Foot Ankle Surg* 2019 doi: 10.1016/j.fas.2019.03.009 [published Online First: 2019/04/30]
8. Henriksen M, Alkjaer T, Lund H, et al. Experimental quadriceps muscle pain impairs knee joint control during walking. *J Appl Physiol (1985)* 2007;103(1):132-9. doi: 10.1152/jappphysiol.01105.2006 [published Online First: 2007/04/07]
9. Tucker K, Butler J, Graven-Nielsen T, et al. Motor unit recruitment strategies are altered during deep-tissue pain. *J Neurosci* 2009;29(35):10820-6. doi: 10.1523/jneurosci.5211-08.2009 [published Online First: 2009/09/04]
10. Haglund-Akerlind Y, Eriksson E. Range of motion, muscle torque and training habits in runners with and without Achilles tendon problems. *Knee Surg Sports Traumatol Arthrosc* 1993;1(3-4):195-9. doi: 10.1007/bf01560205 [published Online First: 1993/01/01]
11. Mahieu NN, Witvrouw E, Stevens V, et al. Intrinsic risk factors for the development of achilles tendon overuse injury: a prospective study. *Am J Sports Med* 2006;34(2):226-35. doi: 10.1177/0363546505279918 [published Online First: 2005/11/02]



12. Henriksen M, Aaboe J, Graven-Nielsen T, et al. Motor responses to experimental Achilles tendon pain. *Br J Sports Med* 2011;45(5):393-8. doi: 10.1136/bjsm.2010.072561 [published Online First: 2010/06/15]
13. Wyndow N, Cowan SM, Wrigley TV, et al. Neuromotor control of the lower limb in Achilles tendinopathy: implications for foot orthotic therapy. *Sports Med* 2010;40(9):715-27. doi: 10.2165/11535920-000000000-00000 [published Online First: 2010/08/24]
14. Arya S, Kulig K. Tendinopathy alters mechanical and material properties of the Achilles tendon. *J Appl Physiol (1985)* 2010;108(3):670-5. doi: 10.1152/jappphysiol.00259.2009 [published Online First: 2009/11/07]
15. Martinez-Valdes E, Negro F, Falla D, et al. Surface electromyographic amplitude does not identify differences in neural drive to synergistic muscles. *J Appl Physiol (1985)* 2018;124(4):1071-79. doi: 10.1152/jappphysiol.01115.2017 [published Online First: 2018/02/09]
16. Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG. *J Appl Physiol (1985)* 2004;96(4):1486-95. doi: 10.1152/jappphysiol.01070.2003 [published Online First: 2004/03/16]
17. Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG: an update. *J Appl Physiol (1985)* 2014;117(11):1215-30. doi: 10.1152/jappphysiol.00162.2014 [published Online First: 2014/10/04]
18. Farina D, Negro F, Muceli S, et al. Principles of Motor Unit Physiology Evolve With Advances in Technology. *Physiology* 2016;31(2):83-94. doi: 10.1152/physiol.00040.2015
19. Alfredson H, Pietila T, Jonsson P, et al. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med* 1998;26(3):360-6. doi: 10.1177/03635465980260030301 [published Online First: 1998/06/09]
20. Maffulli N, Walley G, Sayana MK, et al. Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil* 2008;30(20-22):1677-84. doi: 10.1080/09638280701786427 [published Online First: 2008/07/09]
21. Sayana MK, Maffulli N. Eccentric calf muscle training in non-athletic patients with Achilles tendinopathy. *J Sci Med Sport* 2007;10(1):52-8. doi: 10.1016/j.jsams.2006.05.008 [published Online First: 2006/07/11]
22. Mayer F, Hirschmüller A, Müller S, et al. Effects of short-term treatment strategies over 4 weeks in Achilles tendinopathy. *Br J Sports Med* 2007;41(7):e6. doi: 10.1136/bjsm.2006.031732 [published Online First: 2007/01/31]
23. Jarin I, Backer HC, Vosseller JT. Noninsertional Achilles Tendinopathy: Topical Review. *Foot Ankle Int* 2020;1071100720914605. doi: 10.1177/1071100720914605 [published Online First: 2020/05/01]
24. Mafi N, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 2001;9(1):42-7. doi: 10.1007/s001670000148 [published Online First: 2001/03/28]
25. Yu J, Park D, Lee G. Effect of eccentric strengthening on pain, muscle strength, endurance, and functional fitness factors in male patients with achilles tendinopathy. *American journal of physical medicine & rehabilitation* 2013;92(1):68-76. doi: 10.1097/phm.0b013e31826eda63
26. Lindemann I, Coombes BK, Tucker K, et al. Age-related differences in gastrocnemii muscles and Achilles tendon mechanical properties in vivo. *J Biomech* 2020;112:110067. doi: 10.1016/j.jbiomech.2020.110067 [published Online First: 2020/10/10]
27. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med* 2001;35(5):335-41. doi: 10.1136/bjsm.35.5.335 [published Online First: 2001/10/02]



28. Alghadir AH, Anwer S, Iqbal A, et al. Test-retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal rating scales for measurement of osteoarthritic knee pain. *J Pain Res* 2018;11:851-56. doi: 10.2147/JPR.S158847

29. Beyer R, Kongsgaard M, Hougs Kjær B, et al. Heavy Slow Resistance Versus Eccentric Training as Treatment for Achilles Tendinopathy: A Randomized Controlled Trial. *Am J Sports Med* 2015;43(7):1704-11. doi: 10.1177/0363546515584760 [published Online First: 2015/05/29]

30. Iversen JV, Bartels EM, Langberg H. The victorian institute of sports assessment - achilles questionnaire (visa-a) - a reliable tool for measuring achilles tendinopathy. *Int J Sports Phys Ther* 2012;7(1):76-84.

31. Rompe JD, Furia J, Maffulli N. Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. *Am J Sports Med* 2009;37(3):463-70. doi: 10.1177/0363546508326983 [published Online First: 2008/12/18]

32. Chan A-W, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158(3):200-07. doi: 10.7326/0003-4819-158-3-201302050-00583

33. Chapman JP, Chapman LJ, Allen JJ. The measurement of foot preference. *Neuropsychologia* 1987;25(3):579-84. doi: 10.1016/0028-3932(87)90082-0 [published Online First: 1987/01/01]

34. Faul F, Erdfelder E, Buchner A, et al. Statistical power analyses using G\*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009;41(4):1149-60. doi: 10.3758/brm.41.4.1149 [published Online First: 2009/11/10]

35. Craig CL, Marshall AL, Sjostrom M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35(8):1381-95. doi: 10.1249/01.Mss.0000078924.61453.Fb [published Online First: 2003/08/06]

36. Martin RL, Irrgang JJ, Burdett RG, et al. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int* 2005;26(11):968-83. doi: 10.1177/107110070502601113 [published Online First: 2005/11/29]

37. Sullivan MJL, Bishop S, Pivik J. The Pain Catastrophizing Scale: Development and Validation. *Psychological Assessment* 1996;7:524-32. doi: 10.1037//1040-3590.7.4.524

38. Miller RP, Kori SH, Todd DD. The Tampa Scale: a Measure of Kinisophobia. *The Clinical Journal of Pain* 1991;7(1):51.

39. Del Vecchio A, Negro F, Holobar A, et al. You are as fast as your motor neurons: speed of recruitment and maximal discharge of motor neurons determine the maximal rate of force development in humans. *J Physiol* 2019;597(9):2445-56. doi: 10.1113/jp277396 [published Online First: 2019/02/16]

40. Chen MJ, Fan X, Moe ST. Criterion-related validity of the Borg ratings of perceived exertion scale in healthy individuals: a meta-analysis. *Journal of Sports Sciences* 2002;20(11):873-99. doi: 10.1080/026404102320761787

41. van Poppel MN, Chinapaw MJ, Mokkink LB, et al. Physical activity questionnaires for adults: a systematic review of measurement properties. *Sports Med* 2010;40(7):565-600. doi: 10.2165/11531930-000000000-00000 [published Online First: 2010/06/16]

42. Vlaeyen JW, Kole-Snijders AM, Boeren RG, et al. Fear of movement/(re)injury in chronic low back pain and its relation to behavioral performance. *Pain* 1995;62(3):363-72. doi: 10.1016/0304-3959(94)00279-n [published Online First: 1995/09/01]

43. Goubert L, Crombez G, Van Damme S, et al. Confirmatory factor analysis of the Tampa Scale for Kinesiophobia: invariant two-factor model across low back pain patients and fibromyalgia patients. *Clin J Pain* 2004;20(2):103-10. doi: 10.1097/00002508-200403000-00007 [published Online First: 2004/02/11]

44. Roelofs J, Goubert L, Peters ML, et al. The Tampa Scale for Kinesiophobia: further examination of psychometric properties in patients with chronic low back pain and fibromyalgia. *Eur J Pain* 2004;8(5):495-502. doi: 10.1016/j.ejpain.2003.11.016 [published Online First: 2004/08/25]
45. Swinkels-Meewisse EJ, Swinkels RA, Verbeek AL, et al. Psychometric properties of the Tampa Scale for kinesiophobia and the fear-avoidance beliefs questionnaire in acute low back pain. *Man Ther* 2003;8(1):29-36. doi: 10.1054/math.2002.0484 [published Online First: 2003/02/15]
46. Swinkels-Meewisse IE, Roelofs J, Verbeek AL, et al. Fear of movement/(re)injury, disability and participation in acute low back pain. *Pain* 2003;105(1-2):371-9. doi: 10.1016/s0304-3959(03)00255-0 [published Online First: 2003/09/23]
47. Heuts PH, Vlaeyen JW, Roelofs J, et al. Pain-related fear and daily functioning in patients with osteoarthritis. *Pain* 2004;110(1-2):228-35. doi: 10.1016/j.pain.2004.03.035 [published Online First: 2004/07/28]
48. Bolsterlee B, Gandevia SC, Herbert RD. Effect of Transducer Orientation on Errors in Ultrasound Image-Based Measurements of Human Medial Gastrocnemius Muscle Fascicle Length and Pennation. *PLoS One* 2016;11(6):e0157273. doi: 10.1371/journal.pone.0157273 [published Online First: 2016/06/15]
49. Siu W-I, Chan C-h, Lam C-h, et al. Sonographic evaluation of the effect of long-term exercise on Achilles tendon stiffness using shear wave elastography. *Journal of Science & Medicine in Sport* 2016;19(11):883-87.
50. Coombes BK, Tucker K, Vicenzino B, et al. Achilles and patellar tendinopathy display opposite changes in elastic properties: A shear wave elastography study. *Scand J Med Sci Sports* 2018;28(3):1201-08. doi: 10.1111/sms.12986 [published Online First: 2017/10/04]
51. Arvanitidis M, Falla D, Martinez-Valdes E. Can visual feedback on upper trapezius high-density surface electromyography increase time to task failure of an endurance task? *J Electromyogr Kinesiol* 2019;49:102361. doi: 10.1016/j.jelekin.2019.102361 [published Online First: 2019/10/13]
52. Martinez-Valdes E, Negro F, Farina D, et al. Divergent response of low- versus high-threshold motor units to experimental muscle pain. *J Physiol* 2020;598(11):2093-108. doi: 10.1113/jp279225 [published Online First: 2020/03/19]
53. Negro F, Muceli S, Castronovo AM, et al. Multi-channel intramuscular and surface EMG decomposition by convolutive blind source separation. *J Neural Eng* 2016;13(2):026027. doi: 10.1088/1741-2560/13/2/026027 [published Online First: 2016/03/01]
54. Martinez-Valdes E, Laine CM, Falla D, et al. High-density surface electromyography provides reliable estimates of motor unit behavior. *Clin Neurophysiol* 2016;127(6):2534-41. doi: 10.1016/j.clinph.2015.10.065 [published Online First: 2016/01/19]
55. Boccia G, Martinez-Valdes E, Negro F, et al. Motor unit discharge rate and the estimated synaptic input to the vasti muscles is higher in open compared with closed kinetic chain exercise. *J Appl Physiol (1985)* 2019;127(4):950-58. doi: 10.1152/jappphysiol.00310.2019 [published Online First: 2019/08/02]
56. Martinez-Valdes E, Negro F, Laine CM, et al. Tracking motor units longitudinally across experimental sessions with high-density surface electromyography. *J Physiol* 2017;595(5):1479-96. doi: 10.1113/jp273662 [published Online First: 2016/12/30]
57. Cohen J. Statistical Power Analysis for the Behavioral Sciences. 2nd ed. Hillsdale (NJ) ed: L. Erlbaum Associates 1988:p.567.
58. Baur H, Divert C, Hirschmüller A, et al. Analysis of gait differences in healthy runners and runners with chronic Achilles tendon complaints. *Isokinetics and Exercise Science* 2004;12:111-16. doi: 10.3233/IES-2004-0161

59. Wyndow N, Cowan SM, Wrigley TV, et al. Triceps surae activation is altered in male runners with Achilles tendinopathy. *J Electromyogr Kinesiol* 2013;23(1):166-72. doi: 10.1016/j.jelekin.2012.08.010 [published Online First: 2012/11/13]

60. Masood T, Kalliokoski K, Bojsen-Moller J, et al. Plantarflexor muscle function in healthy and chronic Achilles tendon pain subjects evaluated by the use of EMG and PET imaging. *Clin Biomech (Bristol, Avon)* 2014;29(5):564-70. doi: 10.1016/j.clinbiomech.2014.03.003 [published Online First: 2014/04/10]

61. Reid D, McNair PJ, Johnson S, et al. Electromyographic analysis of an eccentric calf muscle exercise in persons with and without Achilles tendinopathy. *Phys Ther Sport* 2012;13(3):150-5. doi: 10.1016/j.ptsp.2011.08.003 [published Online First: 2012/07/21]

62. Yu J. Comparison of Lower Limb Muscle Activity during Eccentric and Concentric Exercises in Runners with Achilles Tendinopathy. *J Phys Ther Sci* 2014;26(9):1351-3. doi: 10.1589/jpts.26.1351 [published Online First: 2014/10/03]

63. Child S, Bryant AL, Clark RA, et al. Mechanical properties of the achilles tendon aponeurosis are altered in athletes with achilles tendinopathy. *Am J Sports Med* 2010;38(9):1885-93. doi: 10.1177/0363546510366234 [published Online First: 2010/05/29]

64. O'Neill S, Barry S, Watson P. Plantarflexor strength and endurance deficits associated with mid-portion Achilles tendinopathy: The role of soleus. *Phys Ther Sport* 2019;37:69-76. doi: 10.1016/j.ptsp.2019.03.002 [published Online First: 2019/03/19]

65. Washburn N, Onishi K, Wang J. Ultrasound elastography and ultrasound tissue characterisation for tendon evaluation. *Journal of Orthopaedic Translation* 2018;15 doi: 10.1016/j.jot.2018.06.003

66. Maganaris CN, Narici MV, Maffulli N. Biomechanics of the Achilles tendon. *Disabil Rehabil* 2008;30(20-22):1542-7. doi: 10.1080/09638280701785494 [published Online First: 2008/08/23]

67. Leung W, Chu K, Lai C. Sonographic evaluation of the immediate effects of eccentric heel drop exercise on Achilles tendon and gastrocnemius muscle stiffness using shear wave elastography. *PeerJ* 2017;5 doi: 10.7717/peerj.3592

68. Dirrichs T, Quack V, Gatz M, et al. Shear Wave Elastography (SWE) for the Evaluation of Patients with Tendinopathies. *Acad Radiol* 2016;23(10):1204-13. doi: 10.1016/j.acra.2016.05.012 [published Online First: 2016/06/20]

69. Dirrichs T, Quack V, Gatz M, et al. Shear Wave Elastography (SWE) for Monitoring of Treatment of Tendinopathies: A Double-blinded, Longitudinal Clinical Study. *Acad Radiol* 2018;25(3):265-72. doi: 10.1016/j.acra.2017.09.011 [published Online First: 2017/11/21]

70. Grigg NL, Wearing SC, Smeathers JE. Achilles tendinopathy has an aberrant strain response to eccentric exercise. *Med Sci Sports Exerc* 2012;44(1):12-7. doi: 10.1249/MSS.0b013e318227fa8c [published Online First: 2011/06/11]

71. Fredberg U, Bolvig L. Significance of ultrasonographically detected asymptomatic tendinosis in the patellar and achilles tendons of elite soccer players: a longitudinal study. *Am J Sports Med* 2002;30(4):488-91. doi: 10.1177/03635465020300040701 [published Online First: 2002/07/20]



Figure 1. Overview of the study design.

338x190mm (300 x 300 DPI)




Figure 2. Experimental session design.

338x190mm (300 x 300 DPI)




Figure 3. A) Anatomical landmarks used for ultrasonography and B) position of the electrodes in the MG, LG and SO muscles.

338x190mm (300 x 300 DPI)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

CONSORT 2010 checklist of information to include when reporting a randomised trial\*

Section and topic	Items N°	Checklist item	Page N°
Title and abstract			
	1a	Identification as a randomised trial in the title	1
	1b	Structured summary of trial design, methods, results, and conclusions	2
Introduction			
Background and objectives	2a	Scientific background and explanation rationale	4-6
	2b	Specific objectives or hypotheses	6
Methods			
Trial design	3a	Description of trial design (such as parallel, factorial) including allocation ratio	7
	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	N/A
Participants	4a	Eligibility criteria for participants	6, 7
	4b	Settings and locations where the data were collected	7
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were administered	11
Outcomes	6a	Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed	12
	6b	Any changes to trial outcomes after the trial commenced, with reasons	N/A
Sample size	7a	How sample size was determined	8
	7b	When applicable, explanation of any interim analyses and stopping guidelines	N/A
Randomisation:			
Sequence generation	8a	Method used to generate the random allocation sequence	18
	8b	Type of randomisation; details of any restriction (such as blocking and block size)	18
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	18
Implementation	10	Who generated the random allocation sequence, who enrolled the participants, and who assigned participants to interventions	18, 19
Blinding	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how	18, 19
	11b	If relevant, description of the similarity of interventions	N/A
Statistical methods	12a	Statistical methods used to compare groups for primary and secondary outcomes	19
	12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	N/A

136/bmjopen-2021-058683 on 29 July 2022. Downloaded from <http://bmjopen.bmj.com/> on June 13, 2025 at Agence Bibliographique de l'Enseignement Supérieur (A.B.E.S.). All rights reserved. No reuse allowed without permission. AI training, and similar technologies.

<b>Results</b>			
Participant flow (a diagram is strongly recommended)	13a	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome	N/A
	13b	For each group, losses and exclusions after randomisation, together with reasons	N/A
Recruitment	14a	Dates defining the periods of recruitment and follow-up	N/A
	14b	Why the trial ended or was stopped	N/A
Baseline data	15	A table showing baseline demographic and clinical characteristics for each group	N/A
Numbers analysed	16	For each group, number of participants (denominator) included in each analysis and whether analysis was by original assigned groups	N/A
Outcomes and estimation	17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)	N/A
	17b	For binary outcomes, presentation of both absolute and relative effect size is recommended	N/A
Ancillary analysis	18	Results of any other analyses performed, including subgroup analyses and adjusted analyses distinguishing pre-specified from exploratory	N/A
Harms	19	All important harms or unintended effects in each group	N/A
<b>Discussion</b>			
Limitations	20	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant multiplicity of analyses	19-22
Generalisability	21	Generalisability (external validity, applicability) of the trial findings	19-22
Interpretation	22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	19-22
<b>Other information</b>			
Registration	23	Registration number and name of trial registry	2, 7, 22
Protocol	24	Where the full trial protocol can be accessed, if available	N/A
Funding	25	Sources of funding and other support (such as supply of drugs), role of funders	23

\*We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration<sup>13</sup> for important clarifications on all the items. If relevant, we also recommend reading CONSORT extensions for cluster randomised trials,<sup>11</sup> non-inferiority and equivalence trials,<sup>12</sup> non-pharmacological treatments,<sup>32</sup> herbal interventions,<sup>33</sup> and pragmatic trials.<sup>34</sup> Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see [www.consort-statement.org](http://www.consort-statement.org).

From: Schulz K, Altman D, Moher D. CONSORT 2010 Statement: updated guideline for reporting parallel group randomised trials. BMJ 2010; 340:c 332

# BMJ Open

## Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy: Protocol for a randomised controlled trial.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058683.R2
Article Type:	Protocol
Date Submitted by the Author:	30-Jun-2022
Complete List of Authors:	Contreras-Hernandez, Ignacio; University of Birmingham Falla, Deborah; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences Martinez-Valdes, Eduardo; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences
<b>Primary Subject Heading</b>:	Sports and exercise medicine
Secondary Subject Heading:	Rehabilitation medicine
Keywords:	SPORTS MEDICINE, Adult orthopaedics < ORTHOPAEDIC & TRAUMA SURGERY, Foot & ankle < ORTHOPAEDIC & TRAUMA SURGERY, Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY

SCHOLARONE™  
Manuscripts

Title

Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy: Protocol for a randomised controlled trial.

**Authors:** Ignacio Contreras-Hernandez<sup>1</sup>, Deborah Falla<sup>1</sup>, Eduardo Martinez-Valdes<sup>1</sup>

Corresponding author:  
Eduardo Martinez-Valdes<sup>1</sup>  
Assistant Professor in Physiotherapy  
Centre of Precision Rehabilitation for Spinal Pain (CPR Spine)  
School of Sport, Exercise and Rehabilitation Sciences  
College of Life and Environmental Sciences  
University of Birmingham  
Edgbaston B15 2TT, United Kingdom  
T: +44 (0) 121 4158187  
M: +44 (0) 7961 548851  
Email: [e.a.martinezvaldes@bham.ac.uk](mailto:e.a.martinezvaldes@bham.ac.uk)

**Author Affiliations**

<sup>1</sup>Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, UK.

Email: Ignacio Contreras-Hernandez<sup>1</sup> [iac921@student.bham.ac.uk](mailto:iac921@student.bham.ac.uk) – Deborah Falla<sup>1</sup> [d.falla@bham.ac.uk](mailto:d.falla@bham.ac.uk) – Eduardo Martinez-Valdes<sup>1</sup> [e.a.martinezvaldes@bham.ac.uk](mailto:e.a.martinezvaldes@bham.ac.uk)

**WORD COUNT MAIN TEXT: 5943 (excluding title page, abstract, references, figures, tables, and acknowledgments)**

## Abstract

**Introduction.** There is limited evidence on the neural strategies employed by the central nervous system to control muscle force in the presence of non-insertional Achilles tendinopathy (NIAT). Additionally, the neuromuscular mechanisms by which exercise may help to resolve tendon pain remain unclear.

**Objective.** This study aims to firstly establish changes in the gastrocnemius-soleus motor unit firing properties after applying a training protocol of 6-weeks based on either controlled eccentric or concentric contractions in individuals with NIAT. Secondly, we want to determine changes in the level of pain and function and mechanical and structural properties of the Achilles tendon after applying the same training protocol. Additionally, we want to compare these variables at baseline between individuals with NIAT and asymptomatic controls.

**Methods and analysis.** A total of 26 individuals with chronic (> 3 months) NIAT and 13 healthy controls will participate in the study. Individuals with NIAT will be randomised to perform eccentric or concentric training for 6 weeks. Motor unit firing properties of the medial gastrocnemius, lateral gastrocnemius and soleus muscles will be assessed using high-density surface electromyography, as well as Achilles tendon length, cross-sectional area, thickness, and stiffness using B-mode ultrasonography and shear wave elastography. Moreover, participants will complete a battery of questionnaires to document their level of pain and function.

**Ethics and dissemination.** Ethical approval (ERN\_20-0604A) for the study was obtained from the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham. The results of the study will be published in peer-review journals.

**Protocol Registration.** This protocol has been registered on ISRCTN registry (ISRCTN46462385).

**Keywords:** Achilles tendinopathy, HD-sEMG, motor unit, morphological properties, mechanical features.

**WORD COUNT:** 271





## Introduction

Achilles tendinopathy (AT) is a painful overuse injury of the Achilles tendon and it is common among athletes, especially those involved in running and jumping sports<sup>[1-3]</sup>. AT is clinically diagnosed when the patient presents with a combination of localized pain, swelling of the Achilles tendon, and loss of function<sup>[4]</sup>. The essence of tendinopathy is a failed healing response, with degeneration and proliferation of tenocytes, disruption of collagen fibres, and subsequent increase in non-collagenous matrix<sup>[5]</sup>. These structural changes in the tendon result in increased cross-sectional area, reduced tendon stiffness, and altered viscoelastic properties in both symptomatic and asymptomatic tendons<sup>[6]</sup>.

The aetiology of AT remains debated and is likely caused by intrinsic and extrinsic factors<sup>[7]</sup>. One of the most accepted theories is that pain perception during early support loading may trigger inhibition of neuromuscular activity of the calf muscles detected as a reduction in electromyographic (EMG) amplitude<sup>[8-9]</sup>. Thus, the decrease in the ability to generate force in patients with AT could also reflect the decline in neuromuscular activity observed<sup>[10-11]</sup>. Moreover, it has been observed that this motor inhibition also affects synergist and antagonist muscles<sup>[12]</sup>. Additionally, individuals with tendinopathy tend to use movement patterns that place an excessive or abnormal load on their tendons<sup>[6]</sup>, and it is believed that these motor adaptations may generate greater torsional stress in the tendon<sup>[12-13]</sup>. Finally, studies have shown that AT reduces tendon's stiffness<sup>[14]</sup>, which impairs the mechanisms responsible for transmitting force to the bone. Therefore, it is very likely that these alterations in tendon properties may produce changes in the neural drive received by the calf muscles.

Until now, most studies examining the neuromuscular impairments induced by AT have focused on investigating changes in interference EMG amplitude which is an indirect estimate of neural activity with many factors of influence<sup>[15-17]</sup>. Clearer information about the neural strategies employed by the central nervous system to control muscle force in the presence of AT can be obtained through motor unit recordings, since motor unit firing properties represent the direct neural output from the spinal cord to muscles<sup>[18]</sup>. Nevertheless, there are no studies that have measured motor unit firing properties in

individuals with NIAT.

Although eccentric exercise has been widely used for the treatment of NIAT, the mechanisms by which eccentric exercises may help to resolve tendon pain remain unclear<sup>[7]</sup>. Regarding eccentric exercise alone, two prospective studies have reported a significant reduction in pain intensity and change on the Victorian Institute of Sports Assessment-Achilles questionnaire (VISA-A) in recreational athletes following a 12-week exercise program<sup>[19 20]</sup>. In contrast, another study in non-athletic individuals found no significant improvement after a similar 12-week exercise program<sup>[21]</sup>. Concerning eccentric exercises with an adjunctive treatment (e.g., pulsed ultrasound, ice, sensory motor training), a 4-week intervention study resulted in decreased pain and higher plantarflexion peak torque in individuals with NIAT compared to controls<sup>[22]</sup>. However, studies that include eccentric exercises with an adjunctive treatment showed limited evidence of improvement over eccentric exercises alone<sup>[23]</sup>.

There are few studies where the effectiveness of eccentric versus concentric exercises has been compared. Mafi et al. showed that patient satisfaction and return to previous activity were significantly superior after participating in a 12-week rehabilitation protocol based on eccentric exercise compared to concentric exercise. Although pain intensity decreased significantly in both groups, the amount of pain reduction was significantly greater for those that performed eccentric exercise<sup>[24]</sup>. Likewise, Yu et al.<sup>[25]</sup> demonstrated that eight weeks of eccentric exercise was more effective at reducing pain than concentric in individuals with chronic NIAT. Additionally, they found that eccentric exercise was more effective than concentric exercise at increasing muscle strength and endurance, and improving function<sup>[25]</sup>. In these investigations, participants performed the rehabilitation protocols with insufficient control over the load, speed, pain tolerance, or the range of motion in which the exercises were performed. Moreover, it is essential to consider that when participants perform an eccentric plantar flexion exercise without adequate equipment, it is difficult to achieve pure eccentric contractions, which could have influenced the results obtained in these studies.

Based on the above, the aims of this study are to (1) establish changes in the gastrocnemius-soleus motor unit firing properties after applying a training protocol of 6-weeks based on either controlled eccentric or concentric contractions in individuals with NIAT; (2) determine changes in the level of pain and function and mechanical and structural properties of the Achilles tendon after applying the same training protocol; (3) compare these properties at baseline between individuals with NIAT and asymptomatic controls.

## Methods

### Participants

Twenty-six individuals with NIAT and thirteen asymptomatic controls will be recruited from the University of Birmingham staff/student population and the local community via leaflets, e-mail, and social media.

Men or women aged 18 to 55 years old will be recruited. This age range was selected based on previous findings showing lower stiffness and Young's modulus of the Achilles tendon in older than younger population<sup>[26]</sup>. Inclusion criteria are NIAT determined by an experienced physiotherapist based on defined clinical findings (VISA-A<sup>[27]</sup> and NRS (Numerical Rating Scale)<sup>[28]</sup> scores), physical examination, and ultrasound assessment, as well as having pain for at least 3 months<sup>[29]</sup>. VISA-A scores less than 90 will be considered as a reference to identify individuals with NIAT<sup>[30]</sup>. Regarding the NRS scores, previous studies have shown high variability in individuals with NIAT<sup>[29 31]</sup>, thus we will consider individuals with an NRS score  $\geq 2$ . Physical examination will include palpation of the Achilles tendon along its whole length in a proximal to distal direction, and gentle squeezing the tendon between the thumb and the index finger to identify tenderness over the tendon<sup>[20]</sup>. Ultrasound evaluation of the tendon's mid-portion will include identifying local thickening of the tendon and/or irregular tendon structure with hypoechoic areas and/or irregular fiber orientation<sup>[31]</sup>.

The exclusion criteria for both groups will include: (1) systemic or inflammatory conditions including rheumatic, neuromuscular disorders, and malignancy, (2) current or history of chronic respiratory, neurological, or cardiovascular diseases, (3) history of lower limb surgery. Specific exclusion criteria for the participants with NIAT are participation in

any other treatment or rehabilitation program for AT, corticosteroid injections in the previous 12 months, and insertional AT. Additionally, if any participants present non-insertional and insertional AT concurrently in the same limb, they will be excluded. Specific exclusion criteria for the control group are pain/injury in the lower limbs within the previous 6 months, history of AT or lower limb surgery.

## Study Design

This two-arm, parallel-group, randomised controlled trial will be conducted from October 2021 to December 2022 at a laboratory within the Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), University of Birmingham, UK. The Science, Technology, Engineering and Mathematics Ethical Review Committee, University of Birmingham, United Kingdom, approved the study (ERN\_20-0604A). All participants will provide written informed consent prior to participation. The study will be conducted according to the Declaration of Helsinki. This protocol has been designed following the SPIRIT 2013 statement<sup>[32]</sup> (see supplementary file 1). The ISRCTN registry number for this trial is ISRCTN46462385 (trial registration data can be found as supplementary file 2). Time schedule of enrolment, interventions, assessments, and visits can be found as supplementary file 3. The consent forms provided to healthy controls and patients can be also found as supplementary files (supplementary files 4 and 5). Reporting will follow the CONSORT statement and the CONSORT flow diagram will be used to describe the flow of participants throughout the trial (see supplementary file 6).

Participants with NIAT will visit the laboratory over six consecutive weeks for the experimental sessions (at weeks 1, 3, and 6) and training sessions (2-3 sessions per week) (**Fig.1**). We will randomly allocate these participants into two groups: eccentric (ECC) or concentric (CON) training. Healthy participants will visit the laboratory once to allow baseline comparison with ECC and CON groups. Additionally, we will randomise the assessed leg in the healthy control group, and the most symptomatic leg in the ECC and CON groups will be evaluated. Finally, foot preference in specific daily activities (foot dominance) will be determined using a behavioral foot-preference inventory<sup>[33]</sup>. Each experimental session will last 2.5 hours, and each training session will last 40 minutes.

## Sample size

According to power calculations (G\*Power software)<sup>[34]</sup>, a total of 26 individuals with NIAT (ECC group=13, and CON group= 13) and 13 healthy controls will be required for this study. This sample size considers a power=0.80, alpha=0.01, 25% loss of participants and an effect size (d) of 1.7 calculated from the study of Yu et al.<sup>[25]</sup>, where the authors compared reductions in pain after an 8-week concentric and eccentric training protocol in individuals with NIAT.

**Figure 1.** Overview of the study design.

## Experimental sessions

These sessions will involve the completion of questionnaires, ultrasound imaging of the gastrocnemius-soleus muscles and the Achilles tendon, surface electromyography and torque recordings. All the procedures during the experimental sessions will be done by Ignacio Contreras-Hernandez (IC-H) and Joeri van Helden (JVH). IC-H is a PhD student at the University of Birmingham, Master in Physiology, Physiotherapist, and member of the CPR Spine group. Joeri is a PhD student at the University of Birmingham, Master in Neuroscience, Psychologist, and member of the CPR Spine group.

Anthropometric data (age, gender, weight, height, leg dominance, and body mass index) will be obtained, and the participants will then be asked to complete a battery of questionnaires. This includes the International Physical Activity Questionnaire short form (IPAQ-SF),<sup>[35]</sup> VISA-A,<sup>[27]</sup> Foot and Ankle Ability Measure (FAAM),<sup>[36]</sup> Pain Catastrophizing Scale (PCS),<sup>[37]</sup> and Tampa Scale for Kinesiophobia (TSK)<sup>[38]</sup>.

Additionally, participants will be asked to report their current level of pain using the NRS score. After that, participants will lie prone on the chair of a Biodex System 3 dynamometer (Biodex Medical Systems), and ultrasonography (LOGIQ S8 GE Healthcare, Milwaukee, USA) will be used to measure the length, thickness, and cross-sectional area of the Achilles tendon, fascicle length, thickness, and pennation angle of the medial gastrocnemius (MG), and thickness of the lateral gastrocnemius (LG), and soleus (SO) muscles during rest. Then, we will prepare the skin and place the electrodes on the MG, LG, and SO muscles, and using high-density surface electromyography (HD-sEMG); we will ensure minimal electrical activity of these muscles during rest conditions for the measurements of the Achilles tendon stiffness (passive elastography).

Following the ultrasound assessment the maximal voluntary contraction (MVC) will be recorded during three isometric plantarflexion contractions of 5 seconds each<sup>[15]</sup>. Between each MVC, the volunteers will have two minutes of rest<sup>[15]</sup> and all MVCs will be performed at 0° of plantarflexion. The highest MVC value will be used as the reference maximal torque. We will use this MVC value as a reference for the isometric and dynamic plantarflexion contractions during the experimental and training sessions, to avoid multiple MVC measurements that may be produce pain and discomfort in individuals with NIAT. Afterwards, we will measure the stiffness of the tendon during two isometric plantarflexion contractions at 10% MVC (1 s ramp-up, 12 s hold, 1 s ramp-down, and 30 s rest) (active elastography). Subsequently, using HD-sEMG, we will record motor unit activity of the MG, LG, and SO muscles during two isometric plantarflexion contractions at 10%, 40%, and 70% MVC (10% MVC/s ramp-up, 10 s hold, 10% MVC/s ramp-down, and 30 s rest), one concentric-eccentric plantarflexion contraction at 25% and 50% MVC, and one eccentric-concentric plantarflexion contraction at 25% and 50% MVC (the order of the different types of contractions will be randomly selected). Volunteers will have five minutes of rest at the end of each type of contraction (isometric, concentric-eccentric, and eccentric-concentric). After 15 minutes of rest, HD-sEMG will be recorded from the MG, LG, and SO muscles during six explosive (fast force development) isometric plantarflexion contractions at 75% MVC (1 s ramp-up, 3 s hold, 1 s ramp-down, and 10 s rest)<sup>[39]</sup>. Finally, both the rate of perceived exertion and the level of pain will be monitored



regularly throughout the session, using the Borg ratings of perceived exertion scale<sup>[40]</sup> and the NRS. (Fig. 2).

During all contractions, visual feedback of the target torque output will be provided via computer monitor positioned 1 m from the participant. Prior to the contractions, participants will be instructed to match the force output as closely as possible to the target force for the full duration of the contraction. For the dynamic contractions, the range of motion will be set at the total of 30° (neutral position 0° to 30° of plantarflexion) and the angular speed will be set at 3°/s.

**Figure 2.** Experimental session design.

### Training sessions

The training sessions will be done by Michalis Arvanitidis (MA). MA is a PhD student at the University of Birmingham, Master in Advanced Manipulative Physiotherapy, and Specialist Musculoskeletal Physiotherapist (Member of the Musculoskeletal Association of Chartered Physiotherapist) and member of the CPR Spine group.

All the training sessions will be done in prone position on the Biodex System 3 dynamometer.

The participants in the ECC group will be asked to perform a warm-up consisting of three eccentric plantarflexion contractions at 25% MVC; this will be followed by the eccentric training protocol. This protocol consists of 4 x 15 eccentric plantarflexion contractions at 50% MVC, range of motion of 30° (neutral position 0° to 30° of plantarflexion), time under tension of 10 seconds, angular speed of 3°/s and 3 minutes of rest between each series. Visual feedback of the exerted torque will be provided. Participants in the CON group will perform a warm-up consisting of three concentric plantarflexion contractions at 25% MVC, and then, the concentric training protocol. This protocol consists of 4 x 15 concentric plantarflexion contractions at 50% MVC, range of motion of 30° (neutral position 0° to 30° of plantarflexion), time under tension of 10 seconds, angular speed of 3°/s, and 3 minutes of rest between each series.

Preceding the contractions, participants will be instructed to match the torque output as closely as possible to the target torque for the full duration of the contraction.

## Follow-up

Participants with NIAT will be asked to report their level of pain and function at 3 and 6 months after completing the training protocol.

## Outcome measures

### Primary outcomes measure

The primary outcomes for this study will be GM, GL, and SO muscles motor unit firing properties assessed using HD-sEMG and decomposition techniques. These properties include motor unit discharge rate, recruitment and de-recruitment thresholds, and discharge rate variability.

### Secondary outcomes measure

Secondary outcomes will include level of pain and function assessed using the NRS and VISA-A questionnaire, Achilles tendon length, thickness, cross-sectional area, and stiffness using B-mode ultrasonography and shear wave elastography. Additionally, secondary outcomes will include GM, GL, and SO muscles thickness, and GM muscle fascicle length and pennation angle evaluated using B-mode ultrasonography, as well as

the level of physical activity, physical function, pain catastrophizing and fear of movement assessed using the IPAQ-SF, FAAM, PCS, and TSK questionnaires, respectively.

## Questionnaires

In each experimental session, participants will be asked to complete the IPAQ-SF, VISA-A, FAAM, PCS, and TSK questionnaires to measure physical activity level, symptoms in individuals with AT, physical function, pain catastrophizing and fear of movement, respectively. The IPAQ has become the most widely used physical activity questionnaire,<sup>[41]</sup> and it has acceptable measurement properties for monitoring population levels of physical activity among 18 to 65 years old adults in diverse settings<sup>[35]</sup>. The VISA-A was developed with the aim of evaluating the symptoms of AT and their impact on physical activity. This questionnaire is valid, reliable, easy to use, and ideal for comparing patients' progress in clinical settings<sup>[27]</sup>. The FAAM was developed to meet the need for a self-reported evaluative instrument that comprehensively assesses the physical function of individuals with musculoskeletal disorders of the leg, foot, and ankle. The FAAM is a reliable, valid, and responsive measure of self-reported physical function<sup>36</sup>. Additionally, we will use the PCS to understand the psychological processes that lead to heightened physical and emotional distress in response to aversive stimulation. This questionnaire is a reliable and valid measure of catastrophizing<sup>[37]</sup>. Finally, we will apply the TSK to measure the fear of movement/(re)injury. This questionnaire has been validated in patients with chronic back pain <sup>[42-44]</sup>, acute back pain <sup>[45 46]</sup>, osteoarthritis <sup>[47]</sup>, and fibromyalgia <sup>[43 44]</sup>.

## Measurement set-up

For the measurements of the Achilles tendon, MG, LG, and SO muscles, participants will lie prone on the dynamometer, with their knees extended, and their tested foot tightly strapped on the footplate. The pelvis will be stabilized with another strap to minimise compensatory movements. The ankle will be positioned in 0° of plantarflexion and the axis of the dynamometer will be aligned with the inferior tip of the lateral malleolus. The setting and position of the set up (i.e., chair and isokinetic device) will be saved, so the participants' position will be similar in each experimental session.

## Ultrasound measurements

All ultrasound images will be obtained using an ultrasound imaging device equipped with shear wave elastography (SWE) (LOGIQ S8 GE Healthcare, Milwaukee, USA). For the measurements of the length, thickness, and cross-sectional area of the Achilles tendon, and the measurements of the fascicle length, thickness, and pennation angle of the calf muscles, B-mode will be used with a 16-linear array probe (50 mm, 4-15 MHz). Subsequently, for the measurements of the Achilles tendon's stiffness during rest conditions and isometric plantarflexion contraction, the elastography mode will be used with a 9-linear array probe (44 mm, 2-8 MHz).

An adaptation of the protocol developed by Arya and Kulig<sup>[14]</sup> will be used to measure the structural properties of the Achilles tendon. Briefly, to obtain tendon length, the ultrasound transducer will be placed longitudinally over the posterior aspect of the heel, and the distal part of the Achilles tendon will be imaged, and the corresponding point will be marked on the skin with a marker. Then, the ultrasound probe will be moved proximally to locate the musculotendinous junction of the MG, and this point will be marked on the skin. The distance between these two points will be measured with a tape and this distance will represent the resting length of the Achilles tendon. Marks at 2 cm, 4 cm, and 6 cm above the Achilles tendon insertion will then be made on the skin. Later, these marks will be used to place the probe in the transversal plane and determine the cross-sectional area of the Achilles tendon at 2 cm, 4 cm, and 6 cm of its insertion. Additionally, we will use these marks to place the probe in the sagittal plane and determine the thickness of the Achilles tendon at 2 cm, 4 cm, and 6 cm of its insertion.

For muscle ultrasound images, the mid-line of the leg will be marked in the direction of the Achilles tendon. Additionally, a mark will be made on the leg 10 cm above the musculotendinous junction of the MG muscle and 4 cm medial to the mid-line. In this position, we will place the middle point of the HD-sEMG electrode grid and mark the contour of the grid on the skin. We will use these marks to place the probe in the sagittal plane and obtain the images of the MG muscle. Similarly, the leg will be marked 10 cm above the musculotendinous junction of the MG muscle and 4 cm lateral to the mid-line.

Then, we will repeat the procedure mentioned above, but now for the LG muscle. Next, the leg will be marked 5 cm below the musculotendinous junction of the MG and 4 cm lateral to the mid-line. Again, we will repeat the procedure mentioned above, but now for the SO muscle (**Fig. 3**). The middle column of the HD-sEMG electrode grid will be used as a reference (see HD-sEMG and torque section below) to place the probe in the same position during all the experimental sessions and the images will be acquired with the probe oriented in the sagittal plane, and perpendicular to the skin, according to the recommendations of Bolsterlee et al<sup>[48]</sup>. To ensure that we are measuring the exact location of interest, we will use the procedure described above in each experimental session, and we also will mark the middle point of the ultrasound probe. Then, during the acquisition of the ultrasound images, we will align the mark in the ultrasound probe with the marks on the skin at 2 cm, 4 cm, and 6 cm from the Achilles tendon insertion and with the mark of the middle point of the HD-sEMG electrode grid of each muscle. This procedure will allow us to identify the location of interest during the analysis of the ultrasound images since we know that the middle point of the image represents the location of interest. The software Image J (<https://imagej.nih.gov/ij/>) will be used to analyze these images and determine the fascicle length, thickness, and pennation angle.

For tendon stiffness measurements, the probe will be placed in the sagittal plane, with the middle part of the probe located at 4 cm above the Achilles tendon insertion. A probe holder will be used to avoid applying pressure over the tendon and movements that may interfere with the measurements. Then, we will perform a trial SWE to check for possible voids. If voids are detected at this stage, we will remove the probe holder and place the ultrasound probe again. The passive elastography images will be acquired for 12 seconds (twice), and the active elastography images will be acquired during two isometric plantarflexion contractions at 10% MVC (1 s ramp-up, 12 s hold, 1 s ramp down, 1 min rest). We will check the elastography images following each measurement to determine possible voids that may affect our results. If voids in the middle part of the tendon (at 4 cm of the insertion) are detected in this stage, we will repeat the procedure. A shear elastography colour map (height x width, 2.5 cm x 1 cm) will be chosen using elastography ultrasound tools to allow complete visualization of the Achilles tendon width, and a region of interest (ROI) of 3 mm diameter<sup>[49]</sup> will be selected in the middle of each

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

image to determine the shear wave velocity (m/s) and Young’s modulus (kPa). Finally, we will average the mean shear wave velocity and Young’s modulus over the ROIs of consecutive images<sup>[50]</sup>.

**Figure 3.** A) Anatomical landmarks used for ultrasonography and B) position of the electrodes in the MG, LG and SO muscles.

**HD-sEMG and torque recording**

Prior to electrode placement, the skin will be shaved (if necessary), gently abraded (Nuprep, Skin Prep Gel, Weaver and Company, Aurora, Colorado) to reduce skin impedance, and cleaned with water. Three two-dimensional (2D) adhesive grids (SPES Medica, Salerno, Italy) of 13 x 5 equally spaced electrodes (each of 1 mm diameter, with an inter-electrode distance of 8 mm) will be used to record the HD-EMG signals.



Conductive paste (AC-CREAM, SPES Medica, Genova, Italy) will be placed into the cavities of the grid, and the HD-sEMG electrodes will be placed in the exact position described for the ultrasound measurements of the triceps surae (one electrode grid for each muscle).

All signals will be converted from analog-to-digital by a 16-bit converter (Quattrocento- OTBioelettronica, Torino, Italy). The sampling frequency will be 2048 Hz and the amplifier gain will be set to 150. HD-sEMG signals will be digitally filtered with a bandwidth set up to 10 Hz for high pass cut frequency and to 500 Hz for low pass cut frequency<sup>51</sup>. HD-sEMG will be acquired in monopolar mode with reference electrodes (WhiteSensor WS, Ambu A/S, Ballerup, Denmark) positioned in the head of the fibula and with a strip in the thigh of the evaluated leg. All the electrode grids and reference electrodes will be connected to the same bioelectrical amplifier (Quattrocento-OT-Bioelettronica, Torino, Italy).

The torque exerted by the volunteers will be assessed with the isokinetic dynamometer, which will be synchronized with the HD-sEMG signals. Synchronization will be obtained by recording torque signals generated by the isokinetic dynamometer through the auxiliary input of the EMG amplifier<sup>51</sup>.

## Signal Analysis

The torque signal will be low pass filtered at 15 Hz and then used to quantify torque steadiness (coefficient of variation of torque, SD torque/mean torque \* 100) from the stable part of the contractions<sup>52</sup>.

The HD-sEMG signals will be decomposed into motor unit spike trains with an algorithm based on blind source separation, which provides automatic identification of motor unit activity<sup>52</sup>, and the accuracy of the decomposition will be tested with the silhouette measure, which will be set to  $\geq 0.90$ <sup>53</sup>. The signals will be decomposed during the entire duration of the contractions, and the discharge times of the motor units will be transformed in binary spike trains<sup>54</sup>. The mean discharge rate and the discharge rate variability (CoV of the interspike interval [CoV<sub>isi</sub>]) will be determined during the stable plateau of torque signal. Additionally, motor unit recruitment and de-recruitment thresholds will be defined as the ankle plantarflexion torques (%MVC) at the times when

the motor units began and stopped discharging action potentials, respectively<sup>[52]</sup>. Discharges rates at recruitment and de-recruitment will be determined using the first and last six discharges of the motor units.<sup>[52]</sup> Erroneous discharges will be visually inspected and edited using a custom algorithm<sup>[52 55]</sup>. Motor unit activity will be monitored longitudinally with a recent method proposed by Martinez-Valdes et al.<sup>[56]</sup> which allows tracking the same motor units across different experimental sessions.

**Adverse event management**

Participants will be informed that they may experience some pain during or after the experimental and training sessions. Monitoring of participants' pain will be done in each experimental and training sessions using the NRS. Appropriate rest time will be provided throughout the experimental and training sessions, and extra rest periods will be given to the participants at any time if required. If a participant experiences moderate pain (>6 NRS) during the contractions, they will be given additional time to rest. If the pain intensity is maintained or worsens, we will terminate and reschedule the session. The session will be rescheduled in the upcoming three days. If pain intensity is maintained or worsened during these days and rescheduling the session is not possible, the participant will be removed from the study. This will be considered as an adverse effect, and it will be reported to the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham.

**Randomisation and blinding**

Individuals with NIAT will be randomised by an independent researcher (Dr. Eduardo Martinez-Valdes (EM-V)) in a 1:1 allocation ratio to either ECC or CON groups (parallel-groups) using computer-generated simple scheme randomisation (<https://sigdaan.com/randomization/app/randomization-app>). Allocation concealment will be ensured, as EM-V will secure the randomisation code using password-protected files. EM-V will give access to MA to the randomisation code once each participant has completed the first experimental session.

In order to achieve double-blinding, IC-H and JVH will perform the experimental sessions and MA will perform the training sessions. IC-H will use the blindr (<https://github.com/U8NWXD/blindr>) software to encode the results from different participants and will be blinded to the training protocol applied to the participants. MA will be blind to the participant's results, but not to the training protocol applied. EM-V will unmask the results after the data analysis is performed. Due to the nature of the interventions, participants' blinding is not possible.

## Statistical analysis

IBM SPSS Statistics for Windows, Version 25.0 (Armonk, New York, USA) computer software will be used for statistical analysis of the data. Intention-to-treat and per protocol analysis will be performed. Descriptive statistics will be used to interpret the data which will be presented as mean  $\pm$  SD. The Shapiro-Wilk Test will be used to assess data normality. The level of significance for all statistical procedures will be set at  $\alpha=0.05$  and 95% confidence intervals (CI) will be reported. Independent t-test will be used to determine the differences between individuals with NIAT and healthy controls at baseline. If the data are normally distributed, repeated measures analysis of variance (ANOVA) will be used. Factors of group (ECC and CON) and time (at week 1, 3, and 6) will be used to analyze each variable. Bonferroni post hoc analysis will be used if ANOVA is significant. The partial eta-squared ( $\eta_p^2$ ) for ANOVA will be used to examine the effect size of changes after the training intervention. An  $\eta_p^2$  less than 0.06 will be classified as "small", 0.07-0.14 as "moderate", and greater than 0.14 as "large"<sup>[57]</sup>. If data is not normally distributed, appropriate nonparametric tests will be used.

## Discussion

To our knowledge this is the first study aiming to establish changes in motor unit firing properties of the gastrocnemius-soleus muscles after applying a training protocol based on either controlled eccentric or concentric contractions in individuals with NIAT. Additionally, this study will be the first to determine motor unit firing properties of the

gastrocnemius-soleus muscles in individuals with NIAT compared with asymptomatic controls.

Regarding the variables related to the electromyographic activity and motor unit firing properties (i.e., discharge rate, recruitment and discharge rate variability) of the gastrocnemius-soleus muscles; previous studies have estimated the activation of the triceps surae muscles in people with AT during walking,<sup>[58]</sup> running,<sup>[59]</sup> isometric plantarflexion tasks,<sup>[60]</sup> and dynamic plantarflexion tasks<sup>[61 62]</sup>. Currently, there is no agreement in the literature in terms of plantarflexion torque measured during maximal contractions in individuals with NIAT; some authors did not find any difference between groups,<sup>[14 63]</sup> while others found statistically significant differences<sup>[60 64]</sup>. Interestingly, an investigation has observed a significant increase in LG activation during isometric plantarflexion tasks in people with AT following a 12-week training program<sup>[60]</sup>. Despite all of these efforts, currently there are no studies that have evaluated motor unit firing properties of calf muscles in individuals with NIAT.

A strength of this study is that we will perform a detailed assessment of the mechanical and structural properties of the Achilles tendon and the calf muscles. One study showed that tendinopathy alters both the mechanical and material properties of the human Achilles tendon<sup>[14]</sup>. Morphological comparisons of tendinopathic and healthy tendons demonstrated a larger cross-sectional area for the degenerated Achilles tendon. Typically, a larger tendon is considered mechanically stronger due to its ability to dissipate stresses across the tendon and yield lower strain energy. Nonetheless, in the study of Arya and Kulig<sup>[14]</sup>, they demonstrated that despite having a larger cross-sectional area, the degenerated tendon had lower stiffness and Young's modulus compared with healthy tendons. Additionally, our study includes the use of SWE. This procedure has been used to measure tissue elasticity in tendons and might add to a better understanding of the effects of different types of exercises in tendons<sup>[65]</sup>. Furthermore, SWE is able to measure the Young modulus (slope of the stress-strain curve in the linear region<sup>[66]</sup>) of Achilles tendon with high reliability<sup>[67]</sup>. Previous studies suggest that SWE might be a useful tool for diagnosing and monitoring AT<sup>[68]</sup>. For instance, one study demonstrated that symptomatic Achilles tendons had lower Young modulus compared with healthy tendons

and that stiffness increases in correlation with VISA-A scores after 6 months of treatment<sup>[69]</sup>.

Another strength of our study is using an isokinetic dynamometer to perform the training sessions. This device will allow us to control the intensity (50% MVC), range of motion (0° to 30° of plantarflexion), and angular speed (3°/s) of the contractions, enabling us to have close control over the time under tension (10 seconds) of the Achilles tendon. Although the use of isokinetic dynamometers is common to measure peak torque in musculoskeletal research, its use during a training protocol is limited. To our knowledge, there are no studies using isokinetic dynamometers to train individuals with NIAT. This represents an essential aspect of our RCT because previous studies investigating the effects of exercise in individuals with NIAT usually apply training protocols with insufficient control over the load, speed, pain tolerance, or the range of motion, and this could have influenced their results.

Regarding the study's limitations, the relatively short training protocol (6 weeks) might influence the changes expected in the mechanical and structural properties of the Achilles tendon; therefore, longer training interventions might be required to assess changes in these parameters in the long term. Moreover, due to the nature of the training protocol applied (eccentric and concentric exercises), blinding of participants is not possible because we need to explain how to do the different types of exercises on the isokinetic dynamometer. We are aware this introduces bias into the RCT, but unfortunately, it is not possible to achieve participants' blinding. Another study limitation is the inclusion of participants with bilateral symptoms, which could potentially affect the results. Since morphological changes to the asymptomatic tendon are common in this condition<sup>[70]</sup> and 45% of thickened Achilles tendons progress to develop clinical symptoms within 12 months<sup>[71]</sup>, we decided to also include these patients in the study. Additionally, participants' age range is another limitation of our study, as it might affect the reproducibility of our findings in older populations; however, we have decided to recruit participants in this age range based on previous studies showing age-related differences in Achilles tendon's stiffness and Young's modulus, which could confound the results of the intervention.

The study of motor units is an area in continuous development, which in recent years has allowed a more profound understanding of the neural mechanisms involved in muscle contractions. However, much of the research in this area has focused on the normal neurophysiology of muscle rather than its relationship with alterations of the musculoskeletal system.

This research will therefore provide new insights regarding the neuromechanical effects of ECC and CON exercises in the management of individuals with NIAT. A more precise understanding of the mechanisms involved in this pathology is essential to improve the rehabilitation programs commonly used in the management of this condition.

**Patient and Public Involvement**

The research question in this study forms part of a larger discussion about exercise and pain relief within our patient and public involvement meetings. Patients will not be involved in the analysis and data collection but will contribute to data interpretation and production of a lay summary of findings.

**Ethics and Dissemination**

**Ethical approval and trial registration**

The research protocol has been approved by the Science, Technology, Engineering and Mathematics Ethical Review Committee of the University of Birmingham (ERN\_20-0604A). Additionally, this protocol has been registered on ISRCTN registry (ISRCTN46462385).

Researchers will inform all participants of the characteristics of the research and will obtain written consent. Participants will be informed that they are free to withdraw from the study at any time without needing to provide a reason. In any unlikely adverse events, this will be immediately reported by the principal investigator to the ethics committee.

The results of this study will be submitted for publication in a peer review journal and presented at conferences.



## Author Contributions

ICH and EMV are responsible for the conception, design, and development of the protocol. EMV is the lead supervisor of ICH and DF is the co-supervisor. EMV and DF have provided guidance on methodological decisions and critical revision. All authors have read and subsequently approved the final manuscript.

## Funding Statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

## Competing Interests Statement

The authors have no competing interests to report.

## Data Sharing Statement

The datasets generated during the current study will be available from the corresponding author on reasonable request. Datasets will be available from 01/07/2023 to 31/12/2023 (6 months).

## Confidentiality

All information collected will be kept strictly confidential. Personal information will be retained but only available to the researchers using password protected files. In addition, all data for presentations will be anonymized and aggregated, so the participants' identities will not be revealed in any way.

## Sponsor details

Trial Sponsor: University of Birmingham

Sponsor Reference: grid.6572.6

Address: Edgbaston, Birmingham, United Kingdom, B15 2TT

Telephone: +44 (0)1214143344

Email: [researchgovernance@contacts.bham.ac.uk](mailto:researchgovernance@contacts.bham.ac.uk)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Date and version identifier**

Issue date: 26 June 2022  
Protocol amendment number: 2  
Authors: ICH, EMV, and DF

For peer review only

## References

1. Sobhani S, Dekker R, Postema K, et al. Epidemiology of ankle and foot overuse injuries in sports: A systematic review. *Scand J Med Sci Sports* 2013;23(6):669-86. doi: 10.1111/j.1600-0838.2012.01509.x [published Online First: 2012/08/01]
2. Magnussen RA, Dunn WR, Thomson AB. Nonoperative treatment of midportion Achilles tendinopathy: a systematic review. *Clin J Sport Med* 2009;19(1):54-64. doi: 10.1097/JSM.0b013e31818ef090 [published Online First: 2009/01/07]
3. Janssen I, van der Worp H, Hensing S, et al. Investigating Achilles and patellar tendinopathy prevalence in elite athletics. *Research in Sports Medicine* 2018;26(1):1-12. doi: 10.1080/15438627.2017.1393748
4. Kvist M. Achilles tendon injuries in athletes. *Ann Chir Gynaecol* 1991;80(2):188-201. [published Online First: 1991/01/01]
5. Longo UG, Ronga M, Maffulli N. Achilles tendinopathy. *Sports Med Arthrosc Rev* 2009;17(2):112-26. doi: 10.1097/JSA.0b013e3181a3d625 [published Online First: 2009/05/15]
6. Scott A, Backman LJ, Speed C. Tendinopathy: Update on Pathophysiology. *J Orthop Sports Phys Ther* 2015;45(11):833-41. doi: 10.2519/jospt.2015.5884 [published Online First: 2015/09/22]
7. Maffulli N, Longo UG, Kadakia A, et al. Achilles tendinopathy. *Foot Ankle Surg* 2019 doi: 10.1016/j.fas.2019.03.009 [published Online First: 2019/04/30]
8. Henriksen M, Alkjaer T, Lund H, et al. Experimental quadriceps muscle pain impairs knee joint control during walking. *J Appl Physiol (1985)* 2007;103(1):132-9. doi: 10.1152/jappphysiol.01105.2006 [published Online First: 2007/04/07]
9. Tucker K, Butler J, Graven-Nielsen T, et al. Motor unit recruitment strategies are altered during deep-tissue pain. *J Neurosci* 2009;29(35):10820-6. doi: 10.1523/jneurosci.5211-08.2009 [published Online First: 2009/09/04]
10. Haglund-Akerlind Y, Eriksson E. Range of motion, muscle torque and training habits in runners with and without Achilles tendon problems. *Knee Surg Sports Traumatol Arthrosc* 1993;1(3-4):195-9. doi: 10.1007/bf01560205 [published Online First: 1993/01/01]
11. Mahieu NN, Witvrouw E, Stevens V, et al. Intrinsic risk factors for the development of achilles tendon overuse injury: a prospective study. *Am J Sports Med* 2006;34(2):226-35. doi: 10.1177/0363546505279918 [published Online First: 2005/11/02]
12. Henriksen M, Aaboe J, Graven-Nielsen T, et al. Motor responses to experimental Achilles tendon pain. *Br J Sports Med* 2011;45(5):393-8. doi: 10.1136/bjsm.2010.072561 [published Online First: 2010/06/15]
13. Wyndow N, Cowan SM, Wrigley TV, et al. Neuromotor control of the lower limb in Achilles tendinopathy: implications for foot orthotic therapy. *Sports Med* 2010;40(9):715-27. doi: 10.2165/11535920-000000000-00000 [published Online First: 2010/08/24]
14. Arya S, Kulig K. Tendinopathy alters mechanical and material properties of the Achilles tendon. *J Appl Physiol (1985)* 2010;108(3):670-5. doi: 10.1152/jappphysiol.00259.2009 [published Online First: 2009/11/07]
15. Martinez-Valdes E, Negro F, Falla D, et al. Surface electromyographic amplitude does not identify differences in neural drive to synergistic muscles. *J Appl Physiol (1985)* 2018;124(4):1071-79. doi: 10.1152/jappphysiol.01115.2017 [published Online First: 2018/02/09]
16. Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG. *J Appl Physiol (1985)* 2004;96(4):1486-95. doi: 10.1152/jappphysiol.01070.2003 [published Online First: 2004/03/16]

17. Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG: an update. *J Appl Physiol (1985)* 2014;117(11):1215-30. doi: 10.1152/japplphysiol.00162.2014 [published Online First: 2014/10/04]

18. Farina D, Negro F, Muceli S, et al. Principles of Motor Unit Physiology Evolve With Advances in Technology. *Physiology* 2016;31(2):83-94. doi: 10.1152/physiol.00040.2015

19. Alfredson H, Pietila T, Jonsson P, et al. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med* 1998;26(3):360-6. doi: 10.1177/03635465980260030301 [published Online First: 1998/06/09]

20. Maffulli N, Walley G, Sayana MK, et al. Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil* 2008;30(20-22):1677-84. doi: 10.1080/09638280701786427 [published Online First: 2008/07/09]

21. Sayana MK, Maffulli N. Eccentric calf muscle training in non-athletic patients with Achilles tendinopathy. *J Sci Med Sport* 2007;10(1):52-8. doi: 10.1016/j.jsams.2006.05.008 [published Online First: 2006/07/11]

22. Mayer F, Hirschmüller A, Müller S, et al. Effects of short-term treatment strategies over 4 weeks in Achilles tendinopathy. *Br J Sports Med* 2007;41(7):e6. doi: 10.1136/bjsm.2006.031732 [published Online First: 2007/01/31]

23. Jarin I, Backer HC, Vosseller JT. Noninsertional Achilles Tendinopathy: Topical Review. *Foot Ankle Int* 2020;1071100720914605. doi: 10.1177/1071100720914605 [published Online First: 2020/05/01]

24. Mafi N, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 2001;9(1):42-7. doi: 10.1007/s001670000148 [published Online First: 2001/03/28]

25. Yu J, Park D, Lee G. Effect of eccentric strengthening on pain, muscle strength, endurance, and functional fitness factors in male patients with achilles tendinopathy. *American journal of physical medicine & rehabilitation* 2013;92(1):68-76. doi: 10.1097/phm.0b013e31826eda63

26. Lindemann I, Coombes BK, Tucker K, et al. Age-related differences in gastrocnemii muscles and Achilles tendon mechanical properties in vivo. *J Biomech* 2020;112:110067. doi: 10.1016/j.jbiomech.2020.110067 [published Online First: 2020/10/10]

27. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med* 2001;35(5):335-41. doi: 10.1136/bjsm.35.5.335 [published Online First: 2001/10/02]

28. Alghadir AH, Anwer S, Iqbal A, et al. Test-retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal rating scales for measurement of osteoarthritic knee pain. *J Pain Res* 2018;11:851-56. doi: 10.2147/JPR.S158847

29. Beyer R, Kongsgaard M, Hougs Kjær B, et al. Heavy Slow Resistance Versus Eccentric Training as Treatment for Achilles Tendinopathy: A Randomized Controlled Trial. *Am J Sports Med* 2015;43(7):1704-11. doi: 10.1177/0363546515584760 [published Online First: 2015/05/29]

30. Iversen JV, Bartels EM, Langberg H. The victorian institute of sports assessment - achilles questionnaire (visa-a) - a reliable tool for measuring achilles tendinopathy. *Int J Sports Phys Ther* 2012;7(1):76-84.

31. Rompe JD, Furia J, Maffulli N. Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. *Am J Sports Med* 2009;37(3):463-70. doi: 10.1177/0363546508326983 [published Online First: 2008/12/18]

32. Chan A-W, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158(3):200-07. doi: 10.7326/0003-4819-158-3-201302050-00583

33. Chapman JP, Chapman LJ, Allen JJ. The measurement of foot preference. *Neuropsychologia* 1987;25(3):579-84. doi: 10.1016/0028-3932(87)90082-0 [published Online First: 1987/01/01]
34. Faul F, Erdfelder E, Buchner A, et al. Statistical power analyses using G\*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009;41(4):1149-60. doi: 10.3758/brm.41.4.1149 [published Online First: 2009/11/10]
35. Craig CL, Marshall AL, Sjoström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35(8):1381-95. doi: 10.1249/01.Mss.0000078924.61453.Fb [published Online First: 2003/08/06]
36. Martin RL, Irrgang JJ, Burdett RG, et al. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int* 2005;26(11):968-83. doi: 10.1177/107110070502601113 [published Online First: 2005/11/29]
37. Sullivan MJL, Bishop S, Pivik J. The Pain Catastrophizing Scale: Development and Validation. *Psychological Assessment* 1996;7:524-32. doi: 10.1037//1040-3590.7.4.524
38. Miller RP, Kori SH, Todd DD. The Tampa Scale: a Measure of Kinesophobia. *The Clinical Journal of Pain* 1991;7(1):51.
39. Del Vecchio A, Negro F, Holobar A, et al. You are as fast as your motor neurons: speed of recruitment and maximal discharge of motor neurons determine the maximal rate of force development in humans. *J Physiol* 2019;597(9):2445-56. doi: 10.1113/jp277396 [published Online First: 2019/02/16]
40. Chen MJ, Fan X, Moe ST. Criterion-related validity of the Borg ratings of perceived exertion scale in healthy individuals: a meta-analysis. *Journal of Sports Sciences* 2002;20(11):873-99. doi: 10.1080/026404102320761787
41. van Poppel MN, Chinapaw MJ, Mokkink LB, et al. Physical activity questionnaires for adults: a systematic review of measurement properties. *Sports Med* 2010;40(7):565-600. doi: 10.2165/11531930-000000000-00000 [published Online First: 2010/06/16]
42. Vlaeyen JW, Kole-Snijders AM, Boeren RG, et al. Fear of movement/(re)injury in chronic low back pain and its relation to behavioral performance. *Pain* 1995;62(3):363-72. doi: 10.1016/0304-3959(94)00279-n [published Online First: 1995/09/01]
43. Goubert L, Crombez G, Van Damme S, et al. Confirmatory factor analysis of the Tampa Scale for Kinesophobia: invariant two-factor model across low back pain patients and fibromyalgia patients. *Clin J Pain* 2004;20(2):103-10. doi: 10.1097/00002508-200403000-00007 [published Online First: 2004/02/11]
44. Roelofs J, Goubert L, Peters ML, et al. The Tampa Scale for Kinesophobia: further examination of psychometric properties in patients with chronic low back pain and fibromyalgia. *Eur J Pain* 2004;8(5):495-502. doi: 10.1016/j.ejpain.2003.11.016 [published Online First: 2004/08/25]
45. Swinkels-Meewisse EJ, Swinkels RA, Verbeek AL, et al. Psychometric properties of the Tampa Scale for kinesophobia and the fear-avoidance beliefs questionnaire in acute low back pain. *Man Ther* 2003;8(1):29-36. doi: 10.1054/math.2002.0484 [published Online First: 2003/02/15]
46. Swinkels-Meewisse IE, Roelofs J, Verbeek AL, et al. Fear of movement/(re)injury, disability and participation in acute low back pain. *Pain* 2003;105(1-2):371-9. doi: 10.1016/s0304-3959(03)00255-0 [published Online First: 2003/09/23]
47. Heuts PH, Vlaeyen JW, Roelofs J, et al. Pain-related fear and daily functioning in patients with osteoarthritis. *Pain* 2004;110(1-2):228-35. doi: 10.1016/j.pain.2004.03.035 [published Online First: 2004/07/28]
48. Bolsterlee B, Gandevia SC, Herbert RD. Effect of Transducer Orientation on Errors in Ultrasound Image-Based Measurements of Human Medial Gastrocnemius Muscle Fascicle Length and Pennation. *PLoS One* 2016;11(6):e0157273. doi: 10.1371/journal.pone.0157273 [published Online First: 2016/06/15]



49. Siu W-I, Chan C-h, Lam C-h, et al. Sonographic evaluation of the effect of long-term exercise on Achilles tendon stiffness using shear wave elastography. *Journal of Science & Medicine in Sport* 2016;19(11):883-87.

50. Coombes BK, Tucker K, Vicenzino B, et al. Achilles and patellar tendinopathy display opposite changes in elastic properties: A shear wave elastography study. *Scand J Med Sci Sports* 2018;28(3):1201-08. doi: 10.1111/sms.12986 [published Online First: 2017/10/04]

51. Arvanitidis M, Falla D, Martinez-Valdes E. Can visual feedback on upper trapezius high-density surface electromyography increase time to task failure of an endurance task? *J Electromyogr Kinesiol* 2019;49:102361. doi: 10.1016/j.jelekin.2019.102361 [published Online First: 2019/10/13]

52. Martinez-Valdes E, Negro F, Farina D, et al. Divergent response of low- versus high-threshold motor units to experimental muscle pain. *J Physiol* 2020;598(11):2093-108. doi: 10.1113/jp279225 [published Online First: 2020/03/19]

53. Negro F, Muceli S, Castronovo AM, et al. Multi-channel intramuscular and surface EMG decomposition by convolutive blind source separation. *J Neural Eng* 2016;13(2):026027. doi: 10.1088/1741-2560/13/2/026027 [published Online First: 2016/03/01]

54. Martinez-Valdes E, Laine CM, Falla D, et al. High-density surface electromyography provides reliable estimates of motor unit behavior. *Clin Neurophysiol* 2016;127(6):2534-41. doi: 10.1016/j.clinph.2015.10.065 [published Online First: 2016/01/19]

55. Boccia G, Martinez-Valdes E, Negro F, et al. Motor unit discharge rate and the estimated synaptic input to the vasti muscles is higher in open compared with closed kinetic chain exercise. *J Appl Physiol (1985)* 2019;127(4):950-58. doi: 10.1152/jappphysiol.00310.2019 [published Online First: 2019/08/02]

56. Martinez-Valdes E, Negro F, Laine CM, et al. Tracking motor units longitudinally across experimental sessions with high-density surface electromyography. *J Physiol* 2017;595(5):1479-96. doi: 10.1113/jp273662 [published Online First: 2016/12/30]

57. Cohen J. Statistical Power Analysis for the Behavioral Sciences. 2nd ed. Hillsdale (NJ) ed: L. Erlbaum Associates 1988:p.567.

58. Baur H, Divert C, Hirschmüller A, et al. Analysis of gait differences in healthy runners and runners with chronic Achilles tendon complaints. *Isokinetics and Exercise Science* 2004;12:111-16. doi: 10.3233/IES-2004-0161

59. Wyndow N, Cowan SM, Wrigley TV, et al. Triceps surae activation is altered in male runners with Achilles tendinopathy. *J Electromyogr Kinesiol* 2013;23(1):166-72. doi: 10.1016/j.jelekin.2012.08.010 [published Online First: 2012/11/13]

60. Masood T, Kalliokoski K, Bojsen-Moller J, et al. Plantarflexor muscle function in healthy and chronic Achilles tendon pain subjects evaluated by the use of EMG and PET imaging. *Clin Biomech (Bristol, Avon)* 2014;29(5):564-70. doi: 10.1016/j.clinbiomech.2014.03.003 [published Online First: 2014/04/10]

61. Reid D, McNair PJ, Johnson S, et al. Electromyographic analysis of an eccentric calf muscle exercise in persons with and without Achilles tendinopathy. *Phys Ther Sport* 2012;13(3):150-5. doi: 10.1016/j.ptsp.2011.08.003 [published Online First: 2012/07/21]

62. Yu J. Comparison of Lower Limb Muscle Activity during Eccentric and Concentric Exercises in Runners with Achilles Tendinopathy. *J Phys Ther Sci* 2014;26(9):1351-3. doi: 10.1589/jpts.26.1351 [published Online First: 2014/10/03]

63. Child S, Bryant AL, Clark RA, et al. Mechanical properties of the achilles tendon aponeurosis are altered in athletes with achilles tendinopathy. *Am J Sports Med* 2010;38(9):1885-93. doi: 10.1177/0363546510366234 [published Online First: 2010/05/29]



64. O'Neill S, Barry S, Watson P. Plantarflexor strength and endurance deficits associated with mid-portion Achilles tendinopathy: The role of soleus. *Phys Ther Sport* 2019;37:69-76. doi: 10.1016/j.ptsp.2019.03.002 [published Online First: 2019/03/19]
65. Washburn N, Onishi K, Wang J. Ultrasound elastography and ultrasound tissue characterisation for tendon evaluation. *Journal of Orthopaedic Translation* 2018;15 doi: 10.1016/j.jot.2018.06.003
66. Maganaris CN, Narici MV, Maffulli N. Biomechanics of the Achilles tendon. *Disabil Rehabil* 2008;30(20-22):1542-7. doi: 10.1080/09638280701785494 [published Online First: 2008/08/23]
67. Leung W, Chu K, Lai C. Sonographic evaluation of the immediate effects of eccentric heel drop exercise on Achilles tendon and gastrocnemius muscle stiffness using shear wave elastography. *PeerJ* 2017;5 doi: 10.7717/peerj.3592
68. Dirrichs T, Quack V, Gatz M, et al. Shear Wave Elastography (SWE) for the Evaluation of Patients with Tendinopathies. *Acad Radiol* 2016;23(10):1204-13. doi: 10.1016/j.acra.2016.05.012 [published Online First: 2016/06/20]
69. Dirrichs T, Quack V, Gatz M, et al. Shear Wave Elastography (SWE) for Monitoring of Treatment of Tendinopathies: A Double-blinded, Longitudinal Clinical Study. *Acad Radiol* 2018;25(3):265-72. doi: 10.1016/j.acra.2017.09.011 [published Online First: 2017/11/21]
70. Grigg NL, Wearing SC, Smeathers JE. Achilles tendinopathy has an aberrant strain response to eccentric exercise. *Med Sci Sports Exerc* 2012;44(1):12-7. doi: 10.1249/MSS.0b013e318227fa8c [published Online First: 2011/06/11]
71. Fredberg U, Bolvig L. Significance of ultrasonographically detected asymptomatic tendinosis in the patellar and achilles tendons of elite soccer players: a longitudinal study. *Am J Sports Med* 2002;30(4):488-91. doi: 10.1177/03635465020300040701 [published Online First: 2002/07/20]




Figure 1. Overview of the study design.

338x190mm (300 x 300 DPI)




Figure 2. Experimental session design.

338x190mm (300 x 300 DPI)




Figure 3. A) Anatomical landmarks used for ultrasonography and B) position of the electrodes in the MG, LG and SO muscles.

338x190mm (300 x 300 DPI)

# SPIRIT 2013 Checklist: recommended items to address in a clinical trial protocol and related documents\*

Section and topic	Items N°	Checklist item	Page N°
<b>Administrative information</b>			
Title	1	Descriptive title identifying the study design, population, interventions and, if applicable, trial acronym	1
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	2
	2b	All items from the World Health Organization Trial Registration Data Set	Supplementary file 2
Protocol version	3	Date and version identifier	23
Funding	4	Sources and types of financial, material, and other support	22
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	1,22
	5b	Name and contact information for the trial sponsor	22
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities.	N/A
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see item 21a for data monitoring committee)	N/A
<b>Introduction</b>			
Background and rationale	6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention.	4-6
	6b	Explanation for choice of comparators	5
Objectives	7	Specific objectives or hypothesis	6
Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	7
<b>Methods: Participants, Interventions, and outcomes</b>			
Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	7
Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists)	6-7
Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	10-11
	11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease)	17
	11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests)	N/A
	11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	N/A

Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	11-12
Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see fig 1)	Supplementary file 3
Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculation	7-8
Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	6
<b>Methods: Assignment of interventions (for controlled trials)</b>			
Allocation:			
Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	17
Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	17
Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	17-18
Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts) and how	18
	17b	If blinded, circumstances under which unblinding is permissible and procedure for revealing a participant's allocated intervention during the trial	N/A
<b>Methods: Data collection, management, and analysis</b>			
Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	12-17
	18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	N/A
Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	18
Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	18
	20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	18
	20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	N/A



<b>Methods: Monitoring</b>			
Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	N/A
	21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	N/A
Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneous reported adverse events and other unintended effects of trial interventions or trial conduct	17
Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	N/A
<b>Ethics and dissemination</b>			
Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	N/A
Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial regulators, journals, regulators)	N/A
Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	6
	26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	N/A
Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	22
Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	22
Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	N/A
Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	N/A
Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	21
	31b	Authorship eligibility guidelines and any intended use of professional writers	N/A
	31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	22
<b>Appendices</b>			
Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	Supplementary files 4 and 5
Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	N/A



## Trial Registration data

Data category	Information
Primary registry and trial identifying number	ISRCTN registry (ISRCTN46462385).
Date of registration in primary registry	13/08/2021
Secondary identifying numbers	N/A
Source(s) of monetary or material support	This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors
Primary sponsor	University of Birmingham
Secondary sponsor(s)	N/A
Contact for public queries	Ignacio Contreras-Hernandez (iac921@student.bham.ac.uk)
Contact for scientific queries	Eduardo Martinez-Valdes (E.A.MartinezValdes@bham.ac.uk)
Public title	Effects of two different exercise protocols in individuals with non-insertional Achilles tendinopathy
Scientific title	Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy: Protocol for a randomised controlled trial.
Countries of recruitment	United Kingdom
Health condition(s) or problem(s) studied	Non-insertional Achilles tendinopathy
Intervention(s)	Eccentric plantarflexion contractions Concentric plantarflexion contractions
Key inclusion and exclusion criteria	Men or women aged 18 to 55 years old. Inclusion criteria are non-insertional Achilles tendinopathy determined by an experienced physiotherapist based on defined clinical findings, physical examination, and ultrasound assessment, as well as having pain for at least 3 months.  The exclusion criteria for healthy participants and individuals with non-insertional Achilles tendinopathy include: (1) systemic or inflammatory conditions including rheumatic, neuromuscular disorders, and malignancy, (2) current or history of chronic respiratory, neurological, or cardiovascular diseases, (3) history of lower limb surgery. Specific exclusion criteria for the participants with non-insertional Achilles tendinopathy are participation in any other treatment or rehabilitation program for Achilles tendinopathy, corticosteroid injections in the previous 12 months, and insertional Achilles tendinopathy.
Study type	Two-arm, parallel group, randomised controlled trial.
Date of first enrolment	04/10/2021
Target sample size	A total of 26 individuals with non-insertional Achilles tendinopathy and 13 healthy controls
Recruitment status	Recruiting

\* This template is copyrighted by the SPIRIT Group and is reproduced by BMJ with their permission.

From: Chan A, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin JA, Dickersin K, Hróbjartsson A, Schulz KF, Vandenbroucke JP, Vandenbroucke WR, Krleža-Jeric K, Laupacis A, Moher D. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ* 2013;346:f7586

	STUDY PERIOD																		
	Enrolment	Allocation	Post-allocation (6-weeks period)															Close-out	
Timepoint	-t1	0	t1	t2	t3	t4	t5	t6	t7	t8	t9	t10	t11	t12	t13	t14	t15	t16 (follow-up at 3 months)	t17 (follow-up at 6 months)
ENROLMENT:																			
Eligibility screen	x																		
Informed consent	x																		
Allocation		x																	
INTERVENTIONS:																			
Eccentric training				x	x	x	x	x	x		x	x	x	x					
Concentric training				x	x	x	x	x	x		x	x	x	x					
ASSESSMENTS:																			
Anthropometric data			x																
NRS			x							x							x	x	x
VISA-A			x							x							x	x	x
IPAQ-SF			x							x							x		
FAAM			x							x							x		
PCS			x							x							x		
TSK			x							x							x		
Motor unit firing properties			x							x							x		
Morphological and mechanical properties			x							x							x		

\* Schedule of enrolment, interventions, and assessments considering two training sessions per week. This template is copyrighted by the SPIRIT Group and is reproduced by BMJ with their permission.

From: Chan A, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin JA, Dickersin K, Hróbjartsson A, Schulz KF, Parulekar WR, Krleža-Jeric K, Laupacis A, Moher D. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. BMJ 2013;346:e7586.

# Consent Form

Study Title	Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy		
Participant Name:		Date:	
Researcher Name:		Ethics Number:	

This information is being collected as part of a research project, which investigates the relationship between the neuromuscular control of the calf muscles and some characteristics of these muscles and the Achilles tendon in individuals with and without Achilles tendinopathy.

The research will be conducted at the School of Sport, Exercise and Rehabilitation Sciences at the University of Birmingham. The information that you supply and that which may be collected as part of the research project will be entered into a filing system or database and will only be accessed by authorised personnel involved in the project. The information will be retained by the University of Birmingham and will only be used for the purpose of research, and statistical and audit purposes. By supplying this information, you are consenting to the University storing your information for the purposes stated above. The information will be processed by the University of Birmingham in accordance with the provisions of the Data Protection Act 2018. No identifiable personal data will be published.

**This section to be completed by the participant:**  
**Please initial the boxes at the end of each statement if you agree with it.**

1. I confirm that I have read and understood the Participant Information Sheet for the above study. I have had the opportunity to ask questions and these have all been answered satisfactorily

☐

2. I understand that my participation is voluntary and that I am free to withdraw at any time, without giving any reason, up to two weeks after my last visit to the lab.

☐

3. I agree to the storage and use of my data for the purposes of this research study.

☐

4. I confirm that I have read and understand the paragraph relating to COVID-19 related risks in the participant information leaflet for this study and will abide by the measures put in place by the University of Birmingham. I have had the opportunity to ask questions if necessary and have had these answered satisfactorily.

☐

5. Based on the above, I agree to take part in this research study.

☐

Signed:

Name in capitals:

Date:

**This section to be completed by the researcher**  
I certify that this participant has read, properly completed and signed the screening and consent forms, witnessed by myself:

Signed:

Date:

*By supplying this information you are consenting to the University storing your information for the purposes of the stated research study. The information will be processed by the University of Birmingham in accordance with the provisions of the Data Protection Act 2018. No identifiable personal data will be published*



# Consent Form

<b>Study Title</b>	Neuromuscular and structural tendon adaptations after 6-weeks of either concentric or eccentric exercise in individuals with non-insertional Achilles tendinopathy		
<b>Participant Name:</b>		<b>Date:</b>	
<b>Researcher Name:</b>		<b>Ethics Number:</b>	

This information is being collected as part of a research project, which investigates the relationship between the neuromuscular control of the calf muscles and some characteristics of these muscles and the Achilles tendon in individuals with Achilles tendinopathy. Additionally, we want to determine which type of exercise protocol has better results in the treatment of this condition.

The research will be conducted at the School of Sport, Exercise and Rehabilitation Sciences at the University of Birmingham. The information that you supply and that which may be collected as part of the research project will be entered into a filing system or database and will only be accessed by authorised personnel involved in the project. The information will be retained by the University of Birmingham and will only be used for the purpose of research, and statistical and audit purposes. By supplying this information, you are consenting to the University storing your information for the purposes stated above. The information will be processed by the University of Birmingham in accordance with the provisions of the Data Protection Act 2018. No identifiable personal data will be published.

## This section to be completed by the participant:

Please initial the boxes at the end of each statement if you agree with it.

1. I confirm that I have read and understood the Participant Information Sheet for the above study. I have had the opportunity to ask questions and these have all been answered satisfactorily	<input type="checkbox"/>
2. I understand that my participation is voluntary and that I am free to withdraw at any time, without giving any reason, up to two weeks after my last visit to the lab.	<input type="checkbox"/>
3. I agree to the storage and use of my data for the purposes of this research study.	<input type="checkbox"/>
4. I confirm that I have read and understand the paragraph relating to COVID-19 related risks in the participant information leaflet for this study and will abide by the measures put in place by the University of Birmingham. I have had the opportunity to ask questions if necessary and have had these answered satisfactorily.	<input type="checkbox"/>
5. Based on the above, I agree to take part in this research study.	<input type="checkbox"/>
Signed:	
Name in capitals:	
Date:	

## This section to be completed by the researcher

I certify that this participant has read, properly completed and signed the screening and consent forms, witnessed by myself:

Signed:

Date:

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*By supplying this information, you are consenting to the University storing your information for the purposes of the stated research study. The information will be processed by the University of Birmingham in accordance with the provisions of the Data Protection Act 2018. No identifiable personal data will be published*

For peer review only

Enseignement Supérieur (ABES) :  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

## CONSORT 2010 checklist of information to include when reporting a randomised trial\*

Section and topic	Items N°	Checklist item	Page N°
<b>Title and abstract</b>			
	1a	Identification as a randomised trial in the title	1
	1b	Structured summary of trial design, methods, results, and conclusions	2
<b>Introduction</b>			
Background and objectives	2a	Scientific background and explanation rationale	4-6
	2b	Specific objectives or hypotheses	6
<b>Methods</b>			
Trial design	3a	Description of trial design (such as parallel, factorial) including allocation ratio	7
	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	N/A
Participants	4a	Eligibility criteria for participants	6, 7
	4b	Settings and locations where the data were collected	7
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were administered	10,11
Outcomes	6a	Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed	11,12
	6b	Any changes to trial outcomes after the trial commenced, with reasons	N/A
Sample size	7a	How sample size was determined	7,8
	7b	When applicable, explanation of any interim analyses and stopping guidelines	N/A
Randomisation:			
Sequence generation	8a	Method used to generate the random allocation sequence	17
	8b	Type of randomisation; details of any restriction (such as blocking and block size)	17
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	17
Implementation	10	Who generated the random allocation sequence, who enrolled the participants, and who assigned participants to interventions	17, 18
Blinding	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how	17, 18
	11b	If relevant, description of the similarity of interventions	N/A
Statistical methods	12a	Statistical methods used to compare groups for primary and secondary outcomes	18
	12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	N/A

<b>Results</b>			
Participant flow (a diagram is strongly recommended)	13a	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome	N/A
	13b	For each group, losses and exclusions after randomisation, together with reasons	N/A
Recruitment	14a	Dates defining the periods of recruitment and follow-up	N/A
	14b	Why the trial ended or was stopped	N/A
Baseline data	15	A table showing baseline demographic and clinical characteristics for each group	N/A
Numbers analysed	16	For each group, number of participants (denominator) included in each analysis and whether analysis was by original assigned groups	N/A
Outcomes and estimation	17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)	N/A
	17b	For binary outcomes, presentation of both absolute and relative effect size is recommended	N/A
Ancillary analysis	18	Results of any other analyses performed, including subgroup analyses and adjusted analyses distinguishing pre-specified from exploratory	N/A
Harms	19	All important harms or unintended effects in each group	N/A
<b>Discussion</b>			
Limitations	20	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant multiplicity of analyses	18-21
Generalisability	21	Generalisability (external validity, applicability) of the trial findings	18-21
Interpretation	22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	18-21
<b>Other information</b>			
Registration	23	Registration number and name of trial registry	2, 7, 21
Protocol	24	Where the full trial protocol can be accessed, if available	N/A
Funding	25	Sources of funding and other support (such as supply of drugs), role of funders	22

\*We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration<sup>13</sup> for important clarifications on all the items. If relevant, we also recommend reading CONSORT extensions for cluster randomised trials,<sup>11</sup> non-inferiority and equivalence trials,<sup>12</sup> non-pharmacological treatments,<sup>32</sup> herbal interventions,<sup>33</sup> and pragmatic trials.<sup>34</sup> Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see [www.consort-statement.org](http://www.consort-statement.org).

From: Schulz K, Altman D, Moher D. CONSORT 2010 Statement: updated guideline for reporting parallel group randomised trials. BMJ 2010; 340:c 332