



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

A Longitudinal Observational Study Protocol: Preterm Infants: Microbiome Establishment, Neuro-CrossTalk and Origins (PIMENTO)

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-075060
Article Type:	Protocol
Date Submitted by the Author:	25-Apr-2023
Complete List of Authors:	Healy, David; University College Cork, Department of Paediatrics and Child Health Wang, Shuo; University College Cork Grimaud, Ghjuvan; Moorepark Food Research Centre Warda, Alicja; Teagasc Food Research Centre Moorepark Ross, Paul; University College Cork, APC Microbiome Ireland Stanton, Catherine; University College Cork APC Microbiome Institute Dempsey, E; University College Cork, Department of Paediatrics and Child Health
Keywords:	NEONATOLOGY, Microbiology < NATURAL SCIENCE DISCIPLINES, Neonatal intensive & critical care < INTENSIVE & CRITICAL CARE

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A Longitudinal Observational Study Protocol: Preterm Infants: Microbiome Establishment, Neuro-CrossTalk and Origins (PIMENTO)

Authors: David B. Healy ^{1,2}, Shuo Wang ¹, Ghjuvan Grimaud ³, Alicja K. Warda ^{1,3}, R. Paul Ross ^{1,3}, Catherine Stanton ^{1,2,4}, and Eugene M. Dempsey ^{1,2,4,5}

¹ APC Microbiome Ireland, Cork, T12 YT20, Ireland

² Department of Paediatrics and Child Health, University College Cork, T12 YT20, Ireland

³ Teagasc, Moorepark Food Research Centre, Fermoy, P61 C996, Ireland

⁴ University College Cork, Cork, T12 YT20, Ireland

⁵ INFANT Research Centre, University College Cork, T12 DC4A, Ireland

Journal: BMJOpen

Corresponding Author: David B. Healy, APC Microbiome Ireland; Department of Paediatrics and Child Health, david.healy@ucc.ie

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Abstract

Introduction

Very preterm infants are at risk of abnormal microbiome colonisation in the first weeks to months of life. Several important associated factors have been identified including gestational age, mode of delivery, antibiotic exposure and feeding. Preterm infants are at risk of a number of pathologies for which the microbiome may play a central role, including necrotising enterocolitis (NEC) and sepsis. The objective of this study is to determine detailed microbiome changes that occur around implementation of different management practices including empiric antibiotic use, advancement of feeds and administration of probiotics during admission to the neonatal intensive care unit.

Methods and analysis

A single-site, longitudinal observational study of infants born less than 32 weeks gestation, including collection of maternal samples around delivery and breastmilk and infant samples from admission through discharge from the neonatal unit.

Ethics and dissemination

The protocol was approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals (CREC). The trial has been registered at ClinicalTrials.gov (NCT05803577).

The findings from this study will be disseminated in peer-reviewed journals, during scientific conferences, and directly to the study participants. Sequencing data will be deposited in public databases.

Strengths and limitations of this study

- Study design will provide high resolution longitudinal microbiome data that will help to characterise the colonising microbiome of preterm infants, the effect of and progression after treatments.
- An array of samples will be collected including rectal swabs, stool, urine and blood from infants, peripartum maternal samples including stool, oral, skin, vaginal samples and expressed breastmilk from mothers.
- This is a single-site study.
- The study population is heterogenous, and the recruited sample size will be ~200 (mothers and infants).

Abbreviations:

CREC	Clinical Research Ethics Committee of the Cork Teaching Hospitals
CRF	Case report form
CUMH	Cork University Maternity Hospital
GC-MS	Gas chromatography-mass spectrometry
GDPR	General Data Protection Regulation
HSE	Health Service Executive
IMP	Investigational medicinal product
LC-MS	Liquid chromatography-mass spectrometry
NEC	Necrotising enterocolitis
NICU	Neonatal intensive care unit
PCR	Polymerase chain reaction
RBB+C	Repeated bead beating plus column
TLR	Toll-like receptor

Keywords: preterm infants, microbiome, probiotics

1. Introduction

Over the last two decades, the materno-feto-neonatal microbiome has rapidly become an area of budding interest in the specialty of neonatology. While there is much debate as to whether a developing fetus is exposed to microorganisms in utero (1), it is increasingly evident that the gut microbiome does not expand and diversify until the postnatal period at which point microbes from the mother and environment begin colonisation of the infant. Following birth, an initial relatively simple microbial community develops into a diverse microbial ecosystem in a controlled fashion over the first years of life (2). Bacterial colonisation, specifically in very low birth weight infants, may occur in an aberrant fashion for numerous reasons: an immature gut and immune system (3, 4); the environment in which they are nursed(5); frequent exposure to antimicrobial agents(6, 7, 8); and altered feeding practices such as delayed commencement of feeds or using formula in place of mother's own milk(9, 10, 11). It is thought that the colonising microbiome of very preterm infants, and the factors that shape it, play a crucial role in the development of microbiome-mediated complications such as NEC.

NEC is a complex multifactorial condition characterised by intestinal immaturity and dysbiosis in association with altered microvascular blood flow. It is the most commonly acquired serious gastrointestinal complication of prematurity and is rare in babies born above 32 weeks (12). Infants at risk are likely to have deficient intestinal mucus production, immature gut immunity, and reduced endogenously-produced antimicrobial factors. The ability of bacteria to adhere to epithelial cells is thereby enhanced, particularly after gut injury secondary to ischaemia/hypoperfusion. Although not fully understood, the most accepted theory for the pathogenesis of NEC is that in this setting, bacterial endotoxin binds to toll-like receptor 4 (TLR4) found on the intestinal epithelial cells and by so doing, pathogen-associated molecular pattern receptors are activated, resulting in cell apoptosis and epithelial injury (13). This leads to an intense inflammatory cascade in the exposed lamina propria, including activation of inflammatory cytokines such as tumour necrosis factor alpha, interleukin-1 and interleukin-8, and ultimately necrosis of the surrounding tissue. Work by Stewart et al.(14) demonstrated that NEC tends not to occur in infants with a diverse gut microbiome. This may relate to inherent factors that make the individual less susceptible to NEC and simultaneously allow a diverse microbiome or may be due to protective effects associated with having a diverse gut microbiome, such as inflammatory modulation and improved gut epithelial integrity.

With early descriptions of the "ideal" gut microbiome (i.e. that of the vaginally delivered, exclusively breast-fed term infant, and without antibiotic treatment)(15, 16) and the identification of potentially deleterious bacteria harboured in the guts of the hospitalised very preterm infants (15, 17, 18) came the advent of probiotics. Administration of these as a supplement to feeding aimed to lend some "normality" to the gut microbiome of these vulnerable babies by supplanting potentially pathogenic bacteria and attempting to reduce the incidence of NEC (19, 20, 21, 22, 23). While probiotic preparations are widely available, shown to be safe (24, 25) and do, indeed, increase so-called beneficial bacteria present in the

gut(26), the evidence for their benefit is contentious (27) and practice varies from institution to institution.

Work has shown that the full term vaginally delivered and exclusively breastfed baby’s initial gut microbiome is dominated primarily by Actinobacteria (28, 29) while studies looking at preterm infants’ microbiome have shown a Proteobacteria (Bacilli dominance) and Firmicutes predominance (28, 30, 31, 32). Clarification of the initial neonatal gut microbiome in the preterm population is confounded by the low bacterial biomass in early samples and high interindividual variability in composition. Individual medical management of each infant is likely to cause perturbations to the microbiome. Several factors associated with alterations in the neonatal gut microbiome have previously been identified. These include gestational age (28) mode of delivery(28, 33), exposure to antibiotics (both pre- and postnatally) (6) and, importantly, feeding choice (34). Mothers of very low birth weight infants are typically encouraged to express breastmilk for their baby. This is associated with lower rates of sepsis and NEC (35). Some of these effects may be related to the bacteria or supporting molecules received in breastmilk from their mother (36). Correlations between changes in the neonatal gut microbiome and the microbiota of the milk they are being fed are limited and longitudinal analysis of breastmilk in this regard is lacking. Furthermore, interactions within breastmilk between bacteria and bacteriophage are not well characterised.

This work will aim to longitudinally characterise the early and colonising intestinal microbiome of very preterm infants. We will analyse samples from infants less than 32 weeks gestational age at birth, with collection occurring longitudinally from birth to discharge from the neonatal intensive care unit (NICU). Microbiome composition will be evaluated using metagenomics and 16sRNA sequencing of stool. Bacterial function will be assessed longitudinally using untargeted metabolomics of infant urine. We will also interrogate some key perinatal maternal microbiome niches, including oral, vaginal, skin and intestinal, as well as expressed breastmilk to evaluate mother-infant transmission.

2. Methods

2.1. Study design

This single site study is a prospective longitudinal observational study of the intestinal microbiome of infants born between 23 and 32 weeks of gestation. It will investigate the establishment of the very premature infants’ microbiome using culture independent approaches, including the potential initial colonisation process of vertical transmission from mother to preterm infant, the apparent differences in establishment pattern based on feeding type, and the perturbations and recovery associated with different intensive care management strategies such as empiric antibiotic usage.

Study recruitment commenced in April 2020 and finished in December 2022. The primary endpoint of the study will be reached after the last enrolled participant completes 2-year neurodevelopmental follow-up.

2.2. Participant selection

Participants are infants, born between 23+0 weeks gestation and 31+6 weeks gestation, admitted to the Cork University Maternity Hospital (CUMH) NICU and their mothers.

The study was designed as a longitudinal observational study with a time-limited recruitment period of approximately 18 months. With a projected number of 100 infants less than 32 weeks to be admitted to the CUMH NICU annually, and an expected consent rate of 60-70%, a pragmatically pre-defined minimum enrolment number of 80 participants over 18 months of recruitment was set.

2.2.1. Inclusion and exclusion criteria

To be eligible for the study, the participants must meet the terms of the inclusion and exclusion criteria as presented in Table 1.

Table 1. Inclusion and exclusion criteria of the PIMENTO study.

Inclusion criteria	Exclusion criteria
1. Birth at less than 32 weeks gestational age and admitted to the NICU	1. No informed consent
2. Free from antenatal suspicion of major congenital abnormalities	2. Major congenital anomalies
3. Ability of the participant (in the investigator's opinion) to comprehend the full nature and purpose of the study including possible risks and side effects.	3. Gastrointestinal anomalies
4. Consent to participate in the study and willing to comply with the protocol and study restrictions.	4. Inborn errors of metabolism

2.3. Recruitment

Pregnant women will be approached if there is a possibility of delivery before 32 weeks gestation. The study will be explained, and a participant information leaflet will be provided. Written informed consent will be obtained if the pregnant woman is agreeable after a sufficient amount of time to assimilate the information.

Multiple births will be enrolled separately. Samples will be collected from each infant but only one set of maternal samples would be collected. If one twin is delivered vaginally and the other by caesarean section, both skin and vaginal swabs will be collected from the mother if possible.

Written informed consent will be required for all participants. However, in exceptional circumstances (e.g. precipitous delivery) participants may be enrolled with deferred consent with the intent that investigators would obtain written consent as soon as possible thereafter. This would facilitate in the event a mother was not available to provide written informed consent that samples could be collected, i.e. a rectal swab from infant could be taken on admission to the NICU (as additional to the routine clinical swabs) and that, if passed a sample of the first meconium could be retained to ensure that, if subsequently enrolled, completeness of data for that participant could be maintained. Samples would be stored in a freezer on-site and would not be processed or analysed until written informed consent had been obtained. Similarly, out-born infants transferred to the study site may be enrolled with deferred consent and the infant's first meconium and rectal swab retained until the mother is available to provide written informed consent. Again, sample processing or analysis would not be performed without written informed consent and the samples would remain frozen on site until that time.

2.4. Compensation

There is no compensation provided to the participants. There are no cost implications for the Health Service Executive (HSE) or to the participants. The management of patients and investigative tests will comply with current standards of care.

2.5. Study timeline

Each participant will have samples collected during the entirety of their neonatal unit stay (Section 2.9 and Figure 1). Sample collection will not occur if the infant is transferred to another centre for ongoing management.

2.6. Participant withdrawal/exclusion

Under the Declaration of Helsinki, the researcher will explain to the participant that they have the right to withdraw from the study at any time and that this will in no way prejudice their future treatment. The reason for withdrawal will be recorded in the source documents and on the appropriate CRF. Withdrawn participants will be replaced. With consent samples from

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

individuals withdrawing from the study will be stored and may be analysed in the study. Participants excluded from the study after consenting will be replaced.

2.7 Ethics and dissemination

The study is conducted following the version Fortaleza, Brazil, October 2013 of the Declaration of Helsinki 1964. The Protocol and the Informed Consent Form have been approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals (CREC) before commencement (approval letter ECM 4 (t) 10/11/2020 & ECM 3 (mmm) 09/03/2021). On 25th August 2022 Protocol version 5 has been approved by the CREC (approval letter ECM 4 (t) 10/11/2020 & ECM 3 (vvv) 20/09/2022).

If a protocol amendment is necessary, this will be prepared with the agreement of the chief investigator and signed by the relevant parties. If the amendment is substantial, it will be submitted to the CREC and possibly other public bodies according to local requirements for review and approval. The protocol amendment will not be implemented before the required approvals are obtained.

The trial is sponsored by University College Cork (College Road, Cork T12 K8AF, +353 (0)21 490 3000). The sponsor is not involved in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit for publication. The trial has been registered at ClinicalTrials.gov (NCT05803577).

The findings from this study will be disseminated in peer-reviewed journals, during scientific conferences, and directly to the study participants.

2.8. Objectives and Outcomes

2.8.1. Primary Objective

Detailed characterisation of the very preterm microbiome, using culture independent approaches, including maternal origin and initial colonisation, postnatal alterations secondary to medical management (e.g. antibiotic and probiotic administration) and recovery/progression after discontinuation.

2.8.2. Secondary Objectives

To analyse maternal-infant transmission by correlation of the maternal microbiome with the early colonising microbiome of enrolled infants.

To determine changes in the human milk microbiome using culture independent approaches over time with increasing post menstrual age and characterise alterations in microbial content.

2.8.3. Exploratory Objectives

To assess metabolic functioning and systemic indicators in relation to microbiome composition at the time of sampling.

2.9. Sample collection

An overview of the sample collection timeline is presented in Figure 1.

Figure 1. Sample collection timeline. Samples were collected from birth to discharge. Timing of discharge was determined clinically and was variable for each infant so no definitive timeframe was set. Solid bars represent timelines for regular sample collection. Dashed bars represent the “window of opportunity” for sample collection for non-regular samples. Maternal perinatal samples, collected within 4 days of delivery, included stool, oral, skin and vaginal swabs.

2.9.1. Stool samples and rectal swabs

Rectal swabs are taken routinely on admission to the NICU for clinical microbiological screening. These are carried out by the admitting clinical nurse. One extra rectal swab will be taken for microbiome analysis at this time. Stool sampling will be carried out at the cotside during the course of the neonatal stay, a sample of the first meconium passed will be collected from the nappy and thereafter stool will be sampled from the infant’s nappy bi-weekly during routine cares. After 4 weeks of age, samples will be collected fortnightly. Samples are placed in a freezer in the neonatal unit immediately after sample collection. Thereafter samples are transferred to a -20°C freezer on-site for cataloguing prior to transfer to the off-site laboratory (APC Microbiome, Ireland, University College Cork, Ireland) for further processing and storage.

2.9.2. Urine samples

Enseignement Supérieur (ABES) .
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

During the inpatient stay, urine will be collected weekly, if possible, the attending neonatal nurse by placing sterile cotton balls in the infant's nappy which will then be removed at the next occurrence of routine cares. Urine is squeezed from the cotton balls into sterile collection bottles. Urine samples are placed in neonatal unit freezer immediately after collection, subsequently catalogued and transferred to the on-site -20°C freezer, and finally transferred to the off-site laboratory.

2.9.3. Blood samples

Infant blood samples will be collected from infants at times when clinical blood sampling is being performed. These will be collected in Serum Gel Z micro sample tubes. A sample volume of 1% of the estimated blood volume (estimated blood volume = 80mls/kg) will be collected. The sample will be immediately centrifuged on-site for 8 minutes at 9000G and then frozen at -20°C in the on-site freezer. Thereafter, the blood sample will be transferred to the off-site laboratory for further storage and processing. If possible, two to three blood samples will be collected during the entire neonatal stay.

2.9.4. Breastmilk samples

In the case where mothers are expressing milk for their infants, breastmilk samples will be collected once there is enough supply to provide for the infant's intake and surplus for storage. Breastmilk is expressed by mother's either on -site in the neonatal unit or at home into sterile collection bottles. Expressed breastmilk is refrigerated immediately and can be stored refrigerated for up to 24 hours or frozen. 5-15 mls of breastmilk is collected from the refrigerated breastmilk storage bottles using aseptic technique and transferred to sterile cryogenic storage tubes. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter. If possible, three breastmilk samples will be collected during the entire neonatal stay.

2.9.5. Maternal Stool samples

Within three days of delivery, mothers will be approached to provide a sample from their first stool passed after delivery. Once collected, the sample will be brought by the mother to the neonatal unit when it will be placed in the neonatal unit freezer. Samples will be collected from there to be catalogued and transferred to an on-site -20°C freezer, and thereafter, to the off-site laboratory.

2.9.6. Maternal Oral samples

Oral swabs will be collected from mothers within three days after birth. Samples will not be collected within one-hour after mother's have eaten food. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.9.7. Maternal Skin swabs

In the case of infants delivered by caesarean section, skin swabs will be collected from mothers within three days after birth. Swabs are wetted with sterile water for injection prior to swabbing the skin of the left wrist. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.9.8. Maternal Vaginal swabs

In the case of infants born by vaginal delivery mothers will be asked within three days of delivery to provide a vaginal swab. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.10 Sample analysis

Faecal samples will be processed for DNA extraction and metabolomics. DNA will be extracted from stool using the modification of the Repeated Bead Beating Plus Column (RBB+C) method (37) and subjected to 16S rRNA and/or shotgun sequencing. Metabolomics of infant's faeces will be analysed by the gas chromatography–mass spectrometry (GC-MS) technique using methyl chloroformate derivatisation (38). Bioinformatic analysis will identify the microbial composition at the phylum, genus and family levels using both read-based (39) and assembly-based (40) approaches. Metagenomic data will be correlated with physiological and clinical parameters and biomarkers. Advanced statistical and bioinformatics approaches will be applied to identify modifiable environmental factors that influence bacterial groups/consortia. This includes diversity analysis (41), differential abundance analysis (42), strain persistence and vertical transmission (43) and genome-scale metabolic modelling (44).

Breast milk samples will be used for DNA extraction (using modification of Qiagen DNeasy Powerfood Microbial Kit (QIAGEN Ltd, Manchester, United Kingdom) and the milk microbiome composition will be analysed by using state-of-the-art methods, possibly including but not limited to sequencing, and quantitative PCR.

Urine metabolomics analysis will be analysed by the liquid chromatography–mass spectrometry (LC-MS) technique (28).

The maternal oral microbiome composition will be analysed by using state-of-the-art methods after DNA extraction, possibly including but not limited to sequencing, and quantitative PCR. The bioinformatic analysis will identify the microbial composition at the phylum, genus, and family levels.

From vaginal swabs, DNA will be extracted using the modification of the RBB+C method (37) and the vaginal microbiome composition will be analysed by using state-of-the-art

methods, possibly including but not limited to sequencing, and quantitative PCR. The bioinformatic analysis will identify the microbial composition at the phylum, genus, and family levels.

2.11. Adverse events and participant well-being

This is an observational study. No investigational medicinal product (IMP) will be administered to study participants at any stage during the study. The study procedures are not greater than minimal risk, adverse events and serious adverse events are not expected. All self-reported adverse events (AEs) will be listed documenting duration, severity, participant outcome, and if any therapy was required. The study Chief investigator will review these and advance any reports to the local Ethics Committee if it is deemed necessary.

2.12. Data collection and management

2.12.1 Data collection/Case Report Forms (CRFs)

Data will be collected throughout the neonatal stay and from clinic visits after discharge. Data will be extracted from the electronic health records of each subject meeting the eligibility criteria and being included in the study. Data will be recorded in an electronic spreadsheet data collection tool stored on an encrypted password protected computer and backed up to the secure university server. All study staff responsible for entering data into the data collection tool will be trained prior to the start-up of the study.

The data in the CRFs will be consistent with the relevant source documents. The only source documents that will be available will be participants' in-patient electronic health record at CUMH. All data stored will be pseudonymised and treated with strict confidentiality in accordance with the General Data Protection Regulation (GDPR).

2.12.2 Sequencing data

Data types and formats collected and/or produced will include Raw whole genome sequencing (WGS) data (fastq), WGS assembly files and excel file with strains name (fasta), WGS annotation files (fna, gff, faa, ffn), 16S raw data (fastq), 16S ASV table (csv), 16S "phyloseq" file, which includes metadata for each sample (rds), Raw shotgun data (fastq), MAGs/bins from shotgun data (fna), Pangenomes files (newick, fa, csv/tab), and pipeline for the metabolic modelling part and generated data (sbml, mat). Sequencing data will be deposited in public databases.

2.13. Statistical analysis

2.13.1. Sample size justification

This study was designed as a longitudinal observational trial with the primary aim of detailed characterisation of the colonising microbiome of very preterm infants and, additionally, determination of factors relating to perturbations seen therein. As such, there was no statistically predefined sample size but instead the goal to recruit the largest sample population possible in the allotted timeframe for study recruitment, in this case 18-20 months. Post-hoc, we will calculate and state statistical power analysis to give a representation of statistically relevant sample size for this cohort.

3. Discussion

This study aims to longitudinally observe the gut microbiome of very preterm infants during their admission to Cork University Maternity Hospital neonatal unit. We intend to characterise in detail how clinical management practices influence the microbiome of these infants, including but not limited to delivery practices, antibiotic usage, feeding types and regimens, and probiotic usage. Very preterm infants have increased risk of potentially microbiome-mediated pathologies such as NEC and late-onset sepsis. The role that the microbiome plays in these diseases may relate to the overall composition of the microbiome, individual species within the microbiome, or temporal changes that occur. As such, management practices may be influential in the development of microbiome structure and may therefore have an indirect role in the development of these diseases.

At CUMH probiotics (*Bifidobacterim bifidum* and *Lactobacillus acidophilus*) are administered to all infants born less than 32 weeks gestation or less than 1500g birthweight from when feeding is deemed to be tolerated until 34 weeks corrected gestational age. We intend to observe these probiotic strains within the gut microbiome community, how they persist after supplementation is discontinued, and how clinical management may affect their colonisation.

Collected maternal samples will allow us to study transmission of microbes to premature infants and longitudinally collected breastmilk samples will allow us to study how the milk microbiome plays a role in shaping gut microbiome structure of the preterm infant.

This study will inform our understanding of microbial colonisation of very preterm infants, successional patterns of colonisation and how clinical management practices influence these. It will also provide valuable information on probiotic colonisation and practice around probiotic administration and may help development of efficacious probiotic strains.

Status of Study

The trial is ongoing as of 7th April 2021 The minimum number of participants has been reached and recruitment of new participants was completed on 31st December 2022. The primary endpoint will be when 2-year neurodevelopment follow up has been completed for the final participant.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Competing interests

The authors have no known competing financial interests.

Author's contributions

D.B.H. wrote the manuscript. D.B.H., R.P.R., C.S., E.M.D. were involved in study conception and design, acquisition of ethical approval, and review of the manuscript. D.B.H. coordinated participant recruitment, sample collection, on-site sample processing and sample transport. S.W. coordinated laboratory sample processing in conjunction with A.K.W.. G.G. coordinated bioinformatic analysis. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank all the study participants and the neonatal nursing staff in Cork University Maternity Hospital.

Funding

This publication has emanated from research conducted with financial support of Science Foundation Ireland (SFI) under Grant No. 12/RC/2273_P2 and 19/SP/6989.

References

1. Kennedy KM, De Goffau MC, Perez-Muñoz ME, Arrieta M-C, Bäckhed F, Bork P, et al. Questioning the fetal microbiome illustrates pitfalls of low-biomass microbial studies. *Nature*. 2023;613(7945):639-49. doi: 10.1038/s41586-022-05546-8

2. Adlerberth I, Wold AE. Establishment of the gut microbiota in Western infants. *Acta paediatrica*. 2009;98(2):229-38. doi: 10.1111/j.1651-2227.2008.01060.x

3. Fança-Berthon P, Michel C, Pagniez A, Rival M, Van Seuningen I, Darmaun D, et al. Intrauterine Growth Restriction Alters Postnatal Colonic Barrier Maturation in Rats. *Pediatric Research*. 2009;66(1):47-52. doi: 10.1203/pdr.0b013e3181a2047e

4. Zhang W, Ma C, Xie P, Zhu Q, Wang X, Yin Y, et al. Gut microbiota of newborn piglets with intrauterine growth restriction have lower diversity and different taxonomic abundances. *Journal of Applied Microbiology*. 2019;127(2):354-69. doi: 10.1111/jam.14304

5. Brooks B, Firek BA, Miller CS, Sharon I, Thomas BC, Baker R, et al. Microbes in the neonatal intensive care unit resemble those found in the gut of premature infants. *Microbiome*. 2014;2(1):1. doi: 10.1186/2049-2618-2-1

6. Fouhy F, Guinane CM, Hussey S, Wall R, Ryan CA, Dempsey EM, et al. High-throughput sequencing reveals the incomplete, short-term recovery of infant gut microbiota following parenteral antibiotic treatment with ampicillin and gentamicin. *Antimicrobial Agents & Chemotherapy*. 2012;56(11):5811-20. doi: 10.1128/AAC.00789-12

7. Hussey S, Wall R, Gruffman E, O'Sullivan L, Ryan CA, Murphy B, et al. Parenteral antibiotics reduce bifidobacteria colonization and diversity in neonates. *Int J Microbiol*. 2011;2011. doi: 10.1155/2011/130574

8. Zeissig S, Blumberg RS. Life at the beginning: perturbation of the microbiota by antibiotics in early life and its role in health and disease. *Nat Immunol*. 2014;15(4):307-10. doi: 10.1038/ni.2847

9. Cai C, Zhang Z, Morales M, Wang Y, Khafipour E, Friel J. Feeding practice influences gut microbiome composition in very low birth weight preterm infants and the association with oxidative stress: A prospective cohort study. *Free Radic Biol Med*. 2019;142:146-54. doi: 10.1016/j.freeradbiomed.2019.02.032

10. Cong X, Judge M, Xu W, Diallo A, Janton S, Brownell EA, et al. Influence of Feeding Type on Gut Microbiome Development in Hospitalized Preterm Infants. *Nurs Res*. 2017;66(2):123-33. doi: 10.1097/NNR.0000000000000208

11. Parra-Llorca A, Gormaz M, Alcantara C, Cernada M, Nunez-Ramiro A, Vento M, et al. Preterm Gut Microbiome Depending on Feeding Type: Significance of Donor Human Milk. *Front Microbiol*. 2018;9:1376. doi: 10.3389/fmicb.2018.01376

12. Neu J, Mshvildadze M, Mai V. A roadmap for understanding and preventing necrotizing enterocolitis. *Current Gastroenterology Reports*. 2008;10(5):450-7. doi: 10.1007/s11894-008-0084-x

13. Mihi B, Good M. Impact of Toll-Like Receptor 4 Signaling in Necrotizing Enterocolitis. *Clinics in perinatology*. 2019;46(1):145-57. doi: 10.1016/j.clp.2018.09.007

14. Stewart CJ, Embleton ND, Marrs EC, Smith DP, Nelson A, Abdulkadir B, et al. Temporal bacterial and metabolic development of the preterm gut reveals specific signatures in health and disease. *Microbiome*. 2016;4(1):67. doi: 10.1186/s40168-016-0216-8

15. Fanaro S, Chierici R, Guerrini P, Vigi V. Intestinal microflora in early infancy: composition and development. *Acta Paediatr Suppl*. 2003;91(441):48-55. doi: 10.1111/j.1651-2227.2003.tb00646.x

16. Ahrne S, Lonnermark E, Wold AE, Aberg N, Hesselmar B, Saalman R, et al. Lactobacilli in the intestinal microbiota of Swedish infants. *Microbes Infect*. 2005;7(11-12):1256-62. doi: 10.1016/j.micinf.2005.04.011

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

17. Gewolb IH, Schwalbe RS, Taciak VL, Harrison TS, Panigrahi P. Stool microflora in extremely low birthweight infants. *Arch Dis Child Fetal Neonatal Ed.* 1999;80(3):F167-73. doi: 10.1136/fn.80.3.f167
18. Goldmann DA, Leclair J, Macone A. Bacterial colonization of neonates admitted to an intensive care environment. *The Journal of pediatrics.* 1978;93(2):288-93. doi: 10.1016/s0022-3476(78)80523-x
19. Dani C, Biadaoli R, Bertini G, Martelli E, Rubaltelli FF. Probiotics feeding in prevention of urinary tract infection, bacterial sepsis and necrotizing enterocolitis in preterm infants. A prospective double-blind study. *Biology of the neonate.* 2002;82(2):103-8. doi: 10.1159/000063096
20. Lin HC, Su BH, Chen AC, Lin TW, Tsai CH, Yeh TF, et al. Oral probiotics reduce the incidence and severity of necrotizing enterocolitis in very low birth weight infants. *Pediatrics.* 2005;115(1):1-4. doi: 10.1542/peds.2004-1463
21. Bin-Nun A, Bromiker R, Wilschanski M, Kaplan M, Rudensky B, Caplan M, et al. Oral probiotics prevent necrotizing enterocolitis in very low birth weight neonates. *The Journal of pediatrics.* 2005;147(2):192-6. doi: 10.1016/j.jpeds.2005.03.054
22. Lau CS, Chamberlain RS. Probiotic administration can prevent necrotizing enterocolitis in preterm infants: A meta-analysis. *J Pediatr Surg.* 2015;50(8):1405-12. doi: 10.1016/j.jpedsurg.2015.05.008
23. Costeloe K, Hardy P, Juszczak E, Wilks M, Millar MR, Probiotics in Preterm Infants Study Collaborative G. Bifidobacterium breve BBG-001 in very preterm infants: a randomised controlled phase 3 trial. *Lancet.* 2016;387(10019):649-60. doi: 10.1016/S0140-6736(15)01027-2
24. Jacobs SE, Tobin JM, Opie GF, Donath S, Tabrizi SN, Pirotta M, et al. Probiotic effects on late-onset sepsis in very preterm infants: a randomized controlled trial. *Pediatrics.* 2013;132(6):1055-62. doi: 10.1542/peds.2013-1339
25. AlFaleh K, Anabrees J. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev.* 2014(4):CD005496. doi: 10.1002/14651858.CD005496.pub4
26. Plummer EL, Bulach DM, Murray GL, Jacobs SE, Tabrizi SN, Garland SM, et al. Gut microbiota of preterm infants supplemented with probiotics: sub-study of the ProPremis trial. *BMC Microbiol.* 2018;18(1):184. doi: 10.1186/s12866-018-1326-1
27. van den Akker CHP, van Goudoever JB, Shamir R, Domellof M, Embleton ND, Hojsak I, et al. Probiotics and Preterm Infants: A Position Paper by the European Society for Paediatric Gastroenterology Hepatology and Nutrition Committee on Nutrition and the European Society for Paediatric Gastroenterology Hepatology and Nutrition Working Group for Probiotics and Prebiotics. *J Pediatr Gastroenterol Nutr.* 2020;70(5):664-80. doi: 10.1097/MPG.0000000000002655
28. Hill CJ, Lynch DB, Murphy K, Ulaszewska M, Jeffery IB, O'Shea CA, et al. Evolution of gut microbiota composition from birth to 24 weeks in the INFANTMET Cohort. *Microbiome.* 2017;5(1):4. doi: 10.1186/s40168-016-0213-y
29. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, et al. Temporal development of the gut microbiome in early childhood from the TEDDY study. *Nature.* 2018;562(7728):583-8. doi: 10.1038/s41586-018-0617-x
30. Moles L, Gomez M, Heilig H, Bustos G, Fuentes S, de Vos W, et al. Bacterial diversity in meconium of preterm neonates and evolution of their fecal microbiota during the first month of life. *PLoS One.* 2013;8(6):e66986. doi: 10.1371/journal.pone.0066986
31. Cong X, Xu W, Janton S, Henderson WA, Matson A, McGrath JM, et al. Gut Microbiome Developmental Patterns in Early Life of Preterm Infants: Impacts of Feeding and Gender. *PLoS One.* 2016;11(4):e0152751. doi: 10.1371/journal.pone.0152751
32. Young GR, van der Gast CJ, Smith DL, Berrington JE, Embleton ND, Lanyon C. Acquisition and Development of the Extremely Preterm Infant Microbiota Across Multiple Anatomical Sites. *J Pediatr Gastroenterol Nutr.* 2020;70(1):12-9. doi: 10.1097/MPG.0000000000002549

33. Shao Y, Forster SC, Tsaliki E, Vervier K, Strang A, Simpson N, et al. Stunted microbiota and opportunistic pathogen colonization in caesarean-section birth. *Nature*. 2019;574(7776):117-21. doi: 10.1038/s41586-019-1560-1

34. Lyons KE, Ryan CA, Dempsey EM, Ross RP, Stanton C. Breast Milk, a Source of Beneficial Microbes and Associated Benefits for Infant Health. *Nutrients*. 2020;12(4). doi: 10.3390/nu12041039

35. Herrmann K, Carroll K. An exclusively human milk diet reduces necrotizing enterocolitis. *Breastfeed Med*. 2014;9(4):184-90. doi: 10.1089/bfm.2013.0121

36. Masi AC, Embleton ND, Lamb CA, Young G, Granger CL, Najera J, et al. Human milk oligosaccharide DSLNT and gut microbiome in preterm infants predicts necrotising enterocolitis. *Gut*. 2021;70(12):2273-82. doi: 10.1136/gutjnl-2020-322771

37. Yu Z, Morrison M. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques*. 2004;36(5):808-12. doi: 10.2144/04365ST04

38. Smart KF, Aggio RB, Van Houtte JR, Villas-Boas SG. Analytical platform for metabolome analysis of microbial cells using methyl chloroformate derivatization followed by gas chromatography-mass spectrometry. *Nat Protoc*. 2010;5(10):1709-29. doi: 10.1038/nprot.2010.108

39. Beghini F, McIver LJ, Blanco-Miguez A, Dubois L, Asnicar F, Maharjan S, et al. Integrating taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3. *Elife*. 2021;10. doi: 10.7554/eLife.65088

40. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic assembler. *Genome Res*. 2017;27(5):824-34. doi: 10.1101/gr.213959.116

41. Lozupone CA, Hamady M, Kelley ST, Knight R. Quantitative and qualitative beta diversity measures lead to different insights into factors that structure microbial communities. *Appl Environ Microbiol*. 2007;73(5):1576-85. doi: 10.1128/AEM.01996-06

42. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, et al. Multivariable association discovery in population-scale meta-omics studies. *PLoS Comput Biol*. 2021;17(11):e1009442. doi: 10.1371/journal.pcbi.1009442

43. Olm MR, Crits-Christoph A, Bouma-Gregson K, Firek BA, Morowitz MJ, Banfield JF. inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nat Biotechnol*. 2021;39(6):727-36. doi: 10.1038/s41587-020-00797-0

44. Heirendt L, Arreckx S, Pfau T, Mendoza SN, Richelle A, Heinken A, et al. Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v.3.0. *Nat Protoc*. 2019;14(3):639-702. doi: 10.1038/s41596-018-0098-2

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

BMJ Open

A Longitudinal Observational Study Protocol: Preterm Infants: Microbiome Establishment, Neuro-CrossTalk and Origins (PIMENTO)

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-075060.R1
Article Type:	Protocol
Date Submitted by the Author:	03-Jul-2023
Complete List of Authors:	Healy, David; University College Cork, Department of Paediatrics and Child Health Wang, Shuo; University College Cork Grimaud, Ghjuvan; Moorepark Food Research Centre Warda, Alicja; Teagasc Food Research Centre Moorepark Ross, Paul; University College Cork, APC Microbiome Ireland Stanton, Catherine; University College Cork APC Microbiome Institute Dempsey, E; University College Cork, Department of Paediatrics and Child Health
Primary Subject Heading:	Paediatrics
Secondary Subject Heading:	Research methods
Keywords:	Microbiology < NATURAL SCIENCE DISCIPLINES, Neonatal intensive & critical care < INTENSIVE & CRITICAL CARE, NEONATOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A Longitudinal Observational Study Protocol: Preterm Infants: Microbiome Establishment, Neuro-CrossTalk and Origins (PIMENTO)

Authors: David B. Healy ^{1,2}, Shuo Wang ¹, Ghjuvan Grimaud ³, Alicja K. Warda ^{1,3}, R. Paul Ross ^{1,3}, Catherine Stanton ^{1,2,4}, and Eugene M. Dempsey ^{1,2,4,5}

- ¹ APC Microbiome Ireland, Cork, T12 YT20, Ireland
- ² Department of Paediatrics and Child Health, University College Cork, T12 YT20, Ireland
- ³ Teagasc, Moorepark Food Research Centre, Fermoy, P61 C996, Ireland
- ⁴ University College Cork, Cork, T12 YT20, Ireland
- ⁵ INFANT Research Centre, University College Cork, T12 DC4A, Ireland

Journal: BMJOpen

Corresponding Author: David B. Healy, APC Microbiome Ireland; Department of Paediatrics and Child Health, david.healy@ucc.ie

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Abstract

Introduction

Very preterm infants are at risk of abnormal microbiome colonisation in the first weeks to months of life. Several important associated factors have been identified including gestational age, mode of delivery, antibiotic exposure and feeding. Preterm infants are at risk of a number of pathologies for which the microbiome may play a central role, including necrotising enterocolitis (NEC) and sepsis. The objective of this study is to determine detailed microbiome changes that occur around implementation of different management practices including empiric antibiotic use, advancement of feeds and administration of probiotics during admission to the neonatal intensive care unit.

Methods and analysis

A single-site, longitudinal observational study of infants born less than 32 weeks gestation, including collection of maternal samples around delivery and breastmilk and infant samples from admission through discharge from the neonatal unit.

Ethics and dissemination

The protocol was approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals (CREC). The trial has been registered at ClinicalTrials.gov (NCT05803577).

The findings from this study will be disseminated in peer-reviewed journals, during scientific conferences, and directly to the study participants. Sequencing data will be deposited in public databases.

Strengths and limitations of this study

- Study design will provide high resolution longitudinal microbiome data that will help to characterise the colonising microbiome of preterm infants, the effect of and progression after treatments.
- An array of samples will be collected including rectal swabs, stool, urine and blood from infants, peripartum maternal samples including stool, oral, skin, vaginal samples and expressed breastmilk from mothers.
- This is a single-site study.
- The study population is heterogenous, and the recruited sample size will be ~200 (mothers and infants).

Abbreviations:

CREC	Clinical Research Ethics Committee of the Cork Teaching Hospitals
CRF	Case report form
CUMH	Cork University Maternity Hospital
GC-MS	Gas chromatography-mass spectrometry
GDPR	General Data Protection Regulation
HSE	Health Service Executive
IMP	Investigational medicinal product
LC-MS	Liquid chromatography-mass spectrometry
NEC	Necrotising enterocolitis
NICU	Neonatal intensive care unit
PCR	Polymerase chain reaction
RBB+C	Repeated bead beating plus column
TLR	Toll-like receptor

Keywords: preterm infants, microbiome, probiotics

1. Introduction

Over the last two decades, the materno-feto-neonatal microbiome has rapidly become an area of budding interest in the specialty of neonatology. While there is much debate as to whether a developing fetus is exposed to microorganisms in utero (1), it is increasingly evident that the gut microbiome does not expand and diversify until the postnatal period at which point microbes from the mother and environment begin colonisation of the infant. Following birth, an initial relatively simple microbial community develops into a diverse microbial ecosystem in a controlled fashion over the first years of life (2). Bacterial colonisation, specifically in very low birth weight infants, may occur in an aberrant fashion for numerous reasons: an immature gut and immune system (3, 4); the environment in which they are nursed(5); frequent exposure to antimicrobial agents(6, 7, 8); and altered feeding practices such as delayed commencement of feeds or using formula in place of mother's own milk(9, 10, 11). It is thought that the colonising microbiome of very preterm infants, and the factors that shape it, play a crucial role in the development of microbiome-mediated complications such as NEC.

NEC is a complex multifactorial condition characterised by intestinal immaturity and dysbiosis in association with altered microvascular blood flow. It is the most commonly acquired serious gastrointestinal complication of prematurity and is rare in babies born above 32 weeks (12). Infants at risk are likely to have deficient intestinal mucus production, immature gut immunity, and reduced endogenously-produced antimicrobial factors. The ability of bacteria to adhere to epithelial cells is thereby enhanced, particularly after gut injury secondary to ischaemia/hypoperfusion. Although not fully understood, the most accepted theory for the pathogenesis of NEC is that in this setting, bacterial endotoxin binds to toll-like receptor 4 (TLR4) found on the intestinal epithelial cells and by so doing, pathogen-associated molecular pattern receptors are activated, resulting in cell apoptosis and epithelial injury (13). This leads to an intense inflammatory cascade in the exposed lamina propria, including activation of inflammatory cytokines such as tumour necrosis factor alpha, interleukin-1 and interleukin-8, and ultimately necrosis of the surrounding tissue. Work by Stewart et al.(14) demonstrated that NEC tends not to occur in infants with a diverse gut microbiome. This may relate to inherent factors that make the individual less susceptible to NEC and simultaneously allow a diverse microbiome or may be due to protective effects associated with having a diverse gut microbiome, such as inflammatory modulation and improved gut epithelial integrity.

With early descriptions of the "ideal" gut microbiome (i.e. that of the vaginally delivered, exclusively breast-fed term infant, and without antibiotic treatment)(15, 16) and the identification of potentially deleterious bacteria harboured in the guts of the hospitalised very preterm infants (15, 17, 18) came the advent of probiotics. Administration of these as a supplement to feeding aimed to lend some "normality" to the gut microbiome of these vulnerable babies by supplanting potentially pathogenic bacteria and attempting to reduce the incidence of NEC (19, 20, 21, 22, 23). While probiotic preparations are widely available, shown to be safe (24, 25) and do, indeed, increase so-called beneficial bacteria present in the

gut(26), the evidence for their benefit is contentious (27) and practice varies from institution to institution.

Work has shown that the full term vaginally delivered and exclusively breastfed baby’s initial gut microbiome is dominated primarily by Actinobacteria (28, 29) while studies looking at preterm infants’ microbiome have shown a Proteobacteria (Bacilli dominance) and Firmicutes predominance (28, 30, 31, 32). Clarification of the initial neonatal gut microbiome in the preterm population is confounded by the low bacterial biomass in early samples and high interindividual variability in composition. Individual medical management of each infant is likely to cause perturbations to the microbiome. Several factors associated with alterations in the neonatal gut microbiome have previously been identified. These include gestational age (28) mode of delivery(28, 33), exposure to antibiotics (both pre- and postnatally) (6) and, importantly, feeding choice (34). Mothers of very low birth weight infants are typically encouraged to express breastmilk for their baby. This is associated with lower rates of sepsis and NEC (35). Some of these effects may be related to the bacteria or supporting molecules received in breastmilk from their mother (36). Correlations between changes in the neonatal gut microbiome and the microbiota of the milk they are being fed are limited and longitudinal analysis of breastmilk in this regard is lacking. Furthermore, interactions within breastmilk between bacteria and bacteriophage are not well characterised.

Given the rarity of some neonatal diseases, such as NEC, collaborative efforts are necessary to study data from a sufficient number of infants to draw robust conclusions (37). A number of prospective longitudinal repositories have been established, including the Great North Neonatal Biobank in the UK and the NEC Biorepository in the US, with aims including the study and investigation of the preterm neonatal microbiome and it’s role in pathological disease processes such as NEC. Work stemming from these and others has considerably advanced our understanding of the microbiome in term and preterm infants over the past decade. Expanding data and sample collection to other regions will help in future collaborative work, to account for geographical variations in hospital practice and microbiome structure. This work will aim to longitudinally characterise the early and colonising intestinal microbiome of very preterm infants. We will analyse samples from infants less than 32 weeks gestational age at birth, with collection occurring longitudinally from birth to discharge from the neonatal intensive care unit (NICU). Microbiome composition will be evaluated using metagenomics and 16sRNA sequencing of stool. Bacterial function will be assessed longitudinally using untargeted metabolomics of infant urine. We will also interrogate some key perinatal maternal microbiome niches, including oral, vaginal, skin and intestinal, as well as expressed breastmilk to evaluate mother-infant transmission.

2. Methods

2.1. Study design

This single site study is a prospective longitudinal observational study of the intestinal microbiome of infants born between 23 and 32 weeks of gestation. It will investigate the establishment of the very premature infants' microbiome using culture independent approaches, including the potential initial colonisation process of vertical transmission from mother to preterm infant, the apparent differences in establishment pattern based on feeding type, and the perturbations and recovery associated with different intensive care management strategies such as empiric antibiotic usage.

Study recruitment commenced in April 2020 and finished in December 2022. The primary endpoint of the study will be reached after the last enrolled participant completes 2-year neurodevelopmental follow-up.

2.2. Participant selection

Participants are infants, born between 23+0 weeks gestation and 31+6 weeks gestation, admitted to the Cork University Maternity Hospital (CUMH) NICU and their mothers.

The study was designed as a longitudinal observational study with a time-limited recruitment period of approximately 18 months. With a projected number of 100 infants less than 32 weeks to be admitted to the CUMH NICU annually, and an expected consent rate of 60-70%, a pragmatically pre-defined minimum enrolment number of 80 participants over 18 months of recruitment was set.

2.2.1. Inclusion and exclusion criteria

To be eligible for the study, the participants must meet the terms of the inclusion and exclusion criteria as presented in Table 1.

Table 1. Inclusion and exclusion criteria of the PIMENTO study.

Inclusion criteria	Exclusion criteria
1. Birth at less than 32 weeks gestational age and admitted to the NICU	1. No informed consent
2. Free from antenatal suspicion of major congenital abnormalities	2. Major congenital anomalies
3. Ability of the participant (in the investigator's opinion) to comprehend the full nature and purpose of the study including possible risks and side effects.	3. Gastrointestinal anomalies

4. Consent to participate in the study and willing to comply with the protocol and study restrictions.	4. Inborn errors of metabolism
--	--------------------------------

2.3. Recruitment

Pregnant women will be approached if there is a possibility of delivery before 32 weeks gestation. The study will be explained, and a participant information leaflet will be provided. Written informed consent will be obtained if the pregnant woman is agreeable after a sufficient amount of time to assimilate the information.

Multiple births will be enrolled separately. Samples will be collected from each infant but only one set of maternal samples would be collected. If one twin is delivered vaginally and the other by caesarean section, both skin and vaginal swabs will be collected from the mother if possible.

Written informed consent will be required for all participants. However, in exceptional circumstances (e.g. precipitous delivery) participants may be enrolled with deferred consent with the intent that investigators would obtain written consent as soon as possible thereafter. This would facilitate in the event a mother was not available to provide written informed consent that samples could be collected, i.e. a rectal swab from infant could be taken on admission to the NICU (as additional to the routine clinical swabs) and that, if passed a sample of the first meconium could be retained to ensure that, if subsequently enrolled, completeness of data for that participant could be maintained. Samples would be stored in a freezer on-site and would not be processed or analysed until written informed consent had been obtained. Similarly, out-born infants transferred to the study site may be enrolled with deferred consent and the infant’s first meconium and rectal swab retained until the mother is available to provide written informed consent. Again, sample processing or analysis would not be performed without written informed consent and the samples would remain frozen on site until that time.

2.4. Follow up

Two-year neurodevelopmental assessment, using the Bayley Scale for Infant and Toddler Development, 3rd Edition (BSID-III), is routinely performed at CUMH for all infants born at 32 weeks gestational age or less.

2.5. Compensation

There is no compensation provided to the participants. There are no cost implications for the Health Service Executive (HSE) or to the participants. The management of patients and investigative tests will comply with current standards of care.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

2.6. Study timeline

Each participant will have samples collected during the entirety of their neonatal unit stay (Section 2.9 and Figure 1). Sample collection will not occur if the infant is transferred to another centre for ongoing management.

2.7. Participant withdrawal/exclusion

Under the Declaration of Helsinki, the researcher will explain to the participant that they have the right to withdraw from the study at any time and that this will in no way prejudice their future treatment. The reason for withdrawal will be recorded in the source documents and on the appropriate CRF. Withdrawn participants will be replaced. With consent samples from individuals withdrawing from the study will be stored and may be analysed in the study. Participants excluded from the study after consenting will be replaced.

2.8. Ethics and dissemination

The study is conducted following the version Fortaleza, Brazil, October 2013 of the Declaration of Helsinki 1964. The Protocol and the Informed Consent Form have been approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals (CREC) before commencement (approval letter ECM 4 (t) 10/11/2020 & ECM 3 (mmm) 09/03/2021). On 25th August 2022 Protocol version 5 has been approved by the CREC (approval letter ECM 4 (t) 10/11/2020 & ECM 3 (vvv) 20/09/2022).

If a protocol amendment is necessary, this will be prepared with the agreement of the chief investigator and signed by the relevant parties. If the amendment is substantial, it will be submitted to the CREC and possibly other public bodies according to local requirements for review and approval. The protocol amendment will not be implemented before the required approvals are obtained.

The trial is sponsored by University College Cork (College Road, Cork T12 K8AF, +353 (0)21 490 3000). The sponsor is not involved in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit for publication. The trial has been registered at ClinicalTrials.gov (NCT05803577).

The findings from this study will be disseminated in peer-reviewed journals, during scientific conferences, and directly to the study participants.

2.9. Objectives and Outcomes

2.9.1. Primary Objective

Detailed characterisation of the very preterm microbiome from birth to final hospital discharge, using culture independent approaches, including maternal origin and initial colonisation,

postnatal alterations secondary to medical management (e.g. antibiotic and probiotic administration) and recovery/progression after discontinuation.

- a) To describe changes in community structure and function of gut microbiome in preterm infants during stay in the neonatal unit
- b) To compare microbiome successional colonisation in infants requiring varying degrees of initial respiratory support.
- c) To evaluate probiotic strains in the gut microbiome using next generation sequencing during probiotic commencement, colonisation and persistence after discontinuation

2.9.2. Secondary Objectives

- a) To determine changes in the human milk microbiome using culture independent approaches over time with increasing post menstrual age and characterise alterations in microbial content and investigate the influence of breastmilk-derived microbes in the infant gut microbiome.

2.9.3. Exploratory Objectives

- a) To assess metabolic functioning and systemic indicators in relation to microbiome composition at the time of sampling.
- b) To analyse maternal-infant transmission by correlation of the maternal microbiome with the early colonising microbiome of enrolled infants.

2.10. Sample collection

An overview of the sample collection timeline is presented in Figure 1.

2.10.1. Stool samples and rectal swabs

Rectal swabs are taken routinely on admission to the NICU for clinical microbiological screening. These are carried out by the admitting clinical nurse. One extra rectal swab will be taken for microbiome analysis at this time. Stool sampling will be carried out at the cotside during the course of the neonatal stay, a sample of the first meconium passed will be collected from the nappy and thereafter stool will be sampled from the infant’s nappy bi-weekly during routine cares. After 4 weeks of age, samples will be collected fortnightly. Samples are placed in a freezer in the neonatal unit immediately after sample collection. Thereafter samples are transferred to a -20°C freezer on-site for cataloguing prior to transfer to the off-site laboratory (APC Microbiome, Ireland, University College Cork, Ireland) for further processing and storage.

2.10.2. Urine samples

During the inpatient stay, urine will be collected weekly, if possible, the attending neonatal nurse by placing sterile cotton balls in the infant's nappy which will then be removed at the next occurrence of routine cares. Urine is squeezed from the cotton balls into sterile collection bottles. Urine samples are placed in neonatal unit freezer immediately after collection, subsequently catalogued and transferred to the on-site -20°C freezer, and finally transferred to the off-site laboratory.

2.10.3. Blood samples

Infant blood samples will be collected from infants at times when clinical blood sampling is being performed. These will be collected in Serum Gel Z micro sample tubes. A sample volume of 1% of the estimated blood volume (estimated blood volume = 80mls/kg) will be collected. The sample will be immediately centrifuged on-site for 8 minutes at 9000G and then frozen at -20°C in the on-site freezer. Thereafter, the blood sample will be transferred to the off-site laboratory for further storage and processing. If possible, two to three blood samples will be collected during the entire neonatal stay.

2.10.4. Breastmilk samples

In the case where mothers are expressing milk for their infants, breastmilk samples will be collected once there is enough supply to provide for the infant's intake and surplus for storage. Breastmilk is expressed by mother's either on -site in the neonatal unit or at home into sterile collection bottles. Expressed breastmilk is refrigerated immediately and can be stored refrigerated for up to 24 hours or frozen. 5-15 mls of breastmilk is collected from the refrigerated breastmilk storage bottles using aseptic technique and transferred to sterile cryogenic storage tubes. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter. If possible, three breastmilk samples will be collected during the entire neonatal stay.

2.10.5. Maternal Stool samples

Within three days of delivery, mothers will be approached to provide a sample from their first stool passed after delivery. Once collected, the sample will be brought by the mother to the neonatal unit when it will be placed in the neonatal unit freezer. Samples will be collected from there to be catalogued and transferred to an on-site -20°C freezer, and thereafter, to the off-site laboratory.

2.10.6. Maternal Oral samples

Oral swabs will be collected from mothers within three days after birth. Samples will not be collected within one-hour after mother's have eaten food. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.10.7. Maternal Skin swabs

In the case of infants delivered by caesarean section, skin swabs will be collected from mothers within three days after birth. Swabs are wetted with sterile water for injection prior to swabbing the skin of the left wrist. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.10.8. Maternal Vaginal swabs

In the case of infants born by vaginal delivery mothers will be asked within three days of delivery to provide a vaginal swab. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.11 Sample analysis

Faecal samples will be processed for DNA extraction and metabolomics. DNA will be extracted from stool using the modification of the Repeated Bead Beating Plus Column (RBB+C) method (38) and subjected to 16S rRNA and/or shotgun sequencing. Metabolomics of infant's faeces will be analysed by the gas chromatography–mass spectrometry (GC-MS) technique using methyl chloroformate derivatisation (39). Bioinformatic analysis will identify the microbial composition at the phylum, genus and family levels using both read-based (40) and assembly-based (41) approaches. Metagenomic data will be correlated with physiological and clinical parameters and biomarkers. Advanced statistical and bioinformatics approaches will be applied to identify modifiable environmental factors that influence bacterial groups/consortia. This includes diversity analysis (42), differential abundance analysis (43), strain persistence and vertical transmission (44) and genome-scale metabolic modelling (45).

Breast milk samples will be used for DNA extraction (using modification of Qiagen DNeasy Powerfood Microbial Kit (QIAGEN Ltd, Manchester, United Kingdom) and the milk microbiome composition will be analysed by using state-of-the-art methods, possibly including but not limited to sequencing, and quantitative PCR.

Urine metabolomics analysis will be analysed by the liquid chromatography–mass spectrometry (LC-MS) technique (28).

The maternal oral microbiome composition will be analysed by using state-of-the-art methods after DNA extraction, possibly including but not limited to sequencing, and quantitative PCR. The bioinformatic analysis will identify the microbial composition at the phylum, genus, and family levels.

From vaginal swabs, DNA will be extracted using the modification of the RBB+C method (38) and the vaginal microbiome composition will be analysed by using state-of-the-art

methods, possibly including but not limited to sequencing, and quantitative PCR. The bioinformatic analysis will identify the microbial composition at the phylum, genus, and family levels.

2.12. Adverse events and participant well-being

This is an observational study. No investigational medicinal product (IMP) will be administered to study participants at any stage during the study. The study procedures are not greater than minimal risk, adverse events and serious adverse events are not expected. All self-reported adverse events (AEs) will be listed documenting duration, severity, participant outcome, and if any therapy was required. The study Chief investigator will review these and advance any reports to the local Ethics Committee if it is deemed necessary.

2.13. Data collection and management

2.13.1 Data collection/Case Report Forms (CRFs)

Data will be collected throughout the neonatal stay and from clinic visits after discharge. Data will be extracted from the electronic health records of each subject meeting the eligibility criteria and being included in the study. Data will be recorded in an electronic spreadsheet data collection tool stored on an encrypted password protected computer and backed up to the secure university server. All study staff responsible for entering data into the data collection tool will be trained prior to the start-up of the study.

The data in the CRFs will be consistent with the relevant source documents. The only source documents that will be available will be participants' in-patient electronic health record at CUMH. All data stored will be pseudonymised and treated with strict confidentiality in accordance with the General Data Protection Regulation (GDPR).

2.13.2 Sequencing data

Data types and formats collected and/or produced will include Raw whole genome sequencing (WGS) data (fastq), WGS assembly files and excel file with strains name (fasta), WGS annotation files (fna, gff, faa, ffn), 16S raw data (fastq), 16S ASV table (csv), 16S "phyloseq" file, which includes metadata for each sample (rds), Raw shotgun data (fastq), MAGs/bins from shotgun data (fna), Pangenomes files (newick, fa, csv/tab), and pipeline for the metabolic modelling part and generated data (sbml, mat). Sequencing data will be deposited in public databases.

2.14. Statistical analysis

2.14.1. Sample size justification

This study was designed as a longitudinal observational trial with the primary aim of detailed characterisation of the colonising microbiome of very preterm infants and, additionally,

determination of factors relating to perturbations seen therein. As such, there was no statistically predefined sample size but instead the goal to recruit the largest sample population possible in the allotted timeframe for study recruitment, in this case 18-20 months. Post-hoc, we will calculate and state statistical power analysis to give a representation of statistically relevant sample size for this cohort.

3. Discussion

This study aims to longitudinally observe the gut microbiome of very preterm infants during their admission to Cork University Maternity Hospital neonatal unit. We intend to characterise in detail how clinical management practices influence the microbiome of these infants, including but not limited to delivery practices, antibiotic usage, feeding types and regimens, and probiotic usage. Very preterm infants have increased risk of potentially microbiome-mediated pathologies such as NEC and late-onset sepsis. The role that the microbiome plays in these diseases may relate to the overall composition of the microbiome, individual species within the microbiome, or temporal changes that occur. As such, management practices may be influential in the development of microbiome structure and may therefore have an indirect role in the development of these diseases.

At CUMH probiotics (*Bifidobacterim bifidum* and *Lactobacillus acidophilus*) are administered to all infants born less than 32 weeks gestation or less than 1500g birthweight from when feeding is deemed to be tolerated until 34 weeks corrected gestational age. We intend to observe these probiotic strains within the gut microbiome community, how they persist after supplementation is discontinued, and how clinical management may affect their colonisation.

Collected maternal samples will allow us to study transmission of microbes to premature infants and longitudinally collected breastmilk samples will allow us to study how the milk microbiome plays a role in shaping gut microbiome structure of the preterm infant.

The authors acknowledge the limitations of this study design. It will be a single centre observational study and the results may not be generalisable across all other institutions. As all infants that are to be enrolled in the study will receive probiotic supplementation, there will not be a control group without probiotics with which to compare. The study is designed as descriptive and without intervention. Accordingly, the management of the infants enrolled has not been standardised. While the clinical demographics of the enrolled infants will be described in careful detail, all confounders may not be accounted for owing to variability in clinical courses and the variability in management decisions made by the various providers caring for the infants.

We envisage that this work will add to the already existing information on the preterm microbiome. As neonatal care practices can differ markedly from institution to institution we hope that this study will provide novel insights into how the microbiome is established, including maternal transmission of microbes at delivery and through breastmilk. This study

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

will inform our understanding of microbial colonisation of very preterm infants, successional patterns of colonisation and how clinical management practices influence these. It will also provide valuable information on probiotic colonisation and practices around probiotic administration and may help future development of efficacious probiotic strains.

Status of Study

The trial is ongoing as of 7th April 2021. The minimum number of participants has been reached and recruitment of new participants was completed on 31st December 2022. The primary endpoint will be when 2-year neurodevelopment follow up has been completed for the final participant.

Competing interests

The authors have no known competing financial interests.

Author's contributions

D.B.H. wrote the manuscript. D.B.H., R.P.R., C.S., E.M.D. were involved in study conception and design, acquisition of ethical approval, and review of the manuscript. D.B.H. coordinated participant recruitment, sample collection, on-site sample processing and sample transport. S.W. coordinated laboratory sample processing in conjunction with A.K.W.. G.G. coordinated bioinformatic analysis. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank all the study participants and the neonatal nursing staff in Cork University Maternity Hospital.

Funding

This publication has emanated from research conducted with financial support of Science Foundation Ireland (SFI) under Grant No. 12/RC/2273_P2 and 19/SP/6989.

References

1. Kennedy KM, De Goffau MC, Perez-Muñoz ME, Arrieta M-C, Bäckhed F, Bork P, et al. Questioning the fetal microbiome illustrates pitfalls of low-biomass microbial studies. *Nature*. 2023;613(7945):639-49. doi: 10.1038/s41586-022-05546-8

2. Adlerberth I, Wold AE. Establishment of the gut microbiota in Western infants. *Acta paediatrica*. 2009;98(2):229-38. doi: 10.1111/j.1651-2227.2008.01060.x

3. Fança-Berthon P, Michel C, Pagniez A, Rival M, Van Seuningen I, Darmaun D, et al. Intrauterine Growth Restriction Alters Postnatal Colonic Barrier Maturation in Rats. *Pediatric Research*. 2009;66(1):47-52. doi: 10.1203/pdr.0b013e3181a2047e

4. Zhang W, Ma C, Xie P, Zhu Q, Wang X, Yin Y, et al. Gut microbiota of newborn piglets with intrauterine growth restriction have lower diversity and different taxonomic abundances. *Journal of Applied Microbiology*. 2019;127(2):354-69. doi: 10.1111/jam.14304

5. Brooks B, Firek BA, Miller CS, Sharon I, Thomas BC, Baker R, et al. Microbes in the neonatal intensive care unit resemble those found in the gut of premature infants. *Microbiome*. 2014;2(1):1. doi: 10.1186/2049-2618-2-1

6. Fouhy F, Guinane CM, Hussey S, Wall R, Ryan CA, Dempsey EM, et al. High-throughput sequencing reveals the incomplete, short-term recovery of infant gut microbiota following parenteral antibiotic treatment with ampicillin and gentamicin. *Antimicrobial Agents & Chemotherapy*. 2012;56(11):5811-20. doi: 10.1128/AAC.00789-12

7. Hussey S, Wall R, Gruffman E, O'Sullivan L, Ryan CA, Murphy B, et al. Parenteral antibiotics reduce bifidobacteria colonization and diversity in neonates. *Int J Microbiol*. 2011;2011. doi: 10.1155/2011/130574

8. Zeissig S, Blumberg RS. Life at the beginning: perturbation of the microbiota by antibiotics in early life and its role in health and disease. *Nat Immunol*. 2014;15(4):307-10. doi: 10.1038/ni.2847

9. Cai C, Zhang Z, Morales M, Wang Y, Khafipour E, Friel J. Feeding practice influences gut microbiome composition in very low birth weight preterm infants and the association with oxidative stress: A prospective cohort study. *Free Radic Biol Med*. 2019;142:146-54. doi: 10.1016/j.freeradbiomed.2019.02.032

10. Cong X, Judge M, Xu W, Diallo A, Janton S, Brownell EA, et al. Influence of Feeding Type on Gut Microbiome Development in Hospitalized Preterm Infants. *Nurs Res*. 2017;66(2):123-33. doi: 10.1097/NNR.0000000000000208

11. Parra-Llorca A, Gormaz M, Alcantara C, Cernada M, Nunez-Ramiro A, Vento M, et al. Preterm Gut Microbiome Depending on Feeding Type: Significance of Donor Human Milk. *Front Microbiol*. 2018;9:1376. doi: 10.3389/fmicb.2018.01376

12. Neu J, Mshvildadze M, Mai V. A roadmap for understanding and preventing necrotizing enterocolitis. *Current Gastroenterology Reports*. 2008;10(5):450-7. doi: 10.1007/s11894-008-0084-x

13. Mihi B, Good M. Impact of Toll-Like Receptor 4 Signaling in Necrotizing Enterocolitis. *Clinics in perinatology*. 2019;46(1):145-57. doi: 10.1016/j.clp.2018.09.007

14. Stewart CJ, Embleton ND, Marrs EC, Smith DP, Nelson A, Abdulkadir B, et al. Temporal bacterial and metabolic development of the preterm gut reveals specific signatures in health and disease. *Microbiome*. 2016;4(1):67. doi: 10.1186/s40168-016-0216-8

15. Fanaro S, Chierici R, Guerrini P, Vigi V. Intestinal microflora in early infancy: composition and development. *Acta Paediatr Suppl*. 2003;91(441):48-55. doi: 10.1111/j.1651-2227.2003.tb00646.x

16. Ahrne S, Lonnermark E, Wold AE, Aberg N, Hesselmar B, Saalman R, et al. Lactobacilli in the intestinal microbiota of Swedish infants. *Microbes Infect*. 2005;7(11-12):1256-62. doi: 10.1016/j.micinf.2005.04.011

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

17. Gewolb IH, Schwalbe RS, Taciak VL, Harrison TS, Panigrahi P. Stool microflora in extremely low birthweight infants. *Arch Dis Child Fetal Neonatal Ed.* 1999;80(3):F167-73. doi: 10.1136/fn.80.3.f167
18. Goldmann DA, Leclair J, Macone A. Bacterial colonization of neonates admitted to an intensive care environment. *The Journal of pediatrics.* 1978;93(2):288-93. doi: 10.1016/s0022-3476(78)80523-x
19. Dani C, Biadaoli R, Bertini G, Martelli E, Rubaltelli FF. Probiotics feeding in prevention of urinary tract infection, bacterial sepsis and necrotizing enterocolitis in preterm infants. A prospective double-blind study. *Biology of the neonate.* 2002;82(2):103-8. doi: 10.1159/000063096
20. Lin HC, Su BH, Chen AC, Lin TW, Tsai CH, Yeh TF, et al. Oral probiotics reduce the incidence and severity of necrotizing enterocolitis in very low birth weight infants. *Pediatrics.* 2005;115(1):1-4. doi: 10.1542/peds.2004-1463
21. Bin-Nun A, Bromiker R, Wilschanski M, Kaplan M, Rudensky B, Caplan M, et al. Oral probiotics prevent necrotizing enterocolitis in very low birth weight neonates. *The Journal of pediatrics.* 2005;147(2):192-6. doi: 10.1016/j.jpeds.2005.03.054
22. Lau CS, Chamberlain RS. Probiotic administration can prevent necrotizing enterocolitis in preterm infants: A meta-analysis. *J Pediatr Surg.* 2015;50(8):1405-12. doi: 10.1016/j.jpedsurg.2015.05.008
23. Costeloe K, Hardy P, Juszczak E, Wilks M, Millar MR, Probiotics in Preterm Infants Study Collaborative G. *Bifidobacterium breve* BBG-001 in very preterm infants: a randomised controlled phase 3 trial. *Lancet.* 2016;387(10019):649-60. doi: 10.1016/S0140-6736(15)01027-2
24. Jacobs SE, Tobin JM, Opie GF, Donath S, Tabrizi SN, Pirotta M, et al. Probiotic effects on late-onset sepsis in very preterm infants: a randomized controlled trial. *Pediatrics.* 2013;132(6):1055-62. doi: 10.1542/peds.2013-1339
25. AlFaleh K, Anabrees J. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev.* 2014(4):CD005496. doi: 10.1002/14651858.CD005496.pub4
26. Plummer EL, Bulach DM, Murray GL, Jacobs SE, Tabrizi SN, Garland SM, et al. Gut microbiota of preterm infants supplemented with probiotics: sub-study of the ProPremis trial. *BMC Microbiol.* 2018;18(1):184. doi: 10.1186/s12866-018-1326-1
27. van den Akker CHP, van Goudoever JB, Shamir R, Domellof M, Embleton ND, Hojsak I, et al. Probiotics and Preterm Infants: A Position Paper by the European Society for Paediatric Gastroenterology Hepatology and Nutrition Committee on Nutrition and the European Society for Paediatric Gastroenterology Hepatology and Nutrition Working Group for Probiotics and Prebiotics. *J Pediatr Gastroenterol Nutr.* 2020;70(5):664-80. doi: 10.1097/MPG.0000000000002655
28. Hill CJ, Lynch DB, Murphy K, Ulaszewska M, Jeffery IB, O'Shea CA, et al. Evolution of gut microbiota composition from birth to 24 weeks in the INFANTMET Cohort. *Microbiome.* 2017;5(1):4. doi: 10.1186/s40168-016-0213-y
29. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, et al. Temporal development of the gut microbiome in early childhood from the TEDDY study. *Nature.* 2018;562(7728):583-8. doi: 10.1038/s41586-018-0617-x
30. Moles L, Gomez M, Heilig H, Bustos G, Fuentes S, de Vos W, et al. Bacterial diversity in meconium of preterm neonates and evolution of their fecal microbiota during the first month of life. *PLoS One.* 2013;8(6):e66986. doi: 10.1371/journal.pone.0066986
31. Cong X, Xu W, Janton S, Henderson WA, Matson A, McGrath JM, et al. Gut Microbiome Developmental Patterns in Early Life of Preterm Infants: Impacts of Feeding and Gender. *PLoS One.* 2016;11(4):e0152751. doi: 10.1371/journal.pone.0152751
32. Young GR, van der Gast CJ, Smith DL, Berrington JE, Embleton ND, Lanyon C. Acquisition and Development of the Extremely Preterm Infant Microbiota Across Multiple Anatomical Sites. *J Pediatr Gastroenterol Nutr.* 2020;70(1):12-9. doi: 10.1097/MPG.0000000000002549

33. Shao Y, Forster SC, Tsaliki E, Vervier K, Strang A, Simpson N, et al. Stunted microbiota and opportunistic pathogen colonization in caesarean-section birth. *Nature*. 2019;574(7776):117-21. doi: 10.1038/s41586-019-1560-1

34. Lyons KE, Ryan CA, Dempsey EM, Ross RP, Stanton C. Breast Milk, a Source of Beneficial Microbes and Associated Benefits for Infant Health. *Nutrients*. 2020;12(4). doi: 10.3390/nu12041039

35. Herrmann K, Carroll K. An exclusively human milk diet reduces necrotizing enterocolitis. *Breastfeed Med*. 2014;9(4):184-90. doi: 10.1089/bfm.2013.0121

36. Masi AC, Embleton ND, Lamb CA, Young G, Granger CL, Najera J, et al. Human milk oligosaccharide DSLNT and gut microbiome in preterm infants predicts necrotising enterocolitis. *Gut*. 2021;70(12):2273-82. doi: 10.1136/gutjnl-2020-322771

37. Embleton ND, Berrington JE, Dorling J, Ewer AK, Juszczak E, Kirby JA, et al. Mechanisms Affecting the Gut of Preterm Infants in Enteral Feeding Trials. *Frontiers in Nutrition*. 2017;4:14. doi: 10.3389/fnut.2017.00014

38. Yu Z, Morrison M. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques*. 2004;36(5):808-12. doi: 10.2144/04365ST04

39. Smart KF, Aggio RB, Van Houtte JR, Villas-Boas SG. Analytical platform for metabolome analysis of microbial cells using methyl chloroformate derivatization followed by gas chromatography-mass spectrometry. *Nat Protoc*. 2010;5(10):1709-29. doi: 10.1038/nprot.2010.108

40. Beghini F, McIver LJ, Blanco-Miguez A, Dubois L, Asnicar F, Maharjan S, et al. Integrating taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3. *Elife*. 2021;10. doi: 10.7554/eLife.65088

41. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic assembler. *Genome Res*. 2017;27(5):824-34. doi: 10.1101/gr.213959.116

42. Lozupone CA, Hamady M, Kelley ST, Knight R. Quantitative and qualitative beta diversity measures lead to different insights into factors that structure microbial communities. *Appl Environ Microbiol*. 2007;73(5):1576-85. doi: 10.1128/AEM.01996-06


43. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, et al. Multivariable association discovery in population-scale meta-omics studies. *PLoS Comput Biol*. 2021;17(11):e1009442. doi: 10.1371/journal.pcbi.1009442

44. Olm MR, Crits-Christoph A, Bouma-Gregson K, Firek BA, Morowitz MJ, Banfield JF. inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nat Biotechnol*. 2021;39(6):727-36. doi: 10.1038/s41587-020-00797-0

45. Heirendt L, Arreckx S, Pfau T, Mendoza SN, Richelle A, Heinken A, et al. Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v.3.0. *Nat Protoc*. 2019;14(3):639-702. doi: 10.1038/s41596-018-0098-2

Figure Captions:

Figure 1. Sample collection timeline. Samples were collected from birth to discharge. Timing of discharge was determined clinically and was variable for each infant so no definitive timeframe was set. Solid bars represent timelines for regular sample collection. Dashed bars represent the “window of opportunity” for sample collection for non-regular samples. Maternal perinatal samples, collected within 4 days of delivery, included stool, oral, skin and vaginal swabs.



BMJ Open

A Longitudinal Observational Study Protocol: Preterm Infants: Microbiome Establishment, Neuro-CrossTalk and Origins (PIMENTO)

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-075060.R2
Article Type:	Protocol
Date Submitted by the Author:	10-Aug-2023
Complete List of Authors:	Healy, David; University College Cork, Department of Paediatrics and Child Health Wang, Shuo; University College Cork Grimaud, Ghjuvan; Moorepark Food Research Centre Warda, Alicja; Teagasc Food Research Centre Moorepark Ross, Paul; University College Cork, APC Microbiome Ireland Stanton, Catherine; University College Cork APC Microbiome Institute Dempsey, E; University College Cork, Department of Paediatrics and Child Health
Primary Subject Heading:	Paediatrics
Secondary Subject Heading:	Research methods
Keywords:	Microbiology < NATURAL SCIENCE DISCIPLINES, Neonatal intensive & critical care < INTENSIVE & CRITICAL CARE, NEONATOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A Longitudinal Observational Study Protocol: Preterm Infants: Microbiome Establishment, Neuro-CrossTalk and Origins (PIMENTO)

Authors: David B. Healy ^{1,2}, Shuo Wang ¹, Ghjuvan Grimaud ³, Alicja K. Warda ^{1,3}, R. Paul Ross ^{1,3}, Catherine Stanton ^{1,2,4}, and Eugene M. Dempsey ^{1,2,4,5}

- ¹ APC Microbiome Ireland, Cork, T12 YT20, Ireland
- ² Department of Paediatrics and Child Health, University College Cork, T12 YT20, Ireland
- ³ Teagasc, Moorepark Food Research Centre, Fermoy, P61 C996, Ireland
- ⁴ University College Cork, Cork, T12 YT20, Ireland
- ⁵ INFANT Research Centre, University College Cork, T12 DC4A, Ireland

Journal: BMJOpen

Corresponding Author: David B. Healy, APC Microbiome Ireland; Department of Paediatrics and Child Health, david.healy@ucc.ie

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Abstract

Introduction

Very preterm infants are at risk of abnormal microbiome colonisation in the first weeks to months of life. Several important associated factors have been identified including gestational age, mode of delivery, antibiotic exposure and feeding. Preterm infants are at risk of a number of pathologies for which the microbiome may play a central role, including necrotising enterocolitis (NEC) and sepsis. The objective of this study is to determine detailed microbiome changes that occur around implementation of different management practices including empiric antibiotic use, advancement of feeds and administration of probiotics during admission to the neonatal intensive care unit.

Methods and analysis

A single-site, longitudinal observational study of infants born less than 32 weeks gestation, including collection of maternal samples around delivery and breastmilk and infant samples from admission through discharge from the neonatal unit.

Ethics and dissemination

The protocol was approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals (CREC). The trial has been registered at ClinicalTrials.gov (NCT05803577).

The findings from this study will be disseminated in peer-reviewed journals, during scientific conferences, and directly to the study participants. Sequencing data will be deposited in public databases.

Strengths and limitations of this study

- Study design will provide high resolution longitudinal microbiome data that will help to characterise the colonising microbiome of preterm infants, the effect of and progression after treatments.
- An array of samples will be collected including rectal swabs, stool, urine and blood from infants, peripartum maternal samples including stool, oral, skin, vaginal samples and expressed breastmilk from mothers.
- This is a single-site study.
- The study population is heterogenous, and the recruited sample size will be ~200 (mothers and infants).

Abbreviations:

CREC	Clinical Research Ethics Committee of the Cork Teaching Hospitals
CRF	Case report form
CUMH	Cork University Maternity Hospital
GC-MS	Gas chromatography-mass spectrometry
GDPR	General Data Protection Regulation
HSE	Health Service Executive
IMP	Investigational medicinal product
LC-MS	Liquid chromatography-mass spectrometry
NEC	Necrotising enterocolitis
NICU	Neonatal intensive care unit
PCR	Polymerase chain reaction
RBB+C	Repeated bead beating plus column
TLR	Toll-like receptor

Keywords: preterm infants, microbiome, probiotics

1. Introduction

Over the last two decades, the materno-feto-neonatal microbiome has rapidly become an area of budding interest in the specialty of neonatology. While there is much debate as to whether a developing fetus is exposed to microorganisms in utero (1), it is increasingly evident that the gut microbiome does not expand and diversify until the postnatal period at which point microbes from the mother and environment begin colonisation of the infant. Following birth, an initial relatively simple microbial community develops into a diverse microbial ecosystem in a controlled fashion over the first years of life (2). Bacterial colonisation, specifically in very low birth weight infants, may occur in an aberrant fashion for numerous reasons: an immature gut and immune system (3, 4); the environment in which they are nursed(5); frequent exposure to antimicrobial agents(6, 7, 8); and altered feeding practices such as delayed commencement of feeds or using formula in place of mother's own milk(9, 10, 11). It is thought that the colonising microbiome of very preterm infants, and the factors that shape it, play a crucial role in the development of microbiome-mediated complications such as NEC.

NEC is a complex multifactorial condition characterised by intestinal immaturity and dysbiosis in association with altered microvascular blood flow. It is the most commonly acquired serious gastrointestinal complication of prematurity and is rare in babies born above 32 weeks (12). Infants at risk are likely to have deficient intestinal mucus production, immature gut immunity, and reduced endogenously-produced antimicrobial factors. The ability of bacteria to adhere to epithelial cells is thereby enhanced, particularly after gut injury secondary to ischaemia/hypoperfusion. Although not fully understood, the most accepted theory for the pathogenesis of NEC is that in this setting, bacterial endotoxin binds to toll-like receptor 4 (TLR4) found on the intestinal epithelial cells and by so doing, pathogen-associated molecular pattern receptors are activated, resulting in cell apoptosis and epithelial injury (13). This leads to an intense inflammatory cascade in the exposed lamina propria, including activation of inflammatory cytokines such as tumour necrosis factor alpha, interleukin-1 and interleukin-8, and ultimately necrosis of the surrounding tissue. Work by Stewart et al.(14) demonstrated that NEC tends not to occur in infants with a diverse gut microbiome. This may relate to inherent factors that make the individual less susceptible to NEC and simultaneously allow a diverse microbiome or may be due to protective effects associated with having a diverse gut microbiome, such as inflammatory modulation and improved gut epithelial integrity.

With early descriptions of the "ideal" gut microbiome (i.e. that of the vaginally delivered, exclusively breast-fed term infant, and without antibiotic treatment)(15, 16) and the identification of potentially deleterious bacteria harboured in the guts of the hospitalised very preterm infants (15, 17, 18) came the advent of probiotics. Administration of these as a supplement to feeding aimed to lend some "normality" to the gut microbiome of these vulnerable babies by supplanting potentially pathogenic bacteria and attempting to reduce the incidence of NEC (19, 20, 21, 22, 23). While probiotic preparations are widely available, shown to be safe (24, 25) and do, indeed, increase so-called beneficial bacteria present in the

gut(26), the evidence for their benefit is contentious (27) and practice varies from institution to institution.

Work has shown that the full term vaginally delivered and exclusively breastfed baby’s initial gut microbiome is dominated primarily by Actinobacteria (28, 29) while studies looking at preterm infants’ microbiome have shown a Proteobacteria (Bacilli dominance) and Firmicutes predominance (28, 30, 31, 32). Clarification of the initial neonatal gut microbiome in the preterm population is confounded by the low bacterial biomass in early samples and high interindividual variability in composition. Individual medical management of each infant is likely to cause perturbations to the microbiome. Several factors associated with alterations in the neonatal gut microbiome have previously been identified. These include gestational age (28) mode of delivery(28, 33), exposure to antibiotics (both pre- and postnatally) (6) and, importantly, feeding choice (34). Mothers of very low birth weight infants are typically encouraged to express breastmilk for their baby. This is associated with lower rates of sepsis and NEC (35). Some of these effects may be related to the bacteria or supporting molecules received in breastmilk from their mother (36). Correlations between changes in the neonatal gut microbiome and the microbiota of the milk they are being fed are limited and longitudinal analysis of breastmilk in this regard is lacking. Furthermore, interactions within breastmilk between bacteria and bacteriophage are not well characterised.

Given the rarity of some neonatal diseases, such as NEC, collaborative efforts are necessary to study data from a sufficient number of infants to draw robust conclusions (37). A number of prospective longitudinal repositories have been established, including the Great North Neonatal Biobank in the UK and the NEC Biorepository in the US, with aims including the study and investigation of the preterm neonatal microbiome and it’s role in pathological disease processes such as NEC. Work stemming from these and others has considerably advanced our understanding of the microbiome in term and preterm infants over the past decade. Expanding data and sample collection to other regions will help in future collaborative work, to account for geographical variations in hospital practice and microbiome structure. This work will aim to longitudinally characterise the early and colonising intestinal microbiome of very preterm infants. We will analyse samples from infants less than 32 weeks gestational age at birth, with collection occurring longitudinally from birth to discharge from the neonatal intensive care unit (NICU). Microbiome composition will be evaluated using metagenomics and 16sRNA sequencing of stool. Bacterial function will be assessed longitudinally using untargeted metabolomics of infant urine. We will also interrogate some key perinatal maternal microbiome niches, including oral, vaginal, skin and intestinal, as well as expressed breastmilk to evaluate mother-infant transmission.

2. Methods

2.1. Study design

This single site study is a prospective longitudinal observational study of the intestinal microbiome of infants born between 23 and 32 weeks of gestation. It will investigate the establishment of the very premature infants' microbiome using culture independent approaches, including the potential initial colonisation process of vertical transmission from mother to preterm infant, the apparent differences in establishment pattern based on feeding type, and the perturbations and recovery associated with different intensive care management strategies such as empiric antibiotic usage.

Study recruitment commenced in April 2020 and finished in December 2022. The primary endpoint of the study will be reached after the last enrolled participant completes 2-year neurodevelopmental follow-up.

2.2. Participant selection

Participants are infants, born between 23+0 weeks gestation and 31+6 weeks gestation, admitted to the Cork University Maternity Hospital (CUMH) NICU and their mothers.

The study was designed as a longitudinal observational study with a time-limited recruitment period of approximately 18 months. With a projected number of 100 infants less than 32 weeks to be admitted to the CUMH NICU annually, and an expected consent rate of 60-70%, a pragmatically pre-defined minimum enrolment number of 80 participants over 18 months of recruitment was set.

2.2.1. Inclusion and exclusion criteria

To be eligible for the study, the participants must meet the terms of the inclusion and exclusion criteria as presented in Table 1.

Table 1. Inclusion and exclusion criteria of the PIMENTO study.

Inclusion criteria	Exclusion criteria
1. Birth at less than 32 weeks gestational age and admitted to the NICU	1. No informed consent
2. Free from antenatal suspicion of major congenital abnormalities	2. Major congenital anomalies
3. Ability of the participant (in the investigator's opinion) to comprehend the full nature and purpose of the study including possible risks and side effects.	3. Gastrointestinal anomalies

4. Consent to participate in the study and willing to comply with the protocol and study restrictions.	4. Inborn errors of metabolism
--	--------------------------------

2.3. Recruitment

Pregnant women will be approached if there is a possibility of delivery before 32 weeks gestation. The study will be explained, and a participant information leaflet will be provided. Written informed consent will be obtained if the pregnant woman is agreeable after a sufficient amount of time to assimilate the information.

Multiple births will be enrolled separately. Samples will be collected from each infant but only one set of maternal samples would be collected. If one twin is delivered vaginally and the other by caesarean section, both skin and vaginal swabs will be collected from the mother if possible.

Written informed consent will be required for all participants. However, in exceptional circumstances (e.g. precipitous delivery) participants may be enrolled with deferred consent with the intent that investigators would obtain written consent as soon as possible thereafter. This would facilitate in the event a mother was not available to provide written informed consent that samples could be collected, i.e. a rectal swab from infant could be taken on admission to the NICU (as additional to the routine clinical swabs) and that, if passed a sample of the first meconium could be retained to ensure that, if subsequently enrolled, completeness of data for that participant could be maintained. Samples would be stored in a freezer on-site and would not be processed or analysed until written informed consent had been obtained. Similarly, out-born infants transferred to the study site may be enrolled with deferred consent and the infant’s first meconium and rectal swab retained until the mother is available to provide written informed consent. Again, sample processing or analysis would not be performed without written informed consent and the samples would remain frozen on site until that time.

2.4. Follow up

Two-year neurodevelopmental assessment, using the Bayley Scale for Infant and Toddler Development, 3rd Edition (BSID-III), is routinely performed at CUMH for all infants born at 32 weeks gestational age or less.

2.5. Compensation

There is no compensation provided to the participants. There are no cost implications for the Health Service Executive (HSE) or to the participants. The management of patients and investigative tests will comply with current standards of care.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

2.6. Study timeline

Each participant will have samples collected during the entirety of their neonatal unit stay (Section 2.9 and Figure 1). Sample collection will not occur if the infant is transferred to another centre for ongoing management.

2.7. Participant withdrawal/exclusion

Under the Declaration of Helsinki, the researcher will explain to the participant that they have the right to withdraw from the study at any time and that this will in no way prejudice their future treatment. The reason for withdrawal will be recorded in the source documents and on the appropriate CRF. Withdrawn participants will be replaced. With consent samples from individuals withdrawing from the study will be stored and may be analysed in the study. Participants excluded from the study after consenting will be replaced.

2.8. Ethics and dissemination

The study is conducted following the version Fortaleza, Brazil, October 2013 of the Declaration of Helsinki 1964. The Protocol and the Informed Consent Form have been approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals (CREC) before commencement (approval letter ECM 4 (t) 10/11/2020 & ECM 3 (mmm) 09/03/2021). On 25th August 2022 Protocol version 5 has been approved by the CREC (approval letter ECM 4 (t) 10/11/2020 & ECM 3 (vvv) 20/09/2022).

If a protocol amendment is necessary, this will be prepared with the agreement of the chief investigator and signed by the relevant parties. If the amendment is substantial, it will be submitted to the CREC and possibly other public bodies according to local requirements for review and approval. The protocol amendment will not be implemented before the required approvals are obtained.

The trial is sponsored by University College Cork (College Road, Cork T12 K8AF, +353 (0)21 490 3000). The sponsor is not involved in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit for publication. The trial has been registered at ClinicalTrials.gov (NCT05803577).

The findings from this study will be disseminated in peer-reviewed journals, during scientific conferences, and directly to the study participants.

2.9. Patient and Public Involvement

None

2.10. Objectives and Outcomes

2.10.1. Primary Objective

Detailed characterisation of the very preterm microbiome from birth to final hospital discharge, using culture independent approaches, including maternal origin and initial colonisation, postnatal alterations secondary to medical management (e.g. antibiotic and probiotic administration) and recovery/progression after discontinuation.

- a) To describe changes in community structure and function of gut microbiome in preterm infants during stay in the neonatal unit
- b) To compare microbiome successional colonisation in infants requiring varying degrees of initial respiratory support.
- c) To evaluate probiotic strains in the gut microbiome using next generation sequencing during probiotic commencement, colonisation and persistence after discontinuation

2.10.2. Secondary Objectives

- a) To determine changes in the human milk microbiome using culture independent approaches over time with increasing post menstrual age and characterise alterations in microbial content and investigate the influence of breastmilk-derived microbes in the infant gut microbiome.

2.10.3. Exploratory Objectives

- a) To assess metabolic functioning and systemic indicators in relation to microbiome composition at the time of sampling.
- b) To analyse maternal-infant transmission by correlation of the maternal microbiome with the early colonising microbiome of enrolled infants.

2.11. Sample collection

An overview of the sample collection timeline is presented in Figure 1.

2.11.1. Stool samples and rectal swabs

Rectal swabs are taken routinely on admission to the NICU for clinical microbiological screening. These are carried out by the admitting clinical nurse. One extra rectal swab will be taken for microbiome analysis at this time. Stool sampling will be carried out at the cotside during the course of the neonatal stay, a sample of the first meconium passed will be collected from the nappy and thereafter stool will be sampled from the infant’s nappy bi-weekly during routine cares. After 4 weeks of age, samples will be collected fortnightly. Samples are placed in a freezer in the neonatal unit immediately after sample collection. Thereafter samples are transferred to a -20°C freezer on-site for cataloguing prior to transfer to the off-site laboratory (APC Microbiome, Ireland, University College Cork, Ireland) for further processing and storage.

2.11.2. Urine samples

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

During the inpatient stay, urine will be collected weekly, if possible, the attending neonatal nurse by placing sterile cotton balls in the infant's nappy which will then be removed at the next occurrence of routine cares. Urine is squeezed from the cotton balls into sterile collection bottles. Urine samples are placed in neonatal unit freezer immediately after collection, subsequently catalogued and transferred to the on-site -20°C freezer, and finally transferred to the off-site laboratory.

2.11.3. Blood samples

Infant blood samples will be collected from infants at times when clinical blood sampling is being performed. These will be collected in Serum Gel Z micro sample tubes. A sample volume of 1% of the estimated blood volume (estimated blood volume = 80mls/kg) will be collected. The sample will be immediately centrifuged on-site for 8 minutes at 9000G and then frozen at -20°C in the on-site freezer. Thereafter, the blood sample will be transferred to the off-site laboratory for further storage and processing. If possible, two to three blood samples will be collected during the entire neonatal stay.

2.11.4. Breastmilk samples

In the case where mothers are expressing milk for their infants, breastmilk samples will be collected once there is enough supply to provide for the infant's intake and surplus for storage. Breastmilk is expressed by mother's either on -site in the neonatal unit or at home into sterile collection bottles. Expressed breastmilk is refrigerated immediately and can be stored refrigerated for up to 24 hours or frozen. 5-15 mls of breastmilk is collected from the refrigerated breastmilk storage bottles using aseptic technique and transferred to sterile cryogenic storage tubes. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter. If possible, three breastmilk samples will be collected during the entire neonatal stay.

2.11.5. Maternal Stool samples

Within three days of delivery, mothers will be approached to provide a sample from their first stool passed after delivery. Once collected, the sample will be brought by the mother to the neonatal unit when it will be placed in the neonatal unit freezer. Samples will be collected from there to be catalogued and transferred to an on-site -20°C freezer, and thereafter, to the off-site laboratory.

2.11.6. Maternal Oral samples

Oral swabs will be collected from mothers within three days after birth. Samples will not be collected within one-hour after mother's have eaten food. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.11.7. Maternal Skin swabs

In the case of infants delivered by caesarean section, skin swabs will be collected from mothers within three days after birth. Swabs are wetted with sterile water for injection prior to swabbing the skin of the left wrist. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.11.8. Maternal Vaginal swabs

In the case of infants born by vaginal delivery mothers will be asked within three days of delivery to provide a vaginal swab. Samples are catalogued and placed in an on-site -20°C freezer immediately after collection, and subsequently transferred to the off-site laboratory thereafter.

2.12 Sample analysis

Faecal samples will be processed for DNA extraction and metabolomics. DNA will be extracted from stool using the modification of the Repeated Bead Beating Plus Column (RBB+C) method (38) and subjected to 16S rRNA and/or shotgun sequencing. Metabolomics of infant’s faeces will be analysed by the gas chromatography–mass spectrometry (GC-MS) technique using methyl chloroformate derivatisation (39). Bioinformatic analysis will identify the microbial composition at the phylum, genus and family levels using both read-based (40) and assembly-based (41) approaches. Metagenomic data will be correlated with physiological and clinical parameters and biomarkers. Advanced statistical and bioinformatics approaches will be applied to identify modifiable environmental factors that influence bacterial groups/consortia. This includes diversity analysis (42), differential abundance analysis (43), strain persistence and vertical transmission (44) and genome-scale metabolic modelling (45).

Breast milk samples will be used for DNA extraction (using modification of Qiagen DNeasy Powerfood Microbial Kit (QIAGEN Ltd, Manchester, United Kingdom) and the milk microbiome composition will be analysed by using state-of-the-art methods, possibly including but not limited to sequencing, and quantitative PCR.

Urine metabolomics analysis will be analysed by the liquid chromatography–mass spectrometry (LC-MS) technique (28).

The maternal oral microbiome composition will be analysed by using state-of-the-art methods after DNA extraction, possibly including but not limited to sequencing, and quantitative PCR. The bioinformatic analysis will identify the microbial composition at the phylum, genus, and family levels.

From vaginal swabs, DNA will be extracted using the modification of the RBB+C method (38) and the vaginal microbiome composition will be analysed by using state-of-the-art

methods, possibly including but not limited to sequencing, and quantitative PCR. The bioinformatic analysis will identify the microbial composition at the phylum, genus, and family levels.

2.13. Adverse events and participant well-being

This is an observational study. No investigational medicinal product (IMP) will be administered to study participants at any stage during the study. The study procedures are not greater than minimal risk, adverse events and serious adverse events are not expected. All self-reported adverse events (AEs) will be listed documenting duration, severity, participant outcome, and if any therapy was required. The study Chief investigator will review these and advance any reports to the local Ethics Committee if it is deemed necessary.

2.14. Data collection and management

2.14.1 Data collection/Case Report Forms (CRFs)

Data will be collected throughout the neonatal stay and from clinic visits after discharge. Data will be extracted from the electronic health records of each subject meeting the eligibility criteria and being included in the study. Data will be recorded in an electronic spreadsheet data collection tool stored on an encrypted password protected computer and backed up to the secure university server. All study staff responsible for entering data into the data collection tool will be trained prior to the start-up of the study.

The data in the CRFs will be consistent with the relevant source documents. The only source documents that will be available will be participants' in-patient electronic health record at CUMH. All data stored will be pseudonymised and treated with strict confidentiality in accordance with the General Data Protection Regulation (GDPR).

2.14.2 Sequencing data

Data types and formats collected and/or produced will include Raw whole genome sequencing (WGS) data (fastq), WGS assembly files and excel file with strains name (fasta), WGS annotation files (fna, gff, faa, ffn), 16S raw data (fastq), 16S ASV table (csv), 16S "phyloseq" file, which includes metadata for each sample (rds), Raw shotgun data (fastq), MAGs/bins from shotgun data (fna), Pangenomes files (newick, fa, csv/tab), and pipeline for the metabolic modelling part and generated data (sbml, mat). Sequencing data will be deposited in public databases.

2.15. Statistical analysis

2.15.1. Sample size justification

This study was designed as a longitudinal observational trial with the primary aim of detailed characterisation of the colonising microbiome of very preterm infants and, additionally,

determination of factors relating to perturbations seen therein. As such, there was no statistically predefined sample size but instead the goal to recruit the largest sample population possible in the allotted timeframe for study recruitment, in this case 18-20 months. Post-hoc, we will calculate and state statistical power analysis to give a representation of statistically relevant sample size for this cohort.

3. Discussion

This study aims to longitudinally observe the gut microbiome of very preterm infants during their admission to Cork University Maternity Hospital neonatal unit. We intend to characterise in detail how clinical management practices influence the microbiome of these infants, including but not limited to delivery practices, antibiotic usage, feeding types and regimens, and probiotic usage. Very preterm infants have increased risk of potentially microbiome-mediated pathologies such as NEC and late-onset sepsis. The role that the microbiome plays in these diseases may relate to the overall composition of the microbiome, individual species within the microbiome, or temporal changes that occur. As such, management practices may be influential in the development of microbiome structure and may therefore have an indirect role in the development of these diseases.

At CUMH probiotics (*Bifidobacterim bifidum* and *Lactobacillus acidophilus*) are administered to all infants born less than 32 weeks gestation or less than 1500g birthweight from when feeding is deemed to be tolerated until 34 weeks corrected gestational age. We intend to observe these probiotic strains within the gut microbiome community, how they persist after supplementation is discontinued, and how clinical management may affect their colonisation.

Collected maternal samples will allow us to study transmission of microbes to premature infants and longitudinally collected breastmilk samples will allow us to study how the milk microbiome plays a role in shaping gut microbiome structure of the preterm infant.

The authors acknowledge the limitations of this study design. It will be a single centre observational study and the results may not be generalisable across all other institutions. As all infants that are to be enrolled in the study will receive probiotic supplementation, there will not be a control group without probiotics with which to compare. The study is designed as descriptive and without intervention. Accordingly, the management of the infants enrolled has not been standardised. While the clinical demographics of the enrolled infants will be described in careful detail, all confounders may not be accounted for owing to variability in clinical courses and the variability in management decisions made by the various providers caring for the infants.

We envisage that this work will add to the already existing information on the preterm microbiome. As neonatal care practices can differ markedly from institution to institution we hope that this study will provide novel insights into how the microbiome is established, including maternal transmission of microbes at delivery and through breastmilk. This study

will inform our understanding of microbial colonisation of very preterm infants, successional patterns of colonisation and how clinical management practices influence these. It will also provide valuable information on probiotic colonisation and practices around probiotic administration and may help future development of efficacious probiotic strains.

Status of Study

The study is ongoing as of 7th April 2021. Prospective recruitment of new participants was completed on 31st December 2022. The primary endpoint will be when 2-year neurodevelopment follow up has been completed for the final participant. At the time of publication, no sequencing data has been received or analysed.

Competing interests

The authors have no known competing financial interests.

Author's contributions

D.B.H. wrote the manuscript. D.B.H., R.P.R., C.S., E.M.D. were involved in study conception and design, acquisition of ethical approval, and review of the manuscript. D.B.H. coordinated participant recruitment, sample collection, on-site sample processing and sample transport. S.W. coordinated laboratory sample processing in conjunction with A.K.W.. G.G. coordinated bioinformatic analysis. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank all the study participants and the neonatal nursing staff in Cork University Maternity Hospital.

Funding

This publication has emanated from research conducted with financial support of Science Foundation Ireland (SFI) under Grant No. 12/RC/2273_P2 and 19/SP/6989.

References

1. Kennedy KM, De Goffau MC, Perez-Muñoz ME, Arrieta M-C, Bäckhed F, Bork P, et al. Questioning the fetal microbiome illustrates pitfalls of low-biomass microbial studies. *Nature*. 2023;613(7945):639-49. doi: 10.1038/s41586-022-05546-8

2. Adlerberth I, Wold AE. Establishment of the gut microbiota in Western infants. *Acta paediatrica*. 2009;98(2):229-38. doi: 10.1111/j.1651-2227.2008.01060.x

3. Fança-Berthon P, Michel C, Pagniez A, Rival M, Van Seuningen I, Darmaun D, et al. Intrauterine Growth Restriction Alters Postnatal Colonic Barrier Maturation in Rats. *Pediatric Research*. 2009;66(1):47-52. doi: 10.1203/pdr.0b013e3181a2047e

4. Zhang W, Ma C, Xie P, Zhu Q, Wang X, Yin Y, et al. Gut microbiota of newborn piglets with intrauterine growth restriction have lower diversity and different taxonomic abundances. *Journal of Applied Microbiology*. 2019;127(2):354-69. doi: 10.1111/jam.14304

5. Brooks B, Firek BA, Miller CS, Sharon I, Thomas BC, Baker R, et al. Microbes in the neonatal intensive care unit resemble those found in the gut of premature infants. *Microbiome*. 2014;2(1):1. doi: 10.1186/2049-2618-2-1

6. Fouhy F, Guinane CM, Hussey S, Wall R, Ryan CA, Dempsey EM, et al. High-throughput sequencing reveals the incomplete, short-term recovery of infant gut microbiota following parenteral antibiotic treatment with ampicillin and gentamicin. *Antimicrobial Agents & Chemotherapy*. 2012;56(11):5811-20. doi: 10.1128/AAC.00789-12

7. Hussey S, Wall R, Gruffman E, O'Sullivan L, Ryan CA, Murphy B, et al. Parenteral antibiotics reduce bifidobacteria colonization and diversity in neonates. *Int J Microbiol*. 2011;2011. doi: 10.1155/2011/130574

8. Zeissig S, Blumberg RS. Life at the beginning: perturbation of the microbiota by antibiotics in early life and its role in health and disease. *Nat Immunol*. 2014;15(4):307-10. doi: 10.1038/ni.2847

9. Cai C, Zhang Z, Morales M, Wang Y, Khafipour E, Friel J. Feeding practice influences gut microbiome composition in very low birth weight preterm infants and the association with oxidative stress: A prospective cohort study. *Free Radic Biol Med*. 2019;142:146-54. doi: 10.1016/j.freeradbiomed.2019.02.032

10. Cong X, Judge M, Xu W, Diallo A, Janton S, Brownell EA, et al. Influence of Feeding Type on Gut Microbiome Development in Hospitalized Preterm Infants. *Nurs Res*. 2017;66(2):123-33. doi: 10.1097/NNR.0000000000000208

11. Parra-Llorca A, Gormaz M, Alcantara C, Cernada M, Nunez-Ramiro A, Vento M, et al. Preterm Gut Microbiome Depending on Feeding Type: Significance of Donor Human Milk. *Front Microbiol*. 2018;9:1376. doi: 10.3389/fmicb.2018.01376

12. Neu J, Mshvildadze M, Mai V. A roadmap for understanding and preventing necrotizing enterocolitis. *Current Gastroenterology Reports*. 2008;10(5):450-7. doi: 10.1007/s11894-008-0084-x

13. Mihi B, Good M. Impact of Toll-Like Receptor 4 Signaling in Necrotizing Enterocolitis. *Clinics in perinatology*. 2019;46(1):145-57. doi: 10.1016/j.clp.2018.09.007

14. Stewart CJ, Embleton ND, Marrs EC, Smith DP, Nelson A, Abdulkadir B, et al. Temporal bacterial and metabolic development of the preterm gut reveals specific signatures in health and disease. *Microbiome*. 2016;4(1):67. doi: 10.1186/s40168-016-0216-8

15. Fanaro S, Chierici R, Guerrini P, Vigi V. Intestinal microflora in early infancy: composition and development. *Acta Paediatr Suppl*. 2003;91(441):48-55. doi: 10.1111/j.1651-2227.2003.tb00646.x

16. Ahrne S, Lonnermark E, Wold AE, Aberg N, Hesselmar B, Saalman R, et al. Lactobacilli in the intestinal microbiota of Swedish infants. *Microbes Infect*. 2005;7(11-12):1256-62. doi: 10.1016/j.micinf.2005.04.011

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

17. Gewolb IH, Schwalbe RS, Taciak VL, Harrison TS, Panigrahi P. Stool microflora in extremely low birthweight infants. *Arch Dis Child Fetal Neonatal Ed.* 1999;80(3):F167-73. doi: 10.1136/fn.80.3.f167
18. Goldmann DA, Leclair J, Macone A. Bacterial colonization of neonates admitted to an intensive care environment. *The Journal of pediatrics.* 1978;93(2):288-93. doi: 10.1016/s0022-3476(78)80523-x
19. Dani C, Biadaoli R, Bertini G, Martelli E, Rubaltelli FF. Probiotics feeding in prevention of urinary tract infection, bacterial sepsis and necrotizing enterocolitis in preterm infants. A prospective double-blind study. *Biology of the neonate.* 2002;82(2):103-8. doi: 10.1159/000063096
20. Lin HC, Su BH, Chen AC, Lin TW, Tsai CH, Yeh TF, et al. Oral probiotics reduce the incidence and severity of necrotizing enterocolitis in very low birth weight infants. *Pediatrics.* 2005;115(1):1-4. doi: 10.1542/peds.2004-1463
21. Bin-Nun A, Bromiker R, Wilschanski M, Kaplan M, Rudensky B, Caplan M, et al. Oral probiotics prevent necrotizing enterocolitis in very low birth weight neonates. *The Journal of pediatrics.* 2005;147(2):192-6. doi: 10.1016/j.jpeds.2005.03.054
22. Lau CS, Chamberlain RS. Probiotic administration can prevent necrotizing enterocolitis in preterm infants: A meta-analysis. *J Pediatr Surg.* 2015;50(8):1405-12. doi: 10.1016/j.jpedsurg.2015.05.008
23. Costeloe K, Hardy P, Juszczak E, Wilks M, Millar MR, Probiotics in Preterm Infants Study Collaborative G. *Bifidobacterium breve* BBG-001 in very preterm infants: a randomised controlled phase 3 trial. *Lancet.* 2016;387(10019):649-60. doi: 10.1016/S0140-6736(15)01027-2
24. Jacobs SE, Tobin JM, Opie GF, Donath S, Tabrizi SN, Pirotta M, et al. Probiotic effects on late-onset sepsis in very preterm infants: a randomized controlled trial. *Pediatrics.* 2013;132(6):1055-62. doi: 10.1542/peds.2013-1339
25. AlFaleh K, Anabrees J. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev.* 2014(4):CD005496. doi: 10.1002/14651858.CD005496.pub4
26. Plummer EL, Bulach DM, Murray GL, Jacobs SE, Tabrizi SN, Garland SM, et al. Gut microbiota of preterm infants supplemented with probiotics: sub-study of the ProPremis trial. *BMC Microbiol.* 2018;18(1):184. doi: 10.1186/s12866-018-1326-1
27. van den Akker CHP, van Goudoever JB, Shamir R, Domellof M, Embleton ND, Hojsak I, et al. Probiotics and Preterm Infants: A Position Paper by the European Society for Paediatric Gastroenterology Hepatology and Nutrition Committee on Nutrition and the European Society for Paediatric Gastroenterology Hepatology and Nutrition Working Group for Probiotics and Prebiotics. *J Pediatr Gastroenterol Nutr.* 2020;70(5):664-80. doi: 10.1097/MPG.0000000000002655
28. Hill CJ, Lynch DB, Murphy K, Ulaszewska M, Jeffery IB, O'Shea CA, et al. Evolution of gut microbiota composition from birth to 24 weeks in the INFANTMET Cohort. *Microbiome.* 2017;5(1):4. doi: 10.1186/s40168-016-0213-y
29. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, et al. Temporal development of the gut microbiome in early childhood from the TEDDY study. *Nature.* 2018;562(7728):583-8. doi: 10.1038/s41586-018-0617-x
30. Moles L, Gomez M, Heilig H, Bustos G, Fuentes S, de Vos W, et al. Bacterial diversity in meconium of preterm neonates and evolution of their fecal microbiota during the first month of life. *PLoS One.* 2013;8(6):e66986. doi: 10.1371/journal.pone.0066986
31. Cong X, Xu W, Janton S, Henderson WA, Matson A, McGrath JM, et al. Gut Microbiome Developmental Patterns in Early Life of Preterm Infants: Impacts of Feeding and Gender. *PLoS One.* 2016;11(4):e0152751. doi: 10.1371/journal.pone.0152751
32. Young GR, van der Gast CJ, Smith DL, Berrington JE, Embleton ND, Lanyon C. Acquisition and Development of the Extremely Preterm Infant Microbiota Across Multiple Anatomical Sites. *J Pediatr Gastroenterol Nutr.* 2020;70(1):12-9. doi: 10.1097/MPG.0000000000002549

33. Shao Y, Forster SC, Tsaliki E, Vervier K, Strang A, Simpson N, et al. Stunted microbiota and opportunistic pathogen colonization in caesarean-section birth. *Nature*. 2019;574(7776):117-21. doi: 10.1038/s41586-019-1560-1

34. Lyons KE, Ryan CA, Dempsey EM, Ross RP, Stanton C. Breast Milk, a Source of Beneficial Microbes and Associated Benefits for Infant Health. *Nutrients*. 2020;12(4). doi: 10.3390/nu12041039

35. Herrmann K, Carroll K. An exclusively human milk diet reduces necrotizing enterocolitis. *Breastfeed Med*. 2014;9(4):184-90. doi: 10.1089/bfm.2013.0121

36. Masi AC, Embleton ND, Lamb CA, Young G, Granger CL, Najera J, et al. Human milk oligosaccharide DSLNT and gut microbiome in preterm infants predicts necrotising enterocolitis. *Gut*. 2021;70(12):2273-82. doi: 10.1136/gutjnl-2020-322771

37. Embleton ND, Berrington JE, Dorling J, Ewer AK, Juszczak E, Kirby JA, et al. Mechanisms Affecting the Gut of Preterm Infants in Enteral Feeding Trials. *Frontiers in Nutrition*. 2017;4:14. doi: 10.3389/fnut.2017.00014

38. Yu Z, Morrison M. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques*. 2004;36(5):808-12. doi: 10.2144/04365ST04

39. Smart KF, Aggio RB, Van Houtte JR, Villas-Boas SG. Analytical platform for metabolome analysis of microbial cells using methyl chloroformate derivatization followed by gas chromatography-mass spectrometry. *Nat Protoc*. 2010;5(10):1709-29. doi: 10.1038/nprot.2010.108

40. Beghini F, Mclver LJ, Blanco-Miguez A, Dubois L, Asnicar F, Maharjan S, et al. Integrating taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3. *Elife*. 2021;10. doi: 10.7554/eLife.65088

41. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic assembler. *Genome Res*. 2017;27(5):824-34. doi: 10.1101/gr.213959.116

42. Lozupone CA, Hamady M, Kelley ST, Knight R. Quantitative and qualitative beta diversity measures lead to different insights into factors that structure microbial communities. *Appl Environ Microbiol*. 2007;73(5):1576-85. doi: 10.1128/AEM.01996-06

43. Mallick H, Rahnavard A, Mclver LJ, Ma S, Zhang Y, Nguyen LH, et al. Multivariable association discovery in population-scale meta-omics studies. *PLoS Comput Biol*. 2021;17(11):e1009442. doi: 10.1371/journal.pcbi.1009442

44. Olm MR, Crits-Christoph A, Bouma-Gregson K, Firek BA, Morowitz MJ, Banfield JF. inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nat Biotechnol*. 2021;39(6):727-36. doi: 10.1038/s41587-020-00797-0

45. Heirendt L, Arreckx S, Pfau T, Mendoza SN, Richelle A, Heinken A, et al. Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v.3.0. *Nat Protoc*. 2019;14(3):639-702. doi: 10.1038/s41596-018-0098-2

Figure Captions:

Figure 1. Sample collection timeline. Samples were collected from birth to discharge. Timing of discharge was determined clinically and was variable for each infant so no definitive timeframe was set. Solid bars represent timelines for regular sample collection. Dashed bars represent the “window of opportunity” for sample collection for non-regular samples. Maternal perinatal samples, collected within 4 days of delivery, included stool, oral, skin and vaginal swabs.

