



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Effectiveness Of Abdominal Massage For Chronic Constipation In The Elderly: A Protocol For A Systematic Review and Meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-074780
Article Type:	Protocol
Date Submitted by the Author:	17-Apr-2023
Complete List of Authors:	yuan, qiang; Chengdu University of Traditional Chinese Medicine Affiliated Hospital, Department of Tuina Wang, Xiaoyan; Hospital of Chengdu University of Traditional Chinese Medicine, Oncology Department Li, Zhou; Hospital of Chengdu University of Traditional Chinese Medicine, Oncology Department Li, Chuan; Sichuan Acupuncture School Lu, Qunwen; Hospital of Chengdu University of Traditional Chinese Medicine, Tuina Department Wang, Haozhong; Chengdu University of Traditional Chinese Medicine, Luo, Jian; Hospital of Chengdu University of Traditional Chinese Medicine, Tuina Department
Keywords:	Systematic Review, GASTROENTEROLOGY, GERIATRIC MEDICINE, Rehabilitation medicine < INTERNAL MEDICINE, Nursing Care

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Effectiveness Of Abdominal Massage For Chronic Constipation In The Elderly:  
A Protocol For A Systematic Review and Meta-analysis**

Qiang Yuan, MSc<sup>a†</sup>, Xiaoyan Wang, MSc<sup>b†</sup>, Zhou Li, BSc<sup>b†</sup>, Chuan Li,  
MSc<sup>c</sup>, Qunwen Lu, MSc<sup>a</sup>, Haozhong Wang, PhD<sup>d\*</sup>, Jian Luo, PhD<sup>a\*</sup>

<sup>a</sup> Tuina Department, Hospital of Chengdu University of Traditional Chinese Medicine,  
Chengdu, China

<sup>b</sup> Oncology Department, Hospital of Chengdu University of Traditional Chinese  
Medicine, Chengdu, China

<sup>c</sup> Sichuan acupuncture and moxibustion School, Chengdu, China

<sup>d</sup> College of Basic Medicine, Chengdu University of Traditional Chinese Medicine,  
Chengdu, China

\*Correspondence to authors: Haozhong Wang  
wanghaozhong@cdutcm.edu.cn  
Jian Luo  
lj6176@126.com

<sup>†</sup>Co-first authors: Qiang Yuan<sup>1†</sup>, Xiaoyan Wang<sup>2†</sup>, Zhou Li<sup>2†</sup>  
<sup>†</sup>These authors contributed equally to this work and share first authorship.

**ABSTRACT**

**Introduction** Chronic constipation is a health management challenge with high prevalence, difficult treatment, and risk of cardiac and cerebral events in elderly patients. Available laxative medications and lifestyle treatments are ineffective and dependent. Abdominal massage is an economical and effective therapeutic measure to improve chronic constipation in the elderly. The aim of this systematic review and meta-analysis is to evaluate the efficacy and safety of abdominal massage in elderly patients with chronic constipation. An in-depth subgroup analysis will be conducted to explore the factors influencing the efficacy of abdominal massage such as treatment

modality, duration, and frequency to provide credible evidence for future mechanistic studies.

**Methods and analysis** Electronic searches of clinical randomized controlled trials in Web of Science, PubMed, CINAHL, Cochrane Library, Embase, Airiti Library databases, Chinese National Knowledge Infrastructure Databases (CNKI), Chinese Science and Technology Periodical Database (VIP), Chinese Biomedical Literature Database (CBM), and Wan Fang Database will be conducted. Relevant data were extracted, and a meta-analysis was performed using Reviewer Manager 5.4. Quality and risk assessments of the included studies were performed, and the outcome indicators of the trials were observed.

**Results** This meta-analysis will objectively and comprehensively examine the efficacy and safety of abdominal massage in elderly patients with chronic constipation.

**Conclusions** This review will evaluate abdominal massage as a treatment for relieving symptoms and improving quality of life in the elderly with chronic constipation, and will provide additional insight for clinical treatment and mechanistic studies.

**Ethics and dissemination** As there will be no collection or generation of raw data, a statement of ethical approval is not required. We will publish the results of our study in peer-reviewed journals.

**PROSPERO registration number** CRD42023408629

**Keywords:** Abdominal Massage; Chronic constipation; Elderly patients; Systematic review

**Word Count** 2865

**Strengths and limitations of this study**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1. This is the first planned meta-analysis to assess the reliability of the evidence for abdominal massage in the treatment of chronic constipation in the elderly.
  - 2. This study will include elderly patients aged >60 years who meet the diagnostic criteria.
  - 3. Only English and Chinese trials were included, which may lead to an increased risk of bias.
  - 4. There may be a heterogeneity in the different methods of abdominal massage.
- Introduction

Chronic constipation (CC) is a disease characterized by difficult and hard stools, decreased frequency of bowel movements, and a feeling of incomplete defecation. The main diagnostic criteria are the Rome IV criteria and the patient's self-reported symptoms<sup>[1, 2]</sup>. The prevalence of chronic constipation was 16% for men and 26% for women aged 65 years and older, and among those aged 84 years and older, the prevalence could be as high as 26% and 34% for men and women, respectively<sup>[3]</sup>. Decreased contractile motility of the detrusor and smooth muscle of the colonic, and atrophy of the gastrointestinal mucosa with reduced fluid secretion are the mechanisms that make constipation more common in the elderly<sup>[4]</sup>. Aging-related changes in dietary and physical activity lifestyle, arthralgia, osteoporosis, and laxative dependence, long-term use of anticholinergic agents, opioid analgesics<sup>[5]</sup>, calcium supplements, and NSAIDs (non-steroidal anti-inflammatory drugs) are all risk factors for chronic constipation in the elderly<sup>[6]</sup>.

Constipation not only seriously affects the quality of life of the elderly, but can also cause many diseases and consume a large amount of health care resources. Fecal retention leads to gut microbiota disorders and the production of harmful flora metabolites inducing cardiovascular, neurocognitive and other diseases<sup>[7, 8]</sup>. Difficulty in defecation in elderly patients induces cardiovascular and cerebrovascular accidents and increases the risk of sudden death<sup>[9]</sup>. Meanwhile, CC also serves as a prelude to serious diseases such as colon cancer<sup>[10]</sup>. The main treatment for CC is the use of laxatives and lifestyle modification<sup>[11]</sup>. Drug therapy is prone to dependence causing

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

exacerbation of symptoms or complications. Existing treatments have not completely resolved the suffering of CC patients, thus finding economical and effective treatment methods is urgently needed.

Abdominal massage serves as the most convenient and economical alternative therapy for clinical application<sup>[12]</sup>. It has potential mechanisms to improve blood circulation in the gastrointestinal tract and stimulate gastrointestinal motility. Several meta-analyses have explored the effectiveness of abdominal massage in digestive function and bowel disorders<sup>[13-15]</sup>. However, there is still a lack of specific studies on elderly patients with constipation. With the increased prevalence of chronic constipation in the elderly and its specific pathological mechanisms, there is a necessity to explore it in an age-restricted manner.

## 2 Methods and analysis

### OBJECTIVES

To evaluate the efficacy and safety of abdominal massage in the treatment of chronic constipation in the elderly.

### METHODS

#### Study registration

This systematic review protocol was registered with PROSPERO 2023 (registration number: CRD42023408629). And the protocol report is in the base of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols (PRISMA-P) declaration guidelines<sup>[16]</sup>. The review will be performed in line with the PRISMA-P declaration guidelines.

#### Inclusion criteria for study selection

##### Type of study

All randomized controlled trials (RCTs) that fulfilled the inclusion criteria will be included in this systematic evaluation and meta-analysis to evaluate the safety and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

109 efficacy of abdominal massage therapy for the treatment of chronic constipation in the  
110 elderly.

111 **Type of participant**

112 Participants aged 60 years or older diagnosed with chronic constipation will be  
113 included regardless of gender, race, education or economic status. The diagnostic  
114 criteria for chronic constipation are:

- 115 1. Clinical diagnostic criteria (e.g., Rome IV criteria);
- 116 2. Diagnostic criteria defined by the authors or defined by the clinician;
- 117 3. Participants' self-reported constipation.

118 The exclusion criteria contain the following items:1) Cognitive disorder;2) Belonged  
119 to specific clinical population groups (e.g., pregnant women, ICU patients, post  
120 operation patients).

121 **Type of intervention and comparisons.**

122 Abdominal massage, including any type of pressure on the abdominal region, such as  
123 Swedish abdominal massage, acupuncture, and aromatic abdominal massage. There  
124 was no limit to the time or frequency of massage. Comparisons group can include no  
125 intervention, placebo intervention, positive control medicine.

126 **Type of outcome measure**

127 The primary outcomes examined were the frequency of bowel movements, stool  
128 texture and constipation symptoms. Any constipation symptom questionnaire, such as  
129 the Demographic Characteristics Questionnaire, the Rome criteria, and the  
130 Constipation Assessment Scale, measured constipation symptoms. Secondary  
131 outcomes included the influence of constipation on quality of life, laxative use, and  
132 adverse events in older patients.

## Search methods for identification of studies

### Electronic data sources

The following electronic databases will be searched from their respective inception dates to April 21, 2023: Web of Science, PubMed, CINAHL, Cochrane Library, Embase, Airtiti Library databases, Chinese National Knowledge Infrastructure Databases (CNKI), Chinese Science and Technology Periodical Database (VIP), Chinese Biomedical Literature Database (CBM), and Wan Fang Database. Excluding no language or publication restrictions, randomized clinical trials investigating the effectiveness of abdominal massage on chronic constipation in older adults that met eligibility criteria were investigated.

### Searching other resources

The reference lists of potentially missing eligible studies will be scanned and the relevant conference proceedings will be scanned as well.

### Search strategy

The search strategy for PubMed is shown in [Table 1](#). The following search keywords will be used: Massage (e.g., “Tuina” or “Chinese massage” or “Chuna” or “Shiatsu” or “Thai massage” or “Rubbing Abdomen” or “Massage” or “Zone Therapy” or “Therapies, Zone” or “Zone Therapies” or “Therapy, Zone” or “Massage Therapy” or “Massage Therapies” or “Therapies, Massage” or “Therapy, Massage” or “osteopathy” or “Fascial Manipulation”); “Constipation” (e.g., “Dyschezia” or “Colonic Inertia” or “Functional Constipation” or “Primary Constipation” or “Chronic Constipation” or “Idiopathic Constipation” or “Slow Transit Constipation” or “Constipated” or “Defecation Disorder” or “Evacuation Disorder\*” or “Gastrointestinal Transit” or “Gut Transit” or “Slow Transit” or “Hard Stool\*”) ; Randomized Controlled trial (e.g., “Controlled clinical trial” or “Random allocation” or “Randomized” or “Randomly” or

Commented [1]: 具体例子



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

158 “Double-blind Method” or “Single-blind Method” or “Clinical Trial”). The equivalent  
159 search keywords will be used in the Chinese databases.

Search number	Search Details
#1	"Constipation"[MeSH Terms]
#2	"Dyschezia"[Title/Abstract] OR "Colonic inertia"[Title/Abstract] OR "Functional constipation"[Title/Abstract] OR "Primary constipation"[Title/Abstract] OR "Chronic constipation"[Title/Abstract] OR "Idiopathic constipation"[Title/Abstract] OR "Slow transit constipation"[Title/Abstract] OR "Constipated"[Title/Abstract] OR "Defecation disorder"[Title/Abstract] OR "Evacuation Disorder*"[Title/Abstract] OR "Gastrointestinal Transit"[Title/Abstract] OR "Gut Transit"[Title/Abstract] OR "Slow Transit"[Title/Abstract] OR "Hard Stool*"[Title/Abstract] OR "Lumpy Stool*"[Title/Abstract]
#3	#1 OR #2
#4	"Massage"[MeSH Terms]
#5	"Tuina"[Title/Abstract] OR "Chinese massage"[Title/Abstract] OR "Chuna"[Title/Abstract] OR "Shiatsu"[Title/Abstract] OR "Thai massage"[Title/Abstract] OR "Rubbing Abdomen"[Title/Abstract] OR "Massage"[Title/Abstract] OR "Zone Therapy"[Title/Abstract] OR "Therapies, Zone"[Title/Abstract] OR "Zone Therapies"[Title/Abstract] OR "Therapy, Zone"[Title/Abstract] OR "Massage Therapy"[Title/Abstract] OR "Massage Therapies"[Title/Abstract] OR "Therapies, Massage"[Title/Abstract] OR "Therapy, Massage"[Title/Abstract] OR "osteopathy"[Title/Abstract] OR "Fascial Manipulation);
#6	#4 OR #5

#7	"Randomized Controlled Trial" [Publication Type]
#8	"Randomized Controlled Trial"[Title/Abstract] OR "Controlled Clinical Trial"[Title/Abstract] OR "Random Allocation"[Title/Abstract] OR "Randomized"[Title/Abstract] OR "Randomly"[Title/Abstract] OR "Double Blind Method"[Title/Abstract] OR "Single Blind Method"[Title/Abstract] OR "Clinical Trial"[Title/Abstract]
#9	#7 OR #8
#10	#3 AND #6 AND #9

Table 1 Search strategy for the PubMed database.

## Data collection and analysis

### Selection of studies

The titles and abstracts of all searched studies will be reviewed and screened independently by 2 reviewers (YQ and XYW), aiming at identifying eligible trials and eliminating duplicated or irrelevant studies in line with the criteria. The selected research was imported into the document management software EndNote X9. First, an initial screening based on titles and abstracts will be performed and suitable studies will be selected based on inclusion criteria. Secondly, the full text of all potentially eligible studies will be obtained for re-screening. If there are discrepancies in inclusion and exclusion, discussions with the corresponding authors are planned to resolve the differences. The process will use the PRISMA-P flowchart to show the study selection process in [Figure 1](#).

Figure 1. Flow diagram of study selection process.

### Data extraction and management

1  
2  
3  
4  
5  
6  
7  
8  
9  
10 175 The following data will be extracted from the selected studies by 2 independent  
11 176 reviewers using a standard data extraction sheet: year of publication, country, general  
12 177 information, participant characteristics, inclusion and exclusion criteria, sample size,  
13 178 randomization, blinding methods, methods, control, outcome measures, results,  
14 179 adverse reactions, conflicts of interest, ethical approval, and other information. The  
15 180 extracted data were entered into the electronic database, and if a difference occurred,  
16 181 it was verified by corresponding authors (HZW and JI).

17  
18  
19  
20  
21 182 **Management of missing data**

22  
23  
24 183 The cause of the missing data will be determined to solve the problem. And if this is  
25 184 not working, the authors will be contacted for the missing part. This will be  
26 185 documented and the available data will be extracted and analyzed if the missing data  
27 186 cannot be obtained.

28  
29  
30  
31 187 **Risk of bias assessment**

32  
33 188 The Cochrane Collaboration tool<sup>[17]</sup> was used to assess the risk of bias in each study,  
34 189 including the following 6 types of bias: random sequence generation, allocation  
35 190 concealment, participant and personal blinding, outcome assessment blinding,  
36 191 incomplete outcome data, selective reporting, and other sources of bias. The quality of  
37 192 the report was divided into 3 levels: low, unclear, and high-risk. Differences were  
38 193 resolved through group discussion.

39  
40  
41  
42  
43 194 **Statistical analysis**

44  
45 195 This study will be analyzed using RevMan version 5.3. Relative risk (RR) was used  
46 196 when the results were dichotomous variables with 95% confidence intervals. For  
47 197 continuous variables, we used the standardized mean difference and 95% confidence  
48 198 intervals. The chi-square test and  $I^2$  statistic will be used to confirm heterogeneity.  
49 199 The former checks for heterogeneity, whereas the latter reflects the degree of  
50 200 heterogeneity through a specific value. If  $I^2$  was >50%, there was considerable

51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.  
Enseignement Supérieur (ABES).

heterogeneity between the studies; therefore, a subgroup analysis was performed to investigate the potential causes.

### **Subgroup analysis**

If significant heterogeneity exists, subgroup analyses will be conducted to explore its source. Subgroup analyses were conducted according to the characteristics of the included studies, with respect to the time of publication, the geographical scope of the study, the gender of the participants, and the control interventions. In terms of interventions, the effects of massage modality, duration, and frequency on efficacy will be explored in depth.

### **Sensitivity analysis**

Sensitivity analysis will be conducted to test the robustness of the review conclusions if possible. The impacts of sample size, study design, methodological quality, and missing data will be evaluated.

### **Grading the quality of evidence**

The Grading of Recommendations Assessment approach will be used to judge the quality of the evidence for all outcomes. Risk of bias, heterogeneity, indirectness, imprecision and publication bias will be assessed. The assessments will be classified into 4 levels: high, moderate, low, or very low.

### **Ethics and dissemination**

This protocol will not evaluate individual patient information or affect patient rights and therefore does not require ethical approval. Results from this review will be disseminated through peer-reviewed journals and conference reports.

### **DISCUSSION**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

224 This is the first protocol of systematic review and meta-analysis to evaluate the  
225 effectiveness and safety of abdominal massage therapy for chronic constipation in  
226 elderly patients. Our group will objectively and comprehensively evaluate the  
227 therapeutic effects of abdominal massage for constipation in elderly patients. The  
228 results of this review will provide physical therapists, gastroenterologists and patients  
229 with more information on complementary and alternative treatment options for  
230 chronic constipation in the elderly. The credibility of the existing clinical evidence  
231 provides new directions for future research.

232 **Ethics and dissemination** As there will be no collection or generation of raw data, a  
233 statement of ethical approval is not required. We will publish the results of our study in  
234 peer-reviewed journals.

235 **Acknowledgements** Qiang Yuan, Xiaoyan Wang, Zhou Li contributed equally to this  
236 work and share first authorship.

237 **Author Contributions** YQ conceived this study and wrote the manuscript. WXY and  
238 LZ developed the search strategy. WZH and LJ provided methodological advice. LC,  
239 LQW and WHZ revised the manuscript. All authors have reviewed this protocol and  
240 approved the final manuscript.

241 **Funding** This study was supported by Professor Luo Caigui's "Emei injury therapy"  
242 spinal manipulation inheritance research project (Chengdu University of Traditional  
243 Chinese Medicine Foundation No. CCYB2022004) and clinical efficacy study of Luo's  
244 manipulation combined with wax therapy based on shear wave elastography for the  
245 treatment of postpartum low back pain with cold and dampness (Sichuan Provincial  
246 Administration of Traditional Chinese Medicine No. 2021MS075).

247 **Competing interests** None declared.

248 **REFERENCES**

249 1. Bharucha AE, Pemberton JH, Locke GR, 3rd. American Gastroenterological  
250 Association technical review on constipation. **Gastroenterology** 2013,  
251 144(1): 218-238.  
252 2. Siah KTH, Gong X, Yang XJ, Whitehead WE, Chen M, Hou X, *et al.* Rome  
253 Foundation-Asian working team report: Asian functional gastrointestinal

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

- disorder symptom clusters. **Gut** 2018, 67(6): 1071-1077.
3. Gallegos-Orozco JF, Foxx-Orenstein AE, Sterler SM, Stoa JM. Chronic constipation in the elderly. **Am J Gastroenterol** 2012, 107(1): 18-25; quiz 26.
  4. Bharucha AE, Lacy BE. Mechanisms, Evaluation, and Management of Chronic Constipation. **Gastroenterology** 2020, 158(5): 1232-1249.e1233.
  5. Katakami N, Oda K, Tauchi K, Nakata K, Shinozaki K, Yokota T, *et al.* Phase IIb, Randomized, Double-Blind, Placebo-Controlled Study of Naldemedine for the Treatment of Opioid-Induced Constipation in Patients With Cancer. **J Clin Oncol** 2017, 35(17): 1921-1928.
  6. Ness J, Hoth A, Barnett MJ, Shorr RI, Kaboli PJ. Anticholinergic medications in community-dwelling older veterans: prevalence of anticholinergic symptoms, symptom burden, and adverse drug events. **Am J Geriatr Pharmacother** 2006, 4(1): 42-51.
  7. Travagli RA, Browning KN, Camilleri M. Parkinson disease and the gut: new insights into pathogenesis and clinical relevance. **Nat Rev Gastroenterol Hepatol** 2020, 17(11): 673-685.
  8. Sumida K, Molnar MZ, Potukuchi PK, Thomas F, Lu JL, Yamagata K, *et al.* Constipation and risk of death and cardiovascular events. **Atherosclerosis** 2019, 281: 114-120.
  9. Tse G, Lai ET, Lee AP, Yan BP, Wong SH. Electrophysiological Mechanisms of Gastrointestinal Arrhythmogenesis: Lessons from the Heart. **Front Physiol** 2016, 7: 230.
  10. Peng Y, Liu F, Qiao Y, Wang P, Ma B, Li L, *et al.* Association of abnormal bowel health with major chronic diseases and risk of mortality. **Ann Epidemiol** 2022, 75: 39-46.
  11. Vriesman MH, Koppen IJN, Camilleri M, Di Lorenzo C, Benninga MA. Management of functional constipation in children and adults. **Nat Rev Gastroenterol Hepatol** 2020, 17(1): 21-39.
  12. Wang G, Zhang Z, Sun J, Li X, Chu Y, Zhao D, *et al.* Abdominal massage: A review of clinical and experimental studies from 1990 to 2021. **Complement Ther Med** 2022, 70: 102861.
  13. Wang X, Sun J, Li Z, Luo H, Zhao M, Li Z, *et al.* Impact of abdominal massage on enteral nutrition complications in adult critically ill patients: A systematic review and meta-analysis. **Complement Ther Med** 2022, 64: 102796.
  14. Wang J, Chen Y, Xue H, Chen Z, Wang Q, Zhu M, *et al.* Effect of abdominal massage on feeding intolerance in patients receiving enteral nutrition: A systematic review and meta-analysis. **Nurs Open** 2023, 10(5): 2720-2733.
  15. Dai N, He Q, Liu X, Fang M, Xiong M, Li X, *et al.* Therapeutic massage/Tuina for treatment of functional dyspepsia: a systematic review and meta-analysis of randomized controlled trials. **Qual Life Res** 2023, 32(3): 653-667.
  16. Shamseer L, Moher D, Clarke M, Ghera D, Liberati A, Petticrew M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

298 (PRISMA-P) 2015: elaboration and explanation. **Bmj** 2015, 350: g7647.  
299 17. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, *et al.*  
300 The Cochrane Collaboration's tool for assessing risk of bias in randomised  
301 trials. **Bmj** 2011, 343: d5928.  
302

For peer review only

Enseignement Supérieur (ABES).  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.




Figure 1. Flow diagram of study selection process.

423x397mm (72 x 72 DPI)



**PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol\***

Section and topic	Item No	Checklist item	Reported on Page #
<b>ADMINISTRATIVE INFORMATION</b>			
Title:			
Identification	1a	Identify the report as a protocol of a systematic review	1
Update	1b	If the protocol is for an update of a previous systematic review, identify as such	1
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number	2
Authors:			
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author	1
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review	1
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments	NM
Support:			
Sources	5a	Indicate sources of financial or other support for the review	11
Sponsor	5b	Provide name for the review funder and/or sponsor	Not applicable
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol	Not applicable
<b>INTRODUCTION</b>			
Rationale	6	Describe the rationale for the review in the context of what is already known	3
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	3
<b>METHODS</b>			
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review	4
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage	6
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits such that it could be repeated	6

Study records:				
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review		8
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)		8
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently in duplicate), any processes for obtaining and confirming data from investigators		9
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources) and any pre-planned data assumptions and simplifications		8
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale		5
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis		9
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised		9
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as $I^2$ , Kendall's $\tau$ )		9
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression)		10
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned		Not applicable
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)		10
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)		10

**\* It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

*From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.*

# BMJ Open

## Abdominal Massage for Chronic Constipation in the Elderly: A Systematic Review and Meta-analysis Protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-074780.R1
Article Type:	Protocol
Date Submitted by the Author:	01-Nov-2023
Complete List of Authors:	yuan, qiang; Hospital of Chengdu University of Traditional Chinese Medicine, Department of Tuina Wang, Xiaoyan; Hospital of Chengdu University of Traditional Chinese Medicine, Oncology Department Li, Zhou; Hospital of Chengdu University of Traditional Chinese Medicine, Oncology Department Li, Chuan; Chengdu University of Traditional Chinese Medicine School of Acupuncture and Tuina Lu, Qunwen; Hospital of Chengdu University of Traditional Chinese Medicine, Tuina Department Wang, Haozhong; Chengdu University of Traditional Chinese Medicine, Luo, Jian; Hospital of Chengdu University of Traditional Chinese Medicine, Tuina Department
<b>Primary Subject Heading</b>:	Complementary medicine
Secondary Subject Heading:	Gastroenterology and hepatology
Keywords:	COMPLEMENTARY MEDICINE, GASTROENTEROLOGY, Nursing Care

SCHOLARONE™  
Manuscripts



## ABSTRACT

**Introduction:** Chronic constipation is a highly prevalent health challenge, particularly challenging to treat in elderly patients. Although lifestyle guidance and laxative therapy often yield positive outcomes, patients occasionally struggle in maintaining dietary control. Therefore, it is necessary to propose an economical and harmless alternative therapy to the existing treatment methods documented in the international literature. This systematic review and meta-analysis aims to evaluate the efficacy and safety of abdominal massage in elderly CC (Chronic Constipation) patients with chronic constipation. The objective is to investigate an economical and harmless treatment method to provide a basis for future mechanistic research.

**Methods and analysis:** Electronic searches were conducted to identify clinical randomized controlled trials in various databases, including Web of Science, PubMed, CINAHL, Cochrane Library, Embase, Airtiti Library, Chinese National Knowledge Infrastructure Databases (CNKI), Chinese Science and Technology Periodical Database (VIP), Chinese Biomedical Literature Database (CBM), and Wan Fang databases. Relevant data will be extracted, and a meta-analysis will be conducted using Reviewer Manager 5.4. Quality and risk assessments of the studies included were performed, and the outcome indicators of the trials were observed. This review will evaluate abdominal massage as a treatment option for relieving symptoms and improving quality of life in elderly patients with chronic constipation. Moreover, it will provide additional insights for clinical treatment and mechanistic studies. The search will be performed following the publication of this protocol (estimated to occur on January 1, 2023) and will be repeated one month before the submission for publication of the final review (estimated to be March 1, 2024).

**Ethics and dissemination:** As this is a literature review, ethics approval is not required. We will disseminate the findings of this study to publications in peer-reviewed journals as well as presentations at relevant national and international conferences.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

52    **Abbreviations:** CC, chronic constipation; NSAIDs, nonsteroidal anti-inflammatory  
53    drugs.  
54    **Keywords:** Abdominal Massage, Chronic constipation, Elderly patients, Systematic  
55    review  
56

For peer review only

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- All types of Chronic Constipation clinical studies across all elderly age groups will be included without language limitation.
- Adherence to the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) and using Cochrane quality assessment tools ensure a rigorous and standardized methodology.
- The systematic review will rely solely on published data, potentially excluding relevant but unpublished studies.

1

2

3

4

5

65 INTRODUCTION

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Chronic constipation (CC) is characterized by difficult and hard stools, reduced frequency of bowel movements, and a feeling of incomplete defecation. The main diagnostic criteria are the Rome IV criteria and patient's self-reported symptoms.[1, 2] The prevalence of chronic constipation was 16% for men and 26% for women aged  $\geq 65$  years.[3] In individuals aged  $\geq 84$  years, the prevalence could be as high as 26% and 34% for men and women, respectively.[4] Decreased contractile motility of the detrusor and smooth muscle of the colon and atrophy of the gastrointestinal mucosa with reduced fluid secretion are the mechanisms that make constipation more common in the elderly.[5] Aging-related changes in dietary and physical activity, lifestyle factors, arthralgia, osteoporosis, laxative dependence, and long-term use of anticholinergic agents, opioid analgesics,[6] calcium supplements, and nonsteroidal anti-inflammatory drugs (NSAIDs) are risk factors for chronic constipation in elderly individuals.[7]

Constipation not only significantly affects the quality of life of the elderly but can also cause many diseases and consume a substantial amount of health care resources. Fecal retention can result in disruptions of gut microbiota and the production of harmful flora metabolites that induce cardiovascular, neurocognitive, and other diseases.[8, 9] The difficulty in defecation experienced by elderly patients may lead to cardiovascular and cerebrovascular accidents, thereby increasing the risk of sudden death.[10] Concurrently, CC also serves as a risk factor for severe conditions such as colon cancer.[11, 12] The main treatment for CC is the use of laxatives and lifestyle modifications.[13] However, drug therapy may occasionally lead to dependency, potentially exacerbating symptoms or causing complications.[13] While existing treatment methods have received positive feedback, it is still necessary to seek safe, effective, and economical treatments due to the high costs, side effects, and, in some cases, ineffectiveness associated with current interventions.

As an auxiliary alternative therapy, abdominal massage is considered a convenient and cost-effective option for clinical use.[14] It has the potential to improve blood circulation in the gastrointestinal tract and stimulate gastrointestinal motility. Previous studies explored the effectiveness of abdominal massage on digestive function and



bowel disorders.[15-17] However, specific studies on elderly patients with constipation are lacking. Given the increased prevalence of chronic constipation in older adults and its specific pathological mechanisms, it is necessary to explore this condition in an age-restricted manner.

## **METHODS AND ANALYSIS**

### **Study registration**

This systematic review protocol was registered in PROSPERO 2023 (registration number: CRD42023408629). The protocol report adhered to the guidelines outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocol (PRISMA-P) declaration.[18] The review was performed in accordance with the PRISMA-P declaration guidelines.

### **Inclusion criteria for study selection**

#### **Type of study**

All randomized controlled trials that fulfilled the inclusion criteria were included in this systematic evaluation and meta-analysis to assess the safety and efficacy of abdominal massage therapy for the treatment of chronic constipation among the elderly.

#### **Type of participant**

Participants aged 65 years or older diagnosed with chronic constipation will be included, regardless of sex, race, education, or economic status. The diagnostic criteria for chronic constipation were as follows:

- 1) Clinical diagnostic criteria (e.g., Rome IV criteria);[19]
- 2) Diagnostic criteria defined by the authors or defined by the clinician;
- 3) Participants' self-reported constipation.

The exclusion criteria were as follows:

- 1) Severe cognitive impairment that hinders understanding of the trial;
- 2) Belonging to specific clinical population groups (e.g., pregnant women, ICU patients, post-operation patients).

#### **Type of intervention and comparisons**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Abdominal massage includes any type of pressure applied to the abdominal region, such as Swedish abdominal massage, acupressure (abdominal acupoint massage), and aromatic abdominal massage. There were no limits on the time or frequency of the massages. The comparison group consisted of individuals who received either no intervention, a placebo intervention, or other medicinal treatments, such as osmotic or stimulant laxatives, elobixibat, linaclotide, lubiprostone, mizagliflozin, naronapride, plecanatide, prucalopride, tegaserod, tenapanor, or velusetrag.[20]

Type of outcome measure

The primary efficacy outcomes will be the frequency of bowel movements, related scales, and symptoms of constipation. Constipation symptom questionnaires, such as the Demographic Characteristics Questionnaire, Rome criteria,[19] and related Conk Assessment Scales,[21] were used to measure constipation symptoms. Secondary efficacy outcomes will include the influence of constipation on the quality of life and laxative use. The safety outcomes included the monitoring of adverse events.

Search methods for identification of studies

Electronic data sources

The following electronic databases will be searched from their inception dates up to May 30, 2023: Web of Science, PubMed, CINAHL, Cochrane Library, Embase, Airiti Library, Chinese National Knowledge Infrastructure Databases (CNKI), Chinese Science and Technology Periodical Database (VIP), Chinese Biomedical Literature Database (CBM), and Wan Fang Database. After excluding language or publication restrictions, we will conduct randomized clinical trials to investigate the effectiveness of abdominal massage on chronic constipation in older adults who meet the eligibility criteria.

Searching other resources

The reference lists of potentially missing eligible studies will be reviewed, and relevant conference proceedings will also be examined.

Search strategy

The search strategy is summarized in Appendix 1. The following search keywords will be used: abdominal massage (e.g., “tuina,” “massage,” “Chinese massage,”); constipation (e.g., “Dyschezia” or “Colonic Inertia”); randomized controlled trial (e.g., “randomized controlled trial,” “controlled clinical trial,” “random allocation,” “randomized,” “randomly,” “double-blind method,” “single-blind method,” or “clinical trial”). Equivalent search keywords were used in Chinese databases. Additional studies will be sought in the reference lists of the selected articles, and the authors will be contacted for any unclear information.

## **Data collection and analysis**

### **Selection of studies**

The titles and abstracts of all studies retrieved will undergo independent review and screening by two reviewers (YQ and WXY) in order to identify eligible trials and exclude duplicate or irrelevant studies. The selected research was imported into the document management software EndNote X9. The initial screening, based on titles and abstracts, was performed first, and suitable studies were selected based on the inclusion criteria. Subsequently, full texts of all potentially eligible studies were obtained for rescreening. If there are discrepancies between the inclusion and exclusion criteria, discussions with the corresponding authors will be planned to resolve these differences. A PRISMA-P flowchart ( Figure 1) was used to illustrate the study selection process.

### **Data extraction and management**

The following data were extracted from the selected studies by two independent reviewers using a standard data extraction sheet: year of publication, country, general information, participant characteristics, inclusion and exclusion criteria, sample size, randomization, blinding methods, methods, controls, outcome measures, results, adverse reactions, conflicts of interest, ethical approval, and other information. The extracted data were entered into an electronic database, and differences were verified by the corresponding authors (LJ and WHZ).

### **Management of missing data**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

182 The cause of missing data was determined to solve this problem. If investigative  
183 approach proves unsuccessful, the authors will be contacted to request the missing  
184 data. This process will be documented, and in the absence of the missing data, the  
185 available data will be extracted and analyzed.

186 Risk of bias assessment

187 The Cochrane Collaboration tool will be used to assess the risk of bias in each study,  
188 including six types of bias: random sequence generation, allocation concealment,  
189 participant and personal blinding, outcome assessment blinding, incomplete outcome  
190 data, selective reporting, and other sources of bias.[22] The quality of the reports was  
191 divided into three levels: low-, unclear-, and high-risk. Differences were resolved  
192 through group discussion.

193 **Statistical analysis**

194 Data were analyzed using RevMan version 5.3. The relative risk was used when the  
195 results were dichotomous variables, with 95% confidence intervals. For continuous  
196 variables, the standardized mean difference and 95% confidence interval were used.  
197 The chi-square test and  $I^2$  statistic were used to confirm heterogeneity. The former  
198 checks for heterogeneity, whereas the latter reflects the degree of heterogeneity using  
199 a specific value. If  $I^2 > 50\%$ , it indicated considerable heterogeneity among the  
200 studies. Consequently, a subgroup analysis was performed to investigate the potential  
201 causes.

202 **Subgroup analysis**

203 If significant heterogeneity was observed, subgroup analyses were conducted to  
204 explore the sources of heterogeneity. These subgroup analyses were conducted based  
205 on the characteristics of the included studies, including the publication time,  
206 geographical scope of the study, participant sex, and control interventions. Regarding  
207 interventions, we will explore the impact of massage modality, duration, and  
208 frequency on efficacy.

209 **Sensitivity analysis**

A sensitivity analysis will be conducted to test the robustness of the conclusions, examining the impact of sample size, study design, methodological quality, and missing data.

### **Grading the quality of evidence**

The Grading of Recommendations Assessment approach was used to judge the quality of evidence for all outcomes. Risks of bias, heterogeneity, indirectness, imprecision, and publication bias were assessed. The assessments were classified into four levels: high, moderate, low, or very low.

### **Ethics and dissemination**

This protocol will not evaluate individual patient information or affect patient rights; hence, it does not require ethical approval. The outcomes of this review will be shared through peer-reviewed journals and conference reports.

### **Patient and public involvement**

Patients were not involved in the design, conduct, reporting, or dissemination of this research.

## **DISCUSSION**

This is the first protocol for a systematic review and meta-analysis intended to evaluate the effectiveness and safety of abdominal massage therapy for chronic constipation in elderly patients. Our group aimed to objectively and comprehensively evaluate the therapeutic effects of abdominal massage on constipation in elderly patients. The results of this review will provide physical therapists, gastroenterologists, and patients with additional information on complementary and alternative treatment options for chronic constipation in the elderly population. Furthermore, the credibility of existing clinical evidence provides new directions for future research.

## **AUTHORS' CONTRIBUTIONS**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

YQ, WXY and WHZ conceptualised the study. YQ, WXY and ZL wrote the manuscript with support from WHZ and LJ. LQW and LC created the search terms. All authors were responsible for reading and approving this manuscript’s final version.

FUNDING STATEMENT

This project was financially supported by the Program of Sichuan Provincial Administration of Traditional Chinese Medicine (No. 2021MS075 and No. 2023MS534) and the Project of the Key Research Base for Humanities and Social Sciences in the Universities of Sichuan Province- the Coordinated Development of Traditional Chinese Medicine Culture in Sichuan Province ( No. 2023XT105)

COMPETING INTEREST

The authors declare that they have no competing interests.

DATA SHARING STATEMENT

All data relevant to the study are included in the article or uploaded as supplementary information.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

## REFERENCES

- [1] Bharucha AE, Pemberton JH, Locke GR, 3rd. American Gastroenterological Association technical review on constipation. *Gastroenterology* **2013**, 144(1): 218-238.
- [2] Siah KTH, Gong X, Yang XJ, Whitehead WE, Chen M, Hou X, *et al.* Rome Foundation-Asian working team report: Asian functional gastrointestinal disorder symptom clusters. *Gut* **2018**, 67(6): 1071-1077.
- [3] Milosavljevic T, Popovic DD, Mijac DD, Milovanovic T, Krstic S, Krstic MN. Chronic Constipation: Gastroenterohepatologist's Approach. *Dig Dis* **2022**;40(2):175-180.
- [4] Gallegos-Orozco JF, Foxx-Orenstein AE, Sterler SM, Stoa JM. Chronic constipation in the elderly. *Am J Gastroenterol* **2012**, 107(1): 18-25; quiz 26.
- [5] Bharucha AE, Lacy BE. Mechanisms, Evaluation, and Management of Chronic Constipation. *Gastroenterology* **2020**, 158(5): 1232-1249.e1233.
- [6] Katakami N, Oda K, Tauchi K, Nakata K, Shinozaki K, Yokota T, *et al.* Phase IIb, Randomized, Double-Blind, Placebo-Controlled Study of Naldemedine for the Treatment of Opioid-Induced Constipation in Patients With Cancer. *J Clin Oncol* **2017**, 35(17): 1921-1928.
- [7] Ness J, Hoth A, Barnett MJ, Shorr RI, Kaboli PJ. Anticholinergic medications in community-dwelling older veterans: prevalence of anticholinergic symptoms, symptom burden, and adverse drug events. *Am J Geriatr Pharmacother* **2006**, 4(1): 42-51.
- [8] Travagli RA, Browning KN, Camilleri M. Parkinson disease and the gut: new insights into pathogenesis and clinical relevance. *Nat Rev Gastroenterol Hepatol* **2020**, 17(11): 673-685.
- [9] Sumida K, Molnar MZ, Potukuchi PK, Thomas F, Lu JL, Yamagata K, *et al.* Constipation and risk of death and cardiovascular events. *Atherosclerosis* **2019**, 281: 114-120.
- [10] Tse G, Lai ET, Lee AP, Yan BP, Wong SH. Electrophysiological Mechanisms of Gastrointestinal Arrhythmogenesis: Lessons from the Heart. *Front Physiol*



1  
2  
3  
4 286           **2016**, 7: 230.

5  
6 287   [11]   Peng Y, Liu F, Qiao Y, Wang P, Ma B, Li L, *et al.* Association of abnormal  
7  
8 288           bowel health with major chronic diseases and risk of mortality. *Ann Epidemiol*  
9  
10 289           **2022**, 75: 39-46.

11  
12 290   [12]   Staller K, Olén O, Söderling J, *et al.* Chronic Constipation as a Risk Factor for  
13  
14 291           Colorectal Cancer: Results From a Nationwide, Case-Control Study. *Clin*  
15  
16 292           *Gastroenterol Hepatol* **2022 Aug**;20(8):1867-1876.e2.

17  
18 293   [13]   Vriesman MH, Koppen IJN, Camilleri M, Di Lorenzo C, Benninga MA.  
19  
20 294           Management of functional constipation in children and adults. *Nat Rev*  
21  
22 295           *Gastroenterol Hepatol* **2020**, 17(1): 21-39.

23  
24 296   [14]   Wang G, Zhang Z, Sun J, Li X, Chu Y, Zhao D, *et al.* Abdominal massage: A  
25  
26 297           review of clinical and experimental studies from 1990 to 2021. **Complement**  
27  
28 298           **Ther Med** **2022**, 70: 102861.

29  
30 299   [15]   Wang X, Sun J, Li Z, Luo H, Zhao M, Li Z, *et al.* Impact of abdominal  
31  
32 300           massage on enteral nutrition complications in adult critically ill patients: A  
33  
34 301           systematic review and meta-analysis. **Complement Ther Med** **2022**, 64:  
35  
36 302           102796.

37  
38 303   [16]   Wang J, Chen Y, Xue H, Chen Z, Wang Q, Zhu M, *et al.* Effect of abdominal  
39  
40 304           massage on feeding intolerance in patients receiving enteral nutrition: A  
41  
42 305           systematic review and meta-analysis. *Nurs Open* **2023**, 10(5): 2720-2733.

43  
44 306   [17]   Dai N, He Q, Liu X, Fang M, Xiong M, Li X, *et al.* Therapeutic  
45  
46 307           massage/Tuina for treatment of functional dyspepsia: a systematic review and  
47  
48 308           meta-analysis of randomized controlled trials. *Qual Life Res* **2023**, 32(3): 653-  
49  
50 309           667.

51  
52 310   [18]   Shamseer L, Moher D, Clarke M, *et al.* Preferred reporting items for  
53  
54 311           systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration  
55  
56 312           and explanation. *Bmj* **2015**, 350: g7647.

57  
58 313   [19]   Aziz I, Whitehead WE, Palsson OS, *et al.* An approach to the diagnosis and  
59  
60 314           management of Rome IV functional disorders of chronic constipation. *Expert*  
315           *Rev Gastroenterol Hepatol* **2020**;14(1):39-46.



- 316 [20] Luthra P, Camilleri M, Burr NE, *et al.* Efficacy of drugs in chronic idiopathic  
317 constipation: a systematic review and network meta-analysis. *Lancet*  
318 *Gastroenterol Hepatol* **2019**;4(11):831-844.
- 319 [21] Coffin B, Caussé C. Constipation assessment scales in adults: a literature  
320 review including the new Bowel Function Index. *Expert Rev Gastroenterol*  
321 *Hepatol* **2011**;5(5):601-13.
- 322 [22] Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, *et al.*  
323 The Cochrane Collaboration's tool for assessing risk of bias in randomised  
324 trials. *Bmj* **2011**, 343: d5928.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**FIGURE LEGEND**


Figure 1. Flow diagram of study selection process.

**APPENDIX**

**Appendix 1. Search strategies for databases**

For peer review only

Enseignement Supérieur (ABES) .  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.



Flow diagram of study selection process

190x338mm (120 x 120 DPI)

Appendix 1 Search strategies for databases

Web of Science

#1: TS=("constipation")

#2: TS=("dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation")

#3: TS=("Massage")

#4: TS=("massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings"))

#5: TS=("Randomized Controlled Trial" OR "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial")

#6: #1 OR #2

#7: #3 OR #4

#8: #5

#9: #6 AND #7 AND #8

Pubmed

#1 "constipation"[MeSH Terms]

#2 "dyschezia"[Title/Abstract] OR "colonic inertia"[Title/Abstract] OR "functional constipation"[Title/Abstract] OR "primary constipation"[Title/Abstract] OR "chronic constipation"[Title/Abstract] OR "idiopathic constipation"[Title/Abstract] OR "slow transit constipation"[Title/Abstract] OR "constipated"[Title/Abstract] OR "defecation disorder"[Title/Abstract] OR "evacuation disorder\*" [Title/Abstract] OR "gastrointestinal transit"[Title/Abstract] OR "gut transit"[Title/Abstract] OR "slow transit"[Title/Abstract] OR "hard stool\*" [Title/Abstract] OR "lumpy stool\*" [Title/Abstract] OR "hard feces"[Title/Abstract] OR "straining"[Title/Abstract] OR "incomplete evacuation"[Title/Abstract] OR "infrequent bowel movement"[Title/Abstract] OR "Constipation"[Title/Abstract]

#3 #1 OR #2

#4 "Massage"[MeSH Terms]

#5 "massage"[Title/Abstract] OR "tuina"[Title/Abstract] OR "tui na"[Title/Abstract] OR "chinese massage"[Title/Abstract] OR ("rubbed"[Title/Abstract] OR "rubbing"[Title/Abstract] OR "rubbings"[Title/Abstract])

#6 #4 OR #5

#7 "Randomized Controlled Trial" [Publication Type]

#8 "randomized controlled trial"[Title/Abstract] OR "controlled clinical trial"[Title/Abstract] OR "random allocation" [Title/Abstract] OR "randomized" [Title/Abstract] OR

"randomly"[Title/Abstract] OR "double blind method"[Title/Abstract] OR "single blind method"[Title/Abstract] OR "clinical trial"[Title/Abstract]  
 #9 #7 OR #8  
 #10 #3 AND #6 AND #9

### **CINAHL-Plus with full text (EBSCOhost Research Databases )**

#1: MH "Constipation"  
 #2: "dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation"  
 #3: MH "Massage"  
 #4: "massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings")  
 #5: PT "Randomized Controlled Trial" OR "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial"  
 #6: #1 OR #2  
 #7: #3 OR #4  
 #8: #5  
 #9: #6 AND #7 AND #8

### **Cochrane library**

#1: MeSH descriptor: [Constipation]  
 #2: "dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation"  
 #3: MeSH descriptor: [Massage]  
 #4: "massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings")  
 #5: "Randomized Controlled Trial" OR "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial"  
 #6: #1 OR #2  
 #7: #3 OR #4

#8: #5  
#9: #6 AND #7 AND #8

**Embase(Ovid)**

#1: "constipation"/exp  
#2 : "dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation"  
#3: #1 OR #2  
#4: "Massage"/exp  
#5: "massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings")  
#6: #4 OR #5  
#7: "Randomized Controlled Trial"/exp  
#8: "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial"  
#9: #7 OR #8  
#10: #3 AND #6 AND #9

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

# PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol\*

Section and topic	Item No	Checklist item	Reported on Page #
<b>ADMINISTRATIVE INFORMATION</b>			
Title:			
Identification	1a	Identify the report as a protocol of a systematic review	1
Update	1b	If the protocol is for an update of a previous systematic review, identify as such	1
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number	2
Authors:			
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author	1
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review	1
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments	NM
Support:			
Sources	5a	Indicate sources of financial or other support for the review	11
Sponsor	5b	Provide name for the review funder and/or sponsor	Not applicable
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol	Not applicable
<b>INTRODUCTION</b>			
Rationale	6	Describe the rationale for the review in the context of what is already known	3
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	3
<b>METHODS</b>			
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review	4
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage	6
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits such that it could be repeated	6

Study records:			
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review	8
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)	8
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently in duplicate), any processes for obtaining and confirming data from investigators	9
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources) and any pre-planned data assumptions and simplifications	8
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale	5
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis	9
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised	9
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of combining data and methods of combining data from studies, including any planned exploration of consistency (such as I <sup>2</sup> , Kendall's $\tau$ )	9
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression)	10
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned	Not applicable
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)	10
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)	10

**\* It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

*From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.*



# BMJ Open

## Abdominal Massage for Chronic Constipation in the Elderly: A Systematic Review and Meta-analysis Protocol

Journal:	BMJ Open
Manuscript ID	bmjopen-2023-074780.R2
Article Type:	Protocol
Date Submitted by the Author:	08-Dec-2023
Complete List of Authors:	yuan, qiang; Hospital of Chengdu University of Traditional Chinese Medicine, Department of Tuina Wang, Xiaoyan; Hospital of Chengdu University of Traditional Chinese Medicine, Oncology Department Li, Zhou; Hospital of Chengdu University of Traditional Chinese Medicine, Oncology Department Li, Chuan; Chengdu University of Traditional Chinese Medicine School of Acupuncture and Tuina Lu, Qunwen; Hospital of Chengdu University of Traditional Chinese Medicine, Tuina Department Wang, Haozhong; Chengdu University of Traditional Chinese Medicine, Luo, Jian; Hospital of Chengdu University of Traditional Chinese Medicine, Tuina Department
<b>Primary Subject Heading</b>:	Complementary medicine
Secondary Subject Heading:	Gastroenterology and hepatology, Complementary medicine
Keywords:	COMPLEMENTARY MEDICINE, GASTROENTEROLOGY, Nursing Care

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abdominal Massage for Chronic Constipation in the Elderly: A Systematic  
Review and Meta-analysis Protocol**

Qiang Yuan, MSc<sup>a†</sup>; Xiaoyan Wang, MSc<sup>b†</sup>; Li Zhou, BSc<sup>b†</sup>; Chuan Li, MSc<sup>c</sup>;  
Qunwen Lu, MSc<sup>a</sup>; Haozhong Wang, PhD<sup>d\*</sup>; Jian Luo, PhD<sup>a\*</sup>

<sup>a</sup> Tuina Department, Hospital of Chengdu University of Traditional Chinese Medicine,  
Chengdu, China

<sup>b</sup> Oncology Department, Hospital of Chengdu University of Traditional Chinese  
Medicine, Chengdu, China

<sup>c</sup> Sichuan Acupuncture and Moxibustion School, Chengdu, China

<sup>d</sup> Basic Medical College, Chengdu University of Traditional Chinese Medicine,  
Chengdu, China

<sup>†</sup>Co-first authors

<sup>†</sup>These authors contributed equally to this work and share first authorship.

\*Co-corresponding authors: Haozhong Wang and Jian Luo  
Email: [wanghaozhong@cdutcm.edu.cn](mailto:wanghaozhong@cdutcm.edu.cn) (HW); [lj6176@126.com](mailto:lj6176@126.com) (JL)

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

## ABSTRACT

**Introduction:** Chronic constipation (CC) is a highly prevalent health challenge that is particularly challenging to treat in elderly patients. Although lifestyle guidance and laxative therapy often yield positive outcomes, patients occasionally struggle with maintaining dietary control. Therefore, identifying an economical and safe alternative therapy to the existing treatment methods documented in the international literature is necessary. This systematic review and meta-analysis aims to evaluate the efficacy and safety of abdominal massage in elderly patients with CC to provide a basis for future mechanistic research.

**Methods and analysis:** Electronic searches will be conducted to identify clinical randomized controlled trials in various databases, including Web of Science, PubMed, Cumulated Index to Nursing and Allied Health Literature, Cochrane Library, Embase, Airtiti Library, Chinese National Knowledge Infrastructure Databases, Chinese Science and Technology Periodical Database (VIP), Chinese Biomedical Literature Database, and Wan Fang Data. Relevant data will be extracted, and a meta-analysis will be conducted using Reviewer Manager 5.4. Quality and risk assessments of the included studies will be performed, and the outcome indicators of the trials will be observed. This review will evaluate abdominal massage as a treatment option for relieving symptoms and improving quality of life in elderly patients with CC. Moreover, it will provide additional insights for clinical treatment and mechanistic studies. The search will be performed following the publication of this protocol (estimated to occur on December 30, 2023).

**Ethics and dissemination:** As this is a literature review, ethics approval will not be required. We will disseminate the findings of this study to publications in peer-reviewed journals as well as presentations at relevant national and international conferences.

**Abbreviations:** CC, chronic constipation

**Keywords:** Abdominal Massage, Chronic Constipation, Elderly Patients, Systematic Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

50     **STRENGTHS AND LIMITATIONS OF THIS STUDY**

- 51         • All types of clinical studies on chronic constipation across all elderly age groups
- 52             will be included without language limitations.
- 53         • The use of Cochrane quality assessment tools will ensure rigorous and
- 54             standardized methodology.
- 55         • The systematic review will rely solely on published data, potentially excluding
- 56             relevant although unpublished studies.

For peer review only

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.  
Enseignement Supérieur (ABES).

## INTRODUCTION

Chronic constipation (CC) is characterized by difficult to pass and hard stools, reduced frequency of bowel movements, and a feeling of incomplete defecation. The main diagnostic criteria are the Rome IV criteria and self-reported symptoms.[1, 2] The prevalence of CC is 16% for men and 26% for women aged  $\geq 65$  years.[3] In individuals aged  $\geq 84$  years, the prevalence is reportedly as high as 26% and 34% for men and women, respectively.[4] Decreased contractile motility of the detrusor and smooth muscle of the colon and atrophy of the gastrointestinal mucosa with reduced fluid secretion are the mechanisms that make constipation more common in the elderly.[5] Aging-related changes in diet and physical activity, lifestyle factors, arthralgia, osteoporosis, laxative dependence, and long-term use of anticholinergic agents, opioid analgesics,[6] calcium supplements, and nonsteroidal anti-inflammatory drugs are risk factors for CC in elderly individuals.[7]

Constipation not only significantly affects the quality of life of the elderly but can also cause many diseases and consume a substantial amount of health care resources. Fecal retention can result in disruptions of gut microbiota and the production of harmful flora metabolites that induce cardiovascular, neurocognitive, and other diseases.[8, 9] The difficulties with defecation experienced by elderly patients may lead to cardiovascular and cerebrovascular accidents, thereby increasing the risk of sudden death.[10] Concurrently, CC also serves as a risk factor for severe conditions such as colon cancer.[11, 12] The main treatments for CC are laxative use and lifestyle modifications.[13] However, drug therapy may occasionally lead to dependency, potentially exacerbating symptoms or causing complications.[13] Although existing treatment methods have received positive feedback, identifying safe, effective, and economical treatments is necessary due to the high costs, side effects, and, in some cases, the ineffectiveness of current interventions.

As an auxiliary alternative therapy, abdominal massage is considered a convenient and cost-effective option in clinical practice.[14] Abdominal massage has the potential to improve blood circulation in the gastrointestinal tract and stimulate gastrointestinal motility. Previous studies explored the effectiveness of abdominal massage on digestive

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

function and bowel disorders.[15-17] However, specific studies on elderly patients with constipation are lacking. Given the increased prevalence of CC in older adults and its specific pathological mechanisms, exploring this condition in an age-restricted manner is necessary.

**METHODS AND ANALYSIS**

**Study registration**

This systematic review protocol is registered with the International Prospective Register of Systematic Reviews (registration number: CRD42023408629). The reporting of this protocol adheres to the guidelines outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols (PRISMA-P) statement.[18] The review will be reported in accordance with the PRISMA-P guidelines.

**Inclusion criteria for study selection**

Type of study

All randomized controlled trials that fulfill the inclusion criteria will be included in this systematic evaluation and meta-analysis to assess the safety and efficacy of abdominal massage therapy (performed by nurses, nurses aids, caregivers, or doctors) for the treatment of CC among the elderly.

Type of participant

Participants aged 65 years or older diagnosed with CC will be included, regardless of sex, race, education, or economic status. The diagnostic criteria for CC are as follows:

- 1) Clinical diagnostic criteria (e.g., Rome IV criteria);[19]
- 2) Diagnostic criteria defined by the authors or defined by the clinician;
- 3) Participants' self-reported constipation.

The exclusion criteria are as follows:

- 1) Severe cognitive impairment that hinders understanding of the trial;
- 2) Belonging to specific clinical population groups (e.g., pregnant women, intensive care unit patients, post-operation patients).

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

## 116 Type of intervention and comparisons

117 Abdominal massage includes any type of pressure applied to the abdominal region,  
118 such as Swedish abdominal massage, acupressure (abdominal acupoint massage), and  
119 aromatic abdominal massage. No limits will be placed on massage time or frequency.  
120 The comparison group consists of individuals who received either no intervention, a  
121 placebo intervention, or other medicinal treatments, such as osmotic or stimulant  
122 laxatives, elobixibat, linaclotide, lubiprostone, mizagliflozin, naronapride,  
123 plecanatide, prucalopride, tegaserod, tenapanor, or velusetrag.[20]

## 124 Type of outcome measure

125 The primary efficacy outcomes will be the frequency of bowel movements, related  
126 scales, and symptoms of constipation. Constipation symptom questionnaires, such as  
127 the Demographic Characteristics Questionnaire, Rome criteria,[19] and related Conk  
128 Assessment Scales,[21] will be used to measure constipation symptoms. Secondary  
129 efficacy outcomes will include the influence of constipation on the quality of life and  
130 laxative use. The safety outcomes include the monitoring of adverse events.

## 131 Search methods for identification of studies

### 132 Electronic data sources

133 The following electronic databases will be searched from their inception dates up to  
134 December 30, 2023: Web of Science, PubMed, Cumulated Index to Nursing and  
135 Allied Health Literature, Cochrane Library, Embase, Airiti Library, Chinese National  
136 Knowledge Infrastructure Databases, Chinese Science and Technology Periodical  
137 Database (VIP), Chinese Biomedical Literature Database, and Wan Fang Data. After  
138 excluding publication restrictions, we will review randomized clinical trials to  
139 investigate the effectiveness of abdominal massage on CC in older adults who meet  
140 the eligibility criteria.

### 141 Searching other resources

142 The reference lists of potentially missing eligible studies will be reviewed, and  
143 relevant conference proceedings will also be examined.

**Search strategy**

The search strategy is summarized in Appendix 1. The following search keywords will be used: abdominal massage (e.g., “tuina,” “massage,” “Chinese massage,”); constipation (e.g., “Dyschezia” or “Colonic Inertia”); and randomized controlled trial (e.g., “randomized controlled trial,” “controlled clinical trial,” “random allocation,” “randomized,” “randomly,” “double-blind method,” “single-blind method,” or “clinical trial”). Equivalent search keywords were used in Chinese databases. Additional studies will be identified from the reference lists of the selected articles, and the authors will be contacted for any unclear information.

**Data collection and analysis**

**Selection of studies**

The titles and abstracts of all retrieved studies will undergo independent review and screening by two reviewers (YQ and WXY) to identify eligible trials and exclude duplicate or irrelevant studies. The selected studies will be imported using the document management software EndNote X9 (Clarivate, London, UK). The initial screening, based on titles and abstracts, will be performed, and suitable studies will be selected based on the inclusion criteria. Subsequently, full texts of all potentially eligible studies will be obtained for rescreening. In case of discrepancies between the inclusion and exclusion criteria, discussions with the corresponding authors will occur to resolve these differences. A PRISMA-P flowchart (Figure 1) will be used to illustrate the study selection process.

**Data extraction and management**

The following data will be extracted from the selected studies by two independent reviewers (QY and XW) using a standard data extraction sheet: year of publication, country, general information, participant characteristics, inclusion and exclusion criteria, sample size, randomization, blinding methods, methods, controls, outcome measures, results, adverse reactions, conflicts of interest, ethical approval, and other



information. The extracted data will be entered into an electronic database, and differences will be verified by the corresponding authors (JL and HW).

### Management of missing data

The cause of missing data will be determined to solve this problem. If this investigative approach proves unsuccessful, the authors will be contacted to request the missing data. This process will be documented, and in the absence of the missing data, the available data will be extracted and analyzed.

### Risk of bias assessment

The Cochrane Collaboration tool will be used to assess the risk of bias in each study, including six types of bias: random sequence generation, allocation concealment, participant and personal blinding, outcome assessment blinding, incomplete outcome data, selective reporting, as well as other sources of bias.[22] The quality of the reports will be divided into three levels: low, unclear, and high risk. Differences will be resolved through group discussion.

### Statistical analysis

Data will be analyzed using RevMan version 5.3 (The Cochrane Collaboration, London, England). The relative risk will be used to evaluate dichotomous variables, with 95% confidence intervals. The standardized mean difference and 95% confidence interval will be used to evaluate continuous variables. The chi-square test and  $I^2$  statistic will be used to confirm heterogeneity. The former checks for heterogeneity, whereas the latter reflects the degree of heterogeneity using a specific value.  $I^2 > 50\%$ , will indicate considerable heterogeneity among the studies. Consequently, a subgroup analysis will be performed to investigate the potential causes.

### Subgroup analysis

If significant heterogeneity is observed, subgroup analyses will be conducted to explore the sources of heterogeneity. These subgroup analyses will be conducted

based on the characteristics of the included studies, including the publication time, geographical scope of the study, participant sex, and control interventions. Regarding interventions, we will explore the impact of massage modality, duration, and frequency on efficacy.

**Sensitivity analysis**

A sensitivity analysis will be conducted to test the robustness of the conclusions, examining the impact of sample size, study design, methodological quality, and missing data.

**Grading the quality of evidence**

The Grading of Recommendations Assessment approach will be used to judge the quality of evidence for all outcomes. Risks of bias, heterogeneity, indirectness, imprecision, and publication bias were assessed. The assessments were classified into four levels: high, moderate, low, or very low.

**Ethics and dissemination**

Because this study is based exclusively on published literature, ethics approval and informed consent will not be required. The outcomes of this review will be shared through peer-reviewed journals and conference reports.

**Patient and public involvement**

Patients will not be involved in the design, conduct, reporting, or dissemination of this research.

**DISCUSSION**

This is the first protocol for a systematic review and meta-analysis intended to evaluate the effectiveness and safety of abdominal massage therapy for CC in elderly patients. We aim to objectively and comprehensively evaluate the therapeutic effects of abdominal massage on constipation in elderly patients. The results of this review will provide physical therapists, gastroenterologists, and patients with additional

information on complementary and alternative treatment options for CC in the elderly population. Furthermore, the credibility of existing clinical evidence provides new directions for future research.

## **AUTHORS' CONTRIBUTIONS**

QY, XW, and HW conceptualized the protocol. QY, XW, and LZ wrote the manuscript with support from HW and JL. QL and CL created the search terms. All authors were responsible for reading and approving the final version of the manuscript.

## **FUNDING STATEMENT**

This work was supported by the Program of Sichuan Provincial Administration of Traditional Chinese Medicine, grant numbers 2021MS075 and 2023MS534 and the Project of the Key Research Base for Humanities and Social Sciences in the Universities of Sichuan Province-the Coordinated Development of Traditional Chinese Medicine Culture in Sichuan Province, grant number 2023XT105.

## **COMPETING INTEREST**

The authors declare that they have no competing interests.

## **DATA SHARING STATEMENT**

All data relevant to the study are included in the article or as supplementary material.

REFERENCES

[1] Bharucha AE, Pemberton JH, Locke GR, 3rd. American Gastroenterological Association technical review on constipation. *Gastroenterology* 2013;144:218–38.

[2] Siah KTH, Gong X, Yang XJ, et al. Rome Foundation-Asian working team report: Asian functional gastrointestinal disorder symptom clusters. *Gut* 2018;67:1071–7.

[3] Milosavljevic T, Popovic DD, Mijac DD, et al. Chronic constipation: gastroenterohepatologist's approach. *Dig Dis* 2022;40:175–80.

[4] Gallegos-Orozco JF, Foxx-Orenstein AE, Sterler SM, et al. Chronic constipation in the elderly. *Am J Gastroenterol* 2012;107:18–25; quiz 26.

[5] Bharucha AE, Lacy BE. Mechanisms, evaluation, and management of chronic constipation. *Gastroenterology* 2020;158:1232–49.e3.

[6] Katakami N, Oda K, Tauchi K, et al. Phase IIb, randomized, double-blind, placebo-controlled study of naldemedine for the treatment of opioid-induced constipation in patients with cancer. *J Clin Oncol* 2017;35:1921–8.

[7] Ness J, Hoth A, Barnett MJ, et al. Anticholinergic medications in community-dwelling older veterans: prevalence of anticholinergic symptoms, symptom burden, and adverse drug events. *Am J Geriatr Pharmacother* 2006;4:42–51.

[8] Travagli RA, Browning KN, Camilleri M. Parkinson disease and the gut: new insights into pathogenesis and clinical relevance. *Nat Rev Gastroenterol Hepatol* 2020;17:673–85.

[9] Sumida K, Molnar MZ, Potukuchi PK, et al. Constipation and risk of death and cardiovascular events. *Atherosclerosis* 2019;281:114–20.

[10] Tse G, Lai ET, Lee AP, et al. Electrophysiological mechanisms of gastrointestinal arrhythmogenesis: lessons from the heart. *Front Physiol* 2016;7:230.

[11] Peng Y, Liu F, Qiao Y, et al. Association of abnormal bowel health with major chronic diseases and risk of mortality. *Ann Epidemiol* 2022;75:39–46.

[12] Staller K, Olén O, Söderling J, et al. Chronic Constipation as a risk factor for

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

- colorectal cancer: results from a nationwide, case-control study. *Clin Gastroenterol Hepatol* 2022;20:1867–76.e2.
- [13] Vriesman MH, Koppen IJN, Camilleri M, et al. Management of functional constipation in children and adults. *Nat Rev Gastroenterol Hepatol* 2020;17:21–39.
- [14] Wang G, Zhang Z, Sun J, et al. Abdominal massage: a review of clinical and experimental studies from 1990 to 2021. *Complement Ther Med* 2022;70:102861.
- [15] Wang X, Sun J, Li Z, et al. Impact of abdominal massage on enteral nutrition complications in adult critically ill patients: a systematic review and meta-analysis. *Complement Ther Med* 2022;64:102796.
- [16] Wang J, Chen Y, Xue H, et al. Effect of abdominal massage on feeding intolerance in patients receiving enteral nutrition: a systematic review and meta-analysis. *Nurs Open* 2023;10:2720–33.
- [17] Dai N, He Q, Liu X, et al. Therapeutic massage/Tuina for treatment of functional dyspepsia: a systematic review and meta-analysis of randomized controlled trials. *Qual Life Res* 2023;32:653–67.
- [18] Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015;350:g7647.
- [19] Aziz I, Whitehead WE, Palsson OS, et al. An approach to the diagnosis and management of Rome IV functional disorders of chronic constipation. *Expert Rev Gastroenterol Hepatol* 2020;14:39–46.
- [20] Luthra P, Camilleri M, Burr NE, et al. Efficacy of drugs in chronic idiopathic constipation: a systematic review and network meta-analysis. *Lancet Gastroenterol Hepatol* 2019;4:831–44.
- [21] Coffin B, Caussé C. Constipation assessment scales in adults: a literature review including the new Bowel Function Index. *Expert Rev Gastroenterol Hepatol* 2011;5:601–13.
- [22] Higgins JP, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

307 tool for assessing risk of bias in randomised trials. BMJ 2011;343:d5928.  
308  
309

For peer review only

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.  
Enseignement Supérieur (ABES).

For peer review only


310 **FIGURE LEGEND**

311 Figure 1. Flow diagram of study selection process

312

313 **SUPPLEMENTARY MATERIAL**

314 **Appendix 1. Search strategies**



Flow diagram of study selection process

190x338mm (120 x 120 DPI)



## Appendix 1 Search strategies for databases

### Web of Science

#1: TS=("constipation")

#2: TS=("dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation")

#3: TS=("Massage")

#4: TS=("massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings"))

#5: TS=("Randomized Controlled Trial" OR "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial")

#6: #1 OR #2

#7: #3 OR #4

#8: #5

#9: #6 AND #7 AND #8

### Pubmed

#1 "constipation"[MeSH Terms]

#2 "dyschezia"[Title/Abstract] OR "colonic inertia"[Title/Abstract] OR "functional constipation"[Title/Abstract] OR "primary constipation"[Title/Abstract] OR "chronic constipation"[Title/Abstract] OR "idiopathic constipation"[Title/Abstract] OR "slow transit constipation"[Title/Abstract] OR "constipated"[Title/Abstract] OR "defecation disorder"[Title/Abstract] OR "evacuation disorder\*" [Title/Abstract] OR "gastrointestinal transit"[Title/Abstract] OR "gut transit"[Title/Abstract] OR "slow transit"[Title/Abstract] OR "hard stool\*" [Title/Abstract] OR "lumpy stool\*" [Title/Abstract] OR "hard feces"[Title/Abstract] OR "straining"[Title/Abstract] OR "incomplete evacuation"[Title/Abstract] OR "infrequent bowel movement"[Title/Abstract] OR "Constipation"[Title/Abstract]

#3 #1 OR #2

#4 "Massage"[MeSH Terms]

#5 "massage"[Title/Abstract] OR "tuina"[Title/Abstract] OR "tui na"[Title/Abstract] OR "chinese massage"[Title/Abstract] OR ("rubbed"[Title/Abstract] OR "rubbing"[Title/Abstract] OR "rubbings"[Title/Abstract])

#6 #4 OR #5

#7 "Randomized Controlled Trial" [Publication Type]

#8 "randomized controlled trial"[Title/Abstract] OR "controlled clinical trial"[Title/Abstract] OR "random allocation" [Title/Abstract] OR "randomized" [Title/Abstract] OR

"randomly"[Title/Abstract] OR "double blind method"[Title/Abstract] OR "single blind method"[Title/Abstract] OR "clinical trial"[Title/Abstract]  
#9 #7 OR #8  
#10 #3 AND #6 AND #9

**CINAHL-Plus with full text (EBSCOhost Research Databases )**

#1: MH "Constipation"  
#2: "dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation"  
#3: MH "Massage"  
#4: "massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings")  
#5: PT "Randomized Controlled Trial" OR "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial"  
#6: #1 OR #2  
#7: #3 OR #4  
#8: #5  
#9: #6 AND #7 AND #8

**Cochrane library**

#1: MeSH descriptor: [Constipation]  
#2: "dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation"  
#3: MeSH descriptor: [Massage]  
#4: "massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings")  
#5: "Randomized Controlled Trial" OR "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial"  
#6: #1 OR #2  
#7: #3 OR #4

#8: #5

#9: #6 AND #7 AND #8

### **Embase(Ovid)**

#1: "constipation"/exp

#2 : "dyschezia" OR "colonic inertia" OR "functional constipation" OR "primary constipation" OR "chronic constipation" OR "idiopathic constipation" OR "slow transit constipation" OR "constipated" OR "defecation disorder\*" OR "evacuation disorder\*" OR "gastrointestinal transit" OR "gut transit" OR "slow transit" OR "hard stool\*" OR "lumpy stool\*" OR "hard feces" OR "straining" OR "incomplete evacuation" OR "infrequent bowel movement" OR "Constipation"

#3: #1 OR #2

#4: "Massage"/exp

#5: "massage" OR "tuina" OR "tui na" OR "chinese massage" OR ("rubbed" OR "rubbing" OR "rubbings")

#6: #4 OR #5

#7: "Randomized Controlled Trial"/exp

#8: "randomized controlled trial" OR "controlled clinical trial" OR "random allocation" OR "randomized" OR "randomly" OR "double blind method" OR "single blind method" OR "clinical trial"

#9: #7 OR #8

#10: #3 AND #6 AND #9

**PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol\***

Section and topic	Item No	Checklist item	Reported on Page #
<b>ADMINISTRATIVE INFORMATION</b>			
Title:			
Identification	1a	Identify the report as a protocol of a systematic review	1
Update	1b	If the protocol is for an update of a previous systematic review, identify as such	1
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number	2
Authors:			
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author	1
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review	1
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments	NM
Support:			
Sources	5a	Indicate sources of financial or other support for the review	11
Sponsor	5b	Provide name for the review funder and/or sponsor	Not applicable
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol	Not applicable
<b>INTRODUCTION</b>			
Rationale	6	Describe the rationale for the review in the context of what is already known	3
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	3
<b>METHODS</b>			
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review	4
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage	6
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits such that it could be repeated	6

Study records:				
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review		8
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)		8
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently in duplicate), any processes for obtaining and confirming data from investigators		9
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources) and any pre-planned data assumptions and simplifications		8
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale		5
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis		9
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised		9
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of combining data and methods of combining data from studies, including any planned exploration of consistency (such as $I^2$ , Kendall's $\tau$ )		9
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression)		10
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned		Not applicable
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)		10
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)		10

**\* It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

*From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.*