To cite: Shama AT, Terefa DR,

cancer and its determinants in

Ethiopia: a systematic review

and meta-analysis. BMJ Open

2024;14:e080080. doi:10.1136/

Desisa AE. et al. Breast

bmjopen-2023-080080

Prepublication history

and additional supplemental

available online. To view these

online (https://doi.org/10.1136/ bmjopen-2023-080080).

Received 20 September 2023

Accepted 14 October 2024

files, please visit the journal

material for this paper are

BMJ Open Breast cancer and its determinants in Ethiopia: a systematic review and metaanalysis

Adisu Tafari Shama ,¹ Dufera Rikitu Terefa ,¹ Adisu Ewunetu Desisa ,¹ Matiyos Lema ,¹ Melese Chego Cheme ,¹ Edosa Tesfaye Geta ,¹ Jira Wakoya Feyisa ,¹ Bikila Regassa Feyisa ,^{1,2} Bayise Biru ,^{1,3}

ABSTRACT

Objectives Breast cancer is the leading cause of cancer morbidity and mortality among women. Still, there is a paucity of studies to know the magnitude of the problem in Ethiopia. Hence, this review was intended to pool the prevalence and identify the determinants of breast cancer in Ethiopia.

Design A systematic review and meta-analysis was conducted.

Data sources Databases like PubMed/MEDLINE, HINARI, Science Direct, and Google Scholar, as well as websites of organisationsl organizations, rewere searched between 25 February and 6 March 2023.

Eligibility criteria All observational studies in Ethiopia that reported either the magnitude and/or determinants of breast cancer regardless of publication status were included.

Data extraction and synthesis Two authors independently assessed and extracted the data. The Joanna Briggs Institute meta-analysis of statistics assessment and review instrument quality appraisal tool was used to assess the quality of the articles. Effect estimates were done by using the random-effects model. The meta-analysis results were displayed by using forest plots.

Results Seventeen articles were reviewed with 24 435 total participants. The pooled proportion of breast cancer morbidity among patients with cancer was 20. 58% (95% Cl 17.25%, 23.90%) in Ethiopia. Consuming packed foods (POR=2.12, 95% Cl 1.41, 3.17), presence of high cholesterol (POR=4.08; 95% Cl 2.75, 6.07), physical inactivity (POR=3.27; 95% Cl 1.80, 5.94), high body mass index (BMI) (POR=2.27; 95% Cl 0.85, 6.03), postmenopause (POR=2.25; 95% Cl 1.63, 3.10), family history of cancer (POR=3.65; 95% Cl 0.85, 15.71) and lack of breastfeeding (POR=2.76; 95% Cl 0.90, 7.92) were the determinants of breast cancer.

Conclusions One of five patients with cancer is diagnosed with breast cancer in Ethiopia. Furthermore, more than a quarter of women with cancer suffer from breast cancer. Processed food consumption, high cholesterol in the body, lack of physical activity, high BMI, postmenopause, family history of cancer and lack of breastfeeding were the risk factors for breast cancer. The use of healthy food sources such as fruits and vegetables, and homegrown varieties of crops rather than seeking processed foods would help.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The inclusion of prediction interval is a strong aspect of this study, as it is uncommon in many meta-analyses.
- \Rightarrow The review included only observational studies.
- ⇒ The narrow scope of the review (only one country) is a limitation of this study.
- ⇒ A limited number of studies were found to pool the OR for some factors.

PROSPERO registration number CRD42023417733

BACKGROUND

Breast cancer is a diverse disease with numerous morphological and molecular subgroups.¹ It has been found to be the most common cause of cancer deaths in 11 regions of the world.² It is one of the most frequently diagnosed cancers and the leading cause of cancer deaths in women world-wide.³ The recent global burden of cancer statistics (GLOBOCAN 2020) showed that breast cancer has surpassed lung cancer and accounted for 2.3 million (11.7%) of all new cancer cases globally. It affects one in four new cancer cases of women and contributes to one in six deaths of women from cancer.⁴

The cancer burden is increasing worldwide **in** and is estimated to be 28.4 million cases by 2040, which is a 47% increase over the cancer burden in 2020. A higher death rate occurs in developing countries than in developed countries (15 breast cancer deaths in developing countries vs 12.8 in developed countries per 100 000).⁴ In Ethiopia too there were an estimated 5900 incident cases of breast cancer with the highest age-standardised incidence rate of 12.5 per 100 000 and a death rate of 9.7 per 100 000 in 2019.⁵

Previous studies identified that the incidence of breast cancer varies widely across the world due to differences in the level

Check for updates

© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Public Health, Institute of Health Sciences, Wollega University, Nekemte, Ethiopia

²Department of Epidemiology, Faculty of Public Health, Jimma University, Jimma, Ethiopia ³Department of Human Nutrition and Dietetics, Faculty of Public Health, Jimma University, Jimma, Ethiopia

Correspondence to

Adisu Tafari Shama; adisuteferi1906@gmail.com of education, economic status, environmental conditions, eating habits, lifestyle variables and other cultural traditions. Early age at menarche, westernised lifestyles delayed pregnancies/childbirth, reduced (namely breastfeeding, sedentary lifestyles and poor diet), and improving cancer registration and cancer detection are among the factors associated with the breast cancer in low and middle income countries (LMICs).⁶⁻⁹ Lack of knowledge about the disease, improper screening programmes, delayed diagnosis and insufficient medical facilities are also contributing factors to the increasing breast cancer burden in underdeveloped countries.⁶¹⁰¹¹ Widespread urbanisation, shifting patterns of reproductive and environmental risk factors, obesity, decreased physical activity and rising life expectancy are among the major factors contributing to the steady rise in breast cancer incidence in low-income nations. Low socioeconomic level, on the other hand, is related to an increased incidence of aggressive premenopausal breast cancers, as well as late-stage diagnosis and lower survival. Late menopause and early menarche are also among the risk factors that could increase the exposure of breast tissue to oestrogen hormone. In contrast to this, pregnancy and appropriate breastfeeding help to reduce the risk of breast cancer.^{610–13}

Comprehensive identification of the magnitude and determinants of breast cancer is critical for developing nations like Ethiopia, as this will aid in the development and implementation of effective breast cancer prevention initiatives. Breast cancer is not well studied in Ethiopia. Although there is one recently published review, the focus of that study was more on determinants of the problem.¹⁴ Different pocket studies so far may not represent the entire picture of the determinants of breast cancer in Ethiopia. Most of them were limited to small sample sizes, limited portions of populations covered and limited research regions. In this regard, many of the regions were not addressed in the previous studies and our study also helps to show this gap for further study, let alone intervention. It is critical to shed light on the risk factors for breast cancer. As a result, this study aimed to determine the magnitude of breast cancer and its determinants in Ethiopia.

METHODS

Study design

A systematic review and meta-analysis was conducted. The protocol was registered on Prospective Registry of Systematic Reviews with registration number CRD42023417733 and no change was made to the protocol. To conduct this review, the Preferred Reporting Items for Systematic Reviews and Meta-Analysis checklist¹⁵ was used.

Search strategy

A comprehensive search of databases like PubMed/ Medical Literature Analysis and Retrieval System Online (MEDLINE), Health Inter-Newtork Access to Research Initiative (HINARI), Science Direct and Google Scholar

was used to find the relevant articles. The searches were limited to articles written using the English language. In addition to the electronic database search, grey literature was searched using Google search and digital libraries of universities. Finally, the reference lists of the included articles for related studies were searched. To facilitate the article searching process the following keywords were used: 'breast' OR 'mammary gland' AND 'cancer' OR 'tumour' OR 'malignancy' OR 'breast cancer' OR 'breast malignancy' OR 'breast tumour' AND 'Risk factors' OR u 'Associated factors' OR 'Determinants' OR 'predictors' **AND** 'Ethiopia' OR 'Addis Ababa' OR 'Northern Ethi-opia' OR 'North west Ethiopia' OR 'Southern Ethiopia' Š OR 'South Western Ethiopia' OR 'Western Ethiopia' OR 'East Ethiopia' (online supplemental additional file 1). copyright, includi Searching started on 25 February 2023 and the final date of searching was 6 March 2023. **Eligibility criteria** Inclusion criteria

To be included in this review, the study should report bu either the determinants of breast cancer and/or the for magnitude (incidence, prevalence, number) of breast uses rel cancer morbidity.

Study setting

Studies conducted in Ethiopia (both institution-based and population-based) were part of this systematic review.

Study population

The study involved all the human population (male, female, children and adults) in Ethiopia who has been evaluated for cancer and confirmed to be patients with cancer.

Exposure

Those with modifiable or non-modifiable risk factors.

Study design

All observational studies (cross-sectional and case-control) that reported the magnitude of breast cancer morbidity and its determinants were evaluated to be included.

Publication status

Both published and unpublished studies were considered for inclusion.

Exclusion criteria

data mining, AI training, and similar technologies Articles with low quality, unclear methodologies and articles that didn't indicate the outcome of interest were excluded (online supplemental additional file 2). Excluding the studies whose full-text papers were not available after at least two personal email contacts with the corresponding authors was an exclusion criterion but all full texts were available.

Assessment of outcome variables

There were two outcomes in this study: the first outcome was the magnitude of confirmed breast cancer disease/ morbidity among patients with cancer diagnosis. This

ated to text and

Breast cancer

morbidity.

Open access

on the predefined inclusion criteria, two authors (ATS outcome can occur in any population group. Therefore, the population for this outcome was a human population and AED) independently assessed and identified papers of any age evaluated for cancer disease. by their titles, abstracts and full texts. The screened items were then compiled, and any disagreement was handled The diagnosis of breast cancer was used when diagnosis was confirmed by pathological tests in addition to the history and physical examination.¹⁶ Hence, the data were sought if the diagnosis of breast cancer was confirmed by pathological tests. The total number of people who had breast cancer was divided by the total number of people participating in the study and multiplied by 100 which was used to determine the proportion of breast cancer a two-by-two table. The second outcome/variables of this review were **Quality assessment** the determinants of breast cancer. Modifiable and nonmodifiable factors were searched from the literature to pool their value together. For the variables whose categorisation didn't overlap (eg, age category), the category repeatedly reported in the studies or the established categorisation was assumed to get the privilege. Early age at menarche is the starting of menstruation early and mostly before the age of 12 years.^{17 18}

Early age at menarche

Late menopause

Late menopause is delayed age at menopause which is after the age of 55 years in most cases.¹⁷

Benign breast disease and breast injury

These are those breast diseases such as atypical ductal hyperplasia or lobular carcinoma.¹⁷

Menopause status

Menopause status is categorised as postmenopausal if the woman has already stopped menstruation (either absence of menstruation for at least 1 year (any age) or due to bilateral oophorectomy or oestrogen deprivation therapy) and premenopausal otherwise.¹⁶¹⁹²⁰

Body mass index

Body mass index (BMI) is an index which is determined based on the measurement of weight and height and is classified as high if the value is $\geq 25 \text{ kg/m}^2$.¹⁶

Age (<30 years, 30–49 years, >50 years), 17²⁰ residence (rural vs urban), occupation (unemployed vs employed), exposure to smoking dried meat, use of industry processed foods, lack of intake of milk, fruits and sea foods, high cholesterol (total cholesterol > 200 mg/dL),²¹ energy (wood/charcoal/kerosene/animal source;fuel source dung vs electricity), lack of physical activity, contraceptive use, family history of cancer, history of abortion, absence of breastfeeding, benign breast disease and breast injury, exposure to radiation, anaemia and thrombocytosis were also the variables for which data were sought in the literature.

Study selection and data extraction

All the articles searched from the databases were imported into EndNote V.X7, and duplicates were removed. Based

by inviting and discussing with the third author (DRT). Data extraction was performed using the Joanna Briggs Institute (JBI) data extraction format.^{22–24} The data extraction format included the primary author, publication year, study period, region, study area, study setting, study design, study population, publication status, sample size, response rate and the number of cases of breast cancer. For the second outcome, data were extracted into

rotected by copyright JBI meta-analysis of statistics assessment and review instrument quality appraisal tool was used to assess the quality of the articles.²⁴ The JBI parameters included an appropriate sampling frame, proper sampling technique, study subject and setting description, sufficient time to exposure measurement, use of valid methods for the identified conditions, a valid measurement for variables and conditions, using appropriate statistical analysis including rised as low (total score of ≤ 2), moderate (total score of $\leq 3-4$) or high (total score of ≤ 5) in (nseignem es related hood.²⁴ The quality of the included studies was assessed by two independent authors (ATS and DRT). The discrepancy during the quality appraisal of the studies was resolved by the agreement of the two reviewers. Finally, papers with an overall quality score of <37.5%an and/or those not reporting the outcome of interest were excluded from the systematic review and meta-analysis (online supplemental additional file 2).

Data synthesis strategy

, A The data were extracted into Microsoft Excel. Then it was exported to STATA software, V.14, for further analysis. The SEs of the included studies were calculated using the formula $SE = \sqrt{\frac{p(1-p)}{n}}$. The I² statistics and the values of p of the Cochrane Q test were used to identify the heterogeneity problem. The values of p of the Cochrane Q test <0.1 were used to indicate the presence of heterogeneity among the studies. The Higgins I^2 test statistics was used to calculate the percentage of total variance due to heterogeneity across the studies. Heterogeneity was declared for the I^2 value >20%.²³ As a remedial measure for the heterogeneity among the studies by the test $\overset{\circ}{a}$ statistic, the DerSimonian-Laird's impact was evaluated **3** using a random-effects model.²⁵ Moreover, the subgroup analysis by region, study design, study setting and study population was done to identify the possible source of heterogeneity. The effect sizes were expressed as proportions and ORs along with a 95% CIs. Moreover, the 95% prediction interval was computed by using the comprehensive meta-analysis to indicate the true proportion in the comparable population.²⁶ Forest plots were used to display the results of the meta-analysis. Publication bias



Pe

, and



Figure 1 Flow chart of the selection of studies for the systematic review and meta-analysis of breast cancer and its determinants in Ethiopia.

was investigated graphically using a funnel plot and statistically using Egger's weighted regression and/or Begg's rank correlation tests and decided as significant at a value of p<0.05.^{27 28} A leave-one-out sensitivity meta-analysis was used to assess the robustness of the findings.

Patient and public involvement

No patient was involved in this study.

RESULT

Description of included studies in the systematic review and meta-analysis

About 1644 articles were identified through database searching while 17 of them were included in this systematic review and meta-analysis (figure 1). The total number of participants was 16055.

The studies covered the period between 2011 and 2021;^{16 18-21 29-40} 11 were cross-sectional,^{21 29-38} 6 were casecontrol,¹⁶ ^{18–20} ^{39 40} 16 were published,¹⁶ ¹⁸ ¹⁹ ²¹ ^{29–40} 1 was unpublished,²⁰ the majority (15) were institution-based studies, $^{16\,18-21\,29-33\,35\,36\,38-40}$ and 2 of them were populationbased studies.^{34 37} The majority of them (10) were conducted in Addis Ababa,^{16 18 19 29 32 34 35 37-40} followed by southern nations, nationalities and peoples (SNNP) (3 studies)^{20 31 33} and Amhara region (3 studies)^{21 30 36} (table 1).

Prevalence of breast cancer in Ethiopia

From the total of the 17 included studies, 10 articles were useed to pool the prevalence of breast cancer.^{16 18–21 29–40} Accordingly, the pooled proportion of breast cancer morbidity among those patients evaluated for cancer in Ethiopia was found to be 20.58% (95% CI 17.25%, 23.90%; I²=93.8%, p<0.000). The 95% prediction interval is located between 8% and 34%. This indicates that the true magnitude in 95% of all comparable populations falls in the interval between 8% and $34\%^{26}$ (figure 2).

ð the magnitude of breast cancer morbidity, subgroup anal-' uses ysis was done to further check for the source of heterogeneity. For the subgroup analysis by region, the proportion of breast cancer was found to be 21.76%; 95% CI 17.27%, 26.2% in Addis Ababa, 21.64%: 95% CI 15.02%, 28.27% in SNNP, and 14%; 95% CI 11.10%, 16.90% in Amhara (online supplemental additional file 43). The result of **o** subgroup analysis by study setting showed that the pooled magnitude of breast cancer morbidity was high; 22.58%; 95% CI 21.45%, 23.72% for population-based studies while ā 19.84%; 95% CI 15.03%, 24.65% for institution-based studies (online supplemental additional file 4). Furthermore, the subgroup analysis was done by the study population. Accordingly, the pooled magnitude of breast cancer is found to be 26.14%; 95% CI 19.87%, 32.42%, 19.05%; 95% CI 15.91%, 22.18% and 10.0%; 95% CI -8.59%, 28.59% among women with cancer, among general patients with cancer and among children who had cancer, respectively (online supplemental additional file 5).

Publication bias

simi The funnel plot appeared symmetrical indicating the absence of publication bias (online supplemental additional file 6). Egger's test (p=0.533) and Begg's test technologies. (p=0.727) also confirmed this because they both are nonsignificant for a value of p > 0.5.

Sensitivity analysis

A leave-one-out sensitivity analysis was done to test the reliability of the findings. According to the sensitivity analyses output, the random-effects model was robust, and no single study affected the pooled proportion of breast cancer morbidity (online supplemental additional file 7).

Determinants of breast cancer

In individual studies, factors like young age, age at menarche, residence, occupation, exposure to smoking

Author	Age	Study design	Period	Sample size	Proportion (%)	Factors
Abebe <i>et al²⁹</i>	≥18 years	Cross-sectional	1–31 June 2015	112	26.5	
Duche <i>et al¹⁶</i>	>15 years	Case-control	April to September 2017	226		Physical inactivity, postmenopausal, breast feeding, BMI ≥25,
Endalamaw <i>et al³⁰</i>	<15 years	Cross-sectional	1 January 2019 to 30 March 2019	100	10	
Gebretsadik <i>et al³¹</i>	No age limit	Cross-sectional	January 2013 to January 2019	3002	18.6	
Hailu <i>et al³²</i>	No age limit	Cross-sectional	January 2014 to December 2018	2002	29.3	
Hassen <i>et al³⁹</i>	>18 years	Case-control	May 2018 to June 2019	460		Anaemia, thrombocytosis
Hassen <i>et al</i> (2022) ¹⁹	>18 years	Case-control	May 2018 to June 2019	460		Age between 40 years and 49 years, early menarche, unemployment, milk intake, solid oil, use of unclean energy, physical inactivity, breast disease
Kibret <i>et al³³</i>	No age limit	Cross-sectional	January to June 2021	1810	25.4	
Kumie <i>et al²¹</i>	>18 years	Cross-sectional	22 January to 26 May 2020	182		High cholesterol
Mekonen <i>et al⁴⁰</i>	≥18 years	Case-control	February to April 2020	100		
Memirie <i>et al³⁴</i>	≥15 years	Cross-sectional	2012 to 2015	1105	22.9	
Shalamo ²⁰	>15 years	Case-control	1 March to 30 April 2022	408		Age, use of packed food, eating fruits and fish, contraceptive use, history of abortion, radiation exposure, breast injury, history of cancer
Solomon <i>et al³⁵</i>	No age limit	Cross-sectional	1 January 2010 to 15 December 2014	919	14.8	
Tefera <i>et al³⁶</i>	No age limit	Cross-sectional	September 2014 to August 2015	540	14.1	
Timotewos <i>et al³⁷</i>	No age limit	Cross-sectional	2012 to 2013	4139	22.5	
Tolessa <i>et al¹⁸</i>	>20 years	Case-control	1 February to 30 March 2020	348		Early menarche, residence, dried meat, use of packed foods, family history of cancer, lack of breastfeeding overweight
Woldu <i>et al³⁸</i>	No age limit	Cross-sectional	November 2015 to June 2016	142	14.8	
BMI, body mass ind	ex.					
lried meat, use of	processed fo	ods, lack of inta	ke of milk, to be sta	tistically	significant i	n more than one study an

dried meat, use of processed foods, lack of intake of milk, fruits, and eating sea foods, high cholesterol, fuel source (wood, charcoal, kerosene, animal dung), lack of physical activity, menopause, contraceptive use, family history of cancer, history of abortion, benign breast disease and breastinjury, radiation exposure, absence of breastfeeding, high BMI, anaemia and thrombocytosis were found to be the determinants of breast cancer. From these, age, age at menarche, use of processed foods, high cholesterol, lack of physical activity, menopause status, family history of cancer, absence of breastfeeding, and BMI were reported

statistical significance in the meta-analysis. Accordingly, those people who consume processed foods/drinks have 2.12 (pooled odds ratio (POR)=2.12, 95% CI 1.41, 3.17, $I^2=0.0\%$, p=0.826) times more odds of breast cancer than their counterparts (online supplemental additional file 8). This meta-analysis also revealed that the risk of breast cancer is increased by 4 (pooled odds ratio=4.08; 95% CI 2.75, 6.07, I²=0.0%, p=0.888) in the presence of high cholesterol including solid oil as compared with

proportion of breast cancer among cancer patients



Figure 2 Forest plot of the pooled proportion of breast cancer among patients with cancer in Ethiopia. Note: Weights are from the ramdom-effects model.

low cholesterol (online supplemental additional file 9). Those individuals who are physically inactive had 3.27 (pooled odds ratio=3.27; 95% CI 1.80, 5.94, I²=65.2%%, p=0.090) times more odds of breast cancer than their counterparts (online supplemental additional file 10). The pooled odds of breast cancer is 2.25 (pooled odds ratio=2.25; 95% CI 1.63, 3.10, I^2 =0.0%, p=0.433) times more likely in postmenopausal women than premenopausal women (figure 3). In another way, the pooled odds of breast cancer is 3.65 (pooled odds ratio=3.65; 95% CI 0.85, 15.71) times more likely for those who have a family history of cancer as compared with those without a family history of cancer (online supplemental additional file 11). Regarding BMI, when compared with those people having normal BMI, high BMI was associated with 2.27 (pooled odds ratio=2.27; 95% CI 0.85, 6.03) times increased odds of breast cancer (online supplemental additional file 12). Those women who had no history of breastfeeding have 2.76 (pooled odds ratio=2.76; 95% CI 0.90, 7.92) times more odds of breast cancer compared



Figure 3 The pooled OR showing the association between menopausal status and breast cancer in Ethiopia.

Protected by copyright, including with their counterparts (online supplemental additional for uses related file 13).

DISCUSSION

In total, 24 articles were assessed for inclusion and 7 were excluded from this review. The reason for exclusion was q mainly the lack of an intended outcome report and nonsimilarity of the study population. Here are the citations of excluded studies.^{41–47} Data of 17 articles were extracted: 10 for prevalence and 7 for determinants. Accordingly, the pooled proportion of breast cancer morbidity in Ethiopia is found to be 20. 58% (95% CI 17.25%, 23.90%). ∃ Although the result seems to be low when compared with the result in Iranian women (23.6%),⁴⁸ it is still high when ≥ compared with the age-standardised incidence rate of breast cancer in Ethiopia (12.1 per 100000 population).⁵ The observed variation could be due to a difference in ğ the denominator. As shown in the subgroup analysis, the proportion of breast cancer morbidity varies in different situations. For example, the proportion of pooled breast cancer morbidity is higher (26.14%) in women diagnosed with cancer, which is even higher than the one in Iranian women⁴⁸ and 19.05% among general patients with cancer. This shows that breast cancer varies depending on no the study population. In another way, this finding is low as compared with the breast cancer cases (25%) among $\overset{\circ}{\mathbf{G}}$ women newly diagnosed with cancers in the GLOBOCAN 2012 Study and GLOBOCAN 2018 Study^{49 50} and the study in the USA (29%).⁵¹ The difference in socioeconomic and demographic conditions might be the possible reason for this variation. Developed countries have improved cancer detection, registration and reporting as compared with Ethiopia, which could show a difference in breast cancer proportions between countries. Although the prevalence of breast cancer morbidity seems low when compared with developed settings, this result is still high in comparison

e

with the previous estimation. This alerts us to the fact that breast cancer deserves attention, especially in women. The subgroup analysis in this study also indicates a high prevalence of breast cancer disease among women who suffers from different forms of cancer.

In this systematic review and meta-analysis, factors such as the use of processed foods/drinks, high cholesterol, lack of physical activity, postmenopausal status, family history of cancer, absence of breastfeeding and high BMI including obesity were reported as risk factors for breast cancer. Accordingly, a family history of cancer including breast cancer was reported as a risk factor for breast cancer (pooled odds ratio=3.65; 95% CI 0.85, 15.71). This finding is consistent with previous studies conducted in Ethiopia,¹⁴ Iran,^{52 53} the UK,⁵⁴ China⁵⁵ and Malaysia.⁵⁶ This might be due to the presence of some inherited defect that could facilitate the development of the disease. Given that biological exposure is non-modifiable, screening and follow-up of the breast condition would help to get timely treatment that can halt the bad consequences of the disease.¹⁴

This study also revealed non-breastfeeding as a risk factor for breast cancer which is in line with the findings of a previous systematic review,^{14 57} studies done in China,⁵⁵ Iran⁵² and the USA,^{58 59} where studies conducted stated that breastfeeding minimises the risk of breast cancer. The possible reason could be because of the hormonal effect of breastfeeding for the protection or reduction of breast cancer. Both the current study and previous studies revealed the protective effect of breastfeeding for breast cancer. The possible mechanism for the observed protective probability of breast cancer in this study might be attributed to the differentiation induced to the breast lobe by lactation that might transform cancer-prone stem cell 1 to refractive stem cell 2.60 There might also be less exposure of breast tissues to hormones as breastfeeding inhibits ovulation and the hormones from the ovulation cycles.⁶¹ The result is a good indicator for the promotion of breastfeeding which has dual benefits for both the mother and the child.

This current finding also showed that high BMI is a risk factor for breast cancer in which people with high BMI were about 2.27 times more likely to develop breast cancer than their counterparts. This finding is similar to the previous study,^{56 59 62-65} and studies in Iran,^{52 53} but contrasts with the finding of the study conducted in northern California.⁶⁶ High BMI including obesity is found to be a risk factor for breast cancer in postmeno-pausal women.⁶⁷⁻⁶⁹ Increased body fat might increase the level of circulating oestrogens and decrease the levels of sex hormone-binding globulin.⁷⁰ Besides, the inflammation that accompanies obesity might also contribute to breast cancer development.⁷¹

In this study, lack of physical activity was found to be a risk factor for breast cancer and this has also been reported in previous studies.^{52 72 73} The possible explanation for the association between breast cancer and lack of physical activity might be that physical inactivity could increase the probability of fat accumulation in the body as some studies^{52 53} found that obesity is a risk factor for breast cancer. Therefore, adherence to regular physical exercise and healthy food would help to control weight and reduce body fat given that both overweight and physical inactivity are the two modifiable and related risk factors for breast cancer.

This study revealed that the use of processed food and/or drinks was a risk factor for breast cancer. This is consistent with the findings of the individual studies \neg conducted in Latin America,⁷⁴ Iran⁷⁵ and in other reviews.⁶³⁷⁶ According to this study, consumption of packed food/drinks was found to be a risk factor for breast cancer. This result is in line with the study findings of other countries which imply that a decrease in the intake of packed or ultraprocessed food or drink should be encouraged to reduce the incidence of breast cancer among women.^{63 74-76} The possible reason for this association might be due to the presence of different additives to processed foods during the processes that could initiate cancer development.⁷⁷ Other reasons might be that packed foods are rich in energy/added sugar, saturated and transfatty acids, and salt and have low fibre content and vitamins, that would increase the risk of breast cancer.⁷⁸

In this review, a positive association between high cholesterol level and breast cancer was found. However, studies are contradictory in this regard. Some studies found high cholesterol as a risk factor⁷⁹ while some found it as a protective factor.^{80 81} Those studies that found the protective effect of high cholesterol explained it as 'statin-the cholesterol-lowering medication might reduce the breast cancer risk too'.^{82 83} In this study, the total \vec{a} cholesterol including the use of hard oil was used as high \exists . cholesterol and was found to be associated with increased breast cancer risk. The possible reason for the positive 🦉 association between high cholesterol and breast cancer is that cholesterol is the precursor for oestrogen which is the cause of breast cancer.^{84 85} Women with high body fat might have an increased risk of breast cancer though their BMI is normal.

Moreover, the current study revealed that postmenopausal status is one of the risk factors for breast cancer which is consistent with the previous studies^{14 86} which indicated that breast cancer risk is higher in postmenopausal than premenopausal women. However, this result seems to contradict the finding in another meta-analysis in which premenopausal women had about 43% higher & risk of breast cancer than postmenopausal women of the same age. In another way, that study added that postmenopausal women with high body fat had an increased risk of breast cancer than premenopausal women.⁶⁹ Hence, the association between the increased possibility of breast cancer and postmenopausal status in this study might be justified as those postmenopausal women could have high body fat as well. Another possible explanation is that postmenopausal women in this study might have reached menopause at later age commonly after

50 years as late menopause is found to be a risk factor in another study,⁵⁵ because extended menstruation could lead to increased exposure of the breast tissue to hormones like oestrogen.⁶³ Besides, the use of postmenopausal hormone replacement therapy couldn't be ruled out from the possible reasons as this might increase the risk of breast cancer in postmenopausal women.⁵⁷ This implies that the hormonal change in premenopausal and postmenopausal women contributes to a risk or solution for breast cancer. Although there are differences in explanatory models regarding the postmenopausal stage, it is imperative to care for oneself because age extremes are mostly known to have a high risk for disease including chronic conditions like breast cancer.¹⁷

This study has its own implications for research, practice and policy. The practical implication is that a programme on non-communicable diseases like breast cancer in the country should be strengthened to combat the problem given the prevalence in this study is high. On top of that, a protective effect of breastfeeding found in this study implies that programmes that address breastfeeding promotion should incorporate the protective role of breastfeeding in their promotion activities. This research also alerts future research to investigate factors like residence, occupation, smoke-dried meat consumption, unclean energy sources, and the protective effects of foods like milk, seafood, and fruits including their risk and protective mechanisms. The issue of postmenopausal status is still non-conclusive in the literature. It deserves further analysis to put under its appropriate classification (risk or protective factor for breast cancer). Moreover, researchers and policy makers should work together on how to intervene and prevent the consumption of globalised and commercially processed foods as they are contributing to the breast cancer burden. The infection prevention policy of the country should be revisited to incorporate the prevention of non-communicable diseases. Although the disease is partly due to nonmodifiable risk factors, the presence of modifiable factors calls for all concerned bodies to focus on the disease to prevent the disease, diagnose and treat timely, and minimise the risk of death and economic impact. This may include the initiation and strengthening of breast cancer screening in the country.

The strengths of this study are that it is the first systematic review and meta-analysis on breast cancer in Ethiopia which pooled the prevalence of breast cancer. Next, it shares the strengths of systematic review and metaanalysis as the evidence generated from this systematic review and meta-analysis might be more representative of the country's situation than pocket studies. Third, the study estimated the prediction interval for the result obtained which is uncommon in many previous metaanalyses. However, the review was not free of limitations. The first limitation could be the narrow scope of the review in which a single country is covered. Nonetheless, the studied country can use the results to consider their policy decisions. The other drawback was that some of the regions had no primary studies regarding breast cancer and were not included in this review. The majority of the studies were done in Addis Ababa city which is the country's capital. Another drawback is that the evidence pooled together was merely from observational studies (cross-sectional and case-control studies).

Conclusion

In Ethiopia, out of five patients evaluated for cancer disease, one received a diagnosis of breast cancer. Additionally, more than a quarter of cancer disease in women is breast cancer, according to this study. In another way, the use of processed foods, high BMI, high cholesterol, physical inactivity, postmenopausal status, family history of cancer and lack of breastfeeding were the facilitators of breast cancer development.

Postmenopausal women, in particular late menopause **procession**, should stick to the lifestyle modifications that help to control body fat. It would be better if the people of Ethiopia use food sources such as fruits and vegetables, homegrown varieties of crops and the like, rather than seeking to adopt the westernised food culture (processed foods). It is highly recommended to practice regular physical exercise to regulate body weight, and body fat and then to protect against the risk of breast cancer. Appropriate breastfeeding should be practised for at least 2 years after delivery as this contributes to the minimisation of breast cancer risk. Regular breast examination should be practised to detect and control the problem timely.

X Adisu Tafari Shama @adisu Tafari Shama

Contributors ATS conceptualised the study, designed the methods, wrote the protocol, searched, screened and critically evaluated the studies, extracted the data, analysed the data, and wrote the manuscript. DRT was involved in the critical appraisal of the studies and data extraction, and wrote the first draft of the result. AED searched, screened and critically appraised the studies, extracted the data, and interpreted the result. ML searched and screened the studies, drafted the methods and wrote the introduction for the study. MCC and ETG extracted the data and prepared the manuscript. JWF, BRF and BB designed the methods, searched the studies, and extracted and analysed the data. ATS is the guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The data extracted from included studies and analysed in this review are available from the corresponding author upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

9

Open access

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Adisu Tafari Shama http://orcid.org/0000-0002-8963-9390 Dufera Rikitu Terefa http://orcid.org/0000-0002-7064-7163 Adisu Ewunetu Desisa http://orcid.org/0000-0003-4465-6629 Matiyos Lema http://orcid.org/0000-0002-6411-4571 Melese Chego Cheme http://orcid.org/0000-0002-0237-4028 Edosa Tesfaye Geta http://orcid.org/0000-0002-7050-6846 Jira Wakoya Feyisa http://orcid.org/0000-0002-9774-2398 Bikila Regassa Feyisa http://orcid.org/0000-0002-7191-0218 Bayise Biru http://orcid.org/0000-0003-0149-9379

REFERENCES

- 1 Hadgu E, Seifu D, Tigneh W, et al. Breast cancer in Ethiopia: evidence for geographic difference in the distribution of molecular subtypes in Africa. BMC Womens Health 2018;18:40.
- 2 Ferlay J, Colombet M, Soerjomataram I, et al. Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. Int J Cancer 2019;144:1941–53.
- 3 Ruibal A, Benlloch JM, Olmos RV, et al. Molecular imaging in breast cancer. J Oncol 2012;2012:426260.
- 4 Sung H, Ferlay J, Siegel RL, *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021;71:209–49.
- 5 Awedew AF, Asefa Z, Belay WB. National Burden and Trend of Cancer in Ethiopia, 2010-2019: a systemic analysis for Global burden of disease study. *Sci Rep* 2022;12:12736.
- 6 Kashyap D, Pal D, Sharma R, et al. Global Increase in Breast Cancer Incidence: Risk Factors and Preventive Measures. *Biomed Res Int* 2022;2022:9605439.
- 7 Porter P. "Westernizing" women's risks? Breast cancer in lowerincome countries. N Engl J Med 2008;358:213–6.
- 8 Anderson BO, Jakesz R. Breast cancer issues in developing countries: an overview of the Breast Health Global Initiative. *World J Surg* 2008;32:2578–85.
- 9 Kim JH, Lim JS. Early menarche and its consequence in Korean female: reducing fructose intake could be one solution. *Clin Exp Pediatr* 2021;64:12–20.
- 10 da Costa Vieira RA, Biller G, Uemura G, et al. Breast cancer screening in developing countries. *Clinics (Sao Paulo)* 2017;72:244–53.
- 11 Shulman LN, Willett W, Sievers A, et al. Breast cancer in developing countries: opportunities for improved survival. J Oncol 2010;2010:595167.
- 12 Coughlin SS. Social determinants of breast cancer risk, stage, and survival. *Breast Cancer Res Treat* 2019;177:537–48.
- 13 Adebamowo CA, Adekunle OO. Case-controlled study of the epidemiological risk factors for breast cancer in Nigeria. *Br J Surg* 1999;86:665–8.
- 14 Solbana LK, Chaka EE. Determinants of breast cancer in Ethiopia: a systematic review and meta-analysis. *Ecancermedicalscience* 2023;17:1624.
- 15 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009;6:e1000097.
- 16 Duche H, Tsegay AT, Tamirat KS. Identifying Risk Factors of Breast Cancer Among Women Attending Selected Hospitals of Addis Ababa City: Hospital-Based Unmatched Case-Control Study. *Breast Cancer* (*Dove Med Press*) 2021;13:189–97.
- 17 Centers for Disease Control and Prevention (CDC). Breast Cancer Risk Factors, Available: https://www.cdc.gov/breast-cancer/riskfactors/index.html
- 18 Tolessa L, Sendo EG, Dinegde NG, et al. Risk Factors Associated with Breast Cancer among Women in Addis Ababa, Ethiopia: Unmatched Case-Control Study. Int J Womens Health 2021;13:101–10.
- 19 Hassen F, Enquselassie F, Ali A, et al. Association of risk factors and breast cancer among women treated at Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia: a case-control study. BMJ Open 2022;12:e060636.

- 20 Shalamo T. n.d. Predictors of Breast Cancer among Women Attending Hawassa University Comprehensive Specialized Hospital, Hawassa.
- 21 Kumie G, Melak T, Wondifraw Baynes H. The Association of Serum Lipid Levels with Breast Cancer Risks Among Women with Breast Cancer at Felege Hiwot Comprehensive Specialized Hospital, Northwest Ethiopia. *Breast Cancer (Dove Med Press)* 2020;12:279–87.
- 22 Stern C, Lizarondo L, Carrier J, et al. Methodological guidance for the conduct of mixed methods systematic reviews. JBI Evd Synth 2020;18:2108–18.
- 23 Higgins JPT, Altman DG, Gøtzsche PC, *et al.* The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- 24 Aromataris E, Munn Z. JBI manual for evidence synthesis. 2020. Available: https://jbi-global-wiki.refined.site/space/MANUAL
- 25 DerSimonian R, Laird N. Meta-analysis in clinical trials. Control Clin Trials 1986;7:177–88.
- 26 Borenstein M. Research Note: In a meta-analysis, the I² index does not tell us how much the effect size varies across studies. J Physiother 2020;66:135–9.
- 27 Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088–101.
- 28 Egger M, Davey Smith G, Schneider M, *et al.* Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
- 29 Abebe E, Abebe H. Types of Cancers Diagnosed and the Preference of Families of Adult Patients with Cancer about Disclosing Diagnosis to the Patients. *Ethiop J Health Sci* 2017;27:255–62.
- 30 Endalamaw A, Assimamaw NT, Ayele TA, *et al*. Prevalence of childhood Cancer among children attending referral hospitals of outpatient Department in Ethiopia. *BMC Cancer* 2021;21:271.
- 31 Gebretsadik A, Bogale N, Negera DG. Epidemiological Trends of Breast Cancer in Southern Ethiopia: A Seven-Year Retrospective Review. *Cancer Control* 2021;28:10732748211055262.
- 32 Hailu HE, Mondul AM, Rozek LS, et al. Descriptive Epidemiology of breast and gynecological cancers among patients attending Saint Paul's Hospital Millennium Medical College, Ethiopia. PLoS ONE 2020;15:e0230625.
- 33 Kibret YM, Leka YA, Tekle NF, et al. Patterns of cancer in Wolaita Sodo University Hospital: South Ethiopia. PLoS ONE 2022;17:e0274792.
- 34 Memirie ST, Habtemariam MK, Asefa M, et al. Estimates of Cancer Incidence in Ethiopia in 2015 Using Population-Based Registry Data. J Glob Oncol 2018;4:1–11.
- 35 Solomon S, Mulugeta W. Diagnosis and Risk Factors of Advanced Cancers in Ethiopia. *J Cancer Prev* 2019;24:163–72.
- 36 Tefera B, Assefa M, Abebe B, et al. Patterns of Cancer in University of Gondar Hospital: North-West Ethiopia. J Oncol Med Pract 2016;1:106.
- 37 Timotewos G, Solomon A, Mathewos A, et al. First data from a population based cancer registry in Ethiopia. Cancer Epidemiol 2018;53:93–8.
- 38 Woldu M, Legese D, Abamecha F, et al. The prevalence of cancer and its associated risk factors among patients visiting oncology unit, Tikur Anbessa Specialized Hospital, Addis Ababa-Ethiopia. J Cancer Sci Ther 2017;9:1948–5956.
- 39 Hassen F, Enquoselassie F, Ali A, *et al.* Socio-demographic and Haematological Determinants of Breast Cancer in a Tertiary Health Care and Teaching Hospital in Addis Ababa, Ethiopia. *Eth J Health Dev* 2021;35.
- 40 Mekonen S, Ibrahim M, Astatkie H, et al. Exposure to organochlorine pesticides as a predictor to breast cancer: A case-control study among Ethiopian women. PLoS One 2021;16:e0257704.
- 41 Schwartz AD, Adusei A, Tsegaye S, *et al.* Genetic Mutations Associated with Hormone-Positive Breast Cancer in a Small Cohort of Ethiopian Women. *Ann Biomed Eng* 2021;49:1900–8.
- 42 Schwartz AD, Adusei A, Tsegaye S, et al. Genetic mutations associated with hormone-positive breast cancer in ethiopian women. Oncology [Preprint].
- 43 Tesfaw LM, Teshale TA, Muluneh EK. Assessing the Incidence, Epidemiological Description and Associated Risk Factors of Breast Cancer in Western Amhara, Ethiopia. *Breast Cancer Manag* 2020;9:BMT47.
- 44 Tekle G, Dutamo Z. Survival Analysis of Determinants of Breast Cancer Patients at Hossana Queen Elleni Mohammad Memorial Referral Hospital, South Ethiopia: Bayesian Application of Hypertabastic Proportional Hazards Model. 2019.
- 45 Sm A. Trends of Breast Cancer in Ethiopia. *Int J Cancer Res Mol Mech* 2016;2.
- 46 Ayele W, Addissie A, Wienke A, et al. Breast Awareness, Self-Reported Abnormalities, and Breast Cancer in Rural Ethiopia: A

Open access

Survey of 7,573 Women and Predictions of the National Burden. Oncologist 2021;26:e1009–17.

- 47 Ayele W, Führer A, Braun GA, et al. Breast cancer morbidity and mortality in rural Ethiopia: data from 788 verbal autopsies. BMC Womens Health 2022;22:89.
- 48 Kazeminia M, Salari N, Hosseinian-Far A, et al. The Prevalence of Breast Cancer in Iranian Women: A Systematic Review and Meta-Analysis. Ind J Gynecol Oncolog 2022;20.
- 49 Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. Int J Cancer 2015;136:E359–86.
- 50 Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin 2018;68:394–424.
- 51 DeSantis CE, Fedewa SA, Goding Sauer A, *et al.* Breast cancer statistics, 2015: Convergence of incidence rates between black and white women. *CA Cancer J Clin* 2016;66:31–42.
- 52 Khoramdad M, Solaymani-Dodaran M, Kabir A, *et al.* Breast cancer risk factors in Iranian women: a systematic review and meta-analysis of matched case-control studies. *Eur J Med Res* 2022;27:311.
- 53 Ataollahi MR, Sharifi J, Paknahad MR, et al. Breast cancer and associated factors: a review. J Med Life 2015;8:6–11.
- 54 Brewer HR, Jones ME, Schoemaker MJ, *et al.* Family history and risk of breast cancer: an analysis accounting for family structure. *Breast Cancer Res Treat* 2017;165:193–200.
- 55 Liu H, Shi S, Gao J, *et al.* Analysis of risk factors associated with breast cancer in women: a systematic review and meta-analysis. *Transl Cancer Res* 2022;11:1344–53.
- 56 Abubakar M, Sung H, Bcr D, et al. Breast cancer risk factors, survival and recurrence, and tumor molecular subtype: analysis of 3012 women from an indigenous Asian population. Breast Cancer Res 2018;20:114.
- 57 Anothaisintawee T, Wiratkapun C, Lerdsitthichai P, et al. Risk factors of breast cancer: a systematic review and meta-analysis. Asia Pac J Public Health 2013;25:368–87.
- 58 Kwan ML, Bernard PS, Kroenke CH, et al. Breastfeeding, PAM50 Tumor Subtype, and Breast Cancer Prognosis and Survival. J N C I 2015;107:djv087.
- 59 Connor AE, Visvanathan K, Baumgartner KB, *et al.* Pre-diagnostic breastfeeding, adiposity, and mortality among parous Hispanic and non-Hispanic white women with invasive breast cancer: the Breast Cancer Health Disparities Study. *Breast Cancer Res Treat* 2017;161:321–31.
- 60 Russo J, Moral R, Balogh GA, et al. The protective role of pregnancy in breast cancer. Breast Cancer Res 2005;7:131–42.
- Bernstein L. Epidemiology of endocrine-related risk factors for breast cancer. J Mammary Gland Biol Neoplasia 2002;7:3–15.
- 62 Morra A, Jung AY, Behrens S, *et al.* Breast Cancer Risk Factors and Survival by Tumor Subtype: Pooled Analyses from the Breast Cancer Association Consortium. *Cancer Epidemiol Biomarkers Prev* 2021;30:623–42.
- 63 Łukasiewicz S, Czeczelewski M, Forma A, et al. Breast Cancer-Epidemiology, Risk Factors, Classification, Prognostic Markers, and Current Treatment Strategies-An Updated Review. *Cancers (Basel)* 2021;13:4287.
- 64 Barnett GC, Shah M, Redman K, et al. Risk factors for the incidence of breast cancer: do they affect survival from the disease? J Clin Oncol 2008;26:3310–6.
- 65 Reeves GK, Patterson J, Vessey MP, et al. Hormonal and other factors in relation to survival among breast cancer patients. Int J Cancer 2000;89:293–9.
- 66 Cespedes Feliciano EM, Kwan ML, Kushi LH, et al. Body mass index, PAM50 subtype, recurrence, and survival among patients with nonmetastatic breast cancer. Cancer 2017;123:2535–42.

- 67 Greenwald P. Role of dietary fat in the causation of breast cancer: point. Cancer Epidemiol Biomarkers Prev 1999;8:3–7.
- 68 Liu K, Zhang W, Dai Z, et al. Association between body mass index and breast cancer risk: evidence based on a dose-response metaanalysis. Cancer Manag Res 2018;10:143–51.
- 69 Menarche, menopause, and breast cancer risk: individual participant meta-analysis, including 118 964 women with breast cancer from 117 epidemiological studies. *Lancet Oncol* 2012;13:1141–51.
- 70 Khandekar MJ, Cohen P, Spiegelman BM. Molecular mechanisms of cancer development in obesity. *Nat Rev Cancer* 2011;11:886–95.
- 71 Carmichael A. Review article: Obesity as a risk factor for development and poor prognosis of breast cancer. *BJOG* 2006;113:1160–6.
- 72 Friedenreich CM, Stone CR, Cheung WY, et al. Physical Activity and Mortality in Cancer Survivors: A Systematic Review and Meta-Analysis. *JNCI Cancer Spectr* 2020;4:pkz080.
- 73 Lahart IM, Metsios GS, Nevill AM, et al. Physical activity, risk of death and recurrence in breast cancer survivors: A systematic review and meta-analysis of epidemiological studies. Acta Oncol 2015;54:635–54.
- 74 Romieu I, Khandpur N, Katsikari A, et al. Consumption of industrial processed foods and risk of premenopausal breast cancer among Latin American women: the PRECAMA study. BMJNPH 2022;5:1–9.
- 75 Behjat M, Nazari J, Najafi F, et al. n.d. Dietary patterns and risk of breast cancer: case control study in the west of Iran Dietary patterns and risk of breast cancer in Iran. *Epidemiol Health*.
- 76 Kazemi A, Barati-Boldaji R, Soltani S, et al. Intake of Various Food Groups and Risk of Breast Cancer: A Systematic Review and Dose-Response Meta-Analysis of Prospective Studies. Adv Nutr 2021;12:809–49.
- 77 Shu L, Zhang X, Zhu Q, et al. Association between ultra-processed food consumption and risk of breast cancer: a systematic review and dose-response meta-analysis of observational studies. *Front Nutr* 2023;10:1250361.
- 78 Fiolet T, Srour B, Sellem L, et al. Consumption of ultra-processed foods and cancer risk: results from NutriNet-Santé prospective cohort. BMJ 2018;360:k322.
- 79 Ha M, Sung J, Song Y-M. Serum total cholesterol and the risk of breast cancer in postmenopausal Korean women. *Cancer Causes Control* 2009;20:1055–60.
- 80 Garcia-Estevez L, Moreno-Bueno G. Updating the role of obesity and cholesterol in breast cancer. *Breast Cancer Res* 2019;21:35.
- 81 Touvier M, Fassier P, His M, et al. Cholesterol and breast cancer risk: a systematic review and meta-analysis of prospective studies. Br J Nutr 2015;114:347–57.
- 82 Zhong S, Zhang X, Chen L, et al. Statin use and mortality in cancer patients: Systematic review and meta-analysis of observational studies. Cancer Treat Rev 2015;41:554–67.
- 83 Jeong GH, Lee KH, Kim JY, *et al.* Statin and Cancer Mortality and Survival: An Umbrella Systematic Review and Meta-Analysis. *J Clin Med* 2020;9:326.
- 84 Eliassen AH, Missmer SA, Tworoger SS, et al. Endogenous steroid hormone concentrations and risk of breast cancer among premenopausal women. J Natl Cancer Inst 2006;98:1406–15.
- 85 Missmer SA, Eliassen AH, Barbieri RL, et al. Endogenous estrogen, androgen, and progesterone concentrations and breast cancer risk among postmenopausal women. J Natl Cancer Inst 2004;96:1856–65.
- 86 Surakasula A, Nagarjunapu GC, Raghavaiah KV. A comparative study of pre- and post-menopausal breast cancer: Risk factors, presentation, characteristics and management. *J Res Pharm Pract* 2014;3:12–8.