

BMJ Open Enabling participation in community-dwelling children and young people with acquired brain injuries and their families: a theory-based, evidence-based and person-based approach to intervention development

Rachel Keetley ^{1,2}, Joseph C Manning ^{2,3}, Jade Kettlewell^{4,5}, Jane Williams ², Emily Bennett,² Victoria Lyon,⁶ Kate Radford ^{1,5}

To cite: Keetley R, Manning JC, Kettlewell J, *et al.* Enabling participation in community-dwelling children and young people with acquired brain injuries and their families: a theory-based, evidence-based and person-based approach to intervention development. *BMJ Open* 2024;**14**:e088516. doi:10.1136/bmjopen-2024-088516

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-088516>).

Received 08 May 2024
Accepted 25 October 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Rachel Keetley;
rachel.keetley1@nhs.net

ABSTRACT

Objective To codesign and develop an intervention to promote participation and well-being in children and young people (CYP) with acquired brain injury (ABI) and family caregivers.

Design A complex intervention development study including a scoping review, mixed-methods study, co-design workshop and theoretical modelling.

Setting Community-dwelling participants in one geographical region of the UK.

Participants CYP with ABI (5–18 years) and their parents, health, education, social care and voluntary/third-sector practitioners.

Results The intervention development process using a theory-driven and evidence-informed approach, combining the Behaviour Change Wheel and the person-based approach is described. Findings from the scoping review and mixed-methods study were analysed and synthesised using the framework method and the International Classification of Functioning, Health and Disability and the Behaviour Change Wheel. Evidence of identified participation needs, barriers and facilitators was presented at the codesign workshop. The findings demonstrate the significant long-term impact of an ABI on CYP participation and both CYP and parent well-being with significant unmet family needs. Barriers and facilitators were identified, with key barriers being lack of knowledge and understanding, lack of parental and family support and a need for cross-sector collaboration and communication. Stakeholders identified potential solutions and intervention ingredients, such as the need for education for families and schools regarding long-term impact of ABI, and longer-term practical and emotional support for families. Findings from the workshop were analysed using the framework method and synthesised with previous findings using the Behaviour Change Wheel. Theoretical modelling enabled guiding principles to be identified and an intervention logic model to be produced. 'ABI-Participate' is a novel, multifaceted intervention, developed with CYP with ABI, their parents and professionals from across health, education, social care and charity sectors. Using a case

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Involving children and young people, parents and stakeholders from across health, education, social care and voluntary/third sectors at every stage of this research ensured their views and needs remained at the centre of the process and the intervention.
- ⇒ Use of a theory, evidence and person-based approach ensured a detailed and rigorous intervention development process and a theoretically and contextually informed complex intervention.
- ⇒ The findings from this study may not be generalisable, however, understanding the specific needs within the region is important in planning services and delivering care close to home.
- ⇒ Feasibility and effectiveness testing are now required.

coordination model, ABI-Participate aims to address the unmet needs and barriers of this population and includes needs assessment, goal setting, action planning, health coaching, practical and emotional support for families and multiagency liaison and collaboration, adopting an individualised needs-based approach.

Conclusion A systematic process using a theory-based, evidence-based and person-based approach resulted in a novel, codesigned, multifaceted intervention, grounded in an in-depth understanding of CYP with ABI participation needs, barriers and facilitators. Further development and refinement of the individual elements of ABI-Participate and the care pathway to support its implementation are now required prior to feasibility testing.

BACKGROUND

Sustaining an acquired brain injury (ABI) as a child or young person as a result of trauma or non-traumatic causes (eg, infection, stroke, tumour) can lead to an array of

physical, cognitive, emotional and behavioural sequelae.¹ These sequelae can impact on well-being and participation in activities at home, school and the community.^{1 2} Outcomes within this population are heterogeneous with a range of influential factors such as injury severity, location, age at injury, premorbid abilities and personal, socioeconomic and environmental factors (eg, family functioning).^{2 3} Many children and young people (CYP) under the age of 18 experience persistent or life-long effects, which impact on physical and psychological development, quality of life, educational achievement and social inclusion. These continue to emerge months or years after the event as developmental, education and social demands increase.⁴⁻⁷

The International Classification of Functioning, Health and Disability (ICF) defines participation as involvement in life situations, which enhances well-being. It is both a fundamental right and essential part of child development.^{8 9} Research has shown that participation is associated with improved quality of life, social competence, educational success, future life outcomes and overall well-being of CYP with and without disabilities.¹⁰⁻¹² CYP with ABI and their families have reported experiencing participation restrictions, negatively impacting on their well-being, with extensive unmet and unrecognised needs found to persist up to 12 years postinjury.^{1 13-16} The impact of sustaining an ABI on CYP and their family is well documented. Sudden change in roles, routines and lifestyle for families affected by ABI intensifies stress for the entire family, impacting family functioning and well-being.¹⁷ Participation restrictions for a CYP with ABI lead to loss of social interaction, isolation and marginalisation, impacting participation and well-being for the whole family unit.^{18 19} Additionally, sudden health literacy needs, increased caregiver burden, parental stress and financial hardship can impact the mental and emotional health of the whole family, including siblings.^{2 20 21} Rehabilitation interventions must consider the entire family's needs, recognising the interconnectedness of family members' and that addressing parental needs may improve CYP outcomes.^{16 21}

Rehabilitation following an ABI aims to enable individuals to achieve optimal levels of participation by reducing the impact of difficulties and maximising well-being, activities of daily living, functional ability and social integration.^{22 23} However, international variability in paediatric rehabilitation provision and follow-up leads to uncertainty regarding long-term CYP with ABI outcomes and the best way to provide long-term support.^{7 24} Identifying and addressing individual family psychosocial and systemic issues is essential to ensure rehabilitation interventions can be effective.⁴ The literature recommends family-centred rehabilitation care models, collaborative multi-system interventions and long-term regular follow-up.²⁵⁻²⁹ However, it remains unclear what components should be included and how these should be delivered to meet the needs of CYP with ABI and their families, and optimise participation and well-being.

Rehabilitation interventions are typically complex with multiple needs and factors to be addressed.²⁸ Complexity is defined by the number of interacting components, a range of possible outcomes, the need to tailor the intervention to different contexts and dependency on the behaviours of those delivering and receiving the intervention.^{30 31} This complexity makes complex interventions difficult to implement. Factors likely to affect implementation need to be understood and addressed during intervention development. When developing interventions that aim to result in a behaviour change (eg, increasing parental confidence to support their CYP with ABI), we need to understand the target behaviour, its influences, the context for delivery, as well as identify the mechanisms of change and resources required.³²

The Medical Research Council's framework for developing and evaluating complex interventions recommends the use of theory and evidence when designing an intervention.³⁰ Drawing on existing theories, such as the Behaviour Change Wheel (BCW), can help identify important and relevant factors and inform the content and delivery of an intervention.^{33 34} Evidence from previous research can help define the problem, understand context and identify target behaviours. Uncertainties or gaps in the literature can be addressed using primary data collection, such as quantitative surveys to assess outcomes or qualitative interviews and focus groups to gain deeper understanding of needs, barriers and facilitators.³² Engagement with stakeholders through primary data collection is essential to ensure population needs and context are understood and guide intervention design and implementation into real-world practice.³⁵

We describe the intervention development process for the 'ABI-Participate' intervention using an integrated theory-based, evidence-based and person-based approach.³² This approach ensured a pragmatic, systematic, rigorous intervention development process was adhered to. The process necessitates stakeholder engagement and an in-depth understanding of behaviour, its barriers and facilitators, and how implementation of an intervention could change behaviour.^{34 35} The aim was to co-design and develop an intervention to promote participation and well-being in community-dwelling CYP with ABI (all causes and severities) and family caregivers.

Theoretical frameworks

We integrated the BCW and person-based approach in our intervention development process. The BCW was selected as the most appropriate theory for developing our intervention for CYP with ABI as it provides a systematic process using theory and evidence to develop interventions.³⁶ It incorporates the COM-B model of behaviour which aids description of how *capability*, *opportunity* and *motivation* influence *behaviour*, and the Theoretical Domains Framework which subdivides the COM-B components to aid greater understanding of barriers and facilitators at individual, organisational and community levels.^{36 37} Once these have been identified, the BCW leads

Table 1 BCW components and definitions³⁶

COM-B components (for any Behaviour to occur there must be Capability, Opportunity and Motivation to do it)		TDF domains (integrative framework synthesising key theoretical constructs)	BCW intervention functions (broad categories of means by which an intervention can change behaviour)	BCW policy options (types of decisions made by authorities that help to support and enact the interventions)	BCT Taxonomy Groups (16 groups containing 93 BCTs) (active component of an intervention designed to change behaviour)
Capability	Psychological	<ul style="list-style-type: none"> ▶ Knowledge ▶ Skills ▶ Memory, attention and decision processes ▶ Behavioural regulation 	<ul style="list-style-type: none"> ▶ Education ▶ Persuasion ▶ Incentivisation ▶ Coercion ▶ Training ▶ Enablement ▶ Modelling ▶ Environmental Restructuring ▶ Restrictions 	<ul style="list-style-type: none"> ▶ Guidelines ▶ Environmental/ social planning ▶ Communication/ marketing ▶ Legislation ▶ Service provision ▶ Regulation ▶ Fiscal measures 	<ul style="list-style-type: none"> ▶ Goals and planning ▶ Feedback and monitoring ▶ Social support ▶ Shaping knowledge ▶ Natural consequences ▶ Comparison of behaviour ▶ Associations ▶ Repetition and substitution ▶ Comparison of outcomes ▶ Reward and threat ▶ Regulation ▶ Antecedents ▶ Identity ▶ Scheduled consequences ▶ Self-belief ▶ Covert learning
	Physical	<ul style="list-style-type: none"> ▶ Skills 			
Opportunity	Social	<ul style="list-style-type: none"> ▶ Social influences 			
	Physical	<ul style="list-style-type: none"> ▶ Environmental context and resources 			
Motivation	Reflective	<ul style="list-style-type: none"> ▶ Social/professional role and identity ▶ Beliefs about capabilities ▶ Optimism ▶ Beliefs about consequences ▶ Intentions ▶ Goals 			
	Automatic	<ul style="list-style-type: none"> ▶ Social/professional role and identity ▶ Optimism ▶ Reinforcement ▶ Emotion 			

BCT, Behaviour Change Technique; BCW, Behaviour Change Wheel; TDF, Theoretical Domains Framework.

developers through a process identifying the components required for the intervention. It aids identification of *'intervention functions'* to target the behaviour and barriers and *'policies'* to support intervention delivery. This leads to the selection of *'behaviour change techniques'*, specific strategies designed to change behaviour which are the active, observable, replicable and irreducible ingredient of an intervention—that is, the proposed mechanism of change³⁶ (table 1).

The person-based approach, designed for the development of health-related behaviour change interventions, integrates well with the BCW and provides a process for combining stakeholder coproduction with mixed-methods research.³⁸ It ensures that the views of individuals who will interact with the intervention (ie, key stakeholders such as CYP, parents, health, education, social care and charity practitioners) are included throughout, increasing the likelihood of the intervention being successfully implemented in real-world practice. Guiding principles are formulated, describing the key intervention design objectives which can be mapped to

BCW intervention functions and behaviour change techniques (BCTs). This theoretical modelling process facilitated the development of a logic model to describe the intervention, planned mechanisms of change, resources required and impact on outcomes.³⁹

METHODS AND RESULTS

Using the GUIDED framework for reporting intervention development, here we describe the methods and results for each stage of the intervention development process following the process as depicted in figure 1.⁴⁰

Patient and public involvement

CYP with ABI, their parents and stakeholders representing health, education, social care and voluntary/ third sectors were involved throughout the study. Four families were involved in identifying the research question and design of the study. One young adult with ABI has assisted with the data analysis and synthesis and dissemination, including being a coauthor on this paper. Findings from




Figure 1 Intervention development process and methods employed. ABI, acquired brain injury; BCT, behaviour change technique; BCW, Behaviour Change Wheel; CYP, children and young people; TDF, Theoretical Domains Framework.

each stage were disseminated to study participants during the codesign workshop. The findings and recommendations will be disseminated further via the production of a lay summary video.

Stage 1: understanding the target behaviour

To understand the target behaviour, the problem needs to be defined, target behaviour selected and specified and barriers and enablers identified. A scoping review of the literature and primary mixed-methods research was conducted to address this aim.

Synthesis of relevant literature: scoping review

We conducted a scoping review to identify relevant literature regarding the needs of CYP with ABI and their families, and whether needs were met, unmet or unrecognised. The methods and findings of this are reported elsewhere.¹³ Four themes were found regarding needs related to CYP's impairments, parent and family support, return to school and long-term aftercare. Needs were mapped onto the ICF, with a substantial proportion of needs relating to participation and contextual factors. Key findings were the impact on parents and family and the lack of communication, collaboration and long-term follow-up. A lack of awareness and understanding

underpinned all reported needs and led to many needs being unrecognised. Recommendations within the majority of the articles reviewed included the need for specialist follow-up and integrated care pathways that are CYP and family centred. However, there were gaps in the literature. The voices of CYP are limited, there is a lack of focus on personal factors such as psychological and emotional needs for the CYP and families, a lack of focus on community participation, including recreation and leisure activities, and a lack of data regarding outcomes and needs within a UK National Health Service context. These findings led to the development of a mixed-methods research study.

Mixed-methods research study

An exploratory sequential mixed-methods study was designed to explore the long-term participation and well-being needs of CYP with ABI (5–18 years) and their families, 1–4 years after injury, in one geographical region in the UK.

The study consisted of a quantitative cross-sectional survey which explored participation and well-being outcomes and goals of CYP with ABI and their parents. Qualitative interviews and focus groups were conducted with CYP with ABI, parents and stakeholders to explore needs, local context and barriers and facilitators in more depth. We began recruitment in March 2021 with all surveys, interviews and focus groups completed by November 2022. The methods and findings from each of these studies are summarised below and reported more fully elsewhere.^{16 41}

Quantitative study

Survey results demonstrated the significant long-term impact of an ABI on CYP participation and both CYP and parent well-being; 72% of CYP had severely restricted participation, 67% had reduced health-related quality of life (HRQoL).¹⁶ Around half (53%) of parents reported reduced HRQoL and family functioning and 37% of parents screened positive for anxiety/depression. Relationships were found between CYP and parental outcomes. Higher CYP participation and HRQoL was related to higher parental HRQoL and family functioning. Higher levels of parental anxiety/depression were related to lower CYP participation and parental HRQoL and family functioning. CYP and parents reported goals that mapped to the activity and participation domains of the ICF, demonstrating the importance of these activities to their well-being.

Qualitative study

The qualitative study involved CYP with ABI and their parents who had participated in the survey and health, education, social care and voluntary/third-sector stakeholders. Significant unmet participation needs were found, impacting CYP with ABI and family well-being.⁴¹ Barriers and facilitators, mapped to the BCW, spanned 'capability', 'opportunity' and 'motivation'. The greatest

barriers aligned to the TDF domains of knowledge, skills, social influences, environmental context and resources, social identity and emotion. Identified facilitators included increasing awareness and understanding, supporting parents, long-term access to specialist assessment and rehabilitation, peer support and integrated collaborative pathways.

Mapping of current service provision

Current service provision and pathways were mapped out of the information provided by stakeholders and members of the research team and study steering group who work within the clinical service (online supplemental file 1). The mapping demonstrated the complexity of communication and referral routes from acute to community health services, and between health, education and social care providers. Additionally, the lack of provision or capacity of long-term specialist support services and collaborative care pathways was clear.

Synthesis of findings

Following the person-based approach and BCW intervention development process, the findings of the literature review and mixed-methods research were collated and synthesised using the Framework Method of analysis to map the findings and themes to the ICF, COM-B and TDF.^{42 43} This enabled us to define the problem in behavioural terms, identify the target behaviour and identify and specify barriers and facilitators (online supplemental file 2). This also allowed us to consider what needs to change and at what level. Specifying the barriers provided clarity regarding those that were individual factors (CYP/family), external organisational-level (health/education systems) and community-level (society) factors. Although there were individual factors with every CYP with ABI and family experiencing unique circumstances, there were many commonalities and a multitude of external factors that impacted on families in similar ways. Four key issues were identified and used to inform the design of the intervention:

1. Reduced CYP with ABI and family participation and well-being—support needed to enable participation and improve well-being by addressing unmet needs.
2. Lack of practical, psychological and emotional support for parents—support needed for parents to enable them to navigate systems and processes and support their CYP.
3. Lack of understanding and awareness—training and education needed for those who support CYP with ABI across health, education, social care, voluntary/third and community sectors.
4. Lack of cross-sector collaboration—a need to improve communication and collaboration between sectors and access to support in the years after ABI.

A multifaceted intervention was required to target key issues that are common across the CYP with ABI population while also providing individually tailored support to meet the specific needs of CYP and their families.

Stage 2: identify intervention options

The next stage in the BCW process was to link the 'behavioural diagnosis' (online supplemental file 2) with intervention functions likely to be effective and policy categories that can aid implementation.³⁶

As we had identified a large number of barriers and facilitators, there was a need to prioritise which were to be targeted. We therefore consulted CYP, parents and stakeholders on this and asked for their help in generating potential solutions.

Codesign workshop

An experience-based codesign workshop with multiple stakeholders (CYP, parents and health, education and voluntary/third-sector practitioners) was held in person in July 2023. The aim was to codesign and develop an intervention to promote participation and well-being in CYP with ABI and family caregivers. The objectives were to present and confirm the key issues identified, discuss priorities and generate possible solutions and identify local context-specific barriers/enablers to intervention delivery.³⁹ The data collected informed the guiding principles and theoretical modelling of the intervention.

Procedure

Participants were recruited from those who participated in the interview and focus group study with an invitation to attend the workshop sent by email. Additionally, members of the research team, study steering group (healthcare professionals from the acute neurorehabilitation team) and a patient and public involvement representative were present at the workshop to assist with facilitating groups and contribute.

All participants provided written consent/assent prior to the workshop, with parents consenting for CYP under 16 years alongside their CYP's assent. An external facilitator (AH) ran the workshop on the day, to allow the research team to listen and document discussions. The four main issues identified from the previous research, and the aims of the workshop were conveyed to the participants by the researcher (RK). Three break-out groups discussed solutions focused on the main issues, each group was facilitated by a member of the research team and study steering group (KR, JCM, DC, MD). Parents and professionals were divided between two groups, one focusing on parental support and the other on cross-sector collaboration/pathway. A separate group for the CYP was supported by a facilitator, patient and public involvement representative (VL) and play specialist (AP). A range of resources were provided to each group—paper, post-it notes, pens and Lego. The 'draw, write, tell' technique was used within the CYP group with the facilitators ensuring CYP's artwork or verbal contributions were well described in written form.⁴⁴ The groups reconvened and each fed back to the whole group, where potential solutions were discussed.

The workshop findings were collated and analysed by RK using content analysis to code and categorise the data

to the COM-B/TDF using the framework method of analysis.^{42 43} The findings and themes were discussed with the research team and study steering group to ensure rigour by reviewing and triangulating the findings, validate the coding and reduce potential biases.

Findings

In total, 17 participants attended the workshop including four CYP aged 6–17, four parents (mothers), eight health, education, social care and voluntary/third-sector practitioners (including members of study steering group) and one young person PPI representative.

Ten themes emerged regarding possible solutions for the identified issues which were mapped to the COM-B/TDF (table 2). Within these themes the priorities for intervention were identified as ongoing monitoring of CYP needs and goal setting, a single point of contact and support for parents and communication and coordination between sectors. These findings were used to inform the theoretical modelling of the intervention.

Theoretical modelling

Formulate guiding principles

Using the previously synthesised findings and the findings from the codesign workshop, we developed guiding principles. These detail the key issues to be addressed, and the intervention design objectives and distinctive features that are key to successfully addressing the issues (table 3).

Mapping intervention design objectives and features to BCW

Using the BCW, we mapped the intervention design objectives to the nine intervention functions. We identified the corresponding intervention functions that are likely to be effective in addressing the identified barriers and achieving the intervention objectives. Three intervention functions, 'education', 'training' and 'enablement' were identified that could address multiple barriers (online supplemental file 3).

The next step was to consider which of the seven BCW policy options would support the delivery of the identified intervention functions.³⁶ We identified three policy options that would be appropriate for supporting the delivery of the selected intervention functions—'communication/marketing' (using print, electronic, telephonic or broadcast media), 'guidelines' (creating documents that recommend or mandate practice) and 'service provision' (delivering a service) (online supplemental file 3).

Stage 3: identify content and implementation options

Behaviour change techniques

The next step was to identify which 'behaviour change techniques' are most appropriate for the intervention objectives and functions, and which mode of delivery was best suited. Using the BCT taxonomy (v1) we identified the BCTs required for each intervention objective and function, ensuring these also correlated with the COM-B/TDF domains that were originally identified as important to target.³⁶ The guiding principles combined with the identification of BCTs and intervention components

Table 2 Workshop findings mapped to COM-B/TDF

	TDF domain	TDF construct	Theme	Workshop participant comments
Capability	Knowledge /skills	Knowledge (about condition)	Education/training	Parent-directed
		Procedural knowledge		<ul style="list-style-type: none"> ▲ Individualised approach — options — online, face-to-face, one-to-one, groups ▲ Themed educational sessions on aspects of ABI impact, ‘how to’ guides e.g., Education Health Care Plan, where to go for help, looking after yourself, what to expect, what could happen ▲ Understand triggers for seeking help and how with different options — website, charities, nurse specialist/case manager, clinic ▲ Impairment related management strategies e.g., fatigue
		Skills /skills development		School-directed
		Ability /competence		<ul style="list-style-type: none"> ▲ Teacher education re: ABI and impact for return to school and ongoing education e.g., fatigue, cognitive impairments ▲ Peer training e.g., ‘ABI Ambassadors’ in schools, school to school support ▲ Link schools to sources of training and support e.g., charities, local authority and community health training teams ▲ ‘Friends don’t understand’ C—YP peer awareness sessions Professional-directed ▲ Professional education days ▲ Educate primary care — so General Practitioners know where to go for help
Opportunity	Social influences	Social support	Peer support opportunities (whole family)	Whole family
				<ul style="list-style-type: none"> ▲ Family fun/support days ▲ Social opportunities for CYP and siblings — shared activities
				Parent-directed
				<ul style="list-style-type: none"> ▲ Parent support groups (options — local, online, face-to-face, individual, group, therapeutic)
				Professional-directed
				<ul style="list-style-type: none"> ▲ Peer support/networks (health/education/social care/charity)

Continued

Table 2 Continued

TDF domain	TDF construct	Theme	Workshop participant comments
Environmental context and resources	Resources /material resources	Digital resources	<p>Support parents/professionals with advocacy</p> <ul style="list-style-type: none"> ▶ Regional 'hub' —website —co-develop with parents ▶ Information for parents, professionals, schools ▶ Information/signposting to sources of support e.g., charities, local authority services, SENCOS etc. ▶ Idea sharing — e.g. accessible activities — families able to add to ▶ Videos/quotes from other parents ▶ Educational sections — see education/training theme ▶ Timing of accessing information — need to be able to access when need it, even years after
		Practical resources	<ul style="list-style-type: none"> ▶ Financial support —eg, help with benefits ▶ Support with return to school (eg, case manager) ▶ Point of contact and know who can help e.g., specialist nurse/case manager/charities ▶ 1:1 support to access activities (for CYP) e.g., support worker ▶ 'Disability passport' — individualised working document —needs and strategies
		Clinical service provision	<ul style="list-style-type: none"> ▶ Routine check-ups for CYP and parent —4/8 weeks, 3/6/9/12 months with team (prevent getting lost) ▶ Re-access options — need ways back in when need it ▶ Goal setting — reviews and encouragement

Continued

Table 2 Continued

TDF domain	TDF construct	Theme	Workshop participant comments
		School support	<p>Return to school pathway</p> <ul style="list-style-type: none"> ▲ Need for a plan — it's complex and need to reduce delays ▲ Multi-disciplinary team meetings to share information and signpost to support ▲ Special Educational Needs (SEN) Team involvement/Special Educational Needs Register ▲ 'Reasonable adjustments' - e.g., phased return, small groups, reduced timetable ▲ Support with Education Health Care Plan process (see Communication) ▲ Importance of time with friends — restoring friendships ▲ CYP voiced wanting to be 'treated as equal' <p>Transitions</p> <ul style="list-style-type: none"> ▲ MDT meetings — importance of communication and early planning including SEN Team
	Person × environment interaction	Communication strategies	<ul style="list-style-type: none"> ▲ Social media — educational links/ signposting to charities/accessible activities ▲ Central point of access/contact — case manager/key worker role ▲ Website — easy to find information — all in one place ▲ Multi-agency meetings involving parents regarding return to school and transitions ▲ Signposting — professional responsibility to signpost/safety net parents/resources for professionals
Motivation	Beliefs about capabilities	Self-confidence	<p>Confidence building (CYP)</p> <ul style="list-style-type: none"> ▲ Recognition of needs and that going back to school is hard ▲ Strategies to support — CYP know plan, small groups, phased ▲ Support from parents, siblings, school staff, peers
	Goals	Goal/target setting	<p>Goal setting</p> <ul style="list-style-type: none"> ▲ Independence is important (CYP) ▲ Sports and activities — restrictions because of diagnosis but need to 'focus on what can do' (CYP) ▲ Need help to set goals and encouragement to follow them (CYP)

Continued

Table 2 Continued

TDF domain	TDF construct	Theme	Workshop participant comments
Emotion	Affect	Emotional support	<ul style="list-style-type: none"> ▲ Individualised approach — options important (1:1, peers, online, informal groups) ▲ Access to counselling — flexible timing e.g., in hospital, may not be ready until years after ▲ Post traumatic stress disorder support ▲ Targeted support for different family members ▲ CYP support — emotional strategies — how to cope (CYP) ▲ Sibling support — flexible timing ▲ Family support worker (charity provided)
ABI, acquired brain injury; CYP, children and young people; MDT, multidisciplinary team; SENCOs, Special Educational Needs Co-Ordinators; TDF, Theoretical Domains Framework.			

enabled a detailed intervention plan to be added to the guiding principles table (table 3).

Mode of delivery

Deciding on the mode of intervention delivery was important. Considering the workshop findings, participants wanted a range of options, including face-to-face as individuals or in group settings, and 'distance' meetings via telephone or virtual meeting platforms. As the target population have differing needs and reside across a large geographical region, covering five counties, a range of intervention delivery modes were needed. For example, education and training for a school could be delivered virtually or in a group face-to-face. It will be important to assess the acceptability, practicality and affordability of intervention delivery within feasibility testing to ensure it is effective for families living across the region.³⁸

Logic model

Following the systematic and detailed BCW intervention design process enabled the key objectives and active ingredients for the intervention to be identified leading to the production of a logic model that included a description of the core components, necessary resources, mechanisms of change and identification of short-term and long-term impacts and outcomes for the intervention (figure 2). This was developed iteratively, through review with the research team, study steering group and patient and public representative, and refinements made. The TIDieR Checklist informed the description of the intervention.⁴⁵

The 'ABI-Participate' intervention aims to support CYP with ABI and their families to identify and address participation and well-being needs. Using a case coordination model, needs assessments and individualised goal setting and action planning would be completed with CYP and families, considering development stage and family context.^{46–48} This includes supporting participation goals through coaching CYP and parents and ensuring referrals for additional therapy or support are made where necessary. Families and professionals wanted a single point of contact and signposting to sources of information and support. 'ABI-Participate' also includes information sharing and team meeting coordination, liaison with CYP's school/college and supporting parental practical, emotional and psychological needs. The intervention would continue until needs were met or goals achieved, with a single point of contact in the long-term, should families or those working with a CYP with ABI need advice or support. Other features included reassessment of needs at routine reviews in a follow-up clinic, ahead of key educational stage transitions or as new needs emerge. Coordination of transition to adult services would also be provided. The intervention would help to improve understanding and awareness of long-term needs. It would increase support for parents and CYP to improve their participation and well-being by identifying and addressing needs of the whole family. It would help

Table 3 Guiding principles/intervention planning table

	Key issues identified in mixed methods study	Intervention design objectives	Key features of intervention to achieve objective	Mechanisms	
				Behaviour Change Technique taxonomy (v1)	Behaviour Change Wheel intervention functions
1	CYP with ABI experience severely restricted participation as a result of multiple interacting barriers	To identify and address participation restrictions of CYP with ABI and their families	Routine needs assessments Identification of unmet needs Goal setting/coaching (CYP/parents) Action Planning MDT liaison/team around child meeting Referrals/signposting Support parents	Social support (practical) Problem solving Goal setting (behaviour) Action planning Social support (practical) Social support (practical) Social support (practical and emotional)	Enablement
2	Parents and those supporting CYP with ABI lack of awareness and understanding of impact of ABI which leads to under-recognition of needs	To increase understanding about impact of ABI of those supporting CYP with ABI to increase recognition of needs	Support parents to develop health literacy Support parents to understand impact of ABI and recognise needs Provide education regarding impact of ABI to those supporting CYP and family for example, school staff	Instruction on how to perform the behaviour Information about health consequences Information about social and environmental consequences Information about health consequences Information about social and environmental consequences	Training Education
3	CYP with ABI and their families experience reduced HRQoL/well-being which impacts family functioning	To support family well-being	Offer needs-based emotional and practical support Signpost to sources of support/groups, etc MDT liaison/team around child	Social support (practical and emotional) Social support (practical) Social support (practical)	Enablement
4	Parents have a substantial care and advocacy role and experience difficulty navigating systems	To support parents to navigate systems /services effectively	Upskill parents in system navigation Support and empower parents Signposting to resources Advocacy as needed Point of contact for families	Instruction on how to perform the behaviour (impart skills) Social support (practical) Social support (practical) Social support (practical) Problem solving	Training Enablement
5	Parents and professionals experience difficulty with coordinating and communicating across sectors	Facilitate cross-sector collaboration /communication	Liaison/point of contact between services across sectors	Social support (practical)	Enablement

ABI, acquired brain injury; CYP, children and young people; HRQoL, Health-related Quality of Life; MDT, multidisciplinary team.

families and health, education, social care and voluntary/third-sector practitioners working with CYP with ABI and their families to overcome barriers by helping to coordinate cross-sector communication and collaboration. The intervention needs to be situated within and supported by a multidisciplinary neurorehabilitation service. The multidisciplinary team would need capacity to support the assessment of needs and provide targeted rehabilitation interventions, when needs are identified, such as neurocognitive interventions or higher-level physical skills training required for return to sport.

DISCUSSION

This paper has described how we used a theory-based, evidence-based and person-based approach to develop a complex intervention aimed at improving participation in CYP with ABI and their families. As recommended by the Medical Research Council's guidance, underpinning the development of complex interventions with theory and evidence is essential to ensure interventions are evidence informed and grounded in a theoretical understanding of behaviour change.^{30 32} The addition of the person-based approach ensured an in-depth understanding of




Figure 2 ABI-Participate logic model. ABI, acquired brain injury; CYP, children and young people; MDT, multidisciplinary team.

the life experiences of the population and stakeholders, their needs and views on acceptable solutions.³⁹ As this approach focuses specifically on the development of complex behavioural interventions, it was well suited to be combined with the theory and evidence approach in this work.⁴⁹

The intervention we have developed is founded on the biopsychosocial model of the ICF and multidisciplinary teamwork. It is multifaceted, family-centred and tailored to developmental stage, individual needs and contexts. It follows a rehabilitation process, involving assessment, goal setting, intervention delivery, monitoring and review.²³ Based on a case coordination model, ABI-Participate also includes a therapeutic element of goal-oriented coaching, which aims to identify and address the participation needs of the whole family. As found in this study, and recommended by the WHO, there is an increasing acknowledgement of the need for care to be coordinated to support those living with long-term conditions and their family caregivers due to the complexity of coordinating care, ensuring needs are met and undue stress prevented.^{21 48 50–52}

Gagnon *et al*⁴⁷ identified six key supportive roles that family members of adult traumatic brain injury survivors

undertake—researcher, advocate, case manager, coach, activities of daily living supporter and emotional supporter. They concluded that family members require ongoing counselling, support and education about system navigation, accessing community programmes and workplace rights to prevent burnout. A scoping review by Gardiner *et al*⁴⁸ found a diversity of terminology and descriptions used for navigation-type models for children with neuro-disabilities. However, each was characterised by four central domains: *facilitate*—integration/coordination of resources, supports and services; *provide*—information, advice and education; *intended outcomes*—improved health, behaviour and capacity and reduced patient and family distress and *guiding principles*—client-directed, family-centred and collaborative. These findings align with the findings of our study and are incorporated into the proposed ‘ABI-Participate’ intervention.

An additional element identified in our study, and by Gagnon and colleagues,⁴⁷ was coaching and supporting CYP and families in identifying and achieving participation goals. Palisano *et al*⁵³ proposed a conceptual framework for optimal participation of children with physical disabilities that considers the dynamic interaction of determinants (child, family and environment)

and dimensions (physical, social and self-engagement) of participation. Their recommendation that interventions need to be goal-oriented, family-centred, collaborative, strengths-based and ecological also aligns with our findings. ABI-Participate could be used flexibly and at different time points, recognising that needs differ at different developmental stages and points of transition (ie, more intensive support needed during educational transitions, particularly into secondary school and transition to adult services and/or employment).

Health coaching is defined as ‘a goal-oriented, client-centred partnership that is health-focused and occurs through a process of client-enlightenment and empowerment’ (p24).⁵⁴ Coaching can help patients and families identify and achieve their goals and has been shown to be effective in positively influencing health status, health behaviours and costs.²¹ There is increasing evidence supporting coaching in promoting parental self-management and empowerment, addressing parental health literacy and advocacy skills, and CYP participation in parents of CYP with chronic disabilities.^{21 55} Ogourtsova *et al*²¹ systematic review of health coaching for parents of children with developmental disabilities found heterogeneity within the interventions with some being CYP-directed, some parent-directed and some mixed. They recommended further research exploring the outcomes of the different modes of delivery and the effectiveness of these on CYP and parental outcomes, recognising that these are linked with CYP outcomes improving when parent-related outcomes improve.

Existing interventions could be integrated with ABI-Participate. For example, goal-directed interventions (eg, PREP—Pathways and Resources for Engagement and Participation), coaching interventions (eg, CO-OP—Cognitive Orientation to Occupational Performance), parent interventions (eg, Stepping Stones Triple P) or psychological interventions (eg, Acceptance and Commitment Therapy).^{55–57} However further research is recommended to further explore the acceptability and effectiveness of integrating them in this intervention.

The technological advances enabling the widespread use of virtual meeting platforms for health consultations during the COVID-19 pandemic, have made telehealth delivery of some components of ‘ABI-Participate’ possible.^{58 59} Not only are most patients and families now familiar with these platforms, but workshop participants suggested them as plausible modes of delivery. This enables specialist rehabilitation in tertiary settings to be made accessible to people living at long geographical distances and also offers an opportunity to upskill and support local providers with education and training delivered virtually. This delivery mode is in use in Rohrer-Baumgartner *et al*’s ‘Child in Context’ study.²⁸ A future feasibility study should include exploration of both its acceptability and utility, as well as how to deliver this to those without access to the internet.

Strengths and limitations

The strengths and limitations of the scoping review, survey and qualitative study are reported elsewhere.^{13 16 41} Involving CYP, parents and stakeholders from across health, education, social care and voluntary/third sectors at every stage of this research is a strength, ensuring their views and needs remained at the centre of the process and the intervention. To our knowledge, this is the first intervention development study within the CYP with ABI literature to have used a theory, evidence and person-based approach. This novel paper demonstrates the value of combining these approaches to develop a complex intervention for addressing participation needs in CYP with ABI and their families. By following the person-based approach and BCW process, a detailed and rigorous approach was employed to understand needs, identify barriers and facilitators and design the intervention (including the active ingredients/mechanisms). The mixed-methods design enabled us to define the problem further, understand the local context and, using theory, provide a detailed description of the barriers and facilitators to participation. The addition of the codesign workshop with stakeholders to prioritise and generate solutions was invaluable in ensuring the most pertinent targets were selected. The theoretical modelling process and use of guiding principles enabled us to clearly and systematically articulate and document the process of identifying the key objectives, features and functions of the intervention and selection of the policy options for implementing it. Finally, the logic model presented an overview of the intervention, the resources, core components and mechanisms as well as the contextual factors that must be considered and the outcomes that could be measured to assess effectiveness.

This study was conducted in one region within the UK and therefore findings may not generalisable, however, understanding the specific needs within the region is important in planning services and delivering care close to home. While every attempt was made to ensure diverse representation at every stage, this did not occur within the workshop, partly due to some participants being unable to attend on the day. However, the reported needs, barriers and facilitators align with those reported internationally and many of the themes from the qualitative study, which had more diverse representation, were repeated and affirmed at the workshop. This intervention has been developed iteratively, with CYP, parents and stakeholders. Further stakeholder and patient and public representative consultation and expert consensus development workshops are now required to refine, specify and confirm intervention components prior to feasibility testing.

Future directions

There is evidence of effectiveness for the different components of our intervention, but research is needed to test the feasibility of the intervention in our target population and context, and to investigate its acceptability,

deliverability and effectiveness. Within this, identification of standardised outcome measures and methods of determining whether needs are met are required to measure effectiveness. Further consideration also needs to be given to the overall care pathway for CYP with ABI in which this intervention would be situated following hospital discharge. Given the barriers reported and lack of access to rehabilitation, other elements need to be developed and delivered alongside this intervention. For example, specialist rehabilitation and review clinics, particularly for those CYP whose needs cannot be met within primary care, mental health or community therapy services (eg, cognitive, neuropsychological or higher-level motor therapy needs that do not meet referral criteria). For 'ABI-Participate' to be effective, there needs to be referral pathways for CYP with these needs. The mapping of regional service provision demonstrated that these do not exist, except for a very limited regional ABI medical follow-up clinic and neuropsychology service, with long waiting lists, further delaying access to support.

Conclusions

This research has provided an in-depth understanding of the participation and well-being needs of CYP with ABI and their families. The findings demonstrate the significant long-term impact of an ABI on CYP participation and both CYP and parent well-being with significant unmet family needs. Barriers and facilitators that families and stakeholders face in accessing support and rehabilitation are identified. Key barriers identified were a lack of knowledge and understanding of the impact of ABI across every level of society, lack of parental and family support and a need for cross-sector collaboration and communication. Providing parental support, long-term access to specialist assessment and rehabilitation, peer support and integrated collaborative pathways were identified as facilitators.

We have iteratively developed a novel, multifaceted intervention the 'ABI-Participate' intervention with CYP with ABI, their parents and stakeholders from across health, education, social care and voluntary/third sectors with the aim of addressing the unmet needs and barriers of this population. Adopting a case coordination model and an individualised needs-based approach, ABI-Participate includes needs assessment, goal setting, action planning, coaching, practical and emotional support for families and multiagency liaison and collaboration. Further refinement of the components of ABI-Participate and development of the care pathway to support its implementation are now required prior to feasibility testing.

Author affiliations

¹Centre for Rehabilitation and Ageing Research, University of Nottingham, Nottingham, UK

²Nottingham Children's Hospital, Nottingham University Hospitals NHS Trust, Nottingham, UK

³School of Healthcare, University of Leicester, Leicester, UK

⁴School for Primary Care Research, University of Nottingham, Nottingham, UK

⁵NIHR Nottingham Biomedical Research Centre, Nottingham, UK

⁶PPIE, Nottingham University Hospitals NHS Trust, Nottingham, UK

X Rachel Keetley @rachyk77 and Kate Radford @rehab_research

Acknowledgements The authors thank Denise Crozier and Melanie Dean for assisting with facilitating the workshop groups and with the intervention development, Angela Horsley for facilitating the codesign workshop and Alison Price for providing play specialist support to the CYP workshop participants. Our thanks go to the children and young people, their parents/carers and the stakeholders that took part in this study, our patient and public involvement families, the study steering group and clinical team for their ongoing support.

Contributors RK conducted the scoping review, mixed-methods study and workshop, the primary data analysis, wrote the manuscript and is guarantor. KR, JCM, EB, VL assisted with the workshop in facilitating the groups, assisted with the data analysis and contributed to writing the manuscript. KR, JCM, JK, JW supervised the research, assisted with the data analysis, framework mapping and contributed to writing the manuscript. All authors assisted with the intervention development.

Funding This study is funded by the NIHR Applied Research Collaboration East Midlands (ARC EM). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Competing interests RK received PhD studentship funding from NIHR Applied Research Collaboration-East Midlands, University of Nottingham and Health Education England. JM receives funding from NIHR (UK) and NIH (US). KR receives funding from NIHR HTA grant funding (15/130/11), NIHR Programme Grant funding, NIHR HealthTech Research Centres, EPSRC Rehabilitation Technologies Network, NIHR School of Primary Care Research, MS Society. JM is a Data Monitoring Committee member—del Nido versus St. Thomas' blood cardioplegia in the young (DESTINY) trial: a multicentre randomised controlled trial in children undergoing cardiac surgery (funded British Heart Foundation). RK is a Topic Advisor (Children and Young People) NICE Guideline Committee—Rehabilitation for Chronic Neurological Disorders. KR was an unpaid advisor to NHS England in the development of a toolkit for NHS professionals to support return to work after stroke. KR was a member of the NIHR HTA Clinical Evaluation and Trials panel between 2017–2021, JM is current member of the NIHR RfPB East Midlands panel. JK, JW, EB and VL have no competing interests to disclose.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by East Midlands-Nottingham 2 Research Ethics committee (REC-20/EM/0258) and University of Nottingham, Faculty of Medicine and Health Sciences Research Ethics Committee (FMHS 234-0323). Informed consent/assent was gained from all participants via survey completion and written consent forms.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The data that support this study will be shared upon reasonable request to the corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Rachel Keetley <http://orcid.org/0000-0001-6004-1523>

Joseph C Manning <http://orcid.org/0000-0002-6077-4169>

Jane Williams <http://orcid.org/0000-0003-0469-4698>
 Kate Radford <http://orcid.org/0000-0001-6246-3180>

REFERENCES

- Wales L, Davis K, Kelly G, et al. Long Term Participation Outcomes for Severe Acquired Brain Injury in Childhood - An Expanded Scoping Review. *Dev Neurorehabil* 2021;24:379-87.
- Analytis P, Warren N, Crichton A, et al. Coping and wellbeing in families with a child with acquired brain injury compared with families in the community. *Disabil Rehabil* 2022;44:7993-8000.
- Gordon AL, di Maggio A. Rehabilitation for children after acquired brain injury: current and emerging approaches. *Pediatr Neurol* 2012;46:339-44.
- Limond J, Adlam ALR, Cormack M. A model for pediatric neurocognitive interventions: considering the role of development and maturation in rehabilitation planning. *Clin Neuropsychol* 2014;28:181-98.
- Forsyth R, Waugh MC. Paediatric brain injury - getting there from here. *Child Care Health Dev* 2010;36:1-2.
- Anderson V, Godfrey C, Rosenfeld JV, et al. 10 years outcome from childhood traumatic brain injury. *Int J Dev Neurosci* 2012;30:217-24.
- McKinlay A, Linden M, DePompei R, et al. Service provision for children and young people with acquired brain injury: Practice recommendations. *Brain Inj* 2016;30:1656-64.
- World Health Organisation. *Towards a common language for functioning, disability and health - ICF*. Geneva: WHO, 2002.
- Imms C, Green D. *Participation: optimising outcomes in childhood-onset neurodisability*, 1st edn. London: Mac Keith Press, 2020.
- Galvin J, Froude EH, McAleer J. Children's participation in home, school and community life after acquired brain injury. *Aust Occup Ther J* 2010;57:118-26.
- Bedell GM, Dumas HM. Social participation of children and youth with acquired brain injuries discharged from inpatient rehabilitation: a follow-up study. *Brain Inj* 2004;18:65-82.
- Kolehmainen N, Ramsay C, McKee L, et al. Participation in Physical Play and Leisure in Children With Motor Impairments: Mixed-Methods Study to Generate Evidence for Developing an Intervention. *Phys Ther* 2015;95:1374-86.
- Keetley R, Radford K, Manning JC. A scoping review of the needs of children and young people with acquired brain injuries and their families. *Brain Inj* 2019;33:1117-28.
- Allonsius F, de Kloet A, Bedell G, et al. Participation Restrictions among Children and Young Adults with Acquired Brain Injury in a Pediatric Outpatient Rehabilitation Cohort: The Patients' and Parents' Perspective. *Int J Environ Res Public Health* 2021;18:1625.
- van Markus-Doornbosch F, van der Holst M, de Kloet AJ, et al. Fatigue, Participation and Quality of Life in Adolescents and Young Adults with Acquired Brain Injury in an Outpatient Rehabilitation Cohort. *Dev Neurorehabil* 2020;23:328-35.
- Keetley R, Manning JC, Williams J, et al. Child and family health-related quality of life and participation outcomes and goals after acquired brain injury: a cross-sectional survey. *Brain Inj* 2024;38:217-26.
- Rashid M, Goetz HR, Mabood N, et al. The impact of pediatric traumatic brain injury (TBI) on family functioning: A systematic review. *J Pediatr Rehabil Med* 2014;7:241-54.
- Stenberg M, Stålnacke B-M, Saveman B-I. Family experiences up to seven years after a severe traumatic brain injury-family interviews. *Disabil Rehabil* 2022;44:608-16.
- Kwan C, Gitimoghaddam M, Collet J-P. Effects of Social Isolation and Loneliness in Children with Neurodevelopmental Disabilities: A Scoping Review. *Brain Sci* 2020;10:786.
- Allonsius F, De Kloet AJ, Van Markus-Doornbosch F, et al. Parent-reported family impact in children and young adults with acquired brain injury in the outpatient rehabilitation setting. *Brain Inj* 2021;35:563-73.
- Ogourtsova T, O'Donnell M, De Souza Silva W, et al. Health coaching for parents of children with developmental disabilities: a systematic review. *Develop Med Child Neuro* 2019;61:1259-65.
- de Kloet AJ, Gijzen R, Braga LW, et al. Determinants of participation of youth with acquired brain injury: A systematic review. *Brain Inj* 2015;29:1135-45.
- Wade DT. What is rehabilitation? An empirical investigation leading to an evidence-based description. *Clin Rehabil* 2020;34:571-83.
- Haarbauer-Krupa J, Ciccio A, Dodd J, et al. Service Delivery in the Healthcare and Educational Systems for Children Following Traumatic Brain Injury: Gaps in Care. *J Head Trauma Rehabil* 2017;32:367-77.
- Botchway EN, Knight S, Muscara F, et al. Rehabilitation models of care for children and youth living with traumatic brain and/or spinal cord injuries: A focus on family-centred care, psychosocial wellbeing, and transitions. *Neuropsychol Rehabil* 2022;32:537-59.
- Diener ML, Kirby AV, Sumsion F, et al. Community reintegration needs following paediatric brain injury: perspectives of caregivers and service providers. *Disabil Rehabil* 2022;44:5592-602.
- Anaby DR, Avery L, Palisano RJ, et al. Environment-based approaches to improve participation of young people with physical disabilities during COVID -19. *Dev Med Child Neuro* 2024;66:931-8.
- Rohrer-Baumgartner N, Holthe IL, Svendsen EJ, et al. Rehabilitation for children with chronic acquired brain injury in the Child in Context Intervention (CICI) study: study protocol for a randomized controlled trial. *Trials* 2022;23.
- Botchway-Comme E, Muscara F, Greenham M, et al. Rehabilitation models of care for children and youth with traumatic brain and/or spinal cord injuries: A focus on service structure, service organization, and the barriers and facilitators of rehabilitation service provision. *Neuropsychol Rehabil* 2023;33:1697-727.
- Skivington K, Matthews L, Simpson SA, et al. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *BMJ* 2021;374:n2061.
- Rodriguez A, Smith J, Barrett D. Research made simple: developing complex interventions. *Evid Based Nurs* 2020;23:35-7.
- O'Cathain A, Croft L, Duncan E, et al. Guidance on how to develop complex interventions to improve health and healthcare. *BMJ Open* 2019;9:e029954.
- Nilsen P. Making sense of implementation theories, models and frameworks. *Implement Sci* 2015;10:53.
- Atkins L, Francis J, Islam R, et al. A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implement Sci* 2017;12:77.
- Guyatt S, Ferguson M, Beckmann M, et al. Using the Consolidated Framework for Implementation Research to design and implement a perinatal education program in a large maternity hospital. *BMC Health Serv Res* 2021;21:1077.
- Michie S, Atkins L, West R. *The behaviour change wheel: a guide to designing interventions*. London: Silverback Publishing, 2014.
- Cantwell M, Walsh DMJ, Furlong B, et al. The Development of the MedEx IMPACT Intervention: A Patient-Centered, Evidence-Based and Theoretically-Informed Physical Activity Behavior Change Intervention for Individuals Living With and Beyond Cancer. *Cancer Control* 2020;27:1073274820906124.
- Yardley L, Ainsworth B, Arden-Close E, et al. The person-based approach to enhancing the acceptability and feasibility of interventions. *Pilot Feasibility Stud* 2015;1:37.
- Yardley L, Morrison L, Bradbury K, et al. The person-based approach to intervention development: application to digital health-related behavior change interventions. *J Med Internet Res* 2015;17:e30.
- Duncan E, O'Cathain A, Rousseau N, et al. Guidance for reporting intervention development studies in health research (GUIDED): an evidence-based consensus study. *BMJ Open* 2020;10:e033516.
- Keetley R, Manning JC, Williams J, et al. Understanding barriers and facilitators to long-term participation needs in children and young people following acquired brain injuries: a qualitative multi-stakeholder study. *Brain Impair* 2024;25:1B23100.
- Lawton R, Heyhoe J, Louch G, et al. Using the Theoretical Domains Framework (TDF) to understand adherence to multiple evidence-based indicators in primary care: a qualitative study. *Implement Sci* 2016;11:113.
- Ritchie J, Lewis J, McNaughton Nicholls C, et al. *Qualitative research: a guide for social science students and researchers*, 2nd edn. London: SAGE Publications Ltd, 2014.
- Angell C, Alexander J, Hunt JA. 'Draw, write and tell': A literature review and methodological development on the 'draw and write' research method. *J Early Child Res* 2015;13:17-28.
- Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;348:bmj.g1687.
- Fadyl JK, McPherson KM. Approaches to vocational rehabilitation after traumatic brain injury: a review of the evidence. *J Head Trauma Rehabil* 2009;24:195-212.
- Gagnon A, Lin J, Stergiou-Kita M. Family members facilitating community re-integration and return to productivity following traumatic brain injury - motivations, roles and challenges. *Disabil Rehabil* 2016;38:433-41.
- Gardiner E, Wong V, Lin G, et al. Terminology and descriptions of navigation and related practices for children with neurodisability and their families: a scoping review. *BMC Health Serv Res* 2022;22:214.
- Easton S, Ainsworth B, Thomas M, et al. Planning a digital intervention for adolescents with asthma (BREATHE4T): A theory-,

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

- evidence- and Person-Based Approach to identify key behavioural issues. *Pediatr Pulmonol* 2022;57:2589–602.
- 50 Braaf S, Ameratunga S, Christie N, *et al.* Care coordination experiences of people with traumatic brain injury and their family members in the 4-years after injury: a qualitative analysis. *Brain Inj* 2019;33:574–83.
 - 51 Hudon C, Chouinard M-C, Pluye P, *et al.* Characteristics of Case Management in Primary Care Associated With Positive Outcomes for Frequent Users of Health Care: A Systematic Review. *Ann Fam Med* 2019;17:448–58.
 - 52 Budde H, Williams G, Scarpetti G, *et al.* *What are patient navigators and how can they improve integration of care.* Denmark: World Health Organisation, 2022.
 - 53 Palisano RJ, Chiarello LA, King GA, *et al.* Participation-based therapy for children with physical disabilities. *Disabil Rehabil* 2012;34:1041–52.
 - 54 Olsen JM. Health Coaching: A Concept Analysis. *Nurs Forum* 2014;49:18–29.
 - 55 Novak I, Morgan C, Fahey M, *et al.* State of the Evidence Traffic Lights 2019: Systematic Review of Interventions for Preventing and Treating Children with Cerebral Palsy. *Curr Neurol Neurosci Rep* 2020;20:3.
 - 56 Anaby DR, Law M, Feldman D, *et al.* The effectiveness of the Pathways and Resources for Engagement and Participation (PREP) intervention: improving participation of adolescents with physical disabilities. *Dev Med Child Neurol* 2018;60:513–9.
 - 57 Lebrault H, Martini R, Manolov R, *et al.* Cognitive Orientation to daily Occupational Performance to improve occupational performance goals for children with executive function deficits after acquired brain injury. *Dev Med Child Neurol* 2024;66:501–13.
 - 58 Holthe IL, Rohrer-Baumgartner N, Svendsen EJ, *et al.* Feasibility and Acceptability of a Complex Telerehabilitation Intervention for Pediatric Acquired Brain Injury: The Child in Context Intervention (CICI). *J Clin Med* 2022;11:2564.
 - 59 Kettlewell J, Lindley R, Radford K, *et al.* Factors Affecting the Delivery and Acceptability of the ROWTATE Telehealth Vocational Rehabilitation Intervention for Traumatic Injury Survivors: A Mixed-Methods Study. *Int J Environ Res Public Health* 2021;18:9744.