BMJ Open Maternal outcomes of pre-eclampsia with severe features and its determinants at Abebech Gobena Mothers and Childrens Health and Saint Peter's Specialized Hospital, Addis Ababa, Ethiopia: a crosssectional study

Mesfin Tadese ^(D), ¹ Wogene Asefa Damesa, ² Gebeyehu Shumet Solomon, ³ Getu Engida Wakie,¹ Saba Desta Tessema 💿 ,¹ Agizew Endale⁴

ABSTRACT

To cite: Tadese M, Damesa WA, Solomon GS, et al. Maternal outcomes of pre-eclampsia with severe features and its determinants at Abebech Gobena Mothers and Childrens Health and Saint Peter's Specialized Hospital, Addis Ababa, Ethiopia: a crosssectional study. BMJ Open 2024;14:e081901. doi:10.1136/ bmjopen-2023-081901

 Prepublication history for this paper is available online. To view these files, please visit the journal online (https://doi. org/10.1136/bmjopen-2023-081901).

Received 09 November 2023 Accepted 29 February 2024

Check for updates

C Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BM.J.

For numbered affiliations see end of article.

Correspondence to Mr Mesfin Tadese: mesitad031@gmail.com Objective The main objective of this study was to determine the prevalence and factors associated with unfavourable maternal outcomes among pregnant women with pre-eclampsia with severity features (PEWSF) at Abebech Gobena Maternal and Children's Health and St. Peter's Hospital, Addis Ababa, Ethiopia, 2023. **Design** A hospital-based cross-sectional study was conducted from 1 January 2023 to July 2023. The

data was collected using a structured and pre-tested questionnaire through face-to-face interviews and a review clinical chart. Data was entered using EpiData V.4.6 and analysed using SPSS V.26.0 statistical software. Binary logistic regression analysis was run to identify predictors of maternal outcome.

Setting Two hospitals in Addis Ababa, Ethiopia. Participants 348 pregnant women with PEWSF were included.

Outcome measures Unfavourable maternal outcome was defined as mothers with PEWSF that develop at least one complication, that is, eclampsia, abruption placenta, Haemolysis, Elevated Liver Enzymes and Low Platelet Count (HELLP) syndrome, acute renal failure, disseminated intravascular coagulation, cardiac failure, stroke, postpartum haemorrhage, pulmonary oedema and death. Results The overall prevalence of unfavourable maternal outcomes was 33.9% (N=118) (95% CI=28.7 to 38.8). Abruptio placenta (17.2%), HELLP syndrome (15.5%) and postpartum haemorrhage (13.8%) were common complications that occurred among mothers with PEWSF. Age above 35 years (adjusted OR (AOR) (95% CI)=2.70 (1.31 to 5.59)), rural residence (AOR (95% CI)=1.94 (1.07 to 3.53)), unemployment (AOR (95% CI)=0.35 (0.20 to 0.62)), severe blood pressure on admission (AOR (95% CI)=2.32 (1.03 to 5.19)) and complain of severe headache (AOR (95% CI)=1.91 (1.16 to 3.16)) were significant associates of unfavourable maternal outcomes. **Conclusions** The prevalence of unfavourable maternal outcomes was high compared with other studies in

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Interviews and clinical chart reviews were conducted to collect data.
- \Rightarrow One drawback was that, because the research was conducted in a hospital setting, the maternal outcome of home births was not assessed.
- Another limitation was that the study did not include unfavourable maternal outcomes after 24 hours of birth.

Ethiopia. Maternal age, residence, occupation, blood pressure on admission and severe headache have shown a statistically significant association with unfavourable maternal outcomes. Socioeconomic development and early identification of severe signs and symptoms of preeclampsia are needed to reduce unfavourable outcomes.

BACKGROUND

Protected by copyright, including for uses related to text and data mining, AI training, and Pre-eclampsia is a multisystem progressive illness distinguished by the new development of hypertension and either proteinuria or end-organ failure after 20 weeks of gestation, technologi during pregnancy, labour or post partum.¹ A combination of maternal and fetal/ placental factors is most likely the reason. g Relative placental hypoxia, ischaemia or 🖁 under-perfusion can be brought on by abnormalities in the placental vasculature early in pregnancy.² This may then cause the mother's circulation to release antiangiogenic factors, altering the mother's systemic endothelium's function and causing hypertension in addition to other disease manifestations (haematological, neurological, cardiac, pulmonary, renal and hepatic dysfunction). However, the

l simi

and

Open access

reason behind abnormal placental development and the subsequent sequence of events is still unknown.³

Pre-eclampsia complicates between 3% and 5% of pregnancies in high-income countries.⁴ In Africa, hypertension disorders during pregnancy affect 10% of pregnancies.⁵ A Zanzibar study found that pre-eclampsia with severe features (PEWSF) was prevalent in 26.3% of mothers.⁶ Besides, 19.5% of PEWSF was reported in a prospective observational study done at Saint Paul's Hospital Millennium Medical College in Ethiopia.⁷

In the USA, unfavourable maternal outcomes occurred in 10% of women with PEWSF.⁸ According to a prospective cohort study in the Sidama region of Ethiopia, women with PEWSF had a 43% higher risk of unfavourable maternal outcomes.⁹ Similarly, it was shown that 37.7% of mothers with severe pre-eclampsia/eclampsia in referral hospitals in the Amhara region had unfavourable maternal outcomes.¹⁰ Further, in Addis Ababa, Ethiopia, 36% of mothers with PEWSF reported having at least one maternal complication.¹¹

Due to the progressive nature of the disease and the lack of known medical management, delivery is always the definitive treatment, however, there is debate on the best time to deliver for both preterm and term gestations. Extending pregnancy carries a risk of exacerbating endothelial dysfunction in the mother and perpetuating inadequate perfusion of target organs, potentially leading to serious damage to the brain, liver, kidneys, placenta/ fetus, haematological and vascular systems.¹ Thus, there is an increased chance of induction failure and subsequent caesarean birth in pre-eclamptic women.¹² Other potential maternal sequelae include seizure, pulmonary oedema, cerebral haemorrhage, renal detachment or cortical blindness, stroke, hepatic failure, heart failure, renal failure, postpartum haemorrhage, disseminated intravascular coagulation, placental abruption and death.^{1 13} There was also a reported lifetime risk of hypertension.¹⁴ Furthermore, research published in the Lancet Regional Health revealed that pregnant women with hypertensive disorders have an increased risk of developing asthma and chronic obstructive pulmonary diseases.¹⁵

Moreover, pre-eclampsia and eclampsia made a substantial contribution to maternal deaths and severe morbidity.^{4 16} 10% to 15% of all maternal deaths worldwide are attributed to pre-eclampsia and eclampsia.¹¹ In Ethiopia, the five primary direct causes of maternal death were haemorrhage, obstructed labour, pre-eclampsia/ eclampsia, unsafe abortion and sepsis, accounting for 85% of maternal deaths. Pre-eclampsia/eclampsia makes up 11% of these five major causes of maternal mortality.¹⁷

A cross-sectional study in the Amhara region Referral Hospitals, Ethiopia, reported a significant association between residence, level of education, monthly income, parity, history of abortion, booking status, time of drug given and unfavourable maternal outcome.¹⁰ Women admitted at <34 weeks, age 16-24 years, lower wealth quintiles and rural residence had also a positive association with unfavourable maternal outcomes.⁹ Further,

gestational age at admission,¹⁸ onset of the disease and low haemoglobin level¹⁹ were predictors of maternal complication.

Pre-eclampsia causes significant financial losses that affect not just the individual but also the next generation because of the expense of prescription drugs, medical treatment, lost productivity and hindered daily activities. According to a US study, pre-eclampsia during the first 12 months of life is expected to cost US\$2.18 billion (US\$1.03 billion for moms and US\$1.15 billion for infants). The cost burden per infant varies with gestational age, starting at US\$150 000 at 26 weeks and going up to US\$1311 at 36 weeks.²⁰

Limited studies to date have been done to address the unfavourable maternal outcomes among pregnant 8 women with PEWSF in developing countries including pregnant of Ethiopia. The findings could have its own contribution to the local Ethiopian Health Sector Transformation Plan-II targeted to lower maternal mortality ratio from 401 to 140 per 100 000 live births²¹ and global Sustainable Development Goal target plans of less than 70 per 100 000 live births by 2030.²² Hence, the study aimed to determine the prevalence and associated factors of unfavourable uses related to text maternal outcomes among pregnant women with PEWSF in Ethiopia.

Research questions

- 1. What is the magnitude of unfavourable maternal outcomes among pregnant women admitted with PEWSF?
- 2. What are the factors associated with unfavourable maternal outcomes?

METHODS

Study design, period and area

This cross-sectional study was conducted at Abebech Gobena Mothers and Childrens Health (MCH) and St. ⊳ Peter's Specialized Hospital from 1 January 2023 to 30 July 2023, in Addis Ababa, the capital city of Ethiopia. Abebech Gobena MCH Hospital is one of the tertiary referral hospitals directly under the Addis Ababa Health Bureau. Yekatit 12 Hospital Medical College uses it as a teaching hospital as well. The hospital gives service to <u>0</u> more than 200 000 patients annually who were referred by about 18 catchment health centres in the Oromia regional state and Addis Ababa city, as well as one primary hospital. Whereas, St. Peter's Specialized Hospital is a government facility that served as the nation's first tuberculosis referral hospital. The hospital was founded in 1953. Currently providing care for over 100 000 people as a specialised hospital under the supervision of the Federal Ministry of Health. The MCH centre was established in 2006 Ethiopian Calendar and serves 15 catchment health centres and three primary hospitals from the Oromia region and Addis Ababa city.

Population and eligibility criteria

All pregnant mothers who were admitted with a diagnosis of PEWSF in the study area were the source population.

2

Participants were randomly selected from this source population. All pregnant mothers who were diagnosed, admitted and managed for PEWSF were included. Pregnant mothers who were diagnosed with pre-eclampsia but not with severe features and who were not giving birth at the study hospitals with unknown maternal outcomes were excluded.

Sample size and sampling technique

The sample size was determined using OpenEpi V.3.03 statistical software with the assumption of 36% prevalence of unfavourable maternal outcomes in Addis Ababa, Ethiopia,²³ 95% CI, 5% marginal error and 5% non-response rate. Considering, that the final sample size was 372. A total population sampling method was used to select the eligible study participants.

Variables

Maternal outcome was the dependent variable. Independent variables included socio-demographic factors (age, residence, marital status, occupation, educational level and mode of admission), medical and reproductive history (gravidity, parity, history of abortion, antenatal care (ANC); history of gestational hypertension, family history of hypertension, anaemia, chronic hypertension, diabetes and renal disease); clinical features and investigations on admission (headache, dizziness, epigastric pain, visual disturbance, nausea and/or vomiting, convulsion, oedema, haematocrit, liver function test, urea, creatinine, urine protein); and obstetrical factors (onset of labour, mode of delivery, sex of the neonate and duration of hospitalisation).

Outcome measures

Pre-eclampsia with severe features

Is a pre-eclampsia with one of the severity features; including altered mental status, severe headache, altered cerebral or visual disturbance, hepatic abnormality, renal abnormality, severe blood pressure ($\geq 160/110$), thrombocytopenia (platelet count $<100 \times 10^9/L$) and pulmonary oedema.14

Blood pressure at admission

Severe hypertension if blood pressure measurement was ≥160/110 and mild hypertension if 140-159/90-109.⁴¹¹

Severe headache

Incapacitating, 'the worst headache I have ever had' or headache that persists and progresses despite analgesic therapy.

Hepatic abnormality

Severe persistent right upper quadrant or epigastric pain unresponsive to medication and not accounted for by an alternative diagnosis or serum transaminase concentration ≥ 2 times the upper limit of the normal range, or both.¹

Renal abnormality

Progressive renal insufficiency (serum creatinine >1.1 mg/dL (97.2 µmol/L) or a doubling of the serum creatinine concentration in the absence of other renal disease).¹

Unfavourable maternal outcome

Mothers with PEWSF that develop at least one complication, that is, eclampsia, abruption placenta, haemolysis, elevated liver enzymes and low platelet count (HELLP) syndrome, acute renal failure, disseminated intravascular coagulation, cardiac failure, stroke, postpartum haemorrhage, pulmonary oedema and death.⁹¹⁰

Favourable maternal outcome

tected by copyright, Mothers with PEWSF managed and improved without complications.¹⁰

Data collection tool, procedure, quality control

The data was collected using a well-constructed case record form and procedure. The questionnaire was adapted from similar studies.⁷⁻¹⁰ ¹³ The data collection team consisted of two supervisors and four data collectors. The principal investigators gave the supervisors and data collectors a 1-day training on the objectives, methods, procedures and data collection instrument. The ques-Pe tionnaire was translated back and forth from English to Amharic and vice versa to make sure the questions remained true to their original intent. Prior to the real data collection, a pre-test was done on 5% of the samples \mathbf{Z} (19 mothers) at Debre Berhan Comprehensive Specialized Hospital and the necessary adjustments were taken a into account in light of the test results. Over the course of the data collection process, the principal investigators and supervisors closely observed the clarity, consistency and completeness of the data.

Patient and public involvement

There was no patient and/or public involvement in the design and planning of this study.

Data management and analysis

Al training, and Data was entered using EpiData V.4.6 and analysed using S SPSS V.26.0 statistical software. The principal investigator randomly selected a questionnaire for quality control and cross-checked it with the correspondingly entered data and clinical chart. We employed descriptive statistics to <u>lou</u> describe the independent and dependent variables. The results were presented as number, frequency, percentage and comparison of maternal outcomes. Binary logistic 8 regression analysis was run to identify independent predictors of unfavourable maternal outcomes. Variables with a p value of < 0.25 in the bivariable regression analysis were included in the final multivariable logistic regression analysis model. Hosmer and Lemeshow's goodness-of-fit test was employed to evaluate the fitness of the model. The multicollinearity of the explanatory components was also investigated. With a two-sided 95% CI, adjusted ORs (AORs) were used to interpret the strength of the

Socio-demographic characteristics of participants Table 1 admitted with PEWSF at Abebech Gobena Mothers and Childrens Health and St. Peter's Specialized Hospital, Ethiopia, 2023

Variables	Category	Frequency	Per cent (%)
Age in years	20–34	294	84.5
	<20	13	3.7
	≥35	41	11.8
Residence	Urban	272	78.2
	Rural	76	21.8
Level of education	No formal education Primary Secondary Higher education	45 97 134 72	12.9 27.9 38.5 20.7
Marital status	Married	324	93.1
	Others*	24	6.9
Occupation	Employed	204	58.6
	Unemployed	144	41.4
Mode of admission	Self	52	14.9
	Referral	296	85.1

*Single, divorced and widowed.

PEWSF, pre-eclampsia with severe features.

association. A p value of <0.05 was used to declare the level of significance.

RESULTS

Socio-demographic characteristics of participants

A total of 348 mothers participated, giving the survey a 93.5% response rate. The age range of the participants was 18-42 years old, with a mean (SD) of 27.55±5.18 years. Of these, 272 (78.2%) lived in urban, making up more than three-fourths. Furthermore, table 1 shows that 324 (93.1%) of the participants were married, and 134 (38.5%) had completed secondary school.

Medical and obstetrical history

More than half, 179 (51.4%) of mothers, were primigravida and 69 (19.8%) had previously experienced an abortion. Nearly all, 342 (98.3%), of the participants had ANC contact for the current pregnancy. However, only 22 (6.3%) of them had adequate ANC contact. Furthermore, 34 (9.8%) of mothers had a history of gestational hypertension. 27 (7.8%) of participants had a medical history. Of them, chronic hypertension and anaemia were reported in 12 (3.4%) and 8 (2.3%) of cases, respectively (table 2).

Clinical features and investigations on admission

In this study, 180 (51.7%), 119 (34.2%) and 87 (25.0%) of mothers were admitted with a chief issue of headache, epigastric pain and oedema, respectively. Whereas, on an investigation, 38 (10.9%) of the women had deranged

BMJ Open: first published as 10.1136/bmjopen-2023-081901 on 29 March 2024. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES)

Table 2 Medical and obstetrical history of mothers admitted with pre-eclampsia with severe features at Abebech Gobena Mothers and Childrens Health and St. Peter's Specialized Hospital, Ethiopia, 2023

o :	Category	Frequency	(%)
Gravidity	Primigravida Multigravida Grand multipara	179 162 7	51.4 46.6 2.0
Parity	Nulliparous	12	3.4
	1–3	322	92.5
	≥4	14	4.0
History of abortion	Null 1 ≥2	279 61 8	80.2 17.5 2.3
Antenatal care	Yes	342	98.3
(ANC) contact	No	6	1.7
Number of ANC contact	1–3	74	21.3
	4–6	252	72.4
	≥7–8	22	6.3
Number of	Singleton	326	93.7
fetuses	Twin/multiple	22	6.3
History of gestational hypertension	Yes No	34 314	9.8 90.2
Family history of gestational hypertension	Yes No	70 278	20.1 79.9
Medical history	Yes	27	7.8
	No	321	92.2
Anaemia	Yes	8	2.3
	No	340	97.7
Chronic	Yes	12	3.4
hypertension	No	336	96.6
Diabetes	Yes	5	1.4
mellitus	No	343	98.6
Renal disease	Yes	2	0.6
	No	346	99.4

(17.2%), HELLP syndrome (15.5%) and postpartum haemorrhage (13.8%) were the most prevalent complications that occurred among mothers admitted with a diagnosis of PEWSF (figure 1).

Factor of unfavourable outcomes

Variables having a p value of <0.25 in the bivariable analysis were chosen for the multivariable logistic regression analysis model. A crude OR is an OR of univariable

Table 3 Clinical features of participants admitted with preeclampsia with severe features at Abebech Gobena Mothers and Childrens Health and St. Peter's Specialized Hospital, Ethiopia, 2023

Variables	Category	Frequency	Per cent (%)
Headache	Yes	180	51.7
	No	168	48.3
Dizziness	Yes	45	12.9
	No	303	87.1
Epigastric pain	Yes	119	34.2
	No	229	65.8
Visual	Yes	59	17.0
disturbance	No	289	83.0
Nausea and/or vomiting	Yes	15	4.3
	No	333	95.7
Convulsion	Yes	33	9.5
	No	315	90.5
Oedema	Yes	87	25.0
	No	261	75.0
Grade of oedema (n=87)	Grade 1 Grade 2 Grade 3	46 38 3	52.9 43.7 3.4
Blood pressure at admission	Severe range Mild range	297 51	85.3 14.7
Haematocrit	<33%	39	11.2
	≥33%	309	88.8
Liver function test	Normal	310	89.1
	Deranged	38	10.9
Urea	Normal	322	92.5
	Deranged	26	7.5
Creatinine	Normal	319	91.7
	Deranged	29	8.3
Urine protein (dipstick)	Negative 1+ 2+ 3+	105 50 140 53	30.2 14.4 40.2 15.2
Onset of	Spontaneous	104	29.9
labour	Induction	244	70.1
Mode of delivery	Spontaneous vaginal delivery Instrumental Caesarean section	186 14 148	53.4 4.0 42.5
Sex of the neonate	Male	186	53.4
	Female	162	46.6
Duration of	≤3 days	135	38.8
hospital stay	≥4 days	213	61.2

analysis; one independent variable for predicting the dependent variable. Accordingly, age, residence, level of education, occupation, number of fetuses, sex of neonate, blood pressure on admission and headache issue were selected. In the final model, age, residence, occupation, blood pressure on admission and issues of headache were

found to be statistically significantly associated with unfavourable maternal outcomes.

Mothers aged above 35 had approximately threefold increased risk of developing unfavourable outcomes compared with those aged between 20 and 34 (AOR (95% CI)=2.70 (1.31 to 5.59)). Rural residents had a 94% higher chance of experiencing unfavourable outcomes compared with their urban counterparts (AOR (95% CI)=1.94 (1.07 to 3.53)). Unemployed mothers bore a 65% lower risk of unfavourable outcomes in comparison to those who were employed (AOR (95% CI)=0.35 (0.20 to 0.62)). Severe blood pressure measurement on admission increased the risk of unfavourable outcomes by twofold (AOR (95% CI)=2.32 (1.03 to 5.19)). Furthermore, women who were admitted with a headache as 8 pyright, their chief issue had a 91% higher likelihood of having unfavourable outcomes (AOR (95% CI)=1.91 (1.16 to 3.16)) (table 4). including

DISCUSSION

In this study, the overall prevalence of unfavourable maternal outcomes was 33.9% (95% CI: 28.7% to 38.8%). use Age, residence, occupation, blood pressure on admission and headache issues have shown a statistically significant association with unfavourable outcomes among women of PEWSF admitted at Abebech Gobena MCH and St. Peter's Specialized Hospital, Addis Ababa, Ethiopia.

ç Unfavourable maternal outcomes occurred in 33.9% of e mothers with PEWSF. This is comparable with the study findings from Amhara region referral hospitals, where 37.7% of mothers with PEWSF developed unfavourable outcomes.¹⁰ However, it was higher than 10% in the a USA.⁸ This discrepancy could be the result of variations in the study population, time, set-up, sample size and quality and standard of care provided by contemporary, well-≥ equipped maternity hospitals, as well as good prenatal and obstetrical care. On the other hand, it was lower than 43% in the Sidama region of Ethiopia.⁹ Variations in the incidence proportion of unfavourable outcomes between the studies might be attributed to the severity of the disease, differences in clinical features (severity signs and symptoms) on admission and gestational age at diagnosis.

Abruptio placenta (17.2%), HELLP syndrome (15.5%) and postpartum haemorrhage (13.8%) were the most prevalent complications. Similarly, in Thailand, postpartum haemorrhage, placental abruption and heart failure occurred in 9.4%, 1.4% and 0.4% of women with \mathbf{a} PEWSF, respectively.¹³ Further, in the Sidama region, Ethiopia, a higher level of antepartum and postpartum haemorrhage was observed in the mothers of PEWSF.⁹

It was discovered that older mothers were linked to a higher likelihood of unfavourable outcomes. Mothers over 35 were almost three times more likely to experience an adverse outcome. In a similar vein, poor maternal outcome was more common in Indonesia among mothers with pre-eclampsia who were older than 35.²⁴ Because of increased endothelial injuries that lower



Figure 1 Outcomes of pregnant women admitted with pre-eclampsia with severe features at Abebech Gobena Mothers and Childrens Health and St. Peter's Specialized Hospital, Ethiopia, 2023.

renal reserves and the incapacity to adapt to physiological changes during pregnancy, older people may be more susceptible to developing renal insufficiency even if their pre-gestational kidney functions are normal.²⁵ It might also be connected to the extravascular space's increased fluid accumulation during pregnancy. Additionally, older people are more likely to have additional risk factors that increase their likelihood of developing pre-eclampsia, such as diabetes mellitus, obesity and chronic hypertension.

The odds of unfavourable maternal outcomes were 94% higher among rural residents than their urban counterparts. Similarly, in Ethiopia's Sidama region, women who lived in rural regions were more likely to experience unfavourable maternal outcomes.⁹ This could be because women in rural areas may have had a lower socioeconomic level, which may have resulted in a lesser tendency to seek medical attention. Pregnant women with low health-seeking behaviour are less likely to visit ANC clinics, which delays the diagnosis and treatment of pre-eclampsia. In addition, rural women faced significant challenges in getting to health facilities due to transportation issues, which caused delays in receiving medical care. It is improbable that they are aware of the risks and complications associated with pregnancy, labour and delivery. In addition, the cultural practices prevalent in rural areas greatly impact women's nutritional status by preventing them from consuming necessary foods and/ or beverages.²⁶

Unemployed mothers had less risk of unfavourable outcomes compared with those employed. In the Netherlands, when employed women worked longer hours (≥ 40

Protected by copyright, including for uses related hours/week), the mean birth weight of kids decreased by to 45 g.²⁷ Similarly, in South Korea,²⁸ higher risks of early abortive outcomes and stillbirths were more frequent in employed women. The possible explanation might be that unemployed mothers are more likely to have adequate time to care for themselves and listen to updated information regarding gestational hypertension via TV, radio or others. This might help them to have a lower risk of \blacksquare unfavourable outcomes.

A severe blood pressure measurement on admis-≥ sion doubled the likelihood of unfavourable maternal outcomes. Severe blood pressure was also revealed to be a significant predictor of an adverse outcome (eclampsia) among pre-eclamptic mothers in Morocco.²⁹ Hypertension is one of the hallmarks of pre-eclampsia and severe hypertension, defined as a blood pressure of more than 160/110 mm Hg, has been considered a warning indicator of the development of negative outcomes, such as eclampsia.⁴ Thus, severe blood pressure is a symptom of a severe condition, rapid disease progression and a terrible prognosis.

Furthermore, women who were admitted with a chief issue of headache had a 91% increased risk of unfavourable outcomes. In a retrospective chart review of pre-eclamptic patients treated at Ayder Comprehensive Specialized Hospital, Ethiopia, headache and blurring were associated with poor maternal outcomes.³⁰ It has been noted that neurological symptoms indicate an impending negative consequence.²

e

Table 4	Factors associated	with unfavourable	maternal outcome	among mothers	admitted with	pre-eclampsia	with severe
features a	t Abebech Gobena	Mothers and Child	rens Health and St	. Peter's Specia	lized Hospital,	Ethiopia, 2023	

	Maternal outcomes				
Variables	Favourable	Unfavourable	COR (95% CI)	AOR (95% CI)	
Age in years 20–34 <20 ≥35	202 (87.8%) 8 (3.5%) 20 (8.7%)	92 (78.0%) 5 (4.2%) 21 (17.8%)	1 1.37 (0.44 to 4.31) 2.31 (1.19 to 4.46)	1 1.33 (0.39 to 4.52) 2.70 (1.31 to 5.59)*	
Residence Urban Rural	188 (81.7%) 42 (18.3%)	84 (71.2%) 34 (28.8%)	1 1.81 (1.08 to 3.05)	1 1.94 (1.07 to 3.53)*	
Level of education No formal education Primary Secondary Higher education	23 (10.0%) 62 (27.0%) 98 (42.6%) 47 (20.4%)	22 (18.6%) 35 (29.7%) 36 (30.5%) 25 (21.2%)	1.80 (0.84 to 3.84) 1.06 (0.56 to 2.01) 0.69 (0.37 to 1.28) 1	2.15 (0.89 to 5.17) 1.73 (0.82 to 3.67) 1.00 (0.51 to 1.98) 1	
Occupation Employed Unemployed	123 (53.5%) 107 (46.5%)	81 (68.6%) 37 (31.4%)	1 0.53 (0.33 to 0.84)	1 0.35 (0.20 to 0.62)*	
Number of fetuses Singleton Twin/multiple	220 (95.7%) 10 (4.3%)	106 (89.8%) 12 (10.2%)	1 2.49 (1.04 to 5.95)	1 2.04 (0.79 to 5.24)	
Sex of the neonate Male Female	116 (50.4%) 114 (49.6%)	70 (59.3%) 48 (40.7%)	1.43 (0.92 to 2.25) 1	1.43 (0.88 to 2.33) 1	
Blood pressure on admission Severe range Mild range	188 (81.7%) 42 (18.3%)	109 (92.4%) 9 (7.6%)	2.71 (1.27 to 5.77) 1	2.32 (1.03 to 5.19)* 1	
Headache issue Yes No	106 (46.1%) 124 (53.9%)	74 (62.7%) 44 (37.3%)	1.97 (1.25 to 3.10) 1	1.91 (1.16 to 3.16)* 1	
P value≤0.25. *Statistically significant at p value<0.0 AOR, adjusted OR; COR, crude OR.	5				

Conclusion and recommendations

In this study, the prevalence of unfavourable maternal outcomes was high compared with other studies in Ethiopia. Maternal age, residence, occupation, blood pressure on admission and severe headache have shown a statistically significant association with unfavourable maternal outcomes. Socioeconomic development and early identification and treatment of severe signs and symptoms of pre-eclampsia are needed to reduce unfavourable outcomes. Prenatal screening and specialised care for women who are at high risk, such as older mothers, are also recommended. Further, longitudinal studies are recommended to investigate the outcome of mothers with PEWSF.

Limitations

It shares the limitation of a cross-sectional study to draw a causal relationship. Since this study was conducted in

referral hospitals, we are unable to ascertain whether these women delayed visiting the primary health facilities or whether there were delays in referring them. In addiar tion, as this was done in the hospital setting, the maternal technologies. outcome of women delivered at home was not assessed. Further, this study does not include adverse maternal outcomes after 24 hours of birth.

Author affiliations

¹Department of Midwifery, School of Nursing and Midwifery, Asrat Woldeyes Health Science Campus, Debre Berhan University, Debre Berhan, Amhara, Ethiopia ²Department of Medicine, Obstetrics and Gynecology, Abebech Gobena Mothers and Childrens Health Hospital, Addis Ababa, Ethiopia

³Department of Epidemiology, St Peter Specialized Hospital, Addis Ababa, Ethiopia ⁴Department of Nursing, Debre Berhan Health Science College, Debre Berhan, Amhara, Ethiopia

text and data mining, AI training, and similar technologies

Protected by copyright, including for uses related to

Open access

Acknowledgements The authors would like to thank the administrators, data collectors and study participants for providing genuine data, as well as Yekatit 12 Hospital Medical College for approving the project's ethical clearance.

Contributors WAD drafted the topic, designed the proposal and performed data collection. MT critically revised, performed and interpreted the analysis, reviewed and edited, developed the manuscript, and is responsible for the overall content as guarantor. MT, WAD, GSS, GEW, SDT and AE reviewed the proposal, contributed to data collection and analysis and critically revised the manuscript. MT made basic adjustments to the final manuscript and processed publication. All authors approved the manuscript for journal submission.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval The Institutional Review Board (IRB) of Yekatit 12 Hospital Medical College granted ethical clearance (Protocol number 128/23). A formal letter of support was forwarded to the study hospitals. Participants gave their free and informed consent, and they participated willingly. Those who were illiterate were asked to thumbprint the consent form once the content was read. Confidentiality and anonymity were preserved and the client records were returned to their place after the completion of data collection.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. All relevant data sets are incorporated in the article and supplementary files.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Mesfin Tadese http://orcid.org/0000-0001-6288-9771 Saba Desta Tessema http://orcid.org/0000-0003-4983-1086

REFERENCES

August P, Sibai BM. Preeclampsia: clinical features and diagnosis. 2017.22

- 2 Bisson C, Dautel S, Patel E, et al. Preeclampsia pathophysiology and adverse outcomes during pregnancy and postpartum. Front Med 2023;10:485.
- 3 Calvo JP, Rodríguez YP, Figueroa LQ. Update in Preeclampsia. Rev Médica Sinerg 2020;5:01.
- Δ Lisonkova S, Bone JN, Muraca GM, et al. Incidence and risk factors for severe Preeclampsia, hemolysis, elevated liver enzymes, and low platelet count syndrome, and Eclampsia at Preterm and term gestation: a population-based study. Am J Obstet Gynecol 2021-225:538
- Noubiap JJ, Bigna JJ, Nyaga UF, et al. The burden of hypertensive 5 disorders of pregnancy in Africa: a systematic review and Meta-Analysis. J of Clinical Hypertension 2019;21:479-88.
- Machano MM, Joho AA. Prevalence and risk factors associated 6 with severe pre-Eclampsia among postpartum women in Zanzibar: a cross-sectional study. BMC Public Health 2020;20::1347.
- Belay Tolu L, Yigezu E, Urgie T, et al. Maternal and perinatal outcome of Preeclampsia without severe feature among pregnant women managed at a tertiary referral hospital in urban Ethiopia. PLoS One 2020;15:e0230638.

- Venkatesh KK, Strauss RA, Westreich DJ, et al. Adverse maternal 8 and neonatal outcomes among women with Preeclampsia with severe features weeks gestation with versus without Comorbidity. Pregnancy Hypertens 2020;20:75-82.
- Jikamo B, Adefris M, Azale T, et al. The effect of Preeclampsia on adverse maternal outcomes in Sidama region. Ethiopia: a prospective open cohort study. Sci Rep 2022;12:19300.
- 10 Melese MF, Badi MB, Aynalem GL. Maternal outcomes of severe Preeclampsia / Eclampsia and associated factors among mothers admitted in referral hospitals North West Ethiopian context, 2018. BMC Res Notes 2019:12:147
- Chappell LC, Cluver CA, Kingdom J, et al. Pre-Eclampsia. Lancet 11 2021;398:341-54
- 12 Tadesse T, Assefa N, Roba HS, et al. Failed induction of labor and associated factors among women undergoing induction at University of Gondar specialized hospital, Northwest Ethiopia. BMC Pregnancy Childbirth 2022;22:175.
- 13 Kongwattanakul K, Saksiriwuttho P, Chaiyarach S, et al. Incidence, characteristics, maternal complications, and perinatal outcomes associated with Preeclampsia with severe features and HELLP syndrome. Int J Womens Health 2018;10:371-7.
- 14 Turbeville HR, Sasser JM. Preeclampsia beyond pregnancy: longterm consequences for mother and child. Am J Physiol Renal Physiol 2020:318:F1315-26.
- Wang Y-X, Varraso R, Dumas O, et al. Hypertensive disorders of 15 pregnancy and risk of asthma and chronic obstructive pulmonary disease: a prospective cohort study. Lancet Reg Health Am 2023;23:100540
- Tesfay N, Tariku R, Zenebe A, et al. Critical factors associated 16 with postpartum maternal death in Ethiopia. PLoS One 2022;17:e0270495.
- 17 Health FDR of EM of. Maternal Death Surveillance and Response (MDSR) Technical Guideline, Ethiopia: Addis Ababa, 2014.
- 18 Barda S, Yoeli Y, Stav N, et al. Factors associated with progression to Preeclampsia with severe features in pregnancies complicated by mild hypertensive disorders. J Clin Med 2023;12:7022:22:
- 19 Syoum FH, Abreha GF, Teklemichael DM, et al. Fetomaternal outcomes and associated factors among mothers with hypertensive disorders of pregnancy in Suhul hospital, Northwest Tigray, Ethiopia. J Pregnancy 2022;2022:6917009.
- 20 Stevens W, Shih T, Incerti D, et al. Short-term costs of Preeclampsia to the United States health care system. Am J Obstet Gynecol 2017:217:237-48.
- (MOH) EM of H. Health sector transformation plan II: 2020/21-21 2024/25 (2013 EFY-2017 EFY). MOH 2021.
- 22 Boldosser-Boesch A, Brun M, Carvajal L, et al. Setting maternal mortality targets for the Sdgs. Lancet 2017;389:696-7.
- Wagnew M, Dessalegn M, Worku A, et al. Trends of Preeclampsia/ 23 Eclampsia and maternal and neonatal outcomes among women delivering in Addis Ababa selected government hospitals, Ethiopia: a retrospective cross-sectional study. Pan Afr Med J 2016;25:12:12:.
- Tyas BD, Lestari P, Aldika Akbar MI. Maternal perinatal outcomes related to advanced maternal age in Preeclampsia pregnant women. JFRH 2019:13.
- 25 Ronco C, Bellomo R, Kellum J. Understanding renal functional Reserve. Intensive Care Med 2017;43:917-20.
- 26 Bazzano AN, Potts KS, Mulugeta A. How do pregnant and lactating women, and young children, experience religious food restriction at the community level? A qualitative study of fasting traditions and feeding behaviors in four regions of Ethiopia. PLoS One 2018;13:e0208408.
- 27 Jansen PW, Tiemeier H, Verhulst FC, et al. Employment status and the risk of pregnancy complications: the generation R study. Occup Environ Med 2010;67:387-94.
- 28 Kim C-B, Choe S-A, Kim T, et al. Risk of adverse pregnancy outcomes by maternal occupational status: A national populationbased study in South Korea. J Occup Health 2023:65:e12380.
- Rebahi H, Elizabeth Still M, Faouzi Y, et al. Risk factors for Eclampsia 29 in pregnant women with Preeclampsia and positive Neurosensory signs. Turk J Obstet Gynecol 2018;15:227-34.
- 30 Legesse AY, Berhe Y, Mohammednur SA, et al. Prevalence and determinants of maternal and perinatal outcome of Preeclampsia at a tertiary hospital in Ethiopia. Ethiop J Reprod Heal 2019;11.