Protocol

BMJ Open ExtraCorporeal Membrane Oxygenation in the therapy for REfractory Septic shock with Cardiac function Under **Estimated (ECMO-RESCUE): study** protocol for a prospective, multicentre, non-randomised cohort study

Wei-yan Chen 🐵 , Ze-bin Guo, Tian-yu Kong, Wei-xiao Chen, Xiao-hua Chen, Qilin Yang, Yi-chao Wen, Qi-rui Wen, Feng Zhou, Xu-ming Xiong 💩, De-liang Wen. Zhen-hui Zhang 💿

ABSTRACT

To cite: Chen W. Guo Z. Kong T. et al. ExtraCorporeal Membrane Oxygenation in the therapy for REfractory Septic shock with Cardiac function Under Estimated (ECMO-RESCUE): study protocol for a prospective, multicentre, non-randomised cohort study. BMJ Open 2024:14:e079212. doi:10.1136/ bmjopen-2023-079212

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (https://doi.org/10.1136/ bmjopen-2023-079212).

W-yC and Z-bG contributed equally. D-IW and Z-hZ contributed equally.

Received 24 August 2023 Accepted 10 May 2024

Check for updates

C Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Professor Zhen-hui Zhang; yanniskyyx@126.com and Professor De-liang Wen; deliangwen@163.com

Introduction Severe septic cardiomyopathy (SCM) is one of the main causes of refractory septic shock (RSS), with a high mortality. The application of venoarterial extracorporeal membrane oxygenation (ECMO) to support the impaired cardiac function in patients with septic shock remains controversial. Moreover, no prospective studies have been taken to address whether venoarterial ECMO treatment could improve the outcome of patients with sepsis-induced cardiogenic shock. The objective of this study is to assess whether venoarterial ECMO treatment can improve the 30-day survival rate of patients with sepsis-induced refractory cardiogenic shock.

Methods and analysis ExtraCorporeal Membrane Oxygenation in the therapy for REfractory Septic shock with Cardiac function Under Estimated is a prospective, multicentre, non-randomised, cohort study on the application of ECMO in SCM. At least 64 patients with SCM and RSS will be enrolled in an estimated ratio of 1:1.5. Participants taking venoarterial ECMO during the period of study are referred to as cohort 1, and patients receiving only conventional therapy without ECMO belong to cohort 2. The primary outcome is survival in a 30-day follow-up period. Other end points include survival to intensive care unit (ICU) discharge, hospital survival, 6-month survival, quality of life for long-term survival (EQ-5D score), successful rate of ECMO weaning, long-term survivors' cardiac function, the number of days alive without continuous renal replacement therapy, mechanical ventilation and vasopressor, ICU and hospital length of stay, the rate of complications potentially related to ECMO treatment.

Ethics and dissemination The trial has been approved by the Clinical Research and Application Institutional Review Board of the Second Affiliated Hospital of Guangzhou Medical University (2020-hs-51). Participants will be screened and enrolled from ICU patients with septic shock by clinicians, with no public advertisement for recruitment. Results will be disseminated in research journals and through conference presentations. Trial registration number NCT05184296.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- \Rightarrow ExtraCorporeal Membrane Oxygenation in the therapy for REfractory Septic shock with Cardiac function Under Estimated (ECMO-RESCUE) is a prospective, multicentre, non-randomised, cohort study on the application of ECMO in sepsis-induced refractory cardiogenic shock.
- \Rightarrow Patients admitted to intensive care unit and meet diagnostic criteria for cardiogenic shock induced by sepsis will be enrolled in the study and subsequently assigned to one of two cohorts based on their willingness to undergo ECMO.
- \Rightarrow All patients willing to undergo ECMO treatment will initiate venoarterial ECMO within 6 hours of enrolment.
- \Rightarrow Risks of bleeding, thrombosis, leg ischaemia and cannulation-related injuries may be elevated in patients undergoing ECMO treatment; however, this intervention may offer a potential benefit to improve survival.
- \Rightarrow A limitation of the study is its non-randomised design, which would yield bias.

INTRODUCTION

Protected by copyright, including for uses related to text and data mining, Al training, and similar technol Sepsis, a life-threatening syndrome with organ dysfunction caused by infection, is a leading cause of death in intensive care unit (ICU).¹ The global burden of disease **g** arising from sepsis was estimated at 30 million episodes and 6million deaths per year in $2017.^2$ The mortality is as high as 50% when septic shock is present.3 4 Septic cardiomyopathy (SCM) is one of the main causes of septic shock, affecting 20%-65% of patients with sepsis. Severe cardiac function impairment leads to refractory septic shock (RSS), with a mortality up to 70%.^{5–7} However, there are no

evidence-based recommendations for the management of SCM.⁸

In addition to aggressive treatment with antibiotics for infection control, adequate fluid resuscitation and vasoactive drug administration according to the International Guidelines for Management of Sepsis and Septic Shock,⁹ a variety of drugs to improve cardiac function have also been attempted for the treatment of SCM. However, the results were not satisfactory. Studies on dobutamine have demonstrated that the administration of either dobutamine alone or a combination of norepinephrine failed to improve survival rate, microcirculatory perfusion and metabolic despite an increasing cardiac index, heart rate and left ventricular ejection fractions (LVEF).^{10 11} Another positive inotropic drug, levosimendan, also has been tried to apply in the treatment of SCM. However, the therapeutic effect is still controversial.^{12–14} Therefore, there are few recognised effective treatments for sepsisinduced cardiogenic shock at present.

SCM is considered as a sepsis-associated acute syndrome of cardiac dysfunction unrelated to ischaemia.¹⁵ Most studies have suggested that recovery from SCM is prompt. A retrospective study showed that although left ventricular dysfunction may persist in approximately one-third of patients with severe sepsis and septic shock in longterm follow-up, the survival rates did not differ.¹⁶ Therefore, trying new, innovative therapeutic approaches in SCM are valuable and urgently needed.

Venoarterial extracorporeal membrane oxygenation (ECMO) is a circulatory support technology, which can increase cardiac output as well as improve hypoxemia by increasing oxygen delivery. Therefore, the application of venoarterial ECMO in the treatment of RSS has theoretical feasibility. However, sepsis has been considered as a contraindication for ECMO due to its complexity and the unsatisfactory outcomes of earlier studies. As an extracorporeal circulation device, ECMO may be susceptible to pathogen attachment, leading to refractory infections that exacerbate the underlying condition.¹⁷ Furthermore, patients with sepsis often present with thrombocytopenia and abnormal coagulation function, while ECMO necessitates anticoagulation therapy, which can potentially worsen bleeding.^{18 19} In Park *et al*'s study, a total of 32 patients received ECMO support for RSS, only seven patients (21.9%) survived to hospital discharge.²⁰ In Huang et al's study, hospital survival rate was much lower $(15\%).^{2}$

However, as the advancement of ECMO technology, improvements in materials of ECMO²² and the emergence of new research findings, there is a growing reconsideration regarding the utilisation of ECMO in sepsis. It is reported that in neonatal and children with septic shock, the survival rate of ECMO treatment was nearly 80% and 50%, respectively.^{23 24} ECMO treatment of septic shock in neonatal and children has been in guidelines and consensus since 2008.^{25 26} Besides this, a single-centre retrospective study found that venoarterial ECMO rescued more than 70% of the patients who developed refractory

Cardiovascular dysfunction during severe bacterial septic shock.²⁷ Another multicentre cohort study found that survival at 90 days for patients with severe sepsis-induced cardiomyopathy who received venoarterial ECMO was significantly higher than for controls (60% vs 25%).²⁸ Falk *et al* reported a 90% hospital survival rate in septic shock patients with left ventricular failure and 64.7% in patients with distributive shock.²⁹ Cardiac depression, vasoplegia and capillary leakage all lead to circulatory failure in septic shock. Falk *et als* study suggested that patients with sepsis-induced refractory cardiogenic shock may receive high survival benefit from venoarterial ECMO and superior to distributed shock. Moreover, all buryors restored their cardiac function and had good ong-term quality of life.³⁰ A meta-analysis reported by profit (62.0% vs 32.1%), and patients with LVEF between 20% (bas done that perhaps RSS is not an absolute contraindi-cation to ECMO, but rather that the ideal candidates for instrument should be identified. In summary, the use of ECMO in adult patients with RSS is still controversial. More studies are still needed to determine whether the benefit of ECMO outweighs indult patients with sepsis-induced cardiogenic shock have not yet been reported. Therefore, prospective clim-cial studies would encounter numerous challenges. For example, before ECMO is initiated what dose of vasoar tive drug is needed? Are there any other indicators of cardiac function that help to determine the initiation of ECMO in addition to the level of LVEF? This prospective composition to the level of LVEF? This prospective

cardiac function that help to determine the initiation of ECMO in addition to the level of LVEF? This prospective cohort study was designed with reference to indications of ECMO initiation in refractory cardiogenic shock. In this study, we aimed to assess whether venoarterial ECMO treatment can improve the 30-day survival rate of patients with sepsis-induced refractory cardiogenic shock.

METHODS AND ANALYSIS

Study design, setting and patient population

The ExtraCorporeal Membrane Oxygenation in the therapy for REfractory Septic shock with Cardiac function Under Estimated study is a prospective, multicentre, non-randomised, cohort study. All patients admitted to the ICUs of participating centres will be considered as potential candidates for the study. Once the patient is & diagnosed of septic shock, he/she should be screened for **\$** eligibility by the physicians. When the patient fulfils the criterion of recruitment, the researcher provides details on the purpose, specific content and instructions on how to complete the trial. After we obtain written informed consent (online supplemental file 1) from the patient or a responsible surrogate, patients are enrolled in the study.

During the period of study, the participants' legal representative can decide whether to accept ECMO based on their personal conditions. If the participant is confirmed

to receive ECMO treatment, we initiate ECMO within 6 hours. Participants taking ECMO during the period of study are referred to as cohort 1, and patients receiving only conventional therapy without ECMO belong to cohort 2. ECMO is established and managed by a professional team. The teams of each centre have more than 5-year experience.

The study will be conducted in six ICUs located in Guangdong, China. It is anticipated that the study will span a duration of 3 years. Participant recruitment commenced in May 2023, with an anticipated completion date for enrolment set for December 2025. The anticipated completion time for followed-up is projected to be in May 2026.

Patient and public involvement

Patients or the public were not involved in the design, or conduct, or reporting, or dissemination of our research. The results will be available to the public if necessary.

Inclusion criteria

- 1. Age between 18 and 75 years.
- 2. Patients admitted into ICU and diagnosed as septic shock (sepsis-3.0), after adequate fluid resuscitation, high-dose vasoactive drug application (vasoactive inotropic score (VIS) >120) and conventional therapy together with at least one of the following criteria: (1) sustained hypotension (mean arterial pressure (MAP) <65 mm Hg); (2) persistent lactacemia (two consecutive values >5 mmol/L with at least 30 min interval between samples), with non-decreasing trend on steady doses of inotropes and/or vasopressors; (3) persistent low central venous blood oxygen saturation (ScvO_a) (two consecutive values <55% with at least 30 min interval between samples), with non-increasing trend on steady doses of inotropes and/or vasopressors. The above condition lasts more than 5 hours.
- 3. Rapidly deteriorating sepsis-induced myocardial impairment is defined by at least one of the following criteria: (1) rapidly deteriorating ventricular function (LVEF < 35%); (2) CI < 2.0 L/min/m² (>3 hours); (3) emerging refractory arrhythmia.
- 4. Informed consent provided by the patient or person with decisional responsibility.

Exclusion criteria

- 1. Cardiac dysfunction caused by other causes is excluded, such as acute myocardial infarction, chronic heart failure, congenital cardiac disease, myocardial effusion, moderate to severe aortic regurgitation, severe aortic coarctation and so on.
- 2. High suspicion of pulmonary embolism, tension pneumothorax or cardiac tamponade as a cause of shock.
- 3. Prolonged cardiac arrest (>30 min) before ECMO, or cardiopulmonary resuscitation survivors remaining comatose.

- 4. Irreversible condition or meet the inclusion criteria for more than 12 hours.
- 5. Presence of active bleeding or anticoagulant contraindications.
- 6. Peripheral artery disease disabling insertion of outflow cannula to femoral artery.
- 7. Irreversible neurological pathology.
- 8. Severe underlying condition with lift expectancy less than 1 year.
- 9. Special population, such as pregnancy, AIDS.
- Protected by copyright, 10. Patient included in another interventional clinical trial.

Study definitions

Septic shock

Septic shock, a subset of sepsis, can be identified by a vasopressor requirement to maintain an MAP of 65 mm Hg or greater and serum lactate greater than 2mmol/L including despite of resuscitation.

Vasoactive inotropic score

VIS was calculated as ((epinephrine+norepinephrine) ð $\mu g/kg/min$)×100+((dobutamine+dopamine)) $\mu g/kg/$ uses related min)+(milrinone $\mu g/kg/min) \times 10$ +levosimendan μg/ kg/min×50+(vasopressin units/kg/min)×10000.³²

Successful weaning off ECMO

Successful weaning was defined as maintaining stable condition within 24 hours of ECMO weaning.

Study intervention

The study flowchart is detailed in figure 1.

All patients received fluid resuscitation, antibiotic, vasoactive drugs and control of the focus of infection according to the international guidelines of Surviving $\frac{1}{3}$ Sepsis Campaign 2021.⁹ The gold of MAP is $\geq 65 \text{ mm Hg}$. For multiple-organ dysfunction, life support technologies such as mechanical ventilation and continuous renal \triangleright replacement therapy (CRRT) are provided as needed. training, and The patient's primary physicians will determine the management of other comorbidities.

ECMO implantation and management

<u>0</u> All patients in cohort 1 will initiate ECMO as fast as possible. A maximum of 6 hours is allowed between enrolment and the actual initiation of ECMO. ECMO catheterisation and management will be operated by an experienced ECMO team and carried out at the bedside. nologies The initiation and weaning or cessation time, the data of ECMO will be recorded by nurses.

Therapy mode

Venoarterial or venovenoarterial mode will be chosen according to the patient's condition. Intra-aortic balloon pump will be performed simultaneously when necessary to relieve the afterload of left ventricle.

Canulation

All patients will undergo peripheral cannulation. The arterial catheter is placed into the femoral artery, and

to text

an



Figure 1 Study flowchart. ECMO, extracorporeal membrane oxygenation; PiCCO, pulse indicator continuous cardiac output techonology; RSS, refractory septic shock; SCM, septic cardiomyopathy.

the venous catheter is placed into the femoral vein. Ultrasound examination will be performed bedside to select vessels with better conditions and suitable diameter canula before catheterisation. After the arterial cannulation, the distal perfusion catheter is inserted to perform lower limb perfusion.

The blood flow and the goal

The initial flow rate is 80~100 mL/kg of ideal body weight/min. The ECMO blood flow is adjusted to: (1) maintain a MAP >65 mm Hg; (2) reach a preoxygenator oxygen saturation >65%; (3) restore blood lactate to normal level; (4) revert the multiple organ dysfunction syndrome (MODS).

Management

- Inotropes are discontinued or reduced to minimal dosed within a few hours of achieving goal-directed flows (norepinephrine or epinephrine $<0.05 \,\mu g/kg/$ min, and dopamine or dobutamine $<5 \mu g/kg/min$ are suggested).
- During the period of ECMO, maintain a haemoglobin of >70 g/L, a platelet count of $\geq 50 \times 10^9$ /L, antithrombin III (AT3) of >80% and fibrinogen >2 g/L.
- Echocardiography is performed at least daily to monitor cardiac function.

Anticoagulation

Unfractionated heparin (UFH) is recommended for anticoagulation. A bolus dose of UFH is administered at cannulation followed by continuous infusion. Activated

Protected by copyright, including for uses related to text partial thrombin time is targeted between 1.5 and 2 times of normal, or active clotting time (ACT) is targeted between 180 s and 220 s. If the patient is at high risk of bleeding, the target ACT value is lowered to 160s. UFH may be stopped for severe bleeding or coagulation disorders.

Wean off ECMO

ECMO weaning should be considered when patients a exhibit stable haemodynamics and sufficient cardiac recovery.33

Indications for ECMO weaning: (1) adequate upper limb partial pressure of oxygen in arterial blood and **B** saturation with fraction of inspire oxygen <50%, peak inspiratory pressure <30 cm H_oO, positive end expiratory pressure <8 cmH_oO from ventilator when ECMO gas flow at 21%; (2) ECMO flow is reduced to $10\% \sim 25\%$ of normal blood flow or 1.5 L/min; (3) patients exhibit stable haemodynamic (MAP >65 mm Hg, pulse pressure >20 mm Hg), ScvO₂ >70%, LVEF >40%, blood lactate <2.0 mmol/L and without malignant arrhythmia on no/ low doses vasoactive, inotropic support (norepinephrine or epinephrine <0.02 µg/kg/min, and dopamine or dobutamine $<5 \mu g/kg/min$) for more than 2 hours.

The ECMO weaning test is gradually performed according to the patient's systemic haemodynamics and tissue perfusion improvement.^{33 34} During weaning, ECMO flow is decreased progressively by 500 mL every 5-10min. Patients are evaluated after 3-5min of no support (circuit clamped) or alternatively at minimum of 1L/min of support. Successful weaning is defined as

maintaining stable condition within 48 hours of ECMO weaning.

Cessation of ECMO

ECMO will be discontinued with one of the following conditions: (1) brain death; (2) other vital organ dysfunction is difficult to reverse; (3) major bleeding (defined as fatal bleeding, and/or symptomatic bleeding, does ≥ 1 of the following factors apply: (1) bleeding at a critical site, such as intracranial, intraspinal, intraocular, retroperitoneal, intra-articular or pericardial or intramuscular with compartment syndrome; (2) haemodynamic instability; (3) clinically overt bleeding with haemoglobin decrease $\geq 2g/dL$ or administration of ≥ 2 units red blood cells;³⁵ (4) there are no signs of cardiac function recovery and no better therapeutic regimen after 7 to 10 days of ECMO treatment; (5) an uncontrollable infection.

Primary outcome

The primary outcome is 30-day survival measured from the date of enrolled (D0) until death or day 30. For patients who were discharged alive from ICU, information on the primary endpoint will be acquired by a telephone call.

Secondary outcomes

The secondary outcomes include: (1) survival to ICU discharge, hospital survival, 6-month survival and quality of life for long-term survival (EQ-5D score); (2) successful rate of ECMO weaning; (3) long-term survivors' cardiac function was evaluated according to Doppler echocardiography; (4) the number of days alive without CRRT, mechanical ventilation and vasopressor (the numbers of CRRT-free days, mechanical ventilation-free days and vasopressor-free days, between D0 and up to D30); (5) ICU and hospital length of stay.

Safety assessments

In addition to the focus on prognosis of patients with SCM treated with venoarterial ECMO, the safety of the venoarterial ECMO treatment is also a major focus. The complications potentially related to ECMO treatment include: (1) major bleeding associated with anticoagulants; (2) thrombosis (ischaemic stroke, pulmonary embolism, deep venous thrombosis or catheter-associated thrombosis during study confirmed by ultrasound or CT scan); (3) leg ischaemia; (4) cannulation-related injuries (such as arterial laceration, arterial aneurysm and peripheral nerve defect).

Sample size

According to a retrospective study from Bréchot et al, the survival rate of patients with severe sepsis-induced cardiomyopathy was 60% in venoarterial ECMO group and 25% in control group, elevation in survival of 35% in venoarterial ECMO group can be expected. Considering the high cost and uncertain therapeutic effect of ECMO treatment, fewer participants may choose ECMO treatment than conventional treatment, with an estimated

ratio of 1:1.5. The sample size for differences between two independent proportions was calculated by the PASS V.14.0 software to ensure 80% power using a two-sided test with a significance level of α =0.05. We need 23 participants in venoarterial ECMO treatment cohort and 35 participants in conventional treatment cohort. Considering a projected dropout rate of 10%, the sample size of venoarterial ECMO treatment cohort and conventional treatment cohort should be 25 and 39, respectively. The total sample size should be 64.
Data collection and follow-up
Each investigator from the six participating ICUs total sample size should be 64.

9 was trained to the protocol and data collection in the Case Record Form before trial initiation. The 8 data is managed and closed by the Clinical Research centre of the Second Affiliated of Guangzhou Medical University (China).

Flowchart of patient follow-up is shown in table 1. Demographic data and medical history will be collected. Detailed data including reasons for ICU admission, cause of septic shock, focus of infection, acute physiology and chronic health evaluation II (APACHEII) score, dates of hospital and ICU admission will be recorded. Details of mechanical ventilation, vasopressor and CRRT will be documented daily. Sequential organ failure assessment score (SOFA) will be calculated at baseline, day 1 (D1), D3 and D7. Cardiac function will be assessed by echocardiography and recorded at baseline, D1, D3, D5, D7, D10, D14, D30. Blood will be collected at baseline, D1, D3, D5, D7. The following laboratory results will be recorded: white blood cell count and differentials in peripheral circulation, serum electrolyte levels, liver and myocardial enzyme concentrations, arterial blood gas analysis, C **B** reactive protein, procalcitonin and lactate. EQ-5D assess-Bul ment will be acquired through phone call at 6 months. ≥

During ECMO intervention, details of initiation, mode, setting parameters, weaning or cessation, complications will be noted.

ğ All enrolled participants will be followed to determine adverse events, cardiac function recovery and mortality until death or at ICU/hospital discharge, 30 days and 6 months. If the participants survive beyond 30 days and 6 months, they will be requested to revisit the hospital for a cardiac ultrasound examination or provide a cardiac ultrasound report from a local hospital.

Statistical analysis
Data will be double-checked by the clinical research team,

and the database is managed and closed by the Clinical Research centre of the Second Affiliated of Guangzhou Medical University (China).

For each cohort, quantitative variables with normal distribution will be described as mean and SD. Quantitative variables with skewed distribution will be described as median (M) and (IQR, 25th percentile to 75th percentile). Qualitative variables will be described as frequency and percentage.

trair

Elowebart of patient follow up

	Screening	Inclusion D0	Study period	Death/30 days	Follow-up 6 months
Baseline information					
Demographic data and history	\checkmark				
Inclusion and exclusion criteria	\checkmark				
Written informed consent					
Diagnosis and focus of infection	\checkmark				
Vital signs					
APACHEII/SOFA		\checkmark	√D1, 3, 7		
Efficacy observation					
Cardiac function assessment (CI, LVEF)	\checkmark		√D1, 3, 5, 7, 10, 14		
ECMO intervention					
Treatment with vesopressor	\checkmark				
Mechanical ventilation					
CRRT implication					
Laboratory tests	\checkmark		√D1, 3, 5, 7		
Safety observation					
Complications of ECMO					
Adverse event					
Additional observation					
ICU and hospital LOS					
Alive or dead status					
Life quality (EQ-5D score)					

APACHEII, Acute Physiology and Chronic Health Evaluation II; CI, cardiac index; CRRT, continuous renal replacement therapy; ECMO, extracorporeal membrane oxygenation; LOS, length of stay; SOFA, Sequential Organ Failure Assessment.

The effect of venoarterial ECMO treatment versus conventional treatment on 30 days survival will be performed using Fisher's exact test, with secondary analysis by Kaplan-Meier survival analyses, comparison using a log rank test. A propensity score-weighted analysis was done for treatment-effect estimation. Covariate balance between the two groups was assessed after weighting, and we considered an absolute standardised difference of less than 0.1 as evidence of balance. The effect of ECMO on survival at 30 days will be estimated within the weighted pseudopopulation. Adjusted Kaplan-Meier estimator and log-rank test (considering the weighting scheme) will be used. Decision tree analysis will be used to establish a decision tree model to decide whether sepsis-induced cardiogenic shock patients need ECMO treatment.

Safety will be analysed by the frequency of complications in both cohorts and comparing rates using χ^2 or Fisher's exact test, with an alpha risk set at 0.05.

Statistical analyses of the prespecified secondary endpoints will be performed with descriptive and inductive statistical methods. Categorical variables will be compared using the χ^2 or Fisher's exact test, as appropriate. Continuous variables will be compared using Student's t test or the Mann-Whitney test, as appropriate.

Therapeutic efficiency will be analysed using the data of the full analysis set and per protocol set; safety evaluation

will be based on the data of the safety analysis set for statistical analysis.

All analyses were performed with commercially available statistical software SPSS 22.0 and R software version 4.1.0 or later.

ETHICS AND DISSEMINATION

Protected by copyright, including for uses related to text and data mining, AI training The study protocol has been approved by the Clinical Research and Application Institutional Review Board of the Second Affiliated Hospital of Guangzhou Medical University (V.2.0, registration number: 2020-hs-51; date of approval: 17 November 2020). Participants will be screened and enrolled from ICU patients with sepsis-induced cardiogenic shock by clinicians, with no public advertisement for recruitment. When the patient fulfils the criterion of recruitment, written informed consent (online supplemental file 1) should be obtained from the patient or a responsible surrogate before enrolled. After enrolled, the participants can decide whether to accept ECMO based on their personal conditions. All information from the participants will be kept private and will not be provided to any company or institution. Results will be disseminated in research journals and through conference presentations.

Author affiliations

Intensive Care Unit, Guangzhou Medical University Second Affiliated Hospital, Guangzhou, Guangdong, China

Open access

Contributors W-yC and Z-bG designed the trial. T-yK, QY, Y-cW and X-mX participated in the discussion of the plan, D-IW and Z-hZ made a decision to revise the plan. Z-bG, T-yK, W-xC, X-hC, QY, Y-cW, Q-rW and FZ conduct the plan. W-yC draft the manuscript. D-IW and Z-hZ provided critical revision of the manuscript. D-IW and W-yC obtained funding for the trial. All authors discussed and helped to improve the protocol and read and approved the final manuscript.

Funding This study was funded by a Grant from the Second Affiliated Hospital of Guangzhou Medical University (2020-LCYJ-DZX-02) and Runze Medical Research Fund of Wu Jieping Medical Foundation (320.6750.2023-02-6).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer-reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Wei-yan Chen http://orcid.org/0000-0002-5101-8839 Xu-ming Xiong http://orcid.org/0000-0002-6333-7319 Zhen-hui Zhang http://orcid.org/0000-0002-9790-8597

REFERENCES

- Singer M, Deutschman CS, Seymour CW, et al. The third international consensus definitions for sepsis and septic shock (sepsis-3). JAMA 2016;315:801–10.
- 2 Rudd KE, Johnson SC, Agesa KM, *et al.* Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. *Lancet* 2020;395:200–11.
- 3 Fleischmann C, Scherag A, Adhikari NKJ, et al. Assessment of global incidence and mortality of hospital-treated sepsis. Current estimates and limitations. AM J Respir Crit Care Med 2016;193:259–72.
- 4 Reinhart K, Daniels R, Kissoon N, et al. Recognizing sepsis as a global health priority - A WHO resolution. N Engl J Med 2017;377:414–7.
- 5 Narváez I, Canabal A, Martín C, *et al.* Incidence and evolution of sepsis-induced cardiomyopathy in a cohort of patients with sepsis and septic shock. *Med Intensiva (Engl Ed)* 2018;42:283–91.
- 6 Sato R, Kuriyama A, Takada T, *et al.* Prevalence and risk factors of sepsis-induced cardiomyopathy: a retrospective cohort study. *Medicine (Baltimore)* 2016;95:e5031.
- 7 Beesley SJ, Weber G, Sarge T, et al. Septic cardiomyopathy. Crit Care Med 2018;46:625–34.
- 8 Ravikumar N, Sayed MA, Poonsuph CJ, *et al.* Septic cardiomyopathy: from basics to management choices. *Curr Probl Cardiol* 2021;46:100767.
- 9 Evans L, Rhodes A, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. Intensive Care Med 2021;47:1181–247.
- 10 Hernandez G, Bruhn A, Luengo C, *et al.* Effects of dobutamine on systemic, regional and microcirculatory perfusion parameters in septic shock: a randomized, placebo-controlled, double-blind, crossover study. *Intensive Care Med* 2013;39:1435–43.
- 11 Annane D, Vignon P, Renault A, et al. Norepinephrine plus dobutamine versus epinephrine alone for management of septic shock: a randomised trial. *Lancet* 2007;370:676–84.

- 12 Meng J, Hu M, Lai Z, *et al.* Levosimendan versus dobutamine in myocardial injury patients with septic shock: a randomized controlled trial. *Med Sci Monit* 2016;22:1486–96.
- 13 Gordon AC, Perkins GD, Singer M, *et al*. Levosimendan for the prevention of acute organ dysfunction in sepsis. *N Engl J Med* 2016;375:1638–48.
- 14 Pinto BB, Rehberg S, Ertmer C, et al. Role of levosimendan in sepsis and septic shock. Curr Opin Anaesthesiol 2008;21:168–77.
- 15 Martin L, Derwall M, Al Zoubi S, *et al*. The septic heart: current understanding of molecular mechanisms and clinical implications. *Chest* 2019;155:427–37.
- 16 Vallabhajosyula S, Jentzer JC, Geske JB, et al. New-onset heart failure and mortality in hospital survivors of sepsis-related left ventricular dysfunction. *Shock* 2018;49:144–9.
- 17 Han L, Zhang Y, Zhang Y, et al. Risk factors for refractory septic shock treated with VA ECMO. *Ann Transl Med* 2019;7:476.
- 18 Kirkpatrick BV, Krummel TM, Mueller DG, et al. Use of extracorporeal membrane oxygenation for respiratory failure in term infants. *Pediatrics* 1983;72:872–6.
- 19 Riera J, Argudo E, Ruiz-Rodríguez JC, *et al.* Extracorporeal membrane oxygenation for adults with refractory septic shock. *ASAIO J* 2019;65:760–8.
- 20 Park TK, Yang JH, Jeon K, et al. Extracorporeal membrane oxygenation for refractory septic shock in adults. Eur J Cardiothorac Surg 2015;47:e68–74.
- 21 Huang C-T, Tsai Y-J, Tsai P-R, et al. Extracorporeal membrane oxygenation resuscitation in adult patients with refractory septic shock. J Thorac Cardiovasc Surg 2013;146:1041–6.
- 22 Formica F, Avalli L, Martino A, et al. Extracorporeal membrane oxygenation with a poly-methylpentene oxygenator (Quadrox D). The experience of a single Italian centre in adult patients with refractory cardiogenic shock. ASA/O J 2008;54:89–94.
- 23 Tembo M, Harvey C, Duthie M, *et al.* Extracorporeal membrane oxygenation for refractory septic shock in children: one institution's experience. *Pediatr Crit Care Med* 2009;10:534–5.
- 24 McCune S, Short BL, Miller MK, et al. Extracorporeal membrane oxygenation therapy in neonates with septic shock. J Pediatr Surg 1990;25:479–82.
- 25 Brierley J, Carcillo JA, Choong K, et al. Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: 2007 update from the American college of critical care medicine. Crit Care Med 2009;37:666–88.
- 26 Davis AL, Carcillo JA, Aneja RK, et al. American college of critical care medicine clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock. Crit Care Med 2017;45:1061–93.
- 27 Bréchot N, Luyt C-E, Schmidt M, et al. Venoarterial extracorporeal membrane oxygenation support for refractory cardiovascular dysfunction during severe bacterial septic shock*. Crit Care Med 2013;41:1616–26.
- 28 Bréchot N, Hajage D, Kimmoun A, et al. Venoarterial extracorporeal membrane oxygenation to rescue sepsis-induced cardiogenic shock: a retrospective, multicentre, international cohort study. Lancet 2020;396:545–52.
- 29 Falk L, Hultman J, Broman LM. Extracorporeal membrane oxygenation for septic shock. *Crit Care Med* 2019;47:1097–105.
- 30 Barreto BB, Gusmao-Flores D. Long-term mortality in extracorporeal membrane oxygenation. Crit Care Med 2018;46:e348.
- 31 Ling RR, Ramanathan K, Poon WH, et al. Venoarterial extracorporeal membrane oxygenation as mechanical circulatory support in adult septic shock: a systematic review and meta-analysis with individual participant data meta-regression analysis. Crit Care 2021;25:246.
- 32 Koponen T, Karttunen J, Musialowicz T, et al. Vasoactive-inotropic score and the prediction of morbidity and mortality after cardiac surgery. Br J Anaesth 2019;122:428–36.
- 33 Lorusso R, Shekar K, MacLaren G, et al. ELSO interim guidelines for venoarterial extracorporeal membrane oxygenation in adult cardiac patients. ASAIO J 2021;67:827–44.
- 34 Zwischenberger JB, Pitcher HT. Extracorporeal membrane oxygenation management: techniques to liberate from extracorporeal membrane oxygenation and manage post-intensive care unit issues. *Crit Care Clin* 2017;33:843–53.
- 35 Schulman S, Kearon C, Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost* 2005;3:692–4.