



BMJ Open Screening for social anxiety disorder in students of Jordan universities after COVID-19 pandemic: a cross-sectional survey study

Ahlam J Alhemedi ¹, Othman Beni Yonis,¹ Hadeel Allan,¹ Ghalia abu Mohsen,² Ayham Almasri,¹ Haitham Abdulrazzeq,¹ Islam Khasawneh,¹ Osama El-khateeb,¹ Abdallah Y Naser ³

To cite: Alhemedi AJ, Beni Yonis O, Allan H, *et al.* Screening for social anxiety disorder in students of Jordan universities after COVID-19 pandemic: a cross-sectional survey study. *BMJ Open* 2025;**15**:e086066. doi:10.1136/bmjopen-2024-086066

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-086066>).

Received 04 March 2024
Accepted 09 January 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

¹Department of Public Health and Family Medicine, Faculty of Medicine, Jordan University of Science and Technology, P.O. Box 3030, Irbid, 22110, Jordan

²Department of Clinical Medical Sciences, Faculty of Medicine, Yarmouk University, Irbid, Jordan

³Department of Applied Pharmaceutical Sciences and Clinical Pharmacy, Faculty of Pharmacy, Isra University, Amman, Jordan

Correspondence to
Dr Ahlam J Alhemedi;
ajalhmedy@just.edu.jo

ABSTRACT

Objective To examine the prevalence rate of social anxiety disorder (SAD) among university students in Jordan after the COVID-19 pandemic and its associated predictors.

Design A cross-sectional online survey study that was conducted in Jordan between January and December 2023.

Setting: Universities in Jordan.

Participants Healthy university students from any specialty currently enrolled at a Jordanian university.

Primary outcome measure The prevalence rate of SAD, which was assessed using the Social Phobia Inventory.

Results A total of 851 university students participated in this study. More than half of them (65%) were women. The mean age of the study participants was 21.9 (2.7) years. The majority of them (70.6%) were studying medical fields. The median number of times the study participants got infected with COVID-19 was 1.0 (IQR: 0.0–2.0).

The median number of viewing hours spent on social networking sites was 4.0 (IQR: 3.0–6.0). The median SAD score was 19 (IQR: 10–32) out of 68, which represents 27.9% of the maximum attainable score. Up to 45.4% of the study participants were susceptible to SAD, with 12.5% of the study participants reporting severe to very severe SAD symptoms. Students older than 21.9 years were 32% less likely to report SAD symptoms compared with younger students ($p < 0.01$).

Conclusion Jordanian universities students are increasingly likely to report SAD symptoms. The data show how important it is for universities to recognise and deal with this mental health issue. Focused treatments and support networks could help students with social anxiety problems to deal with them.

INTRODUCTION

Mental health constitutes a cornerstone of satisfaction and quality of life, yet among university undergraduate students worldwide, it remains a complex and prevalent concern.¹ Extensive psychiatric and psychological research in various developed and developing nations over the past decades has

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The use of an online survey study design is a cost-effective approach to reach wide and diverse demographic groups of university students.
- ⇒ The cross-sectional study design limited our capacity to track patients over time and investigate causality among the study variables.
- ⇒ Reporting bias frequently occurs in self-administered questionnaire surveys.

consistently highlighted the heightened prevalence of anxiety, depression and stress within this demographic group.^{2–4} In 2019, anxiety disorders affected an estimated 301 million individuals globally, establishing them as the most prevalent mental health condition worldwide.⁵ Notably, social anxiety disorder (SAD), formerly known as social phobia, emerges as a prominent subtype within the spectrum of anxiety disorders.⁶

SAD is a debilitating condition characterised by a profound apprehension towards social interactions or observation, often resulting in hindrance in daily functioning and significant distress.^{7–8} The prevalence of SAD has been established through epidemiological studies, with a lifetime prevalence ranging from 8.4% to 15% and a present-day prevalence ranging from 5% to 10%.^{9–15} Moreover, its impact extends beyond personal well-being, significantly affecting university life and academic persistence and performance.¹⁶ Previous research indicates that SAD is associated with distorted body image and diminished self-esteem, ultimately leading to adverse effects on the scholarly performance of students.^{17–18}

The emergence of the coronavirus disease in the form of the 2019 (COVID-19) pandemic in early 2020 has significantly altered global

educational systems and mental health landscapes.^{19 20} Studies have reported a substantial increase in anxiety disorder cases worldwide by 25.6%, attributed, in part, to the effects of the pandemic,²¹ suggesting a prolonged course for mental health recovery.²² Specifically, on 2 March 2020, Jordan recorded its initial case of COVID-19.²³ The COVID-19 pandemic significantly impacted university students' psychological well-being, with previous studies reporting the high prevalence of a wide range of psychological disorders among university students, including anxiety and depression.^{24–30} A previous study of 4301 individuals in Jordan found that the COVID-19 pandemic affected social relationships markedly, with 30.3% of the participants reporting that the pandemic negatively impacted their social relationships and communication with others.³¹ During the lockdown period, the probability of social anxiety and communication difficulties increased.³² A study by Liang *et al* has reported that online teaching during the pandemic increased the likelihood of social anxiety.³³

In Jordan, higher education is offered by universities and community colleges.³⁴ Tertiary education in Jordan commenced in 1951. Jordan has 29 universities, comprising 10 public and 19 private institutions. In Jordan, higher education institutions are classified into two categories: colleges and universities.³⁴ The prerequisite for university admission is the possession of a general secondary education certificate. The duration of study varies from four to 6 years, depending on the field of study.³⁴

In Jordan, as elsewhere, SAD is considered a significant concern among university students.^{35 36} Risk factors, such as heightened avoidance of social situations³⁷ and reduced social support,^{38 39} have become worse during the COVID-19 pandemic. Nevertheless, research exploring SAD among Jordanian university students, particularly in the post-pandemic context, is limited. Therefore, comprehending the pandemic's impact on the mental well-being of Jordanian university students, especially concerning SAD, is a critical imperative. Hence, the objective of this study is to examine the SAD profile among universities students after the COVID-19 pandemic in Jordan, and its associated predictors.

METHODS

Study design

This is an online cross-sectional study that was conducted in Jordan between January and December 2023.

Participant recruitment

Convenience sampling was used to recruit the participants using an electronic questionnaire linked to a Google form. The study participants were invited through social media platforms such as Facebook, WhatsApp, Snapchat and Instagram. The inclusion criterion of the study population was that the participants should be university students from any specialty currently enrolled at a

Jordanian university. There were no limitations regarding the year or field of study. The cover letter sent to the participants outlined the inclusion criterion. Eligible participants were invited to take part in the study. The cover letter also included the study's goals and objectives. The participants were requested to engage in the study after providing their consent.

Data collection and measures

The Jordanian university students completed the questionnaire after it was distributed through online media. The questionnaire included an English version of the Social Phobia Inventory (SPIN) screening instrument and questions regarding demographic information, including age, gender, student specialty, frequency of COVID-19 infections and time spent on social media (see online supplemental material). The SPIN scale consisted of 17 questions on various anxiety-inducing circumstances, to be rated on a range of 0 (not at all) to 4 (very). Scoring below 20 indicated the normal level, 21–30 indicated mild SAD, 31–40 indicated moderate SAD, 41–50 indicated severe SAD and scores above 51 indicated very severe SAD. The SPIN scale has proven validity, test–retest reliability and internal consistency. This scale can be used for screening for SAD in languages other than English.⁴⁰ The SPIN's psychological characteristics demonstrated strong test–retest reliability over 2 weeks (Spearman's rank correlation coefficient, 0.78–0.89) and internal consistency (Cronbach's alpha coefficient, 0.82–0.94) in an adult community.⁴¹ Convergent validity has been demonstrated through significant correlations between the SPIN and other adult assessment tools for social phobia or anxiety (eg, Liebowitz Social Anxiety Scale and the Brief Social Phobia Scale), while lower correlations with measures less pertinent to social phobia (eg, Medical Outcomes Study Short Form-36, Sheehan Disability Scale, Marks Fear Questionnaire) have evidenced the scale's discriminant validity.⁴⁰

Pilot study

We ran a pilot study with 20 participants who satisfied our criteria for inclusion but were not part of the final study population. The pilot study assessed the participants' comprehension of the questionnaire and inquired about any challenges they had while completing it. The participants in the pilot study confirmed that the questionnaire was easy to understand and no clarification was needed for them to complete it.

Questionnaire validity

Cronbach's alpha was employed to assess the reliability of the questions. The minimum permissible factor loading was set at 0.70.⁴² We computed Cronbach's alpha coefficients to measure the internal consistency of the SAD scale. The Cronbach's alpha coefficient for SAD was calculated to be 0.932, indicating an excellent internal consistency.

Data analysis

The collected data were analysed using the statistical analysis software SPSS V.29. Descriptive analysis was used to consider the data in this study. Categorical variables were presented as percentages and frequencies. Continuous variables were presented as mean and SD or as the median and IQR based on the normality of the variable distribution. The normality of the SAD score was assessed using skewness, kurtosis measurements and a histogram. The predictors of SAD were determined by binary logistic regression analysis, with the dummy variable being the average SAD score of the participants in the study. We used the average SAD score for the study participants as the cut-off to define the dummy variable (the dependent variable) in the regression model, where any individual who had an SAD score above the average for the sample was assigned the number '1' and anyone who had an SAD score below the average score for the sample was assigned the number '0'. The independent variables for the logistic regression analysis were 'participants' demographic characteristics' which were gender, age, field of study, number of viewing hours spent on social networking sites and number of times infected with COVID-19. A 95% CI was used to demonstrate the statistical significance of the results, with a significance threshold set at 5%.

Sample size

Using a CI of 95%, an SD of 0.5 and a 5% error margin, the targeted minimum sample size was 385 students.

Patient and public involvement

Patients and/or the public were not involved in this study.

RESULTS

Participants' demographic characteristics

Table 1 below presents the participants' demographic characteristics. A total of 851 university students participated in this study. More than half of them (65%) were women. The mean age of the study participants was 21.9 (2.7) years. The majority of them (70.6%) were studying medical fields. The median number of times the study participants got infected with COVID-19 was 1.0 (IQR:

Table 1 Participants' demographic characteristics (n=851)

Variable	Frequency	Percentage
Gender		
Females	553	65
Mean age (years) (SD)	21.9 (2.7)	
Field of study		
Medical field	601	70.6
Non-medical field	250	29.4
Median number of viewing hours spent on social networking sites (IQR)	4.0 (3.0–6.0)	

Table 2 SAD profile

SAD level	SAD score	Percentage of participants (frequency)
Normal	0–20	54.6 (n=465)
Mild	21–30	18.8 (n=160)
Moderate	31–40	14.1 (n=120)
Severe	41–50	7.4 (n=63)
Very severe	51 and higher	5.1 (n=43)

SAD, social anxiety disorder.

0.0–2.0). The median number of viewing hours spent on social networking sites was 4.0 (IQR: 3.0–6.0).

Social anxiety disorder profile

The median SAD score was 19.0 (IQR: 10–32) out of 68, which represents 27.9% of the maximum attainable score. Around 45.4% of the study participants were susceptible to SAD and reported its symptoms. Around 12.5% of the study participants reported severe to very severe SAD symptoms (table 2).

Predictors of social anxiety disorder

Table 3 below presents the findings of a binary logistic regression analysis. Younger students were more likely to report severe SAD symptoms compared with older

Table 3 Predictors of reporting SAD symptoms

Variable	OR of reporting SAD symptoms (95% CI)	P value
Gender		
Female (reference category)	1.00	
Male	0.83 (0.62 to 1.01)	0.186
Mean age (category)		
21.9 years or lower	1.00	
Older than 21.9 years	0.68 (0.52 to 0.89)	0.005*
Field of study		
Medical field (reference category)	1.00	
Non-medical field	0.93 (0.69 to 1.25)	0.608
Median number of times infected with COVID-19 (category)		
Once (reference category)	1.00	
More than once	1.14 (0.84 to 1.55)	0.392
Median number of viewing hours you spend on social networking sites (category)		
4 hours or fewer (reference category)	1.00	
More than 4 hours	1.10 (0.82 to 1.47)	0.518

*p<0.01.
SAD, social anxiety disorder.

students (p value <0.01). Other demographic characteristics did not significantly influence the likelihood of reporting SAD symptoms among university students (p value >0.05).

DISCUSSION

The objective of this study was to examine the SAD profile and its associated predictors among university students after the COVID-19 pandemic in Jordan. This was an online cross-sectional study that was conducted among Jordanian university students who completed the SPIN questionnaire, which was distributed through online social media. Up to 45.4% of the study participants reported SAD symptoms, with 12.5% of them reporting severe to very severe symptoms. Younger students showed a higher likelihood of reporting more severe SAD symptoms compared with older students.

Our study highlights the prevalent susceptibility to report SAD symptoms among university students in Jordan after the COVID-19 pandemic, a finding supported by the observation that 45.4% of the participants exhibited susceptibility to report SAD symptoms. A previous study in Jordan examined the prevalence of SAD before the COVID-19 pandemic using the same assessment tool (SPIN) and found that the prevalence rate of SAD among university students was 27.9%.³⁶ The emergence of the COVID-19 pandemic has exacerbated mental health challenges among university students worldwide, including those in Jordan. This study found that over half of Jordanian students reported moderate to severe SAD symptoms during the COVID-19 pandemic. This proportion was significantly higher than the 9% to 16.1% prevalence rate of SAD reported in normal times before the pandemic.^{36 43–46} It has been globally confirmed that COVID-19 increases the risk of various mental disorders.^{27 31 47 48} Although this study used the SPIN to assess the severity of social anxiety and thus lacks clinical or expert diagnosis, it appears that social anxiety among Jordanian students has been significantly exacerbated by the COVID-19 pandemic and associated social isolation.

Consistent with our findings, after the onset of the COVID-19 pandemic, the prevalence of SAD among university students in India was reported to be 42.5%.⁴⁹ Additionally, a study conducted earlier by McLeish *et al*⁵⁰ observed an upsurge in social interaction anxiety among university students following a duration of social isolation when resuming more recurring social meetings. Like numerous other countries, the Jordanian Ministry of Defence implemented stringent public health measures to curb the spread of COVID-19, which might have increased the possibility of developing SAD symptoms. These measures encompassed mandatory mask-wearing in public areas, implementation of lockdowns, enforcement of a nationwide curfew, adherence to social distancing guidelines and the adoption of remote learning.^{51 52} While imperative for public health, these measures had profound implications on individuals' mental well-being,

notably impacting university students⁴⁷ who faced disruptions to their academic, social and personal lives.^{48 53 54}

The prevalence of severe to very severe SAD in a previous study in Jordan that examined the prevalence of SAD before the COVID-19 pandemic was reported to be 2.3%, lower compared with our study findings.³⁶ Another study that was conducted in Jordan after the pandemic reported that the prevalence rate of SAD among university students was 25.8%.⁵⁵ Besides, their study reported that the prevalence of severe to very severe SAD was 14.5%.⁵⁵ The percentage of reporting severe SAD symptoms in this study may align with, surpass, or fall below rates observed in other regions and populations. For instance, the percentage of severe to very severe SAD in Jordan aligns with that found among university students in Saudi Arabia, where 12.4% of students exhibited such levels.⁵⁶ However, the percentage of severe or extremely severe SAD reported in Iran, Nigeria and Malaysia exceeds that in Jordan, ranging from 13.2% to 20%.^{57–59} Conversely, the percentage of reporting severe to very severe SAD symptoms in Jordan is higher than that found among university students in Ethiopia, where 5.2% of students displayed severe to very severe SAD.⁶⁰ These findings are particularly alarming as they indicate a considerable proportion of university students experiencing severe SAD, which can significantly impair their daily functioning and overall well-being.⁷

Our research demonstrates a significant association between age and susceptibility to report SAD symptoms within the university student population in Jordan. Notably, among students aged over 21.9 years, there was a 32% lower likelihood of reporting SAD symptoms compared with their younger peers. This finding is consistent with numerous previous studies. A study on the global prevalence and impact of anxiety disorders during the COVID-19 pandemic in 2020 indicated a higher prevalence of anxiety disorders among younger age groups.²¹ Additionally, existing literature consistently implies an association between age and the onset of SAD, often beginning in childhood or adolescence.⁶¹ Previous studies conducted among undergraduate students at the Faculty of Science and Humanities of the Ekiti State University in Southwestern Nigeria and medical students at Taibah University in Saudi Arabia further support our findings. The Nigerian study revealed that a younger age significantly increased the risk of social anxiety among participants, with those under 21 years being twice as likely to experience SAD.⁶² Likewise, the study among medical students in Saudi Arabia found that SAD is less common among older students.⁶³ These findings collectively support the notion of an inverse association between age and susceptibility to report SAD symptoms, as observed in our study. Accordingly, they emphasise the essentiality of implementing interventions to address the distinctive mental challenges among younger university students.

In clinical settings, older students engage more frequently with patients and refine their interviewing

abilities, potentially leading to a lower prevalence of SAD among this demographic. Poor communication abilities are associated with a higher likelihood of reporting SAD symptoms. At the same time, SAD itself decreases the possibility of effective communication. Previous research indicates that students at advanced academic levels and older age groups typically demonstrate a more favourable perspective toward acquiring communication skills.⁶⁴ Likewise, another study suggests that final-year students exhibit superior communication abilities compared with their first-year peers, indicating a deeper understanding of effective communication.⁶⁵ Conversely, first and second-year students appear to encounter an increased susceptibility to SAD, potentially due to the challenges associated with adapting to university life.^{46 66} The transition to university, especially for those in their initial years, involves newfound independence and exposure to unfamiliar social dynamics, which can contribute to heightened levels of SAD.^{46 66} Research conducted in Turkey has revealed that first and second-year university students often demonstrate elevated anxiety stress scores in comparison to their counterparts.⁶⁷ The onset of SAD at an early stage among undergraduate students raises significant concerns regarding its potential long-term impacts on mental and social well-being.³ Considering that many undergraduates start their university life at a young age, experiencing SAD during this critical period may lead to ongoing unfavourable outcomes.³ Hence, implementing early intervention strategies and establishing healthy support systems are imperative to alleviate these consequences and promote the well-being of university students.

This study is limited. The cross-sectional study design limited our capacity to track patients over time and investigate causality among the study variables. Online survey studies using convenience sampling are criticised as a sample method because they can impact the generalisability of the study results. Therefore, our study sample might have missed other demographic groups. Furthermore, we did not inquire about the participants' specific areas of study; instead, we questioned whether they were pursuing a medical or non-medical field, which limits insights into their specific field of study. Reporting bias frequently occurs in self-administered questionnaire surveys. The diagnosis of SAD in this study is based on a self-administered questionnaire tool, not a clinical diagnosis. Besides, we did not collect information on the medical history of other coexisting disorders. Therefore, we were unable to examine its impact on the prevalence of SAD. Consequently, our study results should be regarded with caution.

Based on our study findings, we recommend that higher education institutions should provide multicultural training for clinicians and other healthcare professionals who provide mental health services and support for their students. Besides, mental health services should include proactive measures that promote the well-being of the students. Confidentiality should be prioritised to

ensure student engagement in such important mental well-being programmes. Furthermore, academic and non-academic staff should be trained to identify and respond to the mental needs of students and provide the necessary support.

Based on the UNESCO recommendations,⁶⁸ policy-makers in the higher education sector have a responsibility to support students' mental health, in alignment with the United Nation's sustainable development goal number 3. Governments have the responsibility to support mental well-being in higher education institutions through policies and funding.⁶⁸ Future research, using longitudinal studies, is recommended to examine SAD among university students from different levels of study and conduct follow-up action for the students across their years of study. Furthermore, future research should examine the influence of cultural background on the prevalence of SAD among university students. Moreover, the role of artificial intelligence and the wide use of social media platforms should be investigated.

CONCLUSION

SAD is becoming a common issue among university students in Jordan. The results here highlight the significance of acknowledging and dealing with this mental health problem in higher education institutions. Mental health support services should be implemented in such institutions to support students' mental health well-being and, ultimately, their academic performance.

Acknowledgements We thank the Jordan University of Science and Technology for its support

Contributors AJA are responsible for conceptualisation, data acquisition, analysis, interpretation, writing original draft, review and editing. OBY, HAI, GaM, AA, HAb, IK, OE and AYN are responsible for writing, review and editing of the draft. AJA acted as the guarantor for this study and accepts full responsibility for the finished work and the conduct of the study, had access to the data and controlled the decision to publish.

Funding This study was supported by Jordan University of Science and Technology (research grant no: 20230176).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval We obtained approval for our study from the Institutional Review Board of the Jordan University of Science and Technology (Reference number: 24/156/2023). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Ahlem J Alhemedi <http://orcid.org/0009-0000-1258-0856>

Abdallah Y Naser <http://orcid.org/0000-0001-8440-7446>

REFERENCES

- Pedrelli P, Nyer M, Yeung A, et al. College Students: Mental Health Problems and Treatment Considerations. *Acad Psychiatry* 2015;39:503–11.
- Brenneisen Mayer F, Souza Santos I, Silveira PSP, et al. Factors associated to depression and anxiety in medical students: a multicenter study. *BMC Med Educ* 2016;16:282.
- Ibrahim AK, Kelly SJ, Adams CE, et al. A systematic review of studies of depression prevalence in university students. *J Psychiatr Res* 2013;47:391–400.
- Mkize LP, Nonkelela NF, Mkize DL. Prevalence of depression in a university population. *Curationis* 1998;21:32–7.
- WHO. Anxiety disorders, 2023. Available: <https://www.who.int/news-room/fact-sheets/detail/anxiety-disorders>
- National Collaborating Centre for Mental Health. National institute for health and care excellence: guidelines, in social anxiety disorder: recognition, assessment and treatment. Leicester (UK) The British Psychological Society & The Royal College of Psychiatrists; 2013.
- Altman M, Sarvaiya N, Neill Epperson C. Sex differences in anxiety and depression clinical perspectives. *Front Neuroendocrinol* 2014;35:320–30.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-5*, Vol 5. American Psychiatric Association, 2013.
- Acarturk C, de Graaf R, van Straten A, et al. Social phobia and number of social fears, and their association with comorbidity, health-related quality of life and help seeking: a population-based study. *Soc Psychiatry Psychiatr Epidemiol* 2008;43:273–9.
- Crome E, Grove R, Baillie AJ, et al. DSM-IV and DSM-5 social anxiety disorder in the Australian community. *Aust N Z J Psychiatry* 2015;49:227–35.
- Kessler RC, Berglund P, Demler O, et al. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005;62:593–602.
- Kessler RC, McGonagle KA, Zhao S, et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry* 1994;51:8–19.
- Ohayon MM, Schatzberg AF. Social phobia and depression: prevalence and comorbidity. *J Psychosom Res* 2010;68:235–43.
- Stein MB. An epidemiologic perspective on social anxiety disorder. *J Clin Psychiatry* 2006;67 Suppl 12:3–8.
- Stein MB, Craske MG. Treating Anxiety in 2017: Optimizing Care to Improve Outcomes. *JAMA* 2017;318:235–6.
- Broaden PN. Lived Experiences of College Students with Social Anxiety Disorder Accessing Support Services. *Walden Dissertations and Doctoral Studies* 2023.
- Izgiç F, Akyüz G, Doğan O, et al. Social phobia among university students and its relation to self-esteem and body image. *Can J Psychiatry* 2004;49:630–4.
- Ahmad RJ, Tahir Faque BH, Seidi PAM, et al. Prevalence of social anxiety in students of college of education–university of Garmian. *Researchers World* 2017;8:79.
- Elbay RY, Kurtulmuş A, Arpacioğlu S, et al. Depression, anxiety, stress levels of physicians and associated factors in Covid-19 pandemics. *Psychiatry Res* 2020;290:113130.
- Das K, Pingali MS, Paital B, et al. A detailed review of the outbreak of COVID-19. *Front Biosci (Landmark Ed)* 2021;26:149–70.
- Santomauro DF, Mantilla Herrera AM, Shadid J, et al. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet* 2021;398:1700–12.
- Quintana-Domeque C, Proto E. On the Persistence of Mental Health Deterioration during the COVID-19 Pandemic by Sex and Ethnicity in the UK: Evidence from Understanding Society. *B E J Econom Anal Policy* 2022;22:361–72.
- Al-Tammemi AB. The Battle Against COVID-19 in Jordan: An Early Overview of the Jordanian Experience. *Front Public Health* 2020;8:188.
- Alsairafi Z, Naser AY, Alsaleh FM, et al. Mental Health Status of Healthcare Professionals and Students of Health Sciences Faculties in Kuwait during the COVID-19 Pandemic. *Int J Environ Res Public Health* 2021;18:2203.
- Abuhamdah SMA, Naser AY, Abdelwahab GM, et al. The Prevalence of Mental Distress and Social Support among University Students in Jordan: A Cross-Sectional Study. *Int J Environ Res Public Health* 2021;18:21.
- Alyami HS, Naser AY, Dahmash EZ, et al. Depression and anxiety during the COVID-19 pandemic in Saudi Arabia: A cross-sectional study. *Int J Clin Pract* 2021;75:e14244.
- Naser AY, Dahmash EZ, Al-Rousan R, et al. Mental health status of the general population, healthcare professionals, and university students during 2019 coronavirus disease outbreak in Jordan: A cross-sectional study. *Brain Behav* 2020;10:e01730.
- Varghese A, George G, Kondaguli SV, et al. Decline in the mental health of nurses across the globe during COVID-19: A systematic review and meta-analysis. *J Glob Health* 2021;11:05009.
- Abukhalaf AH, Naser AY, Cohen SL, et al. Evaluating the mental health of international students in the U.S. during the COVID-19 outbreak: The case of University of Florida. *J Am Coll Health* 2024;72:3464–73.
- Alhemedi AJ, Qasaimeh MG, Abdo N, et al. Depression Among University Students in Jordan After the COVID-19 Pandemic: A Cross-Sectional Study. *Psychol Res Behav Manag* 2023;16:4237–49.
- Naser AY, Al-Hadithi HT, Dahmash EZ, et al. The effect of the 2019 coronavirus disease outbreak on social relationships: A cross-sectional study in Jordan. *Int J Soc Psychiatry* 2021;67:664–71.
- Kindred R, Bates GW. The Influence of the COVID-19 Pandemic on Social Anxiety: A Systematic Review. *Int J Environ Res Public Health* 2023;20:2362.
- Liang Z, Kang D, Zhang M, et al. The Impact of the COVID-19 Pandemic on Chinese Postgraduate Students' Mental Health. *IJERPH* 2021;18:11542.
- Jordan Education. The Jordanian higher education system. 2024. Available: <https://www.jordaneducation.info/higher-education>
- Alsaraireh F. The Psychological Effects of Social Phobia on Undergraduate Students in the south of Jordan. *JPTCP* 2023;30:367–77.
- Mustafa RB, Hamdan-Mansour AM, Hijazeen J, et al. Social phobia among university students in Jordan. *Life Sci J* 2014;11.
- Vriends N, Bolt OC, Kunz SM. Social anxiety disorder, a lifelong disorder? A review of the spontaneous remission and its predictors. *Acta Psychiatr Scand* 2014;130:109–22.
- Barnett MD, Maciel IV, Johnson DM, et al. Social Anxiety and Perceived Social Support: Gender Differences and the Mediating Role of Communication Styles. *Psychol Rep* 2021;124:70–87.
- Reinelt E, Aldinger M, Stopsack M, et al. High social support buffers the effects of 5-HTTLPR genotypes within social anxiety disorder. *Eur Arch Psychiatry Clin Neurosci* 2014;264:433–9.
- Johnson HS, Inderbitzen-Nolan HM, Anderson ER. The Social Phobia Inventory: validity and reliability in an adolescent community sample. *Psychol Assess* 2006;18:269–77.
- Connor KM, Davidson JR, Churchill LE, et al. Psychometric properties of the Social Phobia Inventory (SPIN). New self-rating scale. *Br J Psychiatry* 2000;176:379–86.
- Shelby LB. Beyond Cronbach's Alpha: Considering Confirmatory Factor Analysis and Segmentation. *Human Dimensions of Wildlife* 2011;16:142–8.
- Tillfors M, Furmark T. Social phobia in Swedish university students: prevalence, subgroups and avoidant behavior. *Soc Psychiatry Psychiatr Epidemiol* 2007;42:79–86.
- Baptista CA, Loureiro SR, de Lima Osório F, et al. Social phobia in Brazilian university students: prevalence, under-recognition and academic impairment in women. *J Affect Disord* 2012;136:857–61.
- Bella TT, Omigbodun OO. Social phobia in Nigerian university students: prevalence, correlates and co-morbidity. *Soc Psychiatry Psychiatr Epidemiol* 2009;44:458–63.
- Djidonou A, Tchégnoni FT, Adoukonou T, et al. Associated Factors and Impacts of Social Phobia on Academic Performance among Students from the University of Parakou (UP). *OJPsych* 2016;06:151–7.
- Sahu P. Closure of Universities Due to Coronavirus Disease 2019 (COVID-19): Impact on Education and Mental Health of Students and Academic Staff. *Cureus* 2020;12:e7541.
- Pramukti I, Strong C, Sitthimongkol Y, et al. Anxiety and Suicidal Thoughts During the COVID-19 Pandemic: Cross-Country

- Comparative Study Among Indonesian, Taiwanese, and Thai University Students. *J Med Internet Res* 2020;22:e24487.
- 49 Singh R, Joshi HL. Social Phobia Among College-Going Students Following the Covid-19 Outbreak: Links with Internet Addiction. *Indian J Clin Psychol* 2022;49:01. Available: https://www.researchgate.net/publication/365322156_Social_Phobia_Among_College-Going_Students_Following_the_Covid-19_Outbreak_Links_with_Internet_Addiction#fullTextFileContent
 - 50 McLeish AC, Walker KL, Hart JL. Changes in Internalizing Symptoms and Anxiety Sensitivity Among College Students During the COVID-19 Pandemic. *J Psychopathol Behav Assess* 2022;44:1021–8.
 - 51 Khatatbeh M. Efficacy of Nationwide Curfew to Encounter Spread of COVID-19: A Case From Jordan. *Front Public Health* 2020;8:394.
 - 52 Alqutob R, Al Nsour M, Tarawneh MR, et al. COVID-19 Crisis in Jordan: Response, Scenarios, Strategies, and Recommendations. *JMIR Public Health Surveill* 2020;6:e19332.
 - 53 Wang X, Hegde S, Son C, et al. Investigating Mental Health of US College Students During the COVID-19 Pandemic: Cross-Sectional Survey Study. *J Med Internet Res* 2020;22:e22817.
 - 54 Ma Z, Zhao J, Li Y, et al. Mental health problems and correlates among 746 217 college students during the coronavirus disease 2019 outbreak in China. *Epidemiol Psychiatr Sci* 2020;29:e181.
 - 55 Faris Alsaraireh HA, Leimoon H, Al-Mrayat Y, et al. The Psychological Effects of Social Phobia on Undergraduate Students in the south of Jordan. *JPTCP* 2023;30:367–77.
 - 56 Al-Johani WM, AlShamlan NA, AlAmer NA, et al. Social anxiety disorder and its associated factors: a cross-sectional study among medical students, Saudi Arabia. *BMC Psychiatry* 2022;22:505.
 - 57 Afshari N, Farokhzadian J, Abdi K, et al. Gender Matter, Social Phobia and High-Risk Behaviors in Young Medical Students. *Ethiop J Health Sci* 2021;31:359–70.
 - 58 Uzonwanne FC. Prevalence of social phobia, gender and school type among young adults in Nigerian universities. *Int J Soc Sci Humanit Res* 2014;2:36–45. Available: <https://www.questjournals.org/jrhss/papers/vol2-issue8/E283645.pdf>
 - 59 Norhizan NFA, Ghazi HF, Abdalrazak HA, et al. Social phobia and its association with body shape and internet addiction among private university students in Selangor, Malaysia. *Internati Journ of Medic Toxicol & Lega Medici* 2019;22:106.
 - 60 Desalegn GT, Getinet W, Tadie G. The prevalence and correlates of social phobia among undergraduate health science students in Gondar, Gondar Ethiopia. *BMC Res Notes* 2019;12:438.
 - 61 Nelson EC, Grant JD, Bucholz KK, et al. Social phobia in a population-based female adolescent twin sample: co-morbidity and associated suicide-related symptoms. *Psychol Med* 2000;30:797–804.
 - 62 Obadeji A, Kumolalo BF. Social anxiety disorder among undergraduate students: exploring association with self-esteem and personality traits. *World Social Psychiatry* 2022;4:24–30.
 - 63 Al-Hazmi BH, Sabur SS, Al-Hazmi RH. Social anxiety disorder in medical students at Taibah University, Saudi Arabia. *J Family Med Prim Care* 2020;9:4329–32.
 - 64 Alotaibi FS, Alsaeedi A. Attitudes of medical students toward communication skills learning in Western Saudi Arabia. *Saudi Med J* 2016;37:791–5.
 - 65 Davis H, Nicholaou T. A comparison of the interviewing skills of first- and final-year medical students. *Med Educ* 1992;26:441–7.
 - 66 Sadock BJ. *Kaplan & Sadock's Synopsis of Psychiatry: Behavioral Sciences/Clinical Psychiatry*, Vol 2015. Philadelphia, PA: Wolters Kluwer, 2015.
 - 67 Yen C-F, Ko C-H, Wu Y-Y, et al. Normative Data on Anxiety Symptoms on the Multidimensional Anxiety Scale for Children in Taiwanese Children and Adolescents: Differences in Sex, Age, and Residence and Comparison with an American Sample. *Child Psychiatry Hum Dev* 2010;41:614–23.
 - 68 Supporting the mental health and well-being of higher education students. 2024. Available: <https://unesdoc.unesco.org/ark:/48223/pf0000391501> [Accessed 16 Dec 2024].