

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<u>http://bmjopen.bmj.com</u>).

If you have any questions on BMJ Open's open peer review process please email <u>info.bmjopen@bmj.com</u>

BMJ Open

Association of Occupational Noise Exposure and Shift Work with Non-alcoholic Fatty Liver Disease among Male Workers: a cross-sectional study.

Journal:	BMJ Open
Manuscript ID	bmjopen-2024-085753
Article Type:	Original research
Date Submitted by the Author:	24-Feb-2024
Complete List of Authors:	Jinwei, ZHANG; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Zhang, Yuxia; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Qiu, Cong; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Wenfeng, ZENG; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Yanmei, RUAN; Guangzhou Twelfth People's Hospital, Guangzhou Medical University, Key Laboratory of Occupational Environment and Health Gao , Yun; Guangzhou Medical University, School of Public Health Ma , Wei; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Wu , Kang; Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Wu , Kang; Twelfth People's Hospital Affiliated to Guangzhou Medical University, School of Public Health Zhang , Jing; Guangdong Pharmaceutical University, School of Public Health Cui, Jiaxin; Sun Yat-Sen University, School of public health Ye, Cui; Guangzhou Red Cross Hospital, Department of Preventive Health Liang, Jiabin; Guangzhou Twelfth People's Hospital Wang, Zhi; Guangzhou Twelfth People's Hospital,
Keywords:	Risk Factors, PUBLIC HEALTH, OCCUPATIONAL & INDUSTRIAL MEDICINE, EPIDEMIOLOGIC STUDIES

SCHOLARONE[™] Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our <u>licence</u>.

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which <u>Creative Commons</u> licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

terez oni

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Association of Occupational Noise Exposure and Shift Work with Non-alcoholic Fatty Liver Disease among Male Workers: a crosssectional study.

Jinwei Zhang¹⁺, Yuxia Zhang¹⁺, Congxi Qiu¹, Wenfeng Zeng¹, Yanmei Ruan¹, Yunxia Gao², Weiyu Ma¹, Kangyong Wu², Jingwen Zhang³, Jiaxin Cui⁴, Cuiping Ye⁵, Jiabin Liang¹, Zhi Wang^{1#}

¹Key Laboratory of Occupational Environment and Health, Guangzhou Twelfth People's Hospital, 1 Tianqiang St., Huangpu West Ave., Guangzhou, Guangdong, China 510620

²School of Public Health, Guangzhou Medical University, Guangzhou, Guangdong, China 511436

³School of Public Health, Guangdong Pharmaceutical University, Guangzhou, Guangdong, China 510080

⁴School of Public Health, Sun Yat-sen University, Guangzhou, Guangdong, China 510310

⁵Department of Preventive Health, Guangzhou Red Cross Hospital, Guangzhou Guangdong, China 510220

[#]Corresponding author: Zhi Wang, PhD, MD, Professor

Tel.: +86-20-38981282, Mobile: +86-13434163611

email: zhi_wang@outlook.com

⁺These authors have contributed equally to this work.

Abstract

Objective This study aimed to determine the relationship between occupational noise, shift work, and nonalcoholic fatty liver disease(NAFLD) in male workers in the automobile manufacturing industry.

Design Cross-sectional study.

Setting This study was carried out Guangzhou Twelfth People's Hospital, using data from April to September 2022.

Participants A total of 4672 eligible participants were included in the study.

Primary and secondary outcome measures Diagnosis of NAFLD was made using ultrasound. Noise was detected according to the Measurement of Physical Factors in the Workplace-Part 8: Noise. Environmental noise intensity was assessed using an EDGE personal noise dosimeter manufactured by CASELLA (UK). The working status of workers were investigated by questionnaire.

Results The OR of NAFLD was 1.39(95%CI: 1.02-1.88) in CNE≥95 group compared to CNE<85 group. Improved risk of NAFLD in workers with shift work compared to those without shift work (OR=1.36, 95%CI: 1.10-1.69). As stratified analyses showed, the ORs of NAFLD prevalence related to occupational noise and shift work exposure appears to be increased in young workers. Combined effects analysis revealed that the ORs of NAFLD was 2.02(95%CI: 1.35-3.01) in CNE≥95 and cumulative length of night shifts work>2920 hours.

Conclusion Occupational noise exposure may be an independent risk factor for NAFLD. And it may synergistically affect disease when combined with night shift work, particularly among younger workers. These findings underscore the importance for companies to prioritize the management and training of younger workers, along with targeted occupational health education initiatives, as crucial measures for reducing the incidence of NAFLD.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study contributes to a relatively uncommon exploration of the association between occupational noise exposure, shift work, and their combined effect on NAFLD among male workers in the automotive manufacturing industry. Based on the study findings, we propose several recommendations for management departments.
- Questionnaire may make it difficult to accurately assess hours worked.
- Due to the cross-sectional survey, causal relationships cannot be elucidated.

1 Introduction

In the last decades, there has been a progressive increase in the incidence of NAFLD, leading to a significant burden of morbidity and mortality associated with liver-related disorders^{1 2}. Anticipating future trends, researchers project a continued rise in the prevalence of NAFLD, while expecting a stable or possibly decreasing prevalence of other chronic liver diseases³. In China, the prevalence of NAFLD has experienced a notable increase in the past decade, rising from 18% to 29.2%. Industrialization and changes in lifestyle have contributed to this rise, leading to a 10% increase in national incidence⁴. Recent studies have highlighted the rapid increase in NAFLD prevalence among adolescents ⁵.

The development of non-alcoholic fatty liver disease (NAFLD) is influenced by various factors, including age, gender, type 2 diabetes, metabolic syndrome, its components, and family history of fatty liver⁶⁻¹⁰. Age and gender also play significant roles in the onset of NAFLD. The increasing prevalence of NAFLD can be attributed to population aging, where age-related visceral fat accumulation may contribute to the secretion of inflammatory factors^{7 8}. Unhealthy dietary habits, including high-fat diets, and unhealthy lifestyles such as late-night activities have also been associated with an increased risk of NAFLD¹¹⁻¹³. Furthermore, occupational factors in the workplace, such as noise exposure and shift work, may influence the risk of NAFLD. Previous research has demonstrated a significant dose-response relationship between

cumulative noise exposure (CNE) and the detection rates of fatty liver, lending support to the notion that noise may act as a risk factor for fatty liver¹⁴. Shift work is a prevalent practice in various industries, including manufacturing, pharmaceutical, and service sectors. It not only directly disrupts normal circadian rhythms, but is also closely linked to sleep disorders, leading to insulin resistance and stress reactions, causing physiological disruptions and harm to health¹⁵⁻¹⁷. These disturbances can disrupt glucose and lipid metabolism, which is a significant contributing factor in elevating the risk of NAFLD¹⁸.

Workers in the automotive manufacturing industry frequently endure prolonged occupational noise exposure while may be required to engage in shift work. Despite this, there is a paucity of investigations specifically examining the impact of shift work and noise on NAFLD within the context of this workforce. Consequently, our study aims to investigate the effects of occupational noise exposure and shift work on NAFLD among male workers in the automotive manufacturing industry. By focusing on this specific population, the study intends to provide a theoretical foundation for the implementation of measures to protect against occupational hazards. Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

2 Materials and methods

2.1 Study design and population

From April to September 2022, a cross-sectional study was conducted among male workers at an automobile manufacturing company in Guangzhou for occupational health examination. Participants with incomplete medical examination and questionnaire data, female workers, individuals taking antiretroviral drugs, and those with a history of cancer, excessive alcohol intake, hepatobiliary disease, blood system diseases, or renal failure were excluded. Ultimately, a total of 4672 eligible participants were included in the study. This study was reviewed and approved by the Ethics Committee of Guangzhou Twelfth People's Hospital (No.2021008) and all procedures involving participants were conducted in accordance with the ethical standards of the Institutional Research Board and the 1964 Declaration of Helsinki, and all study participants gave informed consent.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

2.2 NAFLD diagnosis and collection of information on covariates

The occupational health examination was conducted in accordance with a standardized protocol by trained physicians. Diagnosis of NAFLD was made using the criteria proposed by FARRELL et al¹⁹. Blood was drawn from the fasting subjects to measure hemoglobin(HGB), and fasting plasma glucose(FPG) levels. Blood pressure was measured by the omron HEM-7071 electronic blood pressure monitor three times, each time at an interval of 1 min, and the average of the three measurements was taken as the individual blood pressure(BP) value. BP was classified according to 2018 Chinese Guidelines for Prevention and Treatment of Hypertension. FPG normal range is 3.9-6.1mmol/L. According to the Guidelines for the Prevention and Control of Overweight and Obesity in Chinese Adults, our study the inclusion of BMI<24kg/m² and BMI \geq 24kg/m² as dichotomous variables in the multivariate logistic regression model. Long working hours are defined as working more than 40 hours per week²⁰. High-salt foods mainly include pickled vegetables, ham and bacon, etc²¹²². High-fat foods including chips, chocolate, animal offal and fried foods etc. Late evening snack is defined as an additional meal arranged after 21:00. Smoking is defined as a consistent consumption of at least 1 cigarette per day for a duration of more than 6 months. Alcohol consumption is defined as a regular pattern of drinking at least once a week for a period of 1 year or longer.

2.3 Noise exposure assessment Our study was carried out in accordance with the "Measurement of Physical Factors in the Workplace-Part 8: Noise" (GBZ/T189.8-2007) for noise detection. Environmental noise intensity was assessed using an EDGE personal noise dosimeter manufactured by CASELLA (UK). Each monitoring point was measured three times and the average value was taken as the measurement result. In this study, noise exposure was evaluated using the normalization of continuous A-weighted sound pressure level equivalent to an 8h per day (LAeq, 8h), and work exposed to noise is defined as those operations involving the presence of hearing-impairing, health-hazardous or otherwise hazardous sound, and LAeq,8h \geq 80 dB(A) for at least one year. Meanwhile, Cumulative noise exposure (CNE) was used to quantify the noise exposure level of the study participants and it is calculated as:

BMJ Open

CNE= $L_{Aeq, 8h}$ +10logT, where T indicates years of noise exposure and the CNE unit is: dB(A)·year.

2.4 Assessment of shift work The International Labour Organization defines shift work as the working day is extended to 24 hours by alternating between different workers and/or teams²³ ²⁴. And the night shift work is from 0:00 to 5:00 am²⁴ ²⁵. Calculating cumulative length of night shift work in hours in data processing. In this study, the two main types of shift patterns for workers are two shifts system and three shifts system.

2.5 Statistical analyses Daily data verification by subject staff to ensure unique ID codes for each individual and timely identification of missing key data and logical error. R (version 4.2.1) was used for all statistical analyses. Continuous variables that followed normal distribution are presented as mean ±standard deviation(SD), and were compared by t-test. Mann-Whitney U test was used to compare continuous variables with skewed distribution in the form of median with interquartile ranges. The χ^2 test was used to compare categorical variables, which have been expressed as numbers (percentages). The relationship between shift work, occupational noise and NAFLD was analysed using a multivariate logistic regression model, with night shift hours and CNE included in the regression model as categorical variables respectively as well as the joint effect of both was analysed in relation to NAFLD. Referring to the national criteria (GBZT229.4-2012), CNE is divided into four groups <85, 85~, 90~ and 95~. Cumulative length of night shift work are grouped by quartiles, using the first quartile as a reference. Previous studies have considered adolescents and young adults (AYAs) to be the group aged 15 years and older, with the cut-offs at 24, 29 and 39 years being difficult to determine and some studies considering 18-24 years as late adolescence²⁶. We stratified the age into two groups ≤ 29 and ≥ 30 for analysis. To provide more clarity on the effect of CNE on the illness of workers in different age groups, we further divided the youths at ≤ 29 into two groups for analysis(≤ 24 and 25-29 years old)^{27 28}. All test were two-sided, and α =0.05 was considered statistically significant.

3 Results

3.1 Basic information about the study population. A total of 5928 questionnaires were distributed, out of which 4791 valid questionnaires were returned, resulting in a valid response rate of 80.8%. After excluding individuals who did not meet the study's inclusion criteria, and those with incomplete occupational health screening results (n=119), the final sample size included for analysis was determined. Overall, 4672 workers with complete information on occupational health examination and questionnaires were included in the study according to the inclusion and exclusion criteria. The mean age of the study population was (26.8 ± 5.0), with different prevalence rates for workers in the \leq 29 and \geq 30 years age groups (P<0.001), with a total of 1363 workers with NAFLD and a prevalence of 29.2%. There were 3658 (78.3%) shift workers, with different prevalence of NAFLD in different shift work situations (P<0.001). Meanwhile, 2965(63.5%) noise workers had a statistically significant difference in prevalence in different noise exposure scenarios (P<0.05). Cumulative length of night shift work and CNE vary by disease state (p<0.05, Table1 and Supplementary Table 1).

-	-	-		
Characteristics	NA	FLD	Total (N=4672)	<i>P</i> -value
Characteristics	No (N=3309)	Yes (N=1363)	- Total (IV-4072)	
Importance of noise				
protection equipment, n				0.451
(%)				
Unimportant	63 (1.9)	24 (1.8%)	87 (1.9)	
Unknown	72 (2.2)	20 (1.5%)	92 (2.0)	
General	1718 (51.9)	714 (52.4%)	2432 (52.1)	
Extremely	1456 (44.0)	605 (44.4%)	2061 (44.1)	
Long working hours				< 0.001
No	1375 (41.6)	685 (50.3)	2060 (44.1)	
Yes	1934 (58.4)	678 (49.7)	2612 (55.9)	

Table 1 Occupational characteristics of population in different disease states (n, %)

	NA	NAFLD		D 1
Characteristics	No (N=3309)	Yes (N=1363)	- 1 otal (N=46/2)	<i>P</i> -valu
Dust				0.015
No	2990 (90.4)	1262 (92.6)	4252 (91.0)	
Yes	319 (9.6)	101 (7.4)	420 (9.0)	
Aromatic hydrocarbons				0.020
No	2828 (85.5)	1200 (88.0)	4028 (86.2)	
Yes	481 (14.5)	163 (12.0)	644 (13.8)	
Shift work				< 0.001
No	808 (24.4)	206 (15.1)	1014 (21.7)	
Yes	2501 (75.6)	1157 (84.9)	3658 (78.3)	
Work exposed to noise				0.045
No	1239 (37.4)	468 (34.3)	1707 (36.5)	
Yes	2070 (62.6)	895 (65.7)	2965 (63.5)	
Cumulative length of night				<0.001
shift work (hours)				<0.001
≤730	1321 (39.9)	312 (22.9)	1633 (35.0)	
730~	771 (23.3)	240 (17.6)	1011 (21.6)	
1460~	678 (20.5)	290 (21.3)	968 (20.7)	
2920~	539 (16.3)	521 (38.2)	1060 (22.7)	
CNE [dB(A)·year]				< 0.001
<85	1277 (38.6)	330 (24.2)	1607 (34.4)	
85~	992 (30.0)	364 (26.7)	1356 (29.0)	
90~	771 (23.3)	439 (32.2)	1210 (25.9)	
95~	269 (8.1)	230 (16.9)	499 (10.7)	
Seniority in shift work		6.0(3.0,		~0.001
(year), median (IQR)	3.0(1.5, 6.5)	12.0)	4.0(2.0, 8.0)	<0.001
Seniority in work exposed		7 00/2 9 12		
to noise(year), median	3.0(1.3, 7.0)	/.90(2.8,13.	3.8 (1.5, 8.3)	< 0.001
(IQR)		5)		

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

3.2 The relationship between occupational noise exposure and NAFLD. The impact of work exposed to noise on NAFLD was demonstrated in Table 2. Compared with those did not, no statistically significant relationship between work exposed to noise and NAFLD (OR=0.92, 95%CI: 0.78-1.09). Supplementary Table 2 presented the odds ratios (ORs) and 95% confidence intervals (CIs) for the effect of occupational noise exposure on NAFLD. By reference to CNE<85, the OR of NAFLD was 1.39(95%CI: 1.02-1.88) in CNE \geq 95 group. Stratified analysis indicated that the association between CNE and NAFLD was not statistically significant in both those aged \leq 29 and \geq 30 years; Table 3 shows further stratification by age, among those \leq 24 years of age, the OR of NAFLD in the CNE \geq 95 group was 2.95 (95%CI: 1.17-6.91).

1000 2 1	issociation of work e	nposed ic		
work exposed to noise	Model 1		Model 2	
(Yes or No)	OR 95%CI	Р	OR 95%CI	Р
Total	0			
No	Ref			
Yes	0.97(0.83, 1.15)	0.732	0.92(0.78, 1.09)	0.333
Age Group, y				
≤29				
No	Ref			
Yes	1.07(0.87, 1.31)	0.529	0.98(0.79, 1.21)	0.828
<u>≥</u> 30				
No	Ref			
Yes	0.83(0.62, 1.10)	0.196	0.83(0.62, 1.10)	0.197

Table 2 Association of work exposed to noise with NAFLD

Model 1: Adjust for smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB; Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, aromatic hydrocarbons, shift work.

BMJ Open

	1		6.6	1
CNF $[dB(\Lambda))$ vert	Model 1	l	Model	2
	OR 95% <i>CI</i>	Р	OR 95%CI	Р
Age Group, y				
≤24				
<85	Ref			
85~	1.07(0.77, 1.48)	0.683	0.96(0.68, 1.34)	0.792
90~	1.00(0.64, 1.53)	0.997	0.87(0.54, 1.35)	0.532
95~	3.17(1.28, 7.28)	0.009	2.95(1.17, 6.91)	0.016
25~29				
<85	Ref			
85~	1.14(0.80, 1.62)	0.474	1.14(0.78, 1.66)	0.497
90~	1.11(0.75, 1.61)	0.606	1.08(0.73, 1.62)	0.692
95~	1.02(0.55, 1.81)	0.957	1.07(0.58, 1.96)	0.819

Table 3 The relationship between CNE and NAFLD in different age groups

Model 1: Adjust for smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB; Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, aromatic hydrocarbons, shift work.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

3.3 The relationship between shift work and NAFLD. The effect of shift work on NAFLD is shown in the Fig 1. Compared to those without shift work, the OR of NAFLD among workers with shift work was 1.36(95%CI: 1.10-1.69). Stratified analysis showed that shift work was a risk factor for NAFLD among those aged ≤ 29 years (OR=1.46, 95%CI: 1.15-1.87). Similarly, in comparison with cumulative length of night shift work ≤ 730 h, rising cumulative length of night shift work were associated with a higher risk of NAFLD, with ORs(95%CIs) of 1.44(1.15-1.81), 1.45(1.15-1.83) and 1.65(1.24, 2.18), respectively. Among those aged ≤ 29 years, increased cumulative length of night shift work were also a risk factor for NAFLD, with ORs(95%CIs) of 1.66(1.30, 2.12), 1.53(1.17, 1.99) and 2.43(1.50, 3.86), respectively.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

3.4 The combined effect of occupational noise and cumulative length of night shift work in association with NAFLD. The combined effect of occupational noise and cumulative length of night work on NAFLD was shown in Tables 4. After adjusting for variables, the association with NAFLD was statistically significant for all groups at cumulative length of night shift work >2920h and CNE≥85, when compared with CNE<85 and cumulative length of night shift work ≤730h. The OR of NAFLD was greatest at 2.02 (95%CI: 1.35, 3.01) when cumulative length of night shift work >2920h and CNE≥95.

Table 4 Association of CNE and cumulative length of night shift work with NAFLD

CNE	Cu	nulative length of n	night shift work (ho	urs)
$[dB(A) \cdot years]$	≤730	730~	1460~	2920~
<85	Ref	1.51(1.05, 2.14)	1.28(0.82, 1.96)	1.37(0.65, 2.84)
85~	0.87(0.60, 1.26)	1.45(1.05, 1.99)	1.61(1.14, 2.27)	2.00(1.30, 3.09)
90~	1.25(0.77, 1.99)	1.33(0.86, 2.01)	1.42(1.01, 1.99)	1.54(1.08, 2.21)
95~	1.96(0.88, 4.23)	1.60(0.71, 3.47)	1.68(0.96, 2.89)	2.02(1.35, 3.01)

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, aromatic hydrocarbons.

4 Discussion

To summarize, our study included a total of 4672 male workers in the automotive manufacturing industry to investigate the impact of occupational noise and shift work on non-alcoholic fatty liver disease (NAFLD) through a cross-sectional survey. The majority of the company's employees are subjected to a high workload, with 63.5% working in noise and 78.3% working in shifts, which is significantly higher compared to other occupations such as teachers and white-collar workers²⁹. Night work has been categorized by the International Agency for Research on Cancer as "possibly carcinogenic to humans" (Group 2A)³⁰, prompting growing concerns about its adverse effects on health. The prevalence of NAFLD in this survey was 29.2%, a rate close to the combined prevalence estimate of NAFLD reported by Fan

et al^{11 31-33}. Among those who work exposed to noise and those who did not, the prevalence of NAFLD is 30.2% and 27.4%, respectively. After adjusting for relevant variables, multifactorial logistic regression analysis revealed that increased CNE was the significant risk factor associated with NAFLD. This may be because the biological effects of noise are cumulative and the effects of a single noise exposure are not sufficient to cause health damage to the organism. However, the more repeated exposures and the longer the exposure period, the more pronounced the cumulative effects will be when the effects are not eliminated^{34 35}. Associations between CNE and NAFLD vary across age groups. The increase in CNE is a risk factor for NAFLD among workers aged <24 years. This suggests that short-term exposure to highintensity noise may contribute to the development of NAFLD in this population. Possible factors that may contribute to this association include the inadequate use of noise protective equipment among young workers or exposure to noise outside of the occupational setting. The effect of noise on NAFLD in younger workers should be further explored, indicating that enterprises should strengthen training and occupational health education for young workers, which is important to reduce the occurrence of NAFLD. Shins, Saeha et al. have shown that chronic exposure to road traffic noise was associated with increased incidence of diabetes and hypertension in Toronto³⁶. Du et al. found that noise exposure could lead to disorders of glucolipid metabolism in mice through inhibition of the AKT pathway mediated by the biological clock gene BMAL1³⁷. The experiment by Evans et al. also demonstrated significant changes in serum ALT levels across test groups, suggesting that noise can lead to liver dysfunction³⁸. It can be seen that noise may be the basis for the formation of NAFLD by promoting insulin resistance and disorders of glucolipid metabolism in the body, resulting in metabolic abnormalities in the body, and these studies also indirectly support our findings. In this study, the prevalence of NAFLD was 20.3% and 31.6% among auto manufacturing workers not involved in shift work and those who did, respectively, with a higher prevalence among workers in shift work, similar to the findings of Golabi et al¹¹. We observed that shift work and an increase in cumulative length of night shift work were all identified as risk factors for NAFLD.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

We also examined the relationship between shift work and NAFLD across different age groups. Specifically, among workers below 29 years of age, engaging in shift work was found to significantly increase the risk of NAFLD. This indicates the presence of specific physiological adaptations that contribute to this heightened risk. Similarly, we found that an extended cumulative length of night shift work was a significant risk factor for NAFLD in workers below 29 years of age. These findings emphasize the importance of appropriate management strategies to mitigate the adverse effects of shift work on NAFLD among younger workers. Previous epidemiological studies have consistently demonstrated a link between shift work and NAFLD. For instance, a 4-year cohort study conducted among Chinese railway workers reported a higher incidence of NAFLD in individuals engaged in shift work¹¹. Similarly, a cross-sectional survey involving male steelworkers found that night shift workers had an increased risk of NAFLD compared to those working day shifts¹⁵. Wang et al. further supported the association between night shifts and NAFLD by demonstrating the impact of disrupted circadian rhythms, due to delayed sleep, on the incidence of NAFLD³⁹. These findings are in line with our results. Sun et al. also highlighted the association between sleep deprivation, increased cortisol levels, and insulin resistance, further highlighting the possible mechanisms underlying the relationship between sleep disruption and NAFLD⁴⁰. Our study identified a significant association between occupational noise exposure, shift work, and an increased risk of developing NAFLD among workers. This association may be explained by the potential mechanisms through which both noise exposure and shift work contribute to the development of insulin resistance, a key factor in the pathogenesis of NAFLD. Insulin resistance disrupts lipid metabolism, leading to elevated lipase activity and subsequent overproduction of triglycerides and free fatty acids. Consequently, an excessive amount of free fatty acids enters the liver, promoting increased lipid synthesis and deposition within hepatic cells. This lipid accumulation interferes with hepatocyte transport capacity, resulting in hepatocyte apoptosis and the accumulation of fat, ultimately leading to NAFLD development.

This study contributes to a relatively uncommon and innovative exploration of

the association between occupational noise exposure, shift work, and their combined effect on NAFLD among male workers in the automotive manufacturing industry. Based on the study findings, we propose several recommendations for relevant management departments. It is advised to scientifically and reasonably organize the working system, promote the implementation of flexible working arrangements, and minimize the cumulative length of night work. Furthermore, improving the production process to control or eliminate noise sources, such as installing sound insulation and vibration dampers, is recommended. Additionally, establishing and enhancing occupational health supervision, strengthening the management of occupational health, and facilitating regular occupational health check-ups for workers are crucial measures to safeguard worker health. These recommendations hold significant practical importance in protecting the health of workers.

However, it is worth to acknowledge the limitations of our study. Due to the cross-sectional nature of the survey, we were unable to establish a causal relationship between occupational noise exposure, shift work, and the development of illness. To address these limitations, we are in the process of establishing a cohort study among workers in the automotive manufacturing industry. In future studies, we plan to utilize individual noise dosimeters to more accurately measure workers' noise exposure, enabling a more robust analysis of the association between these occupational factors and disease outcomes.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

Acknowledgements We wish to thank the timely help given by workers who participated in this program. we are deeply grateful for Guangzhou Twelfth People's Hospital for their support for collecting data.

Fundings This work was supported by the Key Research and Development Programme of Guangzhou Science and Technology Project (Project Number: 202206010061); Guangzhou Key Medical Discipline(2021-2023); The Science and Technology Program of Guangzhou (Project Number: 2023A03J0502, 202201010816, 2023A03J0979); The Medical Science and Technology Research Fund Project of Guangdong (Project Number: B2023347,

A2023406, A2023396); The Science and Technology Project of Guangzhou Municipal Health Commission (Project Number: 20221A011060, 20211A010041, 20211A010039).

Ethical statements

Patient consent for publication

Consent obtained directly from patient(s).

Ethics approval

The study was approved by Guangzhou Twelfth People's Hospital ethics committee (No.2021008), and all procedures involving participants were conducted in accordance with the ethical standards of the Institutional Research Committee and the 1964 Helsinki Declaration. Participants gave informed consent to participate in the study before taking part.

Data availability statement

Data are available upon reasonable request.

Conflicts of interest

The authors declare that they have no competing interests

Patient and Public Involvement

The study was approved by Guangzhou Twelfth Peoples Hospital ethics committee. The committee believed that the design and plan of the study fully takes into account the principles of safety and fairness, and its research content did not cause any harm to the participants. The recruitment of participants was based on the principle of voluntary and informed consent. The project team protected the rights and privacy of participants in accordance with relevant national regulations, and there was no conflict of interest between the research content and the research results.

4 5

6 7

8

9

10 11

12

13

14

15

16

17

18 19

20

21

22

23 24

25

26

27 28

29

30

31 32

33

34

35

36 37

38

39

40 41

42

43

44

45 46

47

48

49 50

51

52

53 54

55

56

References

- 1. Notarnicola M, Osella AR, Caruso MG, et al. Nonalcoholic Fatty Liver Disease: Focus on New Biomarkers and Lifestyle Interventions. *Int J Mol Sci* 2021;22(8) doi: 10.3390/ijms22083899.
- Masoodi M, Gastaldelli A, Hyotylainen T, et al. Metabolomics and lipidomics in NAFLD: biomarkers and non-invasive diagnostic tests. *Nat Rev Gastroenterol Hepatol* 2021;18(12):835-56. doi: 10.1038/s41575-021-00502-9.
- Younossi ZM, Stepanova M, Afendy M, et al. The changing face of chronic liver disease (CLD) in the United States: The rising epidemic of non-alcoholic fatty liver disease (NAFLD). J Hepatol 2011;54. doi: 10.1016/S0168-8278(11)60018-4.
- 4. Younossi Z, Tacke F, Arrese M, et al. Global Perspectives on Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. *Hepatology* 2019;69(6):2672-82. doi: 10.1002/hep.30251.
- Lin G, Xinhe Z, Haoyu T, et al. Epidemiology and lifestyle survey of non-alcoholic fatty liver disease in school-age children and adolescents in Shenyang, Liaoning. *BMC Pediatr* 2022;22(1):286. doi: 10.1186/s12887-022-03351-w.
- Younossi ZM, Golabi P, de Avila L, et al. The global epidemiology of NAFLD and NASH in patients with type 2 diabetes: A systematic review and meta-analysis. J Hepatol 2019;71(4):793-801. doi: 10.1016/j.jhep.2019.06.021.
- Kim H, Kisseleva T, Brenner DA. Aging and liver disease. *Current opinion in gastroenterology* 2015;31(3):184-91. doi: 10.1097/MOG.00000000000176.
- Barzilai N, Huffman DM, Muzumdar RH, et al. The critical role of metabolic pathways in aging. Diabetes 2012;61(6):1315-22. doi: 10.2337/db11-1300.
- Wong VW, Chu WC, Wong GL, et al. Prevalence of non-alcoholic fatty liver disease and advanced fibrosis in Hong Kong Chinese: a population study using proton-magnetic resonance spectroscopy and transient elastography. *Gut* 2012;61(3):409-15. doi: 10.1136/gutjnl-2011-300342.
- Wang J, Tang Y, Peng K, et al. Development and validation of a nomogram for predicting nonalcoholic fatty liver disease in the non-obese Chinese population. Am J Transl Res 2020;12(10):6149-59.
- Xu J, Ni S, Wang Y, et al. Shift work and nonalcoholic fatty liver disease incidence among Chinese rail workers: a 4-year longitudinal cohort study. *Int Arch Occup Environ Health* 2023;96(1):179-90. doi: 10.1007/s00420-022-01913-0.
- Takahashi A, Anzai Y, Kuroda M, et al. Effects of sleep quality on non-alcoholic fatty liver disease: a cross-sectional survey. *BMJ Open* 2020;10(10):e039947. doi: 10.1136/bmjopen-2020-039947.
- 13. Semmler G, Datz C, Reiberger T, et al. Diet and exercise in NAFLD/NASH: Beyond the obvious. *Liver Int* 2021;41(10):2249-68. doi: 10.1111/liv.15024.
- 14. Guangping B, Huilin Y, Quanzhi Y. Relationship Between Occupational Noise Expose and the Fatty Liver. *HEBEI MEDICINE* 2009;15(01):59-62.
- Zhang S, Wang Y, Wang Z, et al. Rotating night shift work and non-alcoholic fatty liver disease among steelworkers in China: a cross-sectional survey. *Occup Environ Med* 2020;77(5):333-39. doi: 10.1136/oemed-2019-106220.

16. Attarchi M, Dehghan F, Safakhah F, et al. Effect of exposure to occupational noise and shift working on blood pressure in rubber manufacturing company workers. *Ind Health* 2012;50(3):205-13. doi: 10.2486/indhealth.ms1321.

1 2 3

4

5 6

7

8

9

10 11

12

13

14

15

16

17

18 19

20

21

22

23 24

25

26

27 28

29

30

31

32 33

34

35

36 37

38

39

40 41

42

43

44

45 46

47

48

49 50

51

52

53 54

55

56

57

58 59 60

- 17. Dong H, Zhang Q, Zhang Z, et al. Association of sleep disturbance with shift work, occupational stress, and musculoskeletal pain in Chinese public hospital emergency workers: A multicentre cross-sectional study. *Chronobiol Int* 2022;39(6):886-94. doi: 10.1080/07420528.2022.2048662.
- Boivin DB, Boudreau P, Kosmadopoulos A. Disturbance of the Circadian System in Shift Work and Its Health Impact. *J Biol Rhythms* 2022;37(1):3-28. doi: 10.1177/07487304211064218.
- Farrell GC, Chitturi S, Lau GK, et al. Guidelines for the assessment and management of nonalcoholic fatty liver disease in the Asia–Pacific region: Executive summary. J Gastroenterol Hepatol 2007;22(6):775-77. doi: 10.1111/j.1440-1746.2007.05002.x.
- Pega F, Náfrádi B, Momen NC, et al. Global, regional, and national burdens of ischemic heart disease and stroke attributable to exposure to long working hours for 194 countries, 2000-2016: A systematic analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ Int* 2021;154:106595. doi: 10.1016/j.envint.2021.106595.
- Webster JL, Dunford EK, Neal BC. A systematic survey of the sodium contents of processed foods. *Am J Clin Nutr* 2010;91(2):413-20. doi: 10.3945/ajcn.2009.28688.
- 22. Ni Mhurchu C, Capelin C, Dunford EK, et al. Sodium content of processed foods in the United Kingdom: analysis of 44,000 foods purchased by 21,000 households. *Am J Clin Nutr* 2011;93(3):594-600. doi: 10.3945/ajcn.110.004481.
- 23. International Labour Organization. C171-Night Work Convention, 1990(No.171) [Available from: <u>https://www.ilo.org/dyn/normlex/en/f?p=NORMLEXPUB:12100:0::NO::P12100_INSTRUM</u> <u>ENT_ID:312316</u>.
- 24. World Health Organization. IARC Monographs Volume 124: Night Shift Work [Available from: https://www.iarc.who.int/news-events/iarc-monographs-volume-124-night-shift-work/.
- 25. Molzof HE, Peterson CM, Thomas SJ, et al. Nightshift Work and Nighttime Eating Are Associated With Higher Insulin and Leptin Levels in Hospital Nurses. *Front Endocrinol* 2022;13:876752. doi: 10.3389/fendo.2022.876752.
- 26. Arshad T, Paik JM, Biswas R, et al. Nonalcoholic Fatty Liver Disease Prevalence Trends Among Adolescents and Young Adults in the United States, 2007-2016. *Hepatol Commun* 2021;5(10):1676-88. doi: 10.1002/hep4.1760.
- Paik JM, Kabbara K, Eberly KE, et al. Global burden of NAFLD and chronic liver disease among adolescents and young adults. *Hepatology* 2022;75(5):1204-17. doi: 10.1002/hep.32228.
- Zhang X, Wu M, Liu Z, et al. Increasing prevalence of NAFLD/NASH among children, adolescents and young adults from 1990 to 2017: a population-based observational study. *BMJ Open* 2021;11(5):e042843. doi: 10.1136/bmjopen-2020-042843.
- 29. Wang D, Chen H, Chen D, et al. Shift work disorder and related influential factors among shift workers in China. *Sleep Medicine* 2021;81:451-56. doi: 10.1016/j.sleep.2021.03.008.
- 30. Cancer IAfRo. IARC Monographs on the Identification of Carcinogenic Hazards to Humans. Night Shift Work. Lyon (FR): International Agency for Research on Cancer 2020.
- Li H, Wang YJ, Tan K, et al. Prevalence and risk factors of fatty liver disease in Chengdu, Southwest China. *Hepatobiliary Pancreat Dis Int* 2009;8(4):377-82.

BMJ Open

- 32. Golabi P, Paik JM, AlQahtani S, et al. Burden of non-alcoholic fatty liver disease in Asia, the Middle East and North Africa: Data from Global Burden of Disease 2009-2019. J Hepatol 2021;75(4):795-809. doi: 10.1016/j.jhep.2021.05.022.
- Younossi ZM. Non-alcoholic fatty liver disease-A global public health perspective. J Hepatol 2019;70(3):531-44. doi: 10.1016/j.jhep.2018.10.033.
- 34. Ising H, Kruppa B. Health effects caused by noise: evidence in the literature from the past 25 years. *Noise Health* 2004;6(22):5-13.
- 35. Cui B, Li K, Gai Z, et al. Chronic noise exposure acts cumulatively to exacerbate Alzheimer's disease-like amyloid-β pathology and neuroinflammation in the rat hippocampus. *Sci Rep-Uk* 2015;5(1):12943. doi: 10.1038/srep12943.
- 36. Shin S, Bai L, Oiamo TH, et al. Association Between Road Traffic Noise and Incidence of Diabetes Mellitus and Hypertension in Toronto, Canada: A Population-Based Cohort Study. J Am Heart Assoc 2020;9(6):e013021. doi: 10.1161/jaha.119.013021.
- Wen-jing D, Zhi-dan W, Xiang-bin Z, et al. Effects and mechanism of short-term day and night noise exposure on glucose and lipid metabolism in mice. *Modern Preventive Medicine* 2022;49(20):3678-84. doi: 10.20043/j.cnki.MPM.202205073.
- Evans GW, Lercher P, Meis M, et al. Community noise exposure and stress in children. J Acoust Soc Am 2001;109(3):1023-7. doi: 10.1121/1.1340642.
- 39. Wang H, Gu Y, Zheng L, et al. Association between bedtime and the prevalence of newly diagnosed non-alcoholic fatty liver disease in adults. *Liver Int* 2018;38(12):2277-86. doi: 10.1111/liv.13896.
- 40. Sun M, Feng W, Wang F, et al. Meta-analysis on shift work and risks of specific obesity types. *Obes Rev* 2018;19(1):28-40. doi: 10.1111/obr.12621.

iezoni

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.



Fig 1. The relationship between shift work and NAFLD. Adjust for smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, Sleep duration, Physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, aromatic hydrocarbons, work exposed to noise.

Supplementary

 Table 1
 Basic characteristics of population in different disease states (n, %)

Characteristics	NA	FLD	Total	ת
Characteristics	No (N=3309)	Yes (N=1363)	- (N=4672)	P
Age (years), n (%)				< 0.001
≤29	2649 (80.1)	698 (51.2)	3347 (71.6)	
≥30	660 (19.9)	665 (48.8)	1325 (28.4)	
Education, n (%)				0.009
Senior secondary and below	1888 (57.1)	720 (52.8)	2608 (55.8)	
Higher education	1421 (42.9)	643 (47.2)	2064 (44.2)	
Smoking status (cigarettes per day), n (%)				< 0.00
Nonsmokers	1741 (52.6)	729 (53.5)	2470 (52.9)	
1~5	618 (18.7)	198 (14.5)	816 (17.5)	
6~10	599 (18.1)	239 (17.5)	838 (17.9)	
>10	351 (10.6)	197 (14.5)	548 (11.7)	
Drinking status (times/week), n (%)				0.382
Nondrinkers	2124 (64.2)	838 (61.5)	2962 (63.4)	
1~2	1091 (33.0)	484 (35.5)	1575 (33.7)	
3~4	65 (2.0)	28 (2.1)	93 (2.0)	
>4	29 (0.9)	13 (1.0)	42 (0.9)	
Fruit and vegetable (times/week), n (%)				0.002
Seldom	262 (7.9)	89 (6.5)	351 (7.5)	
1~2	1961 (59.3)	752 (55.2)	2713 (58.1)	
3~4	653 (19.7)	297 (21.8)	950 (20.3)	
>4	433 (13.1)	225 (16.5)	658 (14.1)	

There staristics	NAFLD		Total	D
Lnaracteristics	No (N=3309)	Yes (N=1363)	(N=4672)	P
High-salt foods				0.125
(times/week), n (%)				0.155
Seldom	1367 (41.3)	540 (39.6)	1907 (40.8)	
1~2	1749 (52.9)	724 (53.1)	2473 (52.9)	
3~4	144 (4.4)	67 (4.9)	211 (4.5)	
>4	49 (1.5)	32 (2.3)	81 (1.7)	
High-fat foods				0 121
(times/week), n (%)				0.121
Seldom	984 (29.7)	367 (26.9)	1351 (28.9)	
1~2	2138 (64.6)	901 (66.1)	3039 (65.0)	
3~4	152 (4.6)	76 (5.6)	228 (4.9)	
>4	35 (1.1)	19 (1.4)	54 (1.2)	
Late evening snack (times/week), n (%)				0.024
Seldom	747 (22.6)	361 (26.5)	1108 (23.7)	
1~2	1863 (56.3)	735 (53.9)	2598 (55.6)	
3~4	386 (11.7)	137 (10.1)	523 (11.2)	
>4	313 (9.5)	130 (9.5)	443 (9.5)	
BMI (kg/m2), n (%)				< 0.001
<24	3001 (90.7)	555 (40.7)	3556 (76.1)	
≥24	308 (9.3)	808 (59.3)	1116 (23.9)	
Sleep duration (hours),				
n (%)				< 0.001
<6	592 (17.9)	303 (22.2)	895 (19.2)	
6~8	2574 (77.8)	1023 (75.1)	3597 (77.0)	
>8	143 (4.3)	37 (2.7)	180 (3.9)	

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Page 23	of 23
---------	-------

$\begin{array}{l} \text{P} \text{Yes (N=1363)} \\ \text{O} 359 (26.3) \\ \text{O} 915 (67.1) \end{array}$) (N=4672) 1230 (26.3)	0.441
) 359 (26.3) 915 (67.1)	1230 (26.3)	0.441
) 359 (26.3) 915 (67.1)	1230 (26.3)	
915 (67.1)	` '	
, , , , , , , , , , , , , , , , , , , ,	3102 (66.4)	
) 89 (6.5)	340 (7.3)	
		< 0.00
306 (22.5)	1680 (36.0)	
963 (70.7)	2853 (61.1)	
94 (6.9)	139 (3.0)	
		< 0.00
824 (60.5)	2216 (47.4)	
539 (39.5)	2456 (52.6)	
		< 0.00
1001 (73.4)	3888 (83.2)	
362 (26.6)	784 (16.8)	
		0.005
) 1013 (74.3)	3599 (77.0)	
) 350 (25.7)	1073 (23.0)	
2/	Þ	
	 306 (22.5) 963 (70.7) 94 (6.9) 824 (60.5) 539 (39.5) 1001 (73.4) 362 (26.6) 1013 (74.3) 350 (25.7) 	(5) $306 (22.5)$ $1680 (36.0)$ (2) $963 (70.7)$ $2853 (61.1)$ (4) $94 (6.9)$ $139 (3.0)$ (2) $824 (60.5)$ $2216 (47.4)$ (2) $1001 (73.4)$ $3888 (83.2)$ (3) $362 (26.6)$ $784 (16.8)$ (2) $1013 (74.3)$ $3599 (77.0)$ (3) $350 (25.7)$ $1073 (23.0)$

- 58 59
- 60

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

]	Table2 Association of C	CNE with N	IAFLD	
	Model 1	Model 1		
CNE[dB(A)·year]	OR 95%CI	Р	OR 95%CI	Р
Total				
<85	Ref			
85~	1.20(0.97, 1.47)	0.091	1.11(0.89, 1.39)	0.337
90~	1.16(0.93, 1.45)	0.190	1.08(0.85, 1.36)	0.543
95~	1.48(1.10, 1.98)	0.009	1.39(1.02, 1.88)	0.034
Age Group, y				
≤29				
<85	Ref			
85~	1.21(0.96, 1.53)	0.101	1.10(0.86, 1.41)	0.441
90~	1.19(0.90, 1.55)	0.215	1.05(0.79, 1.40)	0.730
95~	1.68(1.04, 2.66)	0.029	1.52(0.93, 2.44)	0.088
≥30				
<85	Ref			
85~	1.03(0.57, 1.88)	0.914	1.02(0.56, 1.86)	0.954
90~	0.99(0.57, 1.75)	0.977	0.96(0.55, 1.71)	0.895
95~	1.25(0.70, 2.25)	0.458	1.23(0.68, 2.23)	0.490

Model 1: Adjust for smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, Sleep duration, Physical activity, BP, ECG, FPG, HGB; Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, aromatic hydrocarbons, shift work.

Association of Occupational Noise Exposure and Shift Work with Non-alcoholic Fatty Liver Disease: A cross-sectional study of male workers in Chinese automobile manufacturing industry.

Journal:	BMJ Open		
Manuscript ID	bmjopen-2024-085753.R1		
Article Type:	Original research		
Date Submitted by the Author:	07-Jan-2025		
Complete List of Authors:	Jinwei, ZHANG; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Guangzhou Zhang, Yuxia; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Qiu, Cong; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Wenfeng, ZENG; Guangzhou Twelfth People's Hospital, Key Laboratory Occupational Environment and Health Yanmei, RUAN; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Gao , Yun; Guangzhou Medical University, School of Public Health Ma , Wei; Guangzhou Medical University, School of Public Health Mu , Kang; Guangzhou Medical University, School of Public Health Zhang , Jing; Guangzhou Medical University, School of Public Health Cui, Jiaxin; Sun Yat-Sen University, School of Public Health Ye, Cui; Guangzhou Red Cross Hospital, Department of Preventive Hea Liang, Jiabin; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Ye, Cui; Guangzhou Red Cross Hospital, Department of Preventive Hea Liang, Jiabin; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Ye, Cui; Guangzhou Red Cross Hospital, Department of Preventive Hea Liang, Jiabin; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health		
Primary Subject Heading :	Occupational and environmental medicine		
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine		
Keywords:	Risk Factors, PUBLIC HEALTH, OCCUPATIONAL & INDUSTRIAL MEDICINE, EPIDEMIOLOGIC STUDIES, Cross-Sectional Studies		

SCHOLARONE[™] Manuscripts

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our <u>licence</u>.

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which <u>Creative Commons</u> licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

terez onz

BMJ Open

Association of Occupational Noise Exposure and Shift Work with Non-alcoholic Fatty Liver Disease: A cross-sectional study of male workers in Chinese automobile manufacturing industry.

Jinwei Zhang¹⁺, Zhang Yuxia¹⁺, Qiu Congxi¹, Wenfeng Zeng¹, Yanmei Ruan¹, Gao Yunxia², Ma Weiyu¹, Wu Kangyong², Zhang Jingwen³, Cui Jiaxin⁴, Ye Cuiping⁵, Liang Jiabin¹, Wang Zhi^{1#}

¹Key Laboratory of Occupational Environment and Health, Guangzhou Twelfth People's Hospital, 1 Tianqiang St., Huangpu West Ave., Guangzhou, Guangdong, China 510620

²School of Public Health, Guangzhou Medical University, Guangzhou, Guangdong, China 511436

³School of Public Health, Guangdong Pharmaceutical University, Guangzhou, Guangdong, China 510080

⁴School of Public Health, Sun Yat-sen University, Guangzhou, Guangdong, China 510310

⁵Department of Preventive Health, Guangzhou Red Cross Hospital, Guangzhou Guangdong, China 510220

[#]Corresponding author: Zhi Wang, PhD, MD, Professor

Tel.: +86-20-38981282, Mobile: +86-13434163611

email: zhi_wang@outlook.com

⁺These authors have contributed equally to this work.

Abstract

Objective This study aimed to determine the relationship between occupational noise, shift work, and nonalcoholic fatty liver disease(NAFLD) in male workers in the automobile manufacturing industry.

Design Cross-sectional study.

Setting This study was carried out Guangzhou Twelfth People's Hospital, using data from April to September 2022.

Participants A total of 4672 eligible participants were included in the study.

Primary and secondary outcome measures Diagnosis of NAFLD was made using ultrasound. Noise was detected according to the Measurement of Physical Factors in the Workplace-Part 8: Noise. Environmental noise intensity was assessed using an EDGE personal noise dosimeter manufactured by CASELLA (UK). The working status of workers were investigated by questionnaire.

Results The OR of NAFLD was 1.39 (1.03, 1.88) in CNE \geq 95 group compared to CNE <85 group. Improved risk of NAFLD in workers with shift work compared to those without shift work (OR=1.35, 95%CI: 1.09-1.68). As stratified analyses showed, the ORs of NAFLD prevalence related to occupational noise and shift work exposure appears to be increased in young workers. When both shift work and noise exposure work are present simultaneously, the synergy index (SI) between them was 0.47 (95%CI: 0.25 to 0.89). Combined effects analysis revealed that the ORs of NAFLD was 2.02 (95%CI: 1.34-2.99) in CNE \geq 95 and cumulative length of night shifts work >2920 hours.

Conclusion Occupational noise exposure may be an independent risk factor for NAFLD. And it may synergistically affect disease when combined with night shift work, particularly among younger workers. These findings underscore the importance for companies to prioritize the management and training of younger workers, along with targeted occupational health education initiatives, as crucial measures for reducing the incidence of NAFLD.

STRENGTHS AND LIMITATIONS OF THIS STUDY

 \Rightarrow In this study, a relatively large sample was used to analyse the association between occupational noise exposure, shift work, and their combined effect on NAFLD using male workers in the automobile manufacturing industry as the study population.

 \Rightarrow The questionnaire items on working hours, lifestyle, diet and so on, were designed simplistically and may be difficult to assess comprehensively.

 \Rightarrow Due to the cross-sectional survey, there may be observational bias and causality cannot be elucidated either.

Introduction

In recent decades, there has been a progressive increase in the incidence of NAFLD, leading to a significant burden of morbidity and mortality associated with liver-related disorders¹². Anticipating future trends, researchers project a continued rise in the prevalence of NAFLD, while expecting a stable or possibly decreasing prevalence of other chronic liver diseases³. In China, the prevalence of NAFLD has notably increased in the past decade, rising from 18% to 29.2%. Industrialization and changes in lifestyle have contributed to this rise, resulting in a 10% increase in national incidence⁴. Recent studies have highlighted the rapid increase in NAFLD prevalence among adolescents⁵.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

The development of non-alcoholic fatty liver disease (NAFLD) is influenced by various factors, including age, gender, type 2 diabetes, metabolic syndrome, its components, and family history of fatty liver⁶⁻¹⁰. Age and gender also play significant roles in the onset of NAFLD. The increasing prevalence of NAFLD can be attributed to population aging, where age-related visceral fat accumulation may contribute to the secretion of inflammatory factors^{7 8}. Unhealthy dietary habits, including high-fat diets, and unhealthy lifestyles such as late-night activities have also been associated with an increased risk of NAFLD¹¹⁻¹³. Furthermore, occupational factors in the workplace, such as noise exposure and shift work, may influence the risk of NAFLD. Previous research has demonstrated a significant dose-response relationship between cumulative noise exposure (CNE) and the detection rates of fatty liver¹⁴. Shift work is a prevalent practice in

various industries, including manufacturing, pharmaceutical, and service sectors. It not only disrupts normal circadian rhythms directly but is also closely linked to sleep disorders, leading to insulin resistance and stress reactions, causing physiological disruptions and harm to health¹⁵⁻¹⁷. These disturbances can disrupt glucose and lipid metabolism, which is a significant contributing factor in elevating the risk of NAFLD¹⁸.

Workers in the automotive manufacturing industry frequently endure prolonged occupational noise exposure and may also be required to engage in shift work. Despite this, few investigations have specifically examined the impact of shift work and noise on NAFLD within the context of this workforce. Consequently, our study aims to investigate the effects of occupational noise exposure and shift work on NAFLD among male workers in the automotive manufacturing industry. By focusing on this specific population, the study intends to provide a theoretical foundation to implement measures to protect against occupational hazards.

Methods

Study design and population

From April to September 2022, a cross-sectional study was conducted among male workers at an automobile manufacturing company in Guangzhou for an occupational health examination. Participants with incomplete medical examination or questionnaire data, female workers, individuals taking antiretroviral drugs, and those with a history of cancer, excessive alcohol intake, hepatobiliary disease, blood system diseases, or renal failure were excluded. Ultimately, a total of 4672 eligible participants were included in the study. This study was reviewed and approved by the Ethics Committee of Guangzhou Twelfth People's Hospital (No.2021008) and all procedures involving participants were conducted in accordance with the ethical standards of the Institutional Research Board and the 1964 Declaration of Helsinki. All study participants gave informed consent.

NAFLD diagnosis and collection of information on covariates

Trained physicians conducted the occupational health examination in accordance with a standardized protocol. Diagnosis of NAFLD was made using the criteria proposed by FARRELL et al¹⁹. Blood was drawn from the fasting subjects to measure

hemoglobin(HGB) and fasting plasma glucose(FPG) levels. Blood pressure was measured by the omron HEM-7071 electronic blood pressure monitor three times, each time at an interval of 1 min, and the average of the three measurements was taken as the individual blood pressure (BP) value. BP was classified according to 2018 Chinese Guidelines for Prevention and Treatment of Hypertension. FPG normal range is 3.9-6.1mmol/L. According to the Guidelines for the Prevention and Control of Overweight and Obesity in Chinese Adults, our study the included BMI<24kg/m² and BMI \geq 24kg/m² as dichotomous variables in the multivariate logistic regression model.

Referring to previous research and consulting multiple experts in the field of occupational health to revise and improve it continuously, we used a self-designed questionnaire to investigate²⁰. The questionnaires were administered face to face by trained investigators to acquire participants' information, including dietary habits, lifestyle, occupational information and past medical history. Finally, investigators checked the completed questionnaires to detect errors promptly. We investigated the average number of working days per week for workers to assess working hours and determine whether it is considered as long working hours. Long working hours are defined as working more than 40 hours per week²¹. Simultaneously, we made inquiries regarding the exposure statuses of other occupational hazard factors, so as to guarantee that the exposures to other occupational factors like dust and benzene series (mainly refers to benzene, toluene and xylene) were beneath the occupational exposure limits (OELs) stipulated in "Occupational exposure limits for hazardous agents in the workplace-Part 1: Chemical hazardous agents" (GBZ 2.1-2019). Classification of dietary habits: high-salt foods mainly include pickled vegetables, ham and bacon, and so on^{22 23}; high-fat foods including chips, chocolate, animal offal and fried foods; late evening snack is defined as an additional meal arranged after 21:00. According to the frequency of eating within a week, they are divided into four groups: seldom, 1~2 times per week, 3~4 times per week, and >4 times per week. Smoking is defined as a consistent consumption of at least 1 cigarette per day for a duration of more than 6 months. Alcohol consumption is defined as a regular pattern of drinking at least once a week for a period of 1 year or longer. According to the duration of sleep, they were Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

divided into three groups: < 6h (short sleep duration), $6 \sim 8h$ (moderate sleep duration), and >8h (long-sleep duration)²⁴.

Noise exposure assessment

Our study followed the "Measurement of Physical Factors in the Workplace - Part 8: Noise" (GBZ/T189.8-2007) for noise detection. Environmental noise intensity was assessed using an EDGE personal noise dosimeter manufactured by CASELLA (UK). Each monitoring point was measured three times and the average value was taken as the measurement result. In this study, noise exposure was evaluated using the normalization of continuous A-weighted sound pressure level equivalent to an 8h per day (LAeq, 8h), and work exposed to noise is defined as those operations involving the presence of hearing-impairing, health-hazardous or otherwise hazardous sound, and LAeq,8h \geq 80 dB(A) for at least one year. Meanwhile, Cumulative noise exposure (CNE) was used to quantify the noise exposure level of the study participants and it is calculated as: CNE=L_{Aeq, 8h}+10logT, where T indicates years of noise exposure and the CNE unit is: dB(A)·year.

Assessment of shift work

The International Labour Organization defines shift work as extending the working day to 24 hours by alternating between different workers and/or teams^{25 26}. Night shift work is defined as work occurring between 0:00 and 5:00 am^{26 27}. The cumulative length of night shift work was calculated in hours during data processing. In this study, the two main types of shift patterns for workers are two shifts system and three shifts system.

In addition, according to the sampling methods and technical requirements in "Specifications of air sampling for hazardous substances monitoring in the workplace" (GBZ 159-2004), collect samples of dust, benzene, toluene, and xylene in the air of workplaces. In accordance with the requirements of "Determination of dust in the air of workplace Part 1: Total dust concentration" (GBZ 192.1-2007) and "Occupational exposure limits for hazardous agents in the workplace-Part 1: Chemical hazardous agents" (GBZ 2.1-2019), we conduct fixed-point sampling and detection on representative operating posts for dust, benzene, toluene, and xylene in the air.

Statistical analyses

Page 9 of 29

BMJ Open

Subject staff verified data daily to ensure unique ID codes for each individual and timely identification of missing key data and logical error. R (version 4.2.1) was used for all statistical analyses. Continuous variables that followed normal distribution were presented as mean ±standard deviation (SD), and were compared by t-test. Mann-Whitney U test was used to compare continuous variables with skewed distribution in the form of median with interquartile ranges. The χ^2 test was used to compare categorical variables, which have been expressed as numbers (percentages). The relationship between shift work, occupational noise and NAFLD was analyzed using a multivariate logistic regression model, with night shift hours and CNE included in the regression model as categorical variables respectively as well as the joint effect of both was analysed in relation to NAFLD. Referring to the national criteria (GBZT229.4-2012), CNE is divided into four groups <85, 85~, 90~ and 95~. Cumulative length of night shift work are grouped by quartiles, using the first quartile as a reference. Previous studies have considered adolescents and young adults (AYAs) to be the group aged 15 years and older, with the cut-offs at 24, 29 and 39 years being difficult to determine and some studies considering 18-24 years as late adolescence²⁸. We stratified the age into two groups ≤ 29 and ≥ 30 for analysis. To provide more clarity on the effect of CNE on the illness of workers in different age groups, we subdivided younger workers at age 29 into two groups (\leq 24 and 25-29 years old)^{29 30}. All test were two-sided, and α =0.05 was considered statistically significant.

Patient and public involvement

Patients and the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Results

Basic information about the study population

A total of 5928 questionnaires were distributed, out of which 4791 valid questionnaires were returned, resulting in a valid response rate of 80.8%. After excluding individuals who did not meet the study's inclusion criteria, and those with incomplete occupational health screening results (n=119), the final sample size included for analysis was determined. Overall, 4672 workers with complete information on occupational health

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

examination and questionnaires were included in the study according to the inclusion and exclusion criteria. The mean age of the study population was (26.8±5.0), with different prevalence rates for workers in the \leq 29 and \geq 30 years age groups (*P*<0.001), with a total of 1363 workers with NAFLD and a prevalence of 29.2%. There were 3658 (78.3%) shift workers, with different prevalence of NAFLD in different shift work situations (*P*<0.001). Meanwhile, 2965 (63.5%) noise workers had a statistically significant difference in prevalence in different noise exposure scenarios (*P*<0.05). Cumulative length of night shift work and CNE vary by disease state (all *p*<0.05). A summary of the sample characteristics is presented in Table 1. The results of univariate analysis of drinking status, high-salt foods, high-fat foods, physical activity and importance of noise protection equipment are presented in Supplementary Table 1.

Table 1 Basic characteristics of population	ulation in different disease states (r	n, %)
---	--	------	---

Characteristics	NAFLD		Total	D
	No (N=3309)	Yes (N=1363)	(N=4672)	1
Age (years), n (%)				< 0.001
≤29	2649 (80.1)	698 (51.2)	3347 (71.6)	
≥30	660 (19.9)	665 (48.8)	1325 (28.4)	
Education, n (%)				0.009
Senior secondary and below	1888 (57.1)	720 (52.8)	2608 (55.8)	
---	-------------	-------------	-------------	---------
Higher education	1421 (42.9)	643 (47.2)	2064 (44.2)	
Smoking status (cigarettes per day), n (%)				< 0.001
Nonsmokers	1741 (52.6)	729 (53.5)	2470 (52.9)	
1~5	618 (18.7)	198 (14.5)	816 (17.5)	
6~10	599 (18.1)	239 (17.5)	838 (17.9)	
>10	351 (10.6)	197 (14.5)	548 (11.7)	
Fruit and vegetable (times per week), n (%)				0.002
Seldom	262 (7.9)	89 (6.5)	351 (7.5)	
1~2	1961 (59.3)	752 (55.2)	2713 (58.1)	
3~4	653 (19.7)	297 (21.8)	950 (20.3)	
>4	433 (13.1)	225 (16.5)	658 (14.1)	
Late evening snack (times per week), n (%)				0.024
Seldom	747 (22.6)	361 (26.5)	1108 (23.7)	
1~2	1863 (56.3)	735 (53.9)	2598 (55.6)	
3~4	386 (11.7)	137 (10.1)	523 (11.2)	
>4	313 (9.5)	130 (9.5)	443 (9.5)	
BMI (kg/m2), n (%)				< 0.001
<24	3001 (90.7)	555 (40.7)	3556 (76.1)	
≥24	308 (9.3)	808 (59.3)	1116 (23.9)	
Sleep duration (hours),n (%)				< 0.001
<6	592 (17.9)	303 (22.2)	895 (19.2)	
6~8	2574 (77.8)	1023 (75.1)	3597 (77.0)	
>8	143 (4.3)	37 (2.7)	180 (3.9)	
BP (mmHg), n (%)				< 0.001
Normal	1374 (41.5)	306 (22.5)	1680 (36.0)	
High-normal	1890 (57.1)	963 (70.7)	2853 (61.1)	
Hypertension	45 (1.4)	94 (6.9)	139 (3.0)	
ECG, n (%)				< 0.001
Normal	1392 (42.1)	824 (60.5)	2216 (47.4)	
Abnormal	1917 (57.9)	539 (39.5)	2456 (52.6)	
FPG (mmol/L), n (%)				< 0.001
Normal	2887 (87.2)	1001 (73.4)	3888 (83.2)	
Abnormal	422 (12.8)	362 (26.6)	784 (16.8)	

Table 1 Continued

Characteristics	NA	FLD	$T_{abal}(N-4(72))$	מ
	No (N=3309)	Yes (N=1363)	-10tal(N-40/2)	P
HGB (g/L), n (%)				0.005
Normal	2586 (78.2)	1013 (74.3)	3599 (77.0)	
Abnormal	723 (21.8)	350 (25.7)	1073 (23.0)	
Long working hours				< 0.001

No	1375 (41.55)	685 (50.26)	2060 (44.09)	
Yes	1934 (58.45)	678 (49.74)	2612 (55.91)	
Dust				0.015
No	2990 (90.36)	1262 (92.60)	4252 (91.01)	
Yes	319 (9.64)	101 (7.41)	420 (8.99)	
Benzene series				0.02
No	2828 (85.46)	1200 (88.04)	4028 (86.22)	
Yes	481 (14.54)	163 (11.96)	644 (13.78)	
Shift work				< 0.001
No	808 (24.42)	206 (15.11)	1014 (21.70)	
Yes	2501 (75.58)	1157 (84.89)	3658 (78.30)	
Work exposed to noise				0.045
No	1239 (37.44)	468 (34.34)	1707 (36.54)	
Yes	2070 (62.56)	895 (65.66)	2965 (63.46)	
Cumulative length of night shift work (hours)	1100 (511-2560)	2190 (1090-4380)	1460 (730-2920)	< 0.001
CNE [dB(A) year]	86.9 (83.1-91.1)	89.8 (85.2-93.3)	87.8 (83.6-91.9)	< 0.001
Seniority in shift work (year), median (IQR)	3.0 (1.50, 6.50)	6.0 (3.95, 12.0)	4.0 (2.0, 8.0)	< 0.001
Seniority in work exposed to noise (year), median (IQR)	3.00 (1.30, 7.00)	7.90 (2.75,13.50)	3.8 (1.50, 8.33)	< 0.001

The relationship between occupational noise exposure and NAFLD

The impact of work exposed to noise on NAFLD was demonstrated in Table 2. Compared with those did not, no statistically significant relationship between work exposed to noise and NAFLD (OR=0.92, 95%CI: 0.78-1.09). Table 3 and Supplementary Table 2 presented the odds ratios (ORs) and 95% confidence intervals (CIs) for the effect of occupational noise exposure on NAFLD. Stratified analysis indicated that the association between CNE and NAFLD was not statistically significant in both those aged \leq 29 and \geq 30 years (see supplemental Table 1); By reference to CNE<85, the OR of NAFLD was 1.39 (95%CI: 1.03, 1.88) in CNE \geq 95 group. Table 3 shows further stratification by age, among those \leq 24 years of age, the OR of NAFLD in the CNE \geq 95 group was 2.97 (95%CI: 1.18-6.95).

Table 2 Association of work exposed to noise with NAFLD

work exposed to	Model 1		Model 2	
noise (Yes or No)	OR 95%CI	Р	OR 95% <i>CI</i>	Р
Total				
No	Ref			
Yes	0.97 (0.83, 1.15)	0.754	0.92 (0.78, 1.09)	0.356

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Age <29 vr				
No	Ref			
Yes	1.07 (0.87, 1.31)	0.537	0.98 (0.79, 1.21)	0.826
≥30 yr				
No	Ref			
Yes	0.84 (0.63, 1.11)	0.212	0.83(0.63, 1.11)	0.212

Model 1: Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB; Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, benzene series, shift work.

CNE [dP(A), wear]	Model 1		Model 2]
	OR 95%CI	Р	OR 95% <i>CI</i>	Р
Total				
<85				
85~	1.20 (0.97, 1.48)	0.087	1.12 (0.90, 1.39)	0.320
90~	1.16 (0.93, 1.45)	0.191	1.08 (0.85, 1.36)	0.536
95~	1.48 (1.10, 1.99)	0.009	1.39 (1.03, 1.88)	0.033
Age				
≤24 yr				
<85	Ref			
85~	1.07 (0.77, 1.48)	0.676	0.96 (0.68, 1.35)	0.810
90~	1.00 (0.64, 1.52)	0.988	0.86 (0.54, 1.34)	0.512
95~	3.18 (1.29, 7.31)	0.008 🧹	2.97 (1.18, 6.95)	0.015
25~29 yr				
<85	Ref			
85~	1.14 (0.80, 1.62)	0.480	1.14 (0.78, 1.66)	0.505
90~	1.11 (0.76, 1.62)	0.603	1.08 (0.73, 1.62)	0.690
95~	1.01 (0.55, 1.81)	0.963	1.07 (0.57, 1.95)	0.825

Table 3 The relationship between CNE and NAFLD in different age groups

Model 1: Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB; Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, benzene series, shift work.

The relationship between shift work and NAFLD

The effect of shift work on NAFLD is shown in the Figure 1. Compared to those without shift work, the OR of NAFLD among workers with shift work was 1.35 (95%CI: 1.09-1.68). Stratified analysis showed that shift work was a risk factor for NAFLD among

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

those aged ≤ 29 years (OR=1.46, 95%CI: 1.15-1.87). Similarly, in comparison with cumulative length of night shift work ≤ 730 h, rising cumulative length of night shift work were associated with a higher risk of NAFLD, with ORs (95%CIs) of 1.43 (1.14, 1.79), 1.44 (1.15, 1.81) and 1.64 (1.23, 2.17), respectively. Among those aged ≤ 29 years, increased cumulative length of night shift work were also a risk factor for NAFLD, with ORs (95%CIs) of 1.66 (1.30, 2.11), 1.52 (1.17, 1.98) and 2.41 (1.50, 3.84) respectively.

The relationship between the interaction of work exposed to noise and shift work and NAFLD

To further explore the combined effects of noise exposure work and shift work, we analyzed the interaction between them. The results were shown in Table 4. When both shift work and noise exposure work are present simultaneously, the excess relative risk of relative excess risk due to interaction (RERI) was -0.43 (95%CI: -1.00 to 0.14), the attributable proportion due to interaction (AP) was -0.31 (-0.69 to 0.07), and the synergy index (SI) between them was 0.47 (95%CI: 0.25 to 0.89).

Table 4	The relationship	between the	interaction	of wo	rk exposec	to noise ar	nd shift v	work and	
NAFLD									

work exposed	shift work			Total		
to noise (Yes or No)	(Yes or No)	OR 95% <i>CI</i>	RERI	AP	SI	$P_{interaction}$
No	No	Ref	-0.43 (-1.00, 0.14)	-0.31 (-0.69, 0.07)	0.47 (0.25, 0.89)	0.112
	Yes	1.60 (1.19, 2.17)				
Yes	No	1.21 (0.83, 1.75)				
	Yes	1.38 (1.05, 1.83)				

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, benzene series.

The combined effect of occupational noise and shift work in association with NAFLD

The combined effect of occupational noise and cumulative length of night work on NAFLD was shown in Table 5. After adjusting for variables, the association with NAFLD was statistically significant for all groups at cumulative length of night shift

BMJ Open

work >2920h and CNE>85, when compared with CNE<85 and cumulative length of night shift work \leq 730h. The OR of NAFLD was greatest at 2.01 (95%CI: 1.34 to 2.99) when cumulative length of night shift work >2920h and CNE \geq 95. Furthermore, we analyzed the relationships between occupational noise, shift work and NAFLD in different states of benzene series exposure as well (see Supplementary Table 3).

CNE		Cumulative length	of night shift work (hou	urs)
CNE [dP(A).woors]	≤730	730~	1460~	2920~
	OR 95%CI	OR 95%CI	OR 95%CI	OR 95% <i>CI</i>
<85	Ref	1.49 (1.04, 2.12)*	1.27 (0.82, 1.94)	1.34 (0.64, 2.76)
85~	0.88 (0.6, 1.27)	1.45 (1.05, 1.99)*	1.61 (1.14, 2.26)**	1.99 (1.29, 3.07)**
90~	1.24 (0.76, 1.97)	1.31 (0.85, 1.99)	1.41 (1.00, 1.98) *	1.54 (1.08, 2.21)*
95~	1.98 (0.89, 4.26)	1.61 (0.72, 3.48)	1.67 (0.95, 2.88)	2.01 (1.34, 2.99)***

Table 5	Association o	f CNE and	d cumulative	length o	of night	shift	work with	NAFLD
---------	---------------	-----------	--------------	----------	----------	-------	-----------	-------

*p<0.05, **p<0.01, ***p<0.001.

Bold values are p<0.05.

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, benzene series.

Discussion

To summarize, our study included a total of 4672 male workers in the automotive manufacturing industry, investigating the impact of occupational noise and shift work on non-alcoholic fatty liver disease (NAFLD) through a cross-sectional survey. The majority of the company's employees are subjected to a high workload, with 63.5% working in noise and 78.3% working in shifts, which is significantly higher compared to other occupations such as teachers and white-collar workers³¹. Night work has been categorized by the International Agency for Research on Cancer as "possibly carcinogenic to humans" (Group 2A)³², prompting growing concerns about its adverse effects on health. The prevalence of NAFLD in this survey was 29.2%, a rate close to the combined prevalence estimate of NAFLD reported by Fan et al^{11 33-35}. Among those exposed to noise and those not exposed, the prevalence of NAFLD is 30.2% and 27.4%, respectively. After adjusting for relevant variables, multifactorial logistic regression analysis revealed that increased CNE was the significant risk factor associated with NAFLD. This may be because the biological effects of noise are cumulative and the

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

effects of a single noise exposure are not sufficient to cause health damage to the organism. However, the more repeated exposures and the longer the exposure period, the more pronounced the cumulative effects will be when the effects are not eliminated^{36 37}. Associations between CNE and NAFLD vary across age groups. An increase in CNE is a risk factor for NAFLD among workers under 24 years of age. This suggests that short-term exposure to high-intensity noise may contribute to the development of NAFLD in this population. Possible factors that may contribute to this association include the inadequate use of noise protective equipment among young workers or exposure to noise outside of the occupational setting. The effect of noise on NAFLD in younger workers should be further explored. This indicates that enterprises should strengthen training and occupational health education for young workers, which is important to reduce the occurrence of NAFLD. Shins, Saeha et al. have shown that chronic exposure to road traffic noise was associated with an increased incidence of diabetes and hypertension in Toronto³⁸. Du et al. found that noise exposure could lead to disorders of glucolipid metabolism in mice through inhibition of the AKT pathway mediated by the biological clock gene BMAL1³⁹. The experiment by Evans et al. also demonstrated significant changes in serum ALT levels across test groups, suggesting that noise can lead to liver dysfunction⁴⁰. It can be seen that noise may be the basis for the formation of NAFLD by promoting insulin resistance and disorders of glucolipid metabolism in the body, resulting in metabolic abnormalities in the body. These studies also indirectly support our findings. In this study, the prevalence of NAFLD was 20.3% and 31.6% among auto manufacturing workers not involved in shift work and those who did, respectively, with a higher prevalence among workers in shift work, similar to the findings of Golabi et al¹¹. We observed that shift work and an increase in cumulative length of night shift work were identified as risk factors for NAFLD. We also examined the relationship between shift work and NAFLD across different age groups. Specifically, among workers below 29 years of age, engaging in shift work was found to significantly increase the risk of NAFLD. This indicates the presence of specific physiological adaptations that contribute to this heightened risk. Similarly, we found that an extended cumulative length of night shift work was a significant risk

Page 17 of 29

BMJ Open

factor for NAFLD in workers below 29 years of age. These findings emphasize the importance of appropriate management strategies to mitigate the adverse effects of shift work on NAFLD among younger workers. Previous epidemiological studies have consistently demonstrated a link between shift work and NAFLD. For instance, a 4year cohort study conducted among Chinese railway workers reported a higher incidence of NAFLD in individuals engaged among shift work¹¹. Similarly, a crosssectional survey involving male steelworkers found that night shift workers had an increased risk of NAFLD compared to those working day shifts¹⁵. Wang et al. further supported the association between night shifts and NAFLD by demonstrating the impact of disrupted circadian rhythms, due to delayed sleep, on the incidence of NAFLD⁴¹. These findings are in line with our results. Sun et al. also revealed the association between sleep deprivation, increased cortisol levels, and insulin resistance, further highlighting the possible mechanisms underlying the relationship between sleep disruption and NAFLD⁴². Our study identified a significant association between occupational noise exposure, shift work, and an increased risk of developing NAFLD among workers. This association may be explained by the potential mechanisms through which both noise exposure and shift work contribute to the development of insulin resistance, a key factor in the pathogenesis of NAFLD. Insulin resistance disrupts lipid metabolism, leading to elevated lipase activity and subsequent overproduction of triglycerides and free fatty acids. Consequently, an excessive amount of free fatty acids enters the liver, promoting increased lipid synthesis and deposition within hepatic cells. This lipid accumulation interferes with hepatocyte transport capacity, resulting in hepatocyte apoptosis and the accumulation of fat, ultimately leading to NAFLD development.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

It should be noted that some participants in this study were simultaneously exposed to benzene series (mainly benzene, toluene, and xylene). The study by Yun - Hee Cho et al. showed that exposure to chemicals such as benzene, toluene, and xylene could significantly increase the risk of NAFLD⁴³ ⁴⁴. In obese populations, benzene exposure is positively correlated with liver injury, and systemic inflammation increases with the exposure to benzene and toluene, indicating that benzene and toluene exposure

 have certain potential impacts on steatohepatitis⁴⁵. These effects may result from the activation of cell death, the induction of oxidative and endoplasmic reticulum stress, and the disruption of metabolism, which are caused by the dysfunction of xenobiotic and endobiotic receptors⁴⁶. After detection, we found that the concentrations of benzene series in the researchers' workplaces were all lower than the occupational exposure limits (OELs) stipulated in "Occupational exposure limits for hazardous agents in the workplace - Part 1: Chemical hazardous agents". To ensure the reliability of the results, we also analyzed the relationships between occupational noise, shift work and NAFLD under different benzene series exposure, occupational noise exposure and shift work were still risk factors for NAFLD.

This study contributes to a relatively uncommon and innovative exploration of the association between occupational noise exposure, shift work, and their combined effect on NAFLD among male workers in the automotive manufacturing industry. Based on the study findings, we propose several recommendations for relevant management departments. It is advised to scientifically and reasonably organize the working system, promote the implementation of flexible working arrangements, and minimize the cumulative length of night work. Furthermore, improving the production process to control or eliminate noise sources, such as installing sound insulation and vibration dampers, is recommended. Additionally, establishing and enhancing occupational health supervision, strengthening the management of occupational health, and facilitating regular occupational health check-ups for workers are crucial measures to safeguard worker health. These recommendations hold significant practical importance in protecting the health of workers.

However, it has to acknowledge the limitations of our study. Due to the crosssectional nature of the survey, there may be observational bias and causality cannot be elucidated either. To address these limitations, we are in the process of establishing a cohort study among workers in the automotive manufacturing industry. In future studies, we plan to utilize individual noise dosimeters to more accurately measure workers' noise exposure, enabling a more robust analysis of the association between these occupational factors and disease outcomes.

Acknowledgements

We wish to thank the timely help given by workers who participated in this program. we are deeply grateful for Guangzhou Twelfth People's Hospital for their support for collecting data.

Contributors

Jinwei Zhang, Zhang Yuxia and Wang Zhi conceived and designed the study. Wang Zhi provided administrative support. Jinwei Zhang, Qiu Congxi, Ma Weiyu, Wenfeng Zeng and Wang Zhi provided study materials. Jinwei Zhang, Zhang Yuxia, Qiu Congxi, Wu Kangyong, Zhang Jingwen, and Cui Jiaxin collected and assembled data. Zhang Yuxia, Yanmei Ruan, Gao Yunxia, Liang Jiabin and Ye Cuiping analyzed and interpreted the data. Zhang Yuxia, Jinwei Zhang and Gao Yunxia wrote the manuscript. Wang Zhi and Jinwei Zhang approved the final manuscript. The guarantor of the study is Wang Zhi , accepts full responsibility for the finished work and the conduct of the study, has access to the data and controls the decision to publish.

Fundings

This work was supported by the Key Research and Development Program of Guangzhou Science and Technology Project (Project Number: 202206010061); Guangzhou Key Medical Discipline (2025-2027); The Science and Technology Program of Guangzhou (Project Number: 2023A03J0502, 2023A03J0979, 2024A03J0432, 2024A03J0505, 2025A03J3591); The Medical Science and Technology Research Fund Project of Guangdong (Project Number: B2023347, A2023406, A2023396); The Science and Technology Project of Guangzhou Municipal Health Commission (Project Number: 20251A010042);

Ethics statements

Patient consent for publication

Consent obtained directly from patient(s).

Ethics approval

The study was approved by Guangzhou Twelfth People's Hospital ethics committee(No.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

BMJ Open

> 2021008), and all procedures involving participants were conducted in accordance with the ethical standards of the Institutional Research Committee and the 1964 Helsinki Declaration. The informed consent form was sent in electronic form to each participant before the survey began, and participants could also request a paper copy.

> > ion peer teries only

Data availability statement

Data are available upon reasonable request.

Conflicts of interest

The authors declare that they have no competing interests

References

 Notarnicola M, Osella AR, Caruso MG, et al. Nonalcoholic Fatty Liver Disease: Focus on New Biomarkers and Lifestyle Interventions. *Int J Mol Sci* 2021;22(8):3899.
 Masoodi M, Gastaldelli A, Hyötyläinen T, et al. Metabolomics and lipidomics in NAFLD: biomarkers and non-invasive diagnostic tests. *Nat Rev Gastroenterol Hepatol* 2021;18(12):835-56. doi: 10.1038/s41575-021-00502-9 [published Online First: 2021/09/12]

3. Younossi ZM, Stepanova M, Afendy M, et al. The changing face of chronic liver disease (CLD) in the United States: The rising epidemic of non-alcoholic fatty liver disease (NAFLD). *J Hepatol* 2011;54

 Younossi Z, Tacke F, Arrese M, et al. Global Perspectives on Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. *Hepatology* 2019;69(6):2672-82. doi: 10.1002/hep.30251 [published Online First: 2018/09/05]

5. Lin G, Xinhe Z, Haoyu T, et al. Epidemiology and lifestyle survey of non-alcoholic fatty liver disease in school-age children and adolescents in Shenyang, Liaoning. *BMC Pediatr* 2022;22(1):286. doi: 10.1186/s12887-022-03351-w [published Online First: 2022/05/18]

6. Younossi ZM, Golabi P, de Avila L, et al. The global epidemiology of NAFLD and NASH in patients with type 2 diabetes: A systematic review and meta-analysis. *J Hepatol* 2019;71(4):793-801. doi: 10.1016/j.jhep.2019.06.021 [published Online First: 2019/07/08]

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

7. Kim IH, Kisseleva T, Brenner DA. Aging and liver disease. *Curr Opin Gastroenterol* 2015;31(3):184-91. doi: 10.1097/mog.00000000000176 [published Online First: 2015/04/09]

8. Barzilai N, Huffman DM, Muzumdar RH, et al. The critical role of metabolic pathways in aging. *Diabetes* 2012;61(6):1315-22. doi: 10.2337/db11-1300 [published Online First: 2012/05/24]

9. Wong VW, Chu WC, Wong GL, et al. Prevalence of non-alcoholic fatty liver disease and advanced fibrosis in Hong Kong Chinese: a population study using protonmagnetic resonance spectroscopy and transient elastography. *Gut* 2012;61(3):409-15. doi: 10.1136/gutjnl-2011-300342 [published Online First: 2011/08/19]

 10. Wang J, Tang Y, Peng K, et al. Development and validation of a nomogram for predicting nonalcoholic fatty liver disease in the non-obese Chinese population. *Am J Transl Res* 2020;12(10):6149-59. [published Online First: 2020/11/17]

11. Xu J, Ni S, Wang Y, et al. Shift work and nonalcoholic fatty liver disease incidence among Chinese rail workers: a 4-year longitudinal cohort study. *Int Arch Occup Environ Health* 2023;96(1):179-90. doi: 10.1007/s00420-022-01913-0

Takahashi A, Anzai Y, Kuroda M, et al. Effects of sleep quality on non-alcoholic fatty liver disease: a cross-sectional survey. *BMJ Open* 2020;10(10):e039947. doi: 10.1136/bmjopen-2020-039947 [published Online First: 2020/10/31]

13. Semmler G, Datz C, Reiberger T, et al. Diet and exercise in NAFLD/NASH: Beyond the obvious. *Liver Int* 2021;41(10):2249-68. doi: 10.1111/liv.15024 [published Online First: 2021/07/31]

14. Bai G-p. Relationship Between Occupational Noise Expose and the Fatty Liver. *Hebei Medicine* 2009

15. Zhang S, Wang Y, Wang Z, et al. Rotating night shift work and non-alcoholic fatty liver disease among steelworkers in China: a cross-sectional survey. *Occup Environ Med* 2020;77(5):333-39. doi: 10.1136/oemed-2019-106220 [published Online First: 2020/02/06]

16. Attarchi M, Dehghan F, Safakhah F, et al. Effect of exposure to occupational noise and shift working on blood pressure in rubber manufacturing company workers. *Ind Health* 2012;50(3):205-13. doi: 10.2486/indhealth.ms1321 [published Online First: 2012/03/29]

17. Dong H, Zhang Q, Zhang Z, et al. Association of sleep disturbance with shift work, occupational stress, and musculoskeletal pain in Chinese public hospital emergency workers: A multicentre cross-sectional study. *Chronobiol Int* 2022;39(6):886-94. doi: 10.1080/07420528.2022.2048662 [published Online First: 2022/03/08]

18. Boivin DB, Boudreau P, Kosmadopoulos A. Disturbance of the Circadian System in Shift Work and Its Health Impact. *J Biol Rhythms* 2022;37(1):3-28. doi: 10.1177/07487304211064218 [published Online First: 2022/01/01]

19. Farrell GC, Chitturi S, Lau GK, et al. Guidelines for the assessment and management of non-alcoholic fatty liver disease in the Asia-Pacific region: executive summary. *J Gastroenterol Hepatol* 2007;22(6):775-7. doi: 10.1111/j.1440-1746.2007.05002.x [published Online First: 2007/06/15]

20. Ruan Y, Huang G, Zhang J, et al. Risk analysis of noise-induced hearing loss of workers in the automobile manufacturing industries based on back-propagation neural network model: a cross-sectional study in Han Chinese population. *BMJ Open* 2024;14(5):e079955. doi: 10.1136/bmjopen-2023-079955 [published Online First: 2024/05/18]

21. Pega F, Náfrádi B, Momen NC, et al. Global, regional, and national burdens of ischemic heart disease and stroke attributable to exposure to long working hours for 194 countries, 2000-2016: A systematic analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ Int* 2021;154:106595. doi: 10.1016/j.envint.2021.106595 [published Online First: 2021/05/21]

22. Webster JL, Dunford EK, Neal BC. A systematic survey of the sodium contents of processed foods. *Am J Clin Nutr* 2010;91(2):413-20. doi: 10.3945/ajcn.2009.28688 [published Online First: 2009/12/04]

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

23. Ni Mhurchu C, Capelin C, Dunford EK, et al. Sodium content of processed foods in the United Kingdom: analysis of 44,000 foods purchased by 21,000 households. *Am J Clin Nutr* 2011;93(3):594-600. doi: 10.3945/ajcn.110.004481 [published Online First: 2010/12/31]

24. Imaizumi H, Takahashi A, Tanji N, et al. The Association between Sleep Duration and Non-Alcoholic Fatty Liver Disease among Japanese Men and Women. *Obes Facts* 2015;8(4):234-42. doi: 10.1159/000436997

25. International Labour Organization. C171-Night Work Convention, 1990(No.171)1990 [Available from:

https://www.ilo.org/dyn/normlex/en/f?p=NORMLEXPUB:12100:0::NO::P12100_IN STRUMENT_ID:312316.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

26. Organization, WH. IARC Monographs Volume 124: Night Shift Work

[Available from: <u>https://www.iarc.who.int/news-events/iarc-monographs-volume-</u> 124-night-shift-work/.

27. Molzof HE, Peterson CM, Thomas SJ, et al. Nightshift Work and Nighttime Eating Are Associated With Higher Insulin and Leptin Levels in Hospital Nurses. *Front Endocrinol (Lausanne)* 2022;13 doi: 10.3389/fendo.2022.876752

28. Arshad T, Paik JM, Biswas R, et al. Nonalcoholic Fatty Liver Disease Prevalence Trends Among Adolescents and Young Adults in the United States, 2007-2016. *Hepatol Commun* 2021;5(10):1676-88. doi: 10.1002/hep4.1760 [published Online First: 2021/09/25]

29. Paik JM, Kabbara K, Eberly KE, et al. Global burden of NAFLD and chronic liver disease among adolescents and young adults. *Hepatology* 2022;75(5):1204-17. doi: 10.1002/hep.32228 [published Online First: 2021/11/07]

30. Zhang X, Wu M, Liu Z, et al. Increasing prevalence of NAFLD/NASH among children, adolescents and young adults from 1990 to 2017: a population-based observational study. *BMJ Open* 2021;11(5):e042843. doi: 10.1136/bmjopen-2020-042843 [published Online First: 2021/05/06]

31. Wang D, Chen H, Chen D, et al. Shift work disorder and related influential factors among shift workers in China. *Sleep Med* 2021;81:451-56. doi: 10.1016/j.sleep.2021.03.008 [published Online First: 2021/04/14]

32. Hum IMICH. Night shift work. Geneva, Switzerland: WHO Press 2020:1–371.

33. Li H, Wang YJ, Tan K, et al. Prevalence and risk factors of fatty liver disease in Chengdu, Southwest China. *Hepatobiliary Pancreat Dis Int* 2009;8(4):377-82. [published Online First: 2009/08/12]

34. Golabi P, Paik JM, AlQahtani S, et al. Burden of non-alcoholic fatty liver disease in Asia, the Middle East and North Africa: Data from Global Burden of Disease 2009-2019. *J Hepatol* 2021;75(4):795-809. doi: 10.1016/j.jhep.2021.05.022 [published Online First: 2021/06/04]

BMJ Open

35. Younossi ZM. Non-alcoholic fatty liver disease - A global public health perspective. *J Hepatol* 2019;70(3):531-44. doi: 10.1016/j.jhep.2018.10.033 [published Online First: 2018/11/12]

36. Ising H, Kruppa B. Health effects caused by noise: evidence in the literature from the past 25 years. *Noise Health* 2004;6(22):5-13. [published Online First: 2004/04/09] 37. Cui B, Li K, Gai Z, et al. Chronic Noise Exposure Acts Cumulatively to Exacerbate Alzheimer's Disease-Like Amyloid- β Pathology and Neuroinflammation in the Rat Hippocampus. *Sci Rep* 2015;5:12943. doi: 10.1038/srep12943 [published Online First: 2015/08/08]

38. Shin S, Bai L, Oiamo TH, et al. Association Between Road Traffic Noise and Incidence of Diabetes Mellitus and Hypertension in Toronto, Canada: A Population-Based Cohort Study. *J Am Heart Assoc* 2020;9(6):e013021. doi: 10.1161/jaha.119.013021 [published Online First: 2020/03/10]

39. Wen-jing D Z-dW, Xiang-bin Z, et al. Effects and mechanism of short-term day and night noise exposure on glucose and lipid metabolism in mice. *Modern Preventive Medicine* 2022;49(20):3678-84. doi: 10.20043/j.cnki.MPM.202205073

40. Evans GW, Lercher P, Meis M, et al. Community noise exposure and stress in children. *J Acoust Soc Am* 2001;109(3):1023-7. doi: 10.1121/1.1340642 [published Online First: 2001/04/17]

41. Wang H, Gu Y, Zheng L, et al. Association between bedtime and the prevalence of newly diagnosed non-alcoholic fatty liver disease in adults. *Liver Int* 2018;38(12):2277-86. doi: 10.1111/liv.13896 [published Online First: 2018/06/01]

42. Sun M, Feng W, Wang F, et al. Meta-analysis on shift work and risks of specific obesity types. *Obes Rev* 2018;19(1):28-40. doi: 10.1111/obr.12621 [published Online First: 2017/10/05]

43. Choi Y-H, Lee J-Y, Moon KW. Exposure to volatile organic compounds and polycyclic aromatic hydrocarbons is associated with the risk of non-alcoholic fatty liver disease in Korean adolescents: Korea National Environmental Health Survey (KoNEHS) 2015–2017. *Ecotoxicol Environ Saf* 2023;251:114508. doi: https://doi.org/10.1016/j.ecoenv.2023.114508

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

44. VoPham T. Environmental Risk Factors for Liver Cancer and Nonalcoholic Fatty Liver Disease. *Current Epidemiology Reports* 2019;6(1):50-66. doi: 10.1007/s40471-019-0183-2

45. Werder EJ, Beier JI, Sandler DP, et al. Blood BTEXS and heavy metal levels are associated with liver injury and systemic inflammation in Gulf states residents. *Food Chem Toxicol* 2020;139:111242. doi: <u>https://doi.org/10.1016/j.fct.2020.111242</u>
46. Lang AL, Beier JI. Interaction of volatile organic compounds and underlying liver disease: a new paradigm for risk. *Biol Chem* 2018;399(11):1237-48. doi:

doi:10.1515/hsz-2017-0324

Figure 1. The relationship between shift work and NAFLD.

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, highfat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, benzene series, work exposed to noise.

Page 28 of 29

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

BMJ Open

3
4
5
6
7
/
8
9
10
11
12
13
14
15
15
16
17
18
19
20
21
22
∠∠ 22
23
24
25
26
27
28
29
20
20
31
32
33
34
35
36
37
38
20
27
40
41
42
43
44
45
46
47
47
40
49
50
51
52
53
54
55
55
50
5/
58
59

60



Supplementary

Table 1 Basic characteristics of population in different disease states (n, %)

Chamatanistica	NA	Total	ת	
Characteristics	No (N=3309)	Yes (N=1363)	(N=4672)	P
Drinking status (times per week), n (%)				0.382
Nondrinkers	2124 (64.2)	838 (61.5)	2962 (63.4)	
1~2	1091 (33.0)	484 (35.5)	1575 (33.7)	
3~4	65 (2.0)	28 (2.1)	93 (2.0)	
>4	29 (0.9)	13 (1.0)	42 (0.9)	
High-salt foods (times per week), n (%)				0.135
Seldom	1367 (41.3)	540 (39.6)	1907 (40.8)	
1~2	1749 (52.9)	724 (53.1)	2473 (52.9)	
3~4	144 (4.4)	67 (4.9)	211 (4.5)	
>4	49 (1.5)	32 (2.3)	81 (1.7)	
High-fat foods (times per week), n (%)				0.121
Seldom	984 (29.7)	367 (26.9)	1351 (28.9)	
1~2	2138 (64.6)	901 (66.1)	3039 (65.0)	
3~4	152 (4.6)	76 (5.6)	228 (4.9)	
>4	35 (1.1)	19 (1.4)	54 (1.2)	
Physical activity (times per week), n (%)				0.441
Never	871 (26.3)	359 (26.3)	1230 (26.3)	
1~2	2187 (66.1)	915 (67.1)	3102 (66.4)	
≥3	251 (7.6)	89 (6.5)	340 (7.3)	
Importance of noise protection				0 451
equipment, n (%)				0.451
Unimportant	63 (1.90)	24 (1.76)	87 (1.86)	
Unknown	72 (2.18)	20 (1.47)	92 (1.97)	
General	1718(51.92)	714 (52.38)	2432(52.05)	
Extremely	1456(44.00)	605 (44.39)	2061(44.11)	

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

	^				
CNE [dB(A)·year]	Model 1		Model 2		
	OR 95%CI	Р	OR 95%CI	Р	
Age					
≤29 yr					
<85	Ref				
85~	1.21(0.96, 1.53)	0.103	1.10 (0.86, 1.41)	0.443	
90~	1.19(0.91, 1.56)	0.209	1.05 (0.79, 1.40)	0.728	
95~	1.68(1.04, 2.66)	0.029	1.52 (0.93, 2.44)	0.088	
≥30 yr					
<85	Ref				
85~	1.07 (0.59, 1.94)	0.824	1.05 (0.58, 1.93)	0.863	
90~	1.03 (0.59, 1.82)	0.906	1.00 (0.57, 1.78)	0.989	
95~	1.29 (0.72, 2.33)	0.389	1.28 (0.71, 2.31)	0.419	

 Table 2
 The relationship between CNE and NAFLD in different age groups

Model 1: Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB;

Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, benzene series, shift work.

, shn w

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

		Benzene series (Yes or No)				
Variable	No		Yes			
	OR 95%CI	Р	OR 95%CI	Р		
Work exposed to noise						
(Yes or No)						
No	Ref					
Yes	0.96 (0.80, 1.16)	0.695	0.71 (0.43, 1.17)	0.182		
CNE [dB(A)·year]						
<85	Ref					
85~	1.19 (0.94, 1.52)	0.147	0.78 (0.42, 1.42)	0.419		
90~	1.14 (0.88, 1.48)	0.304	0.84 (0.42, 1.67)	0.624		
95~	1.44 (1.04, 2.00)	0.029	1.64 (0.66, 3.98)	0.279		
Shift work (Yes or No)						
No	Ref					
Yes	1.42 (1.13, 1.79)	0.003	1.07 (0.59, 2)	0.820		
Cumulative length of						
night shift work (hours)					
≤730	Ref					
730~	1.43 (1.12, 1.84)	0.004	1.37 (0.73, 2.56)	0.323		
1460~	1.55 (1.21, 1.98)	< 0.001	0.94 (0.46, 1.88)	0.852		
2920~	1.74 (1.29, 2.35)	< 0.001	1.24 (0.45, 3.28)	0.675		

 Table 3
 The relationship between occupational noise exposure, shift work and NAFLD under different benzene series exposure states

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, shift work/ work exposed to noise.

Association of Occupational Noise Exposure and Shift Work with Non-alcoholic Fatty Liver Disease: A cross-sectional study of male workers in Chinese automobile manufacturing industry.

Journal:	BMJ Open
Manuscript ID	bmjopen-2024-085753.R2
Article Type:	Original research
Date Submitted by the Author:	03-Feb-2025
Complete List of Authors:	Jinwei, ZHANG; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Guangzhou Zhang, Yuxia; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Qiu, Cong; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Wenfeng, ZENG; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Yanmei, RUAN; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Yanmei, RUAN; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Gao , Yun; Guangzhou Medical University, School of Public Health Ma , Wei; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Wu , Kang; Guangzhou Medical University, School of Public Health Xhang , Jing; Guangdong Pharmaceutical University, School of Public Health Zhang , Jing; Guangdong Pharmaceutical University, School of Public Health Cui, Jiaxin; Sun Yat-Sen University, School of public health Ye, Cui; Guangzhou Red Cross Hospital, Department of Preventive Health Liang, Jiabin; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health Wang, Zhi; Guangzhou Twelfth People's Hospital, Key Laboratory of Occupational Environment and Health
Primary Subject Heading :	Occupational and environmental medicine
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine
Keywords:	Risk Factors, PUBLIC HEALTH, OCCUPATIONAL & INDUSTRIAL MEDICINE, EPIDEMIOLOGIC STUDIES, Cross-Sectional Studies



Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our <u>licence</u>.

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which <u>Creative Commons</u> licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

terez onz

BMJ Open

Association of Occupational Noise Exposure and Shift Work with Non-alcoholic Fatty Liver Disease: A cross-sectional study of male workers in Chinese automobile manufacturing industry.

Jinwei Zhang¹⁺, Zhang Yuxia¹⁺, Qiu Congxi¹, Wenfeng Zeng¹, Yanmei Ruan¹, Gao Yunxia², Ma Weiyu¹, Wu Kangyong², Zhang Jingwen³, Cui Jiaxin⁴, Ye Cuiping⁵, Liang Jiabin¹, Wang Zhi^{1#}

¹Key Laboratory of Occupational Environment and Health, Guangzhou Twelfth People's Hospital, 1 Tianqiang St., Huangpu West Ave., Guangzhou, Guangdong, China 510620

²School of Public Health, Guangzhou Medical University, Guangzhou, Guangdong, China 511436

³School of Public Health, Guangdong Pharmaceutical University, Guangzhou, Guangdong, China 510080

⁴School of Public Health, Sun Yat-sen University, Guangzhou, Guangdong, China 510310

⁵Department of Preventive Health, Guangzhou Red Cross Hospital, Guangzhou Guangdong, China 510220

[#]Corresponding author: Zhi Wang, PhD, MD, Professor

Tel.: +86-20-38981282, Mobile: +86-13434163611

email: zhi_wang@outlook.com

⁺These authors have contributed equally to this work.

Abstract

Objective This study aimed to determine the relationship between occupational noise, shift work, and nonalcoholic fatty liver disease(NAFLD) in male workers in the automobile manufacturing industry.

Design Cross-sectional study.

Setting This study was carried out Guangzhou Twelfth People's Hospital, using data from April to September 2022.

Participants A total of 4672 eligible participants were included in the study.

Primary and secondary outcome measures Diagnosis of NAFLD was made using ultrasound. Noise was detected according to the Measurement of Physical Factors in the Workplace-Part 8: Noise. Environmental noise intensity was assessed using an EDGE personal noise dosimeter manufactured by CASELLA (UK). The working status of workers were investigated by questionnaire.

Results The OR of NAFLD was 1.39 (1.03, 1.88) in CNE \geq 95 group compared to CNE <85 group. Improved risk of NAFLD in workers with shift work compared to those without shift work (OR=1.35, 95%CI: 1.09-1.68). As stratified analyses showed, the ORs of NAFLD prevalence related to occupational noise and shift work exposure appears to be increased in young workers. When both shift work and noise exposure work are present simultaneously, the synergy index (SI) between them was 0.47 (95%CI: 0.25 to 0.89). Combined effects analysis revealed that the ORs of NAFLD was 2.02 (95%CI: 1.34-2.99) in CNE \geq 95 and cumulative length of night shifts work >2920 hours.

Conclusion Occupational noise exposure may be an independent risk factor for NAFLD. And it may synergistically affect disease when combined with night shift work, particularly among younger workers. These findings underscore the importance for companies to prioritize the management and training of younger workers, along with targeted occupational health education initiatives, as crucial measures for reducing the incidence of NAFLD.

STRENGTHS AND LIMITATIONS OF THIS STUDY

 \Rightarrow In this study, a cross-sectional survey was used to analyze the effects of occupational noise exposure and shift work on NAFLD. Data were collected by trained investigators, and health examinations were provided to the participants by professional physicians.

 \Rightarrow The questionnaire items on working hours, lifestyle, diet and so on, were designed simplistically and may be difficult to assess comprehensively.

 \Rightarrow Due to the cross-sectional survey, there may be observational bias and causality cannot be elucidated either.

Introduction

In recent decades, there has been a progressive increase in the incidence of NAFLD, leading to a significant burden of morbidity and mortality associated with liver-related disorders¹². Anticipating future trends, researchers project a continued rise in the prevalence of NAFLD, while expecting a stable or possibly decreasing prevalence of other chronic liver diseases³. In China, the prevalence of NAFLD has notably increased in the past decade, rising from 18% to 29.2%. Industrialization and changes in lifestyle have contributed to this rise, resulting in a 10% increase in national incidence⁴. Recent studies have highlighted the rapid increase in NAFLD prevalence among adolescents⁵.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

The development of non-alcoholic fatty liver disease (NAFLD) is influenced by various factors, including age, gender, type 2 diabetes, metabolic syndrome, its components, and family history of fatty liver⁶⁻¹⁰. Age and gender also play significant roles in the onset of NAFLD. The increasing prevalence of NAFLD can be attributed to population aging, where age-related visceral fat accumulation may contribute to the secretion of inflammatory factors^{7 8}. Unhealthy dietary habits, including high-fat diets, and unhealthy lifestyles such as late-night activities have also been associated with an increased risk of NAFLD¹¹⁻¹³. Furthermore, occupational factors in the workplace, such as noise exposure and shift work, may influence the risk of NAFLD. Previous research has demonstrated a significant dose-response relationship between cumulative noise exposure (CNE) and the detection rates of fatty liver¹⁴. Shift work is a prevalent practice in

various industries, including manufacturing, pharmaceutical, and service sectors. It not only disrupts normal circadian rhythms directly but is also closely linked to sleep disorders, leading to insulin resistance and stress reactions, causing physiological disruptions and harm to health¹⁵⁻¹⁷. These disturbances can disrupt glucose and lipid metabolism, which is a significant contributing factor in elevating the risk of NAFLD¹⁸.

Workers in the automotive manufacturing industry frequently endure prolonged occupational noise exposure and may also be required to engage in shift work. Despite this, few investigations have specifically examined the impact of shift work and noise on NAFLD within the context of this workforce. Consequently, our study aims to investigate the effects of occupational noise exposure and shift work on NAFLD among male workers in the automotive manufacturing industry. By focusing on this specific population, the study intends to provide a theoretical foundation to implement measures to protect against occupational hazards.

Methods

Study design and population

From April to September 2022, a cross-sectional study was conducted among male workers at an automobile manufacturing company in Guangzhou for an occupational health examination. Participants with incomplete medical examination or questionnaire data, female workers, individuals taking antiretroviral drugs, and those with a history of cancer, excessive alcohol intake, hepatobiliary disease, blood system diseases, or renal failure were excluded. Ultimately, a total of 4672 eligible participants were included in the study. This study was reviewed and approved by the Ethics Committee of Guangzhou Twelfth People's Hospital (No.2021008) and all procedures involving participants were conducted in accordance with the ethical standards of the Institutional Research Board and the 1964 Declaration of Helsinki. All study participants gave informed consent.

NAFLD diagnosis and collection of information on covariates

Trained physicians conducted the occupational health examination in accordance with a standardized protocol. Diagnosis of NAFLD was made using the criteria proposed by FARRELL et al¹⁹. Blood was drawn from the fasting subjects to measure

hemoglobin(HGB) and fasting plasma glucose(FPG) levels. Blood pressure was measured by the omron HEM-7071 electronic blood pressure monitor three times, each time at an interval of 1 min, and the average of the three measurements was taken as the individual blood pressure (BP) value. BP was classified according to 2018 Chinese Guidelines for Prevention and Treatment of Hypertension. FPG normal range is 3.9-6.1mmol/L. According to the Guidelines for the Prevention and Control of Overweight and Obesity in Chinese Adults, our study the included BMI<24kg/m² and BMI \geq 24kg/m² as dichotomous variables in the multivariate logistic regression model.

Referring to previous research and consulting multiple experts in the field of occupational health to revise and improve it continuously, we used a self-designed questionnaire to investigate²⁰. The questionnaires were administered face to face by trained investigators to acquire participants' information, including dietary habits, lifestyle, occupational information and past medical history. Finally, investigators checked the completed questionnaires to detect errors promptly. We investigated the average number of working days per week for workers to assess working hours and determine whether it is considered as long working hours. Long working hours are defined as working more than 40 hours per week²¹. Simultaneously, we made inquiries regarding the exposure statuses of other occupational hazard factors, so as to guarantee that the exposures to other occupational factors like dust and benzene series (mainly refers to benzene, toluene and xylene) were beneath the occupational exposure limits (OELs) stipulated in "Occupational exposure limits for hazardous agents in the workplace-Part 1: Chemical hazardous agents" (GBZ 2.1-2019). Classification of dietary habits: high-salt foods mainly include pickled vegetables, ham and bacon, and so on^{22 23}; high-fat foods including chips, chocolate, animal offal and fried foods; late evening snack is defined as an additional meal arranged after 21:00. According to the frequency of eating within a week, they are divided into four groups: seldom, 1~2 times per week, 3~4 times per week, and >4 times per week. Smoking is defined as a consistent consumption of at least 1 cigarette per day for a duration of more than 6 months. Alcohol consumption is defined as a regular pattern of drinking at least once a week for a period of 1 year or longer. According to the duration of sleep, they were Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

divided into three groups: < 6h (short sleep duration), $6 \sim 8h$ (moderate sleep duration), and >8h (long-sleep duration)²⁴.

Noise exposure assessment

Our study followed the "Measurement of Physical Factors in the Workplace - Part 8: Noise" (GBZ/T189.8-2007) for noise detection. Environmental noise intensity was assessed using an EDGE personal noise dosimeter manufactured by CASELLA (UK). Each monitoring point was measured three times and the average value was taken as the measurement result. In this study, noise exposure was evaluated using the normalization of continuous A-weighted sound pressure level equivalent to an 8h per day (LAeq, 8h), and work exposed to noise is defined as those operations involving the presence of hearing-impairing, health-hazardous or otherwise hazardous sound, and LAeq,8h \geq 80 dB(A) for at least one year. Meanwhile, Cumulative noise exposure (CNE) was used to quantify the noise exposure level of the study participants and it is calculated as: CNE=L_{Aeq, 8h}+10logT, where T indicates years of noise exposure and the CNE unit is: dB(A)·year.

Assessment of shift work

The International Labour Organization defines shift work as extending the working day to 24 hours by alternating between different workers and/or teams^{25 26}. Night shift work is defined as work occurring between 0:00 and 5:00 am^{26 27}. The cumulative length of night shift work was calculated in hours during data processing. In this study, the two main types of shift patterns for workers are two shifts system and three shifts system.

In addition, according to the sampling methods and technical requirements in "Specifications of air sampling for hazardous substances monitoring in the workplace" (GBZ 159-2004), collect samples of dust, benzene, toluene, and xylene in the air of workplaces. In accordance with the requirements of "Determination of dust in the air of workplace Part 1: Total dust concentration" (GBZ 192.1-2007) and "Occupational exposure limits for hazardous agents in the workplace-Part 1: Chemical hazardous agents" (GBZ 2.1-2019), we conduct fixed-point sampling and detection on representative operating posts for dust, benzene, toluene, and xylene in the air.

Statistical analyses

Page 9 of 29

BMJ Open

Subject staff verified data daily to ensure unique ID codes for each individual and timely identification of missing key data and logical error. R (version 4.2.1) was used for all statistical analyses. Continuous variables that followed normal distribution were presented as mean ±standard deviation (SD), and were compared by t-test. Mann-Whitney U test was used to compare continuous variables with skewed distribution in the form of median with interquartile ranges. The χ^2 test was used to compare categorical variables, which have been expressed as numbers (percentages). The relationship between shift work, occupational noise and NAFLD was analyzed using a multivariate logistic regression model, with night shift hours and CNE included in the regression model as categorical variables respectively as well as the joint effect of both was analysed in relation to NAFLD. Referring to the national criteria (GBZT229.4-2012), CNE is divided into four groups <85, 85~, 90~ and 95~. Cumulative length of night shift work are grouped by quartiles, using the first quartile as a reference. Previous studies have considered adolescents and young adults (AYAs) to be the group aged 15 years and older, with the cut-offs at 24, 29 and 39 years being difficult to determine and some studies considering 18-24 years as late adolescence²⁸. We stratified the age into two groups ≤ 29 and ≥ 30 for analysis. To provide more clarity on the effect of CNE on the illness of workers in different age groups, we subdivided younger workers at age 29 into two groups (\leq 24 and 25-29 years old)^{29 30}. All test were two-sided, and α =0.05 was considered statistically significant.

Patient and public involvement

Patients and the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Results

Basic information about the study population

A total of 5928 questionnaires were distributed, out of which 4791 valid questionnaires were returned, resulting in a valid response rate of 80.8%. After excluding individuals who did not meet the study's inclusion criteria, and those with incomplete occupational health screening results (n=119), the final sample size included for analysis was determined. Overall, 4672 workers with complete information on occupational health

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

examination and questionnaires were included in the study according to the inclusion and exclusion criteria. The mean age of the study population was (26.8±5.0), with different prevalence rates for workers in the \leq 29 and \geq 30 years age groups (*P*<0.001), with a total of 1363 workers with NAFLD and a prevalence of 29.2%. There were 3658 (78.3%) shift workers, with different prevalence of NAFLD in different shift work situations (*P*<0.001). Meanwhile, 2965 (63.5%) noise workers had a statistically significant difference in prevalence in different noise exposure scenarios (*P*<0.05). Cumulative length of night shift work and CNE vary by disease state (all *p*<0.05). A summary of the sample characteristics is presented in Table 1. The results of univariate analysis of drinking status, high-salt foods, high-fat foods, physical activity and importance of noise protection equipment are presented in Supplementary Table 1.

Table 1 Basic characteristics of population	pulation in different disease states (r	1, %)
---	---	------	---

Characteristics	NAI	NAFLD			
	No (N=3309)	Yes (N=1363)	(N=4672)	Г	
Age (years), n (%)				< 0.001	
≤29	2649 (80.1)	698 (51.2)	3347 (71.6)		
≥30	660 (19.9)	665 (48.8)	1325 (28.4)		
Education, n (%)				0.009	

Senior secondary and below	1888 (57.1)	720 (52.8)	2608 (55.8)	
Higher education	1421 (42.9)	643 (47.2)	2064 (44.2)	
Smoking status (cigarettes per day), n (%)				< 0.001
Nonsmokers	1741 (52.6)	729 (53.5)	2470 (52.9)	
1~5	618 (18.7)	198 (14.5)	816 (17.5)	
6~10	599 (18.1)	239 (17.5)	838 (17.9)	
>10	351 (10.6)	197 (14.5)	548 (11.7)	
Fruit and vegetable (times per week), n (%)				0.002
Seldom	262 (7.9)	89 (6.5)	351 (7.5)	
1~2	1961 (59.3)	752 (55.2)	2713 (58.1)	
3~4	653 (19.7)	297 (21.8)	950 (20.3)	
>4	433 (13.1)	225 (16.5)	658 (14.1)	
Late evening snack (times per week), n (%)				0.024
Seldom	747 (22.6)	361 (26.5)	1108 (23.7)	
1~2	1863 (56.3)	735 (53.9)	2598 (55.6)	
3~4	386 (11.7)	137 (10.1)	523 (11.2)	
>4	313 (9.5)	130 (9.5)	443 (9.5)	
BMI (kg/m2), n (%)				< 0.001
<24	3001 (90.7)	555 (40.7)	3556 (76.1)	
≥24	308 (9.3)	808 (59.3)	1116 (23.9)	
Sleep duration (hours),n (%)				< 0.001
<6	592 (17.9)	303 (22.2)	895 (19.2)	
6~8	2574 (77.8)	1023 (75.1)	3597 (77.0)	
>8	143 (4.3)	37 (2.7)	180 (3.9)	
BP (mmHg), n (%)				< 0.001
Normal	1374 (41.5)	306 (22.5)	1680 (36.0)	
High-normal	1890 (57.1)	963 (70.7)	2853 (61.1)	
Hypertension	45 (1.4)	94 (6.9)	139 (3.0)	
ECG, n (%)				< 0.001
Normal	1392 (42.1)	824 (60.5)	2216 (47.4)	
Abnormal	1917 (57.9)	539 (39.5)	2456 (52.6)	
FPG (mmol/L), n (%)				< 0.001
Normal	2887 (87.2)	1001 (73.4)	3888 (83.2)	
Abnormal	422 (12.8)	362 (26.6)	784 (16.8)	

Table 1 Continued

Characteristics	NA	FLD	$T_{abal}(N - 4(72))$	מ
	No (N=3309)	Yes (N=1363)	-10tal(N-40/2)	P
HGB (g/L), n (%)				0.005
Normal	2586 (78.2)	1013 (74.3)	3599 (77.0)	
Abnormal	723 (21.8)	350 (25.7)	1073 (23.0)	
Long working hours				< 0.001

No	1375 (41.55)	685 (50.26)	2060 (44.09)	
Yes	1934 (58.45)	678 (49.74)	2612 (55.91)	
Dust				0.015
No	2990 (90.36)	1262 (92.60)	4252 (91.01)	
Yes	319 (9.64)	101 (7.41)	420 (8.99)	
Benzene series				0.02
No	2828 (85.46)	1200 (88.04)	4028 (86.22)	
Yes	481 (14.54)	163 (11.96)	644 (13.78)	
Shift work				< 0.001
No	808 (24.42)	206 (15.11)	1014 (21.70)	
Yes	2501 (75.58)	1157 (84.89)	3658 (78.30)	
Work exposed to noise				0.045
No	1239 (37.44)	468 (34.34)	1707 (36.54)	
Yes	2070 (62.56)	895 (65.66)	2965 (63.46)	
Cumulative length of night shift work (hours)	1100 (511-2560)	2190 (1090-4380)	1460 (730-2920)	< 0.001
CNE [dB(A) year]	86.9 (83.1-91.1)	89.8 (85.2-93.3)	87.8 (83.6-91.9)	< 0.001
Seniority in shift work (year), median (IQR)	3.0 (1.50, 6.50)	6.0 (3.95, 12.0)	4.0 (2.0, 8.0)	< 0.001
Seniority in work exposed to noise (year), median (IQR)	3.00 (1.30, 7.00)	7.90 (2.75,13.50)	3.8 (1.50, 8.33)	< 0.001

The relationship between occupational noise exposure and NAFLD

The impact of work exposed to noise on NAFLD was demonstrated in Table 2. Compared with those did not, no statistically significant relationship between work exposed to noise and NAFLD (OR=0.92, 95%CI: 0.78-1.09). Table 3 and Supplementary Table 2 presented the odds ratios (ORs) and 95% confidence intervals (CIs) for the effect of occupational noise exposure on NAFLD. Stratified analysis indicated that the association between CNE and NAFLD was not statistically significant in both those aged \leq 29 and \geq 30 years (see supplemental Table 1); By reference to CNE<85, the OR of NAFLD was 1.39 (95%CI: 1.03, 1.88) in CNE \geq 95 group. Table 3 shows further stratification by age, among those \leq 24 years of age, the OR of NAFLD in the CNE \geq 95 group was 2.97 (95%CI: 1.18-6.95).

Table 2 Association of work exposed to noise with NAFLD

work exposed to noise (Yes or No)	Model 1		Model 2		
	OR 95%CI	Р	OR 95% <i>CI</i>	Р	
Total					
No	Ref				
Yes	0.97 (0.83, 1.15)	0.754	0.92 (0.78, 1.09)	0.356	

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Age <29 vr				
No	Ref			
Yes	1.07 (0.87, 1.31)	0.537	0.98 (0.79, 1.21)	0.826
≥30 yr				
No	Ref			
Yes	0.84 (0.63, 1.11)	0.212	0.83(0.63, 1.11)	0.212

Model 1: Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB; Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, benzene series, shift work.

CNE [dP(A), wear]	Model 1		Model 2]
	OR 95%CI	Р	OR 95% <i>CI</i>	Р
Total				
<85				
85~	1.20 (0.97, 1.48)	0.087	1.12 (0.90, 1.39)	0.320
90~	1.16 (0.93, 1.45)	0.191	1.08 (0.85, 1.36)	0.536
95~	1.48 (1.10, 1.99)	0.009	1.39 (1.03, 1.88)	0.033
Age				
≤24 yr				
<85	Ref			
85~	1.07 (0.77, 1.48)	0.676	0.96 (0.68, 1.35)	0.810
90~	1.00 (0.64, 1.52)	0.988	0.86 (0.54, 1.34)	0.512
95~	3.18 (1.29, 7.31)	0.008 🧹	2.97 (1.18, 6.95)	0.015
25~29 yr				
<85	Ref			
85~	1.14 (0.80, 1.62)	0.480	1.14 (0.78, 1.66)	0.505
90~	1.11 (0.76, 1.62)	0.603	1.08 (0.73, 1.62)	0.690
95~	1.01 (0.55, 1.81)	0.963	1.07 (0.57, 1.95)	0.825

Table 3 The relationship between CNE and NAFLD in different age groups

Model 1: Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB; Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, benzene series, shift work.

The relationship between shift work and NAFLD

The effect of shift work on NAFLD is shown in the Figure 1. Compared to those without shift work, the OR of NAFLD among workers with shift work was 1.35 (95%CI: 1.09-1.68). Stratified analysis showed that shift work was a risk factor for NAFLD among

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

those aged ≤ 29 years (OR=1.46, 95%CI: 1.15-1.87). Similarly, in comparison with cumulative length of night shift work ≤ 730 h, rising cumulative length of night shift work were associated with a higher risk of NAFLD, with ORs (95%CIs) of 1.43 (1.14, 1.79), 1.44 (1.15, 1.81) and 1.64 (1.23, 2.17), respectively. Among those aged ≤ 29 years, increased cumulative length of night shift work were also a risk factor for NAFLD, with ORs (95%CIs) of 1.66 (1.30, 2.11), 1.52 (1.17, 1.98) and 2.41 (1.50, 3.84) respectively.

The relationship between the interaction of work exposed to noise and shift work and NAFLD

To further explore the combined effects of noise exposure work and shift work, we analyzed the interaction between them. The results were shown in Table 4. When both shift work and noise exposure work are present simultaneously, the excess relative risk of relative excess risk due to interaction (RERI) was -0.43 (95%CI: -1.00 to 0.14), the attributable proportion due to interaction (AP) was -0.31 (-0.69 to 0.07), and the synergy index (SI) between them was 0.47 (95%CI: 0.25 to 0.89).

Table 4	The relationship	between the	interaction	of wo	rk exposec	to noise ar	nd shift v	work and	
NAFLD									

work exposed	shift work	Total					
to noise (Yes or No)	(Yes or No)	OR 95% <i>CI</i>	RERI	AP	SI	$P_{interaction}$	
No	No	Ref	-0.43 (-1.00, 0.14)	-0.31 (-0.69, 0.07)	0.47 (0.25, 0.89)	0.112	
	Yes	1.60 (1.19, 2.17)					
Yes	No	1.21 (0.83, 1.75)					
	Yes	1.38 (1.05, 1.83)					

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, benzene series.

The combined effect of occupational noise and shift work in association with NAFLD

The combined effect of occupational noise and cumulative length of night work on NAFLD was shown in Table 5. After adjusting for variables, the association with NAFLD was statistically significant for all groups at cumulative length of night shift

BMJ Open

work >2920h and CNE>85, when compared with CNE<85 and cumulative length of night shift work \leq 730h. The OR of NAFLD was greatest at 2.01 (95%CI: 1.34 to 2.99) when cumulative length of night shift work >2920h and CNE \geq 95. Furthermore, we analyzed the relationships between occupational noise, shift work and NAFLD in different states of benzene series exposure as well (see Supplementary Table 3).

CNE [dB(A)∙years]	Cumulative length of night shift work (hours)					
	≤730	730~	1460~	2920~		
	OR 95%CI	OR 95%CI	OR 95%CI	OR 95% <i>CI</i>		
<85	Ref	1.49 (1.04, 2.12)*	1.27 (0.82, 1.94)	1.34 (0.64, 2.76)		
85~	0.88 (0.6, 1.27)	1.45 (1.05, 1.99)*	1.61 (1.14, 2.26)**	1.99 (1.29, 3.07)**		
90~	1.24 (0.76, 1.97)	1.31 (0.85, 1.99)	1.41 (1.00, 1.98) *	1.54 (1.08, 2.21)*		
95~	1.98 (0.89, 4.26)	1.61 (0.72, 3.48)	1.67 (0.95, 2.88)	2.01 (1.34, 2.99)***		

Table 5	Association o	f CNE and	d cumulative	length o	of night	shift	work with	NAFLD
---------	---------------	-----------	--------------	----------	----------	-------	-----------	-------

*p<0.05, **p<0.01, ***p<0.001.

Bold values are p<0.05.

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, benzene series.

Discussion

To summarize, our study included a total of 4672 male workers in the automotive manufacturing industry, investigating the impact of occupational noise and shift work on non-alcoholic fatty liver disease (NAFLD) through a cross-sectional survey. The majority of the company's employees are subjected to a high workload, with 63.5% working in noise and 78.3% working in shifts, which is significantly higher compared to other occupations such as teachers and white-collar workers³¹. Night work has been categorized by the International Agency for Research on Cancer as "possibly carcinogenic to humans" (Group 2A)³², prompting growing concerns about its adverse effects on health. The prevalence of NAFLD in this survey was 29.2%, a rate close to the combined prevalence estimate of NAFLD reported by Fan et al^{11 33-35}. Among those exposed to noise and those not exposed, the prevalence of NAFLD is 30.2% and 27.4%, respectively. After adjusting for relevant variables, multifactorial logistic regression analysis revealed that increased CNE was the significant risk factor associated with NAFLD. This may be because the biological effects of noise are cumulative and the
BMJ Open

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

effects of a single noise exposure are not sufficient to cause health damage to the organism. However, the more repeated exposures and the longer the exposure period, the more pronounced the cumulative effects will be when the effects are not eliminated^{36 37}. Associations between CNE and NAFLD vary across age groups. An increase in CNE is a risk factor for NAFLD among workers under 24 years of age. This suggests that short-term exposure to high-intensity noise may contribute to the development of NAFLD in this population. Possible factors that may contribute to this association include the inadequate use of noise protective equipment among young workers or exposure to noise outside of the occupational setting. The effect of noise on NAFLD in younger workers should be further explored. This indicates that enterprises should strengthen training and occupational health education for young workers, which is important to reduce the occurrence of NAFLD. Shins, Saeha et al. have shown that chronic exposure to road traffic noise was associated with an increased incidence of diabetes and hypertension in Toronto³⁸. Du et al. found that noise exposure could lead to disorders of glucolipid metabolism in mice through inhibition of the AKT pathway mediated by the biological clock gene BMAL1³⁹. The experiment by Evans et al. also demonstrated significant changes in serum ALT levels across test groups, suggesting that noise can lead to liver dysfunction⁴⁰. It can be seen that noise may be the basis for the formation of NAFLD by promoting insulin resistance and disorders of glucolipid metabolism in the body, resulting in metabolic abnormalities in the body. These studies also indirectly support our findings. In this study, the prevalence of NAFLD was 20.3% and 31.6% among auto manufacturing workers not involved in shift work and those who did, respectively, with a higher prevalence among workers in shift work, similar to the findings of Golabi et al¹¹. We observed that shift work and an increase in cumulative length of night shift work were identified as risk factors for NAFLD. We also examined the relationship between shift work and NAFLD across different age groups. Specifically, among workers below 29 years of age, engaging in shift work was found to significantly increase the risk of NAFLD. This indicates the presence of specific physiological adaptations that contribute to this heightened risk. Similarly, we found that an extended cumulative length of night shift work was a significant risk

Page 17 of 29

BMJ Open

factor for NAFLD in workers below 29 years of age. These findings emphasize the importance of appropriate management strategies to mitigate the adverse effects of shift work on NAFLD among younger workers. Previous epidemiological studies have consistently demonstrated a link between shift work and NAFLD. For instance, a 4year cohort study conducted among Chinese railway workers reported a higher incidence of NAFLD in individuals engaged among shift work¹¹. Similarly, a crosssectional survey involving male steelworkers found that night shift workers had an increased risk of NAFLD compared to those working day shifts¹⁵. Wang et al. further supported the association between night shifts and NAFLD by demonstrating the impact of disrupted circadian rhythms, due to delayed sleep, on the incidence of NAFLD⁴¹. These findings are in line with our results. Sun et al. also revealed the association between sleep deprivation, increased cortisol levels, and insulin resistance, further highlighting the possible mechanisms underlying the relationship between sleep disruption and NAFLD⁴². Our study identified a significant association between occupational noise exposure, shift work, and an increased risk of developing NAFLD among workers. This association may be explained by the potential mechanisms through which both noise exposure and shift work contribute to the development of insulin resistance, a key factor in the pathogenesis of NAFLD. Insulin resistance disrupts lipid metabolism, leading to elevated lipase activity and subsequent overproduction of triglycerides and free fatty acids. Consequently, an excessive amount of free fatty acids enters the liver, promoting increased lipid synthesis and deposition within hepatic cells. This lipid accumulation interferes with hepatocyte transport capacity, resulting in hepatocyte apoptosis and the accumulation of fat, ultimately leading to NAFLD development.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

It should be noted that some participants in this study were simultaneously exposed to benzene series (mainly benzene, toluene, and xylene). The study by Yun - Hee Cho et al. showed that exposure to chemicals such as benzene, toluene, and xylene could significantly increase the risk of NAFLD⁴³ ⁴⁴. In obese populations, benzene exposure is positively correlated with liver injury, and systemic inflammation increases with the exposure to benzene and toluene, indicating that benzene and toluene exposure

BMJ Open

 have certain potential impacts on steatohepatitis⁴⁵. These effects may result from the activation of cell death, the induction of oxidative and endoplasmic reticulum stress, and the disruption of metabolism, which are caused by the dysfunction of xenobiotic and endobiotic receptors⁴⁶. After detection, we found that the concentrations of benzene series in the researchers' workplaces were all lower than the occupational exposure limits (OELs) stipulated in "Occupational exposure limits for hazardous agents in the workplace - Part 1: Chemical hazardous agents". To ensure the reliability of the results, we also analyzed the relationships between occupational noise, shift work and NAFLD under different benzene series exposure, occupational noise exposure and shift work were still risk factors for NAFLD.

This study contributes to a relatively uncommon and innovative exploration of the association between occupational noise exposure, shift work, and their combined effect on NAFLD among male workers in the automotive manufacturing industry. Based on the study findings, we propose several recommendations for relevant management departments. It is advised to scientifically and reasonably organize the working system, promote the implementation of flexible working arrangements, and minimize the cumulative length of night work. Furthermore, improving the production process to control or eliminate noise sources, such as installing sound insulation and vibration dampers, is recommended. Additionally, establishing and enhancing occupational health supervision, strengthening the management of occupational health, and facilitating regular occupational health check-ups for workers are crucial measures to safeguard worker health. These recommendations hold significant practical importance in protecting the health of workers.

However, it has to acknowledge the limitations of our study. Due to the crosssectional nature of the survey, there may be observational bias and causality cannot be elucidated either. To address these limitations, we are in the process of establishing a cohort study among workers in the automotive manufacturing industry. In future studies, we plan to utilize individual noise dosimeters to more accurately measure workers' noise exposure, enabling a more robust analysis of the association between these occupational factors and disease outcomes.

Acknowledgements

We wish to thank the timely help given by workers who participated in this program. we are deeply grateful for Guangzhou Twelfth People's Hospital for their support for collecting data.

Contributors

Jinwei Zhang, Zhang Yuxia and Wang Zhi conceived and designed the study. Wang Zhi provided administrative support. Jinwei Zhang, Qiu Congxi, Ma Weiyu, Wenfeng Zeng and Wang Zhi provided study materials. Jinwei Zhang, Zhang Yuxia, Qiu Congxi, Wu Kangyong, Zhang Jingwen, and Cui Jiaxin collected and assembled data. Zhang Yuxia, Yanmei Ruan, Gao Yunxia, Liang Jiabin and Ye Cuiping analyzed and interpreted the data. Zhang Yuxia, Jinwei Zhang and Gao Yunxia wrote the manuscript. Wang Zhi and Jinwei Zhang approved the final manuscript. The guarantor of the study is Wang Zhi , accepts full responsibility for the finished work and the conduct of the study, has access to the data and controls the decision to publish.

Fundings

This work was supported by the Key Research and Development Program of Guangzhou Science and Technology Project (Project Number: 202206010061); Guangzhou Key Medical Discipline (2025-2027); The Science and Technology Program of Guangzhou (Project Number: 2023A03J0502, 2023A03J0979, 2024A03J0432, 2024A03J0505, 2025A03J3591); The Medical Science and Technology Research Fund Project of Guangdong (Project Number: B2023347, A2023406, A2023396); The Science and Technology Project of Guangzhou Municipal Health Commission (Project Number: 20251A010042);

Ethics statements

Patient consent for publication

Consent obtained directly from patient(s).

Ethics approval

The study was approved by Guangzhou Twelfth People's Hospital ethics committee(No.

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

2021008), and all procedures involving participants were conducted in accordance with the ethical standards of the Institutional Research Committee and the 1964 Helsinki Declaration. The informed consent form was sent in electronic form to each participant before the survey began, and participants could also request a paper copy. All participants have signed written informed consent.

toppt tellewony

Data availability statement

Data are available upon reasonable request.

Conflicts of interest

The authors declare that they have no competing interests

References

 Notarnicola M, Osella AR, Caruso MG, et al. Nonalcoholic Fatty Liver Disease: Focus on New Biomarkers and Lifestyle Interventions. *Int J Mol Sci* 2021;22(8):3899.
 Masoodi M, Gastaldelli A, Hyötyläinen T, et al. Metabolomics and lipidomics in NAFLD: biomarkers and non-invasive diagnostic tests. *Nat Rev Gastroenterol Hepatol* 2021;18(12):835-56. doi: 10.1038/s41575-021-00502-9 [published Online First: 2021/09/12]

3. Younossi ZM, Stepanova M, Afendy M, et al. The changing face of chronic liver disease (CLD) in the United States: The rising epidemic of non-alcoholic fatty liver disease (NAFLD). *J Hepatol* 2011;54

 Younossi Z, Tacke F, Arrese M, et al. Global Perspectives on Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. *Hepatology* 2019;69(6):2672-82. doi: 10.1002/hep.30251 [published Online First: 2018/09/05]

5. Lin G, Xinhe Z, Haoyu T, et al. Epidemiology and lifestyle survey of non-alcoholic fatty liver disease in school-age children and adolescents in Shenyang, Liaoning. *BMC Pediatr* 2022;22(1):286. doi: 10.1186/s12887-022-03351-w [published Online First: 2022/05/18]

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

6. Younossi ZM, Golabi P, de Avila L, et al. The global epidemiology of NAFLD and NASH in patients with type 2 diabetes: A systematic review and meta-analysis. *J Hepatol* 2019;71(4):793-801. doi: 10.1016/j.jhep.2019.06.021 [published Online First: 2019/07/08]

7. Kim IH, Kisseleva T, Brenner DA. Aging and liver disease. *Curr Opin Gastroenterol* 2015;31(3):184-91. doi: 10.1097/mog.000000000000176 [published Online First: 2015/04/09]

8. Barzilai N, Huffman DM, Muzumdar RH, et al. The critical role of metabolic pathways in aging. *Diabetes* 2012;61(6):1315-22. doi: 10.2337/db11-1300 [published Online First: 2012/05/24]

9. Wong VW, Chu WC, Wong GL, et al. Prevalence of non-alcoholic fatty liver disease and advanced fibrosis in Hong Kong Chinese: a population study using proton-

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

magnetic resonance spectroscopy and transient elastography. Gut 2012;61(3):409-15.

doi: 10.1136/gutjnl-2011-300342 [published Online First: 2011/08/19]

10. Wang J, Tang Y, Peng K, et al. Development and validation of a nomogram for predicting nonalcoholic fatty liver disease in the non-obese Chinese population. *Am J Transl Res* 2020;12(10):6149-59. [published Online First: 2020/11/17]

11. Xu J, Ni S, Wang Y, et al. Shift work and nonalcoholic fatty liver disease incidence among Chinese rail workers: a 4-year longitudinal cohort study. *Int Arch Occup Environ Health* 2023;96(1):179-90. doi: 10.1007/s00420-022-01913-0

Takahashi A, Anzai Y, Kuroda M, et al. Effects of sleep quality on non-alcoholic fatty liver disease: a cross-sectional survey. *BMJ Open* 2020;10(10):e039947. doi: 10.1136/bmjopen-2020-039947 [published Online First: 2020/10/31]

13. Semmler G, Datz C, Reiberger T, et al. Diet and exercise in NAFLD/NASH: Beyond the obvious. *Liver Int* 2021;41(10):2249-68. doi: 10.1111/liv.15024 [published Online First: 2021/07/31]

14. Bai G-p. Relationship Between Occupational Noise Expose and the Fatty Liver. *Hebei Medicine* 2009

15. Zhang S, Wang Y, Wang Z, et al. Rotating night shift work and non-alcoholic fatty liver disease among steelworkers in China: a cross-sectional survey. *Occup Environ Med* 2020;77(5):333-39. doi: 10.1136/oemed-2019-106220 [published Online First: 2020/02/06]

16. Attarchi M, Dehghan F, Safakhah F, et al. Effect of exposure to occupational noise and shift working on blood pressure in rubber manufacturing company workers. *Ind Health* 2012;50(3):205-13. doi: 10.2486/indhealth.ms1321 [published Online First: 2012/03/29]

17. Dong H, Zhang Q, Zhang Z, et al. Association of sleep disturbance with shift work, occupational stress, and musculoskeletal pain in Chinese public hospital emergency workers: A multicentre cross-sectional study. *Chronobiol Int* 2022;39(6):886-94. doi: 10.1080/07420528.2022.2048662 [published Online First: 2022/03/08]

BMJ Open

18. Boivin DB, Boudreau P, Kosmadopoulos A. Disturbance of the Circadian System
in Shift Work and Its Health Impact. J Biol Rhythms 2022;37(1):3-28. doi:
10.1177/07487304211064218 [published Online First: 2022/01/01]

19. Farrell GC, Chitturi S, Lau GK, et al. Guidelines for the assessment and management of non-alcoholic fatty liver disease in the Asia-Pacific region: executive summary. *J Gastroenterol Hepatol* 2007;22(6):775-7. doi: 10.1111/j.1440-1746.2007.05002.x [published Online First: 2007/06/15]

20. Ruan Y, Huang G, Zhang J, et al. Risk analysis of noise-induced hearing loss of workers in the automobile manufacturing industries based on back-propagation neural network model: a cross-sectional study in Han Chinese population. *BMJ Open* 2024;14(5):e079955. doi: 10.1136/bmjopen-2023-079955 [published Online First: 2024/05/18]

21. Pega F, Náfrádi B, Momen NC, et al. Global, regional, and national burdens of ischemic heart disease and stroke attributable to exposure to long working hours for 194 countries, 2000-2016: A systematic analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ Int* 2021;154:106595. doi: 10.1016/j.envint.2021.106595 [published Online First: 2021/05/21]

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

22. Webster JL, Dunford EK, Neal BC. A systematic survey of the sodium contents of processed foods. *Am J Clin Nutr* 2010;91(2):413-20. doi: 10.3945/ajcn.2009.28688 [published Online First: 2009/12/04]

23. Ni Mhurchu C, Capelin C, Dunford EK, et al. Sodium content of processed foods in the United Kingdom: analysis of 44,000 foods purchased by 21,000 households. *Am J Clin Nutr* 2011;93(3):594-600. doi: 10.3945/ajcn.110.004481 [published Online First: 2010/12/31]

24. Imaizumi H, Takahashi A, Tanji N, et al. The Association between Sleep Duration and Non-Alcoholic Fatty Liver Disease among Japanese Men and Women. *Obes Facts* 2015;8(4):234-42. doi: 10.1159/000436997

25. International Labour Organization. C171-Night Work Convention, 1990(No.171)1990 [Available from:

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

https://www.ilo.org/dyn/normlex/en/f?p=NORMLEXPUB:12100:0::NO::P12100_IN STRUMENT_ID:312316.

26. Organization, WH. IARC Monographs Volume 124: Night Shift Work [Available from: <u>https://www.iarc.who.int/news-events/iarc-monographs-volume-124-night-shift-work/</u>.

27. Molzof HE, Peterson CM, Thomas SJ, et al. Nightshift Work and Nighttime Eating Are Associated With Higher Insulin and Leptin Levels in Hospital Nurses. *Front Endocrinol (Lausanne)* 2022;13 doi: 10.3389/fendo.2022.876752

28. Arshad T, Paik JM, Biswas R, et al. Nonalcoholic Fatty Liver Disease Prevalence Trends Among Adolescents and Young Adults in the United States, 2007-2016. *Hepatol Commun* 2021;5(10):1676-88. doi: 10.1002/hep4.1760 [published Online First: 2021/09/25]

29. Paik JM, Kabbara K, Eberly KE, et al. Global burden of NAFLD and chronic liver disease among adolescents and young adults. *Hepatology* 2022;75(5):1204-17. doi: 10.1002/hep.32228 [published Online First: 2021/11/07]

30. Zhang X, Wu M, Liu Z, et al. Increasing prevalence of NAFLD/NASH among children, adolescents and young adults from 1990 to 2017: a population-based observational study. *BMJ Open* 2021;11(5):e042843. doi: 10.1136/bmjopen-2020-042843 [published Online First: 2021/05/06]

31. Wang D, Chen H, Chen D, et al. Shift work disorder and related influential factors among shift workers in China. *Sleep Med* 2021;81:451-56. doi: 10.1016/j.sleep.2021.03.008 [published Online First: 2021/04/14]

32. Hum IMICH. Night shift work. Geneva, Switzerland: WHO Press 2020:1-371.

33. Li H, Wang YJ, Tan K, et al. Prevalence and risk factors of fatty liver disease in Chengdu, Southwest China. *Hepatobiliary Pancreat Dis Int* 2009;8(4):377-82. [published Online First: 2009/08/12]

34. Golabi P, Paik JM, AlQahtani S, et al. Burden of non-alcoholic fatty liver disease in Asia, the Middle East and North Africa: Data from Global Burden of Disease 2009-2019. *J Hepatol* 2021;75(4):795-809. doi: 10.1016/j.jhep.2021.05.022 [published Online First: 2021/06/04]

BMJ Open

35. Younossi ZM. Non-alcoholic fatty liver disease - A global public health perspective. *J Hepatol* 2019;70(3):531-44. doi: 10.1016/j.jhep.2018.10.033 [published Online First: 2018/11/12]

36. Ising H, Kruppa B. Health effects caused by noise: evidence in the literature from the past 25 years. *Noise Health* 2004;6(22):5-13. [published Online First: 2004/04/09] 37. Cui B, Li K, Gai Z, et al. Chronic Noise Exposure Acts Cumulatively to Exacerbate Alzheimer's Disease-Like Amyloid- β Pathology and Neuroinflammation in the Rat Hippocampus. *Sci Rep* 2015;5:12943. doi: 10.1038/srep12943 [published Online First: 2015/08/08]

38. Shin S, Bai L, Oiamo TH, et al. Association Between Road Traffic Noise and Incidence of Diabetes Mellitus and Hypertension in Toronto, Canada: A Population-Based Cohort Study. *J Am Heart Assoc* 2020;9(6):e013021. doi: 10.1161/jaha.119.013021 [published Online First: 2020/03/10]

39. Wen-jing D Z-dW, Xiang-bin Z, et al. Effects and mechanism of short-term day and night noise exposure on glucose and lipid metabolism in mice. *Modern Preventive Medicine* 2022;49(20):3678-84. doi: 10.20043/j.cnki.MPM.202205073

40. Evans GW, Lercher P, Meis M, et al. Community noise exposure and stress in children. *J Acoust Soc Am* 2001;109(3):1023-7. doi: 10.1121/1.1340642 [published Online First: 2001/04/17]

41. Wang H, Gu Y, Zheng L, et al. Association between bedtime and the prevalence of newly diagnosed non-alcoholic fatty liver disease in adults. *Liver Int* 2018;38(12):2277-86. doi: 10.1111/liv.13896 [published Online First: 2018/06/01]

42. Sun M, Feng W, Wang F, et al. Meta-analysis on shift work and risks of specific obesity types. *Obes Rev* 2018;19(1):28-40. doi: 10.1111/obr.12621 [published Online First: 2017/10/05]

43. Choi Y-H, Lee J-Y, Moon KW. Exposure to volatile organic compounds and polycyclic aromatic hydrocarbons is associated with the risk of non-alcoholic fatty liver disease in Korean adolescents: Korea National Environmental Health Survey (KoNEHS) 2015–2017. *Ecotoxicol Environ Saf* 2023;251:114508. doi: https://doi.org/10.1016/j.ecoenv.2023.114508

Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

44. VoPham T. Environmental Risk Factors for Liver Cancer and Nonalcoholic Fatty Liver Disease. *Current Epidemiology Reports* 2019;6(1):50-66. doi: 10.1007/s40471-019-0183-2

45. Werder EJ, Beier JI, Sandler DP, et al. Blood BTEXS and heavy metal levels are associated with liver injury and systemic inflammation in Gulf states residents. *Food Chem Toxicol* 2020;139:111242. doi: <u>https://doi.org/10.1016/j.fct.2020.111242</u>
46. Lang AL, Beier JI. Interaction of volatile organic compounds and underlying liver disease: a new paradigm for risk. *Biol Chem* 2018;399(11):1237-48. doi:

doi:10.1515/hsz-2017-0324

Figure 1. The relationship between shift work and NAFLD.

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, highfat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, benzene series, work exposed to noise.

Page 28 of 29

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

BMJ Open

3
4
5
6
7
/
8
9
10
11
12
13
14
15
15
16
17
18
19
20
21
22
∠∠ วว
23
24
25
26
27
28
29
20
20
31
32
33
34
35
36
37
38
20
27
40
41
42
43
44
45
46
47
77 10
40
49
50
51
52
53
54
55
55
50
5/
58
59

60



Supplementary

Table 1 Basic characteristics of population in different disease states (n, %)

Chamatariatian	NA	FLD	Total	D
Characteristics	No (N=3309)	Yes (N=1363)	(N=4672)	P
Drinking status (times per week), n (%)				0.382
Nondrinkers	2124 (64.2)	838 (61.5)	2962 (63.4)	
1~2	1091 (33.0)	484 (35.5)	1575 (33.7)	
3~4	65 (2.0)	28 (2.1)	93 (2.0)	
>4	29 (0.9)	13 (1.0)	42 (0.9)	
High-salt foods (times per week), n (%)				0.135
Seldom	1367 (41.3)	540 (39.6)	1907 (40.8)	
1~2	1749 (52.9)	724 (53.1)	2473 (52.9)	
3~4	144 (4.4)	67 (4.9)	211 (4.5)	
>4	49 (1.5)	32 (2.3)	81 (1.7)	
High-fat foods (times per week), n (%)				0.121
Seldom	984 (29.7)	367 (26.9)	1351 (28.9)	
1~2	2138 (64.6)	901 (66.1)	3039 (65.0)	
3~4	152 (4.6)	76 (5.6)	228 (4.9)	
>4	35 (1.1)	19 (1.4)	54 (1.2)	
Physical activity (times per week), n (%)				0.441
Never	871 (26.3)	359 (26.3)	1230 (26.3)	
1~2	2187 (66.1)	915 (67.1)	3102 (66.4)	
≥3	251 (7.6)	89 (6.5)	340 (7.3)	
Importance of noise protection				0 451
equipment, n (%)				0.451
Unimportant	63 (1.90)	24 (1.76)	87 (1.86)	
Unknown	72 (2.18)	20 (1.47)	92 (1.97)	
General	1718(51.92)	714 (52.38)	2432(52.05)	
Extremely	1456(44.00)	605 (44.39)	2061(44.11)	

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

	*			
	Model	l	Model 2	·
CIVE [db(A)·year]	OR 95%CI	Р	OR 95% <i>CI</i>	Р
Age				
≤29 yr				
<85	Ref			
85~	1.21(0.96, 1.53)	0.103	1.10 (0.86, 1.41)	0.443
90~	1.19(0.91, 1.56)	0.209	1.05 (0.79, 1.40)	0.728
95~	1.68(1.04, 2.66)	0.029	1.52 (0.93, 2.44)	0.088
≥30 yr				
<85	Ref			
85~	1.07 (0.59, 1.94)	0.824	1.05 (0.58, 1.93)	0.863
90~	1.03 (0.59, 1.82)	0.906	1.00 (0.57, 1.78)	0.989
95~	1.29 (0.72, 2.33)	0.389	1.28 (0.71, 2.31)	0.419

 Table 2
 The relationship between CNE and NAFLD in different age groups

Model 1: Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, Late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB;

Model 2: Based on model 1, adjust for importance of noise protection equipment, long working hours, dust, benzene series, shift work.

, shin ...

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

		Benzene se	eries (Yes or No)	
Variable	No		Yes	
	OR 95%CI	Р	OR 95%CI	Р
Work exposed to noise				
(Yes or No)				
No	Ref			
Yes	0.96 (0.80, 1.16)	0.695	0.71 (0.43, 1.17)	0.182
CNE [dB(A)∙year]				
<85	Ref			
85~	1.19 (0.94, 1.52)	0.147	0.78 (0.42, 1.42)	0.419
90~	1.14 (0.88, 1.48)	0.304	0.84 (0.42, 1.67)	0.624
95~	1.44 (1.04, 2.00)	0.029	1.64 (0.66, 3.98)	0.279
Shift work (Yes or No)				
No	Ref			
Yes	1.42 (1.13, 1.79)	0.003	1.07 (0.59, 2)	0.820
Cumulative length of				
night shift work (hours)			
≤730	Ref			
730~	1.43 (1.12, 1.84)	0.004	1.37 (0.73, 2.56)	0.323
1460~	1.55 (1.21, 1.98)	< 0.001	0.94 (0.46, 1.88)	0.852
2920~	1.74 (1.29, 2.35)	< 0.001	1.24 (0.45, 3.28)	0.67

 Table 3
 The relationship between occupational noise exposure, shift work and NAFLD under different benzene series exposure states

Adjust for age, smoking and drinking status, fruit and vegetable, high-salt foods, high-fat foods, late evening snack, BMI, sleep duration, physical activity, BP, ECG, FPG, HGB, importance of noise protection equipment, long working hours, dust, shift work/ work exposed to noise.