BMJ Open Effects of pre-CABG program on discharge readiness and surgery outcomes for patients undergoing elective CABG surgery: a study protocol for a randomised control trial

Pouya Dolat Abadi 💿 ,¹ Masoumeh Zakerimoghadam 💿 ,¹ Zahra Abbasi Dolat Abadi,¹ Mehrzad Rahmanian,² Seyed Mohammad Riahi,³ Ali Khanipour-Kencha¹

ABSTRACT

To cite: Dolat Abadi P. Zakerimoghadam M, Abadi ZAD, et al. Effects of pre-CABG program on discharge readiness and surgery outcomes for patients undergoing elective CABG surgery: a study protocol for a randomised control trial. BMJ Open 2025;15:e090256. doi:10.1136/ bmjopen-2024-090256

 Prepublication history for this paper is available online. To view these files, please visit the journal online (https://doi. org/10.1136/bmjopen-2024-090256).

Received 20 June 2024 Accepted 03 April 2025



C Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Masoumeh Zakerimoghadam; zakerimo@tums.ac.ir

Introduction Cardiovascular diseases, a leading cause of death globally, impose significant health and economic burdens, particularly in countries like Iran. Coronary artery bypass grafting (CABG) is a common intervention for ischaemic heart disease, yet it entails a long recovery process with potential complications and psychological impacts. This study aims to evaluate the effectiveness of a prehabilitation programme (pre-CABG) on postoperative outcomes and discharge readiness in patients undergoing elective CABG.

Methods and analysis This randomised controlled trial involves 60 patients diagnosed with coronary artery disease at Imam Khomeini Hospital Complex, Tehran. Participants will be randomly assigned to either the intervention group, receiving the pre-CABG programme, or the control group, receiving standard care. The pre-CABG programme includes patient education, stress management techniques, respiratory muscle training and nutritional guidance. Primary outcomes include discharge readiness, duration of intubation, Intensive Care Uniy (ICU) stay, occurrence of atelectasis, onset of mobility, hospital stay and levels of anxiety and depression. Secondary outcomes include the rate of 30-day readmissions. Data collection will involve standardised scales and checklists administered at various stages preoperation and postoperation.

Ethics and dissemination The research study has received approval from the Research Ethics Committee at Tehran University of Medical Sciences' School of Nursing and Midwifery and Rehabilitation. All participants must provide written consent for their involvement in this study. The findings will be shared with appropriate groups and published in peer-reviewed journals.

Trial registration number The study is registered with the Iranian Registry of Clinical Trials under the ID IBCT20231019059768N1.

INTRODUCTION

Cardiovascular diseases (CVDs) are a leading cause of death and a significant global health

STRENGTHS AND LIMITATIONS OF THIS STUDY

- \Rightarrow The prehabilitation programme is well-defined, including specific components such as stress management techniques, nutritional guidance, respiratory exercises and patient education.
- \Rightarrow Emphasising discharge readiness as a primary outcome is significant, as it is directly related to patient recovery, hospital resource utilisation and overall healthcare costs.
- \Rightarrow Single-centre design may limit the generalisability of findinas.
- \Rightarrow Unblinded participants and interventionists may introduce bias.
- \Rightarrow Relatively small sample size may limit the detection of minor differences between groups.

ng, issue.¹ It is predicted that in 2030, about 23 million deaths in the world will be due to CVD.² CVD is one of the leading causes of a significant reduction in quality of life and life expectancy and also imposes enormous costs **g** on health systems in different countries.¹ Several risk factors have been identified, including physical inactivity, obesity, hypertension, hyperlipidaemia, hyperglycaemia and stress. These factors also influence the recovery process from major cardiac events.³ Studies conducted in Iran also show that coronary artery disease is the leading cause of **G** death in Iran, and Iran has the highest disease burden caused by this disease in the region.⁴ Based on the studies, CVDs are the leading cause of 46% of deaths and about 20%–23% of the total burden of disease in Iran, and it is one of the main problems of Iran's health system.⁵ ⁶ Current therapeutic interventions for ischaemic heart disease include drug therapy (antiplatelet drugs, beta-blockers or statins) to stabilise the disease and reduce

≥

Protected by copyright, including for uses related to text and data mini

BMJ Group

acute events (such as myocardial infarction or sudden death) or immediate restoration of blood flow through surgical revascularisation treatment, such as coronary artery bypass grafting (CABG) or percutaneous coronary intervention.⁷⁸ The recovery process for these surgeries is long and involves caregiving challenges.9 Nevertheless, this method is a common choice for heart patients and surgeons. Complications of this surgery include increased hospitalisation time, delirium, anxiety related to treatment and their ability to adhere to recommended physical activity, depression, stroke, bleeding and cardiac tamponade, myocardial dysfunction, sternal wound infection and pulmonary complications.^{10–12}

Despite all the advantages, CABG is a stressful and traumatic event and usually has negative psychological consequences in the preoperative period that may persist after the surgery.¹³¹⁴ When a person is diagnosed with coronary artery disease, it causes high levels of anxiety, which is exceptionally high in the preoperative period and when the client is waiting for major surgery.¹³ ¹⁵ High anxiety reduces quality of life and worsens long-term psychological consequences.¹¹ Persistent stress can negatively affect the prognosis and physiological parameters of patients (preoperatively and intraoperatively or during anaesthesia), which may prolong recovery and length of stay and negatively impact the quality of life.¹⁶¹⁷ Also, the relationship between depression and CVD and its effect on the outcome of patients hospitalised for acute coronary artery disease, as well as before and after vascular surgery, is well known.¹⁸ The presence of depressive symptoms during or shortly after hospitalisation increases the risk of death or non-fatal cardiac events by two to three times and significantly increases the morbidity and mortality of these clients.^{19 20} Studies have shown a lower healthrelated functional status and quality of life, as well as a higher rate of readmission after discharge and increased mortality in these clients.²¹

Surgeries that require a thoracotomy, such as coronary artery bypass surgery, carry a high risk of pulmonary complications.²² Patients undergoing cardiac surgery are more vulnerable to developing these pulmonary complications after surgery.² Today, in most hospitals, chest physiotherapy is used to minimise postoperative pulmonary complications after CABG.^{24 25} Various studies have reported that early rehabilitation significantly reduces long-term mortality in patients undergoing coronary artery bypass graft surgery.²⁶⁻²⁸ Despite the well-documented importance of postoperative rehabilitation, insufficient information on the value of preoperative interventions is available.^{23 29}

Prehabilitation is a method of preparing the patient for heart surgery.³⁰ Patients admitted to the hospital for CABG surgery may have to wait up to a week for the procedure, which may lead to regression and physical weakness and negatively affect the outcome of the surgery.³¹ To better use this waiting period, patients can participate in prehabilitation programmes. These

<page-header><text><text><text><text><text><text><text><text><text><text>



Figure 1 Flowchart showing patient allocation to intervention and control groups. HADS, Hospital Anxiety and Depression Scale; RHDS, Readiness for Hospital Discharge Scale.** COPD Chronic Obstructive Pulmonary Disease.** RHDS-RN Readiness for Hospital Discharge Scale based on Registered Nurses' Score

elective coronary artery bypass surgery list; (3) experiencing heart surgery for the first time; (4) the ability to speak and understand Persian; (5) not having been diagnosed with anxiety or other psychological disorders (self-report) and (6) non-concurrency of CABG surgery with valve replacement or other surgeries.

Exclusion criteria

Exclusion criteria for the study include (1) lack of consent to continue participating in the study; (2) postoperative haemodynamic instability; (3) aortic dissection or aneurysm; (4) haemodynamic instability leading to surgical and (10) patients with cognitive disorders and Alzheimer's disease.

Intervention

This study involves a prehabilitation programme led by a nurse. A cardiac surgeon and nurse interventionist will determine patients' eligibility during a face-to-face visit. Participants who meet the eligibility criteria and agree to participate will be randomly

divided into intervention and control groups. The control group will receive standard care, while patients in the intervention group will receive the pre-CABG intervention. This includes three visits from the nurse interventionist-the first during the surgery clinic meeting with the surgeon and the next two while in the hospital waiting for surgery. During the first session, patients in the intervention group will receive an educational booklet with four sections and an incentive spirometry.

The pre-CABG intervention

This intervention programme will encompass the following steps:

- 1. A meeting between the interventional nurse, the patient and the surgeon during clinic visits.
- 2. Delivering a comprehensive explanation of the surgical procedure in a manner designed to minimise patient anxiety.
- 3. Encouraging patients to discuss anxiety triggers and rectifying any misconceptions with the intervention team's assistance.
- 4. Introducing stress management techniques, such as progressive muscle relaxation, deep breathing, guided imagery and recommended practice repetition. Patients will choose their preferred stress management method, which will be taught and practised.
- 5. Providing preoperative nutritional guidance, emphasising reduced carbohydrate intake and increased protein, vitamins and micronutrient consumption.
- 6. Advise patients to quit smoking if necessary and offer respiratory muscle strengthening exercises, such as the pursed lip breathing method, incentive spirometry before the operation and diaphragmatic breathing with hands-on practice until proficiency is achieved. The accuracy of the patient's technique will be assessed, and if necessary, additional teaching sessions will be provided until the patient is proficient.
- 7. Implementing a logbook in which patients will record the number of training sessions completed and the duration of each session. Given that patients with cardiovascular conditions often experience high levels of anxiety and fear of death, they tend to follow the instructions with great attention to detail.

Participants will complete a demographic information questionnaire on expressing interest and providing informed consent. For the intervention group, the intervention will commence approximately 7 days before surgery, with the first session during the surgery clinic visit. The subsequent two sessions will occur during hospitalisation in the cardiac surgery department 2 days before the operation, focusing on face-to-face training, continued prehabilitation activities and answering questions.

The first session will involve the steps mentioned earlier. The second and third sessions will be held in the heart surgery ward during the patient's hospitalisation. The researcher and the patient will review previous topics, address patient concerns and reinforce stress

management techniques and breathing exercises. Each session will last 45-60 min, catering to individual patient needs.

Both control and intervention groups will receive routine care, including drug therapy and haemodynamic monitoring. The Hospital Anxiety and Depression Scale (HADS) will be completed at the end of the third session. Additional assessments, such as intubation duration, intensive care unit stay, occurrence of atelectasis, overall hospital stay and discharge readiness (assessed by adult form Readiness for hospital discharge scale patient version and nurse version), will be conducted before **g** patient discharge. 30-day re-hospitalisation checklist will ŝ be completed 4weeks postdischarge for both control and copyright, includ intervention groups, ensuring a comprehensive evaluation of the intervention's impact.

Outcome measures

The study will evaluate the intervention's effectiveness based on the following primary and secondary outcomes:

Primary outcomes

for uses r The primary outcome (measured variables) of this study will include discharge readiness; discharge readiness will be assessed using the short form of the Readiness for Hospital Discharge Scale (RHDS) for both patients and nurses. The RHDS evaluates a patient's preparedness to transition home from $\overline{\mathbf{a}}$ acute care, typically administered 4 hours before ŧ discharge. The RHDS includes eight items that cover four domains: personal status, knowledge, perceived coping ability and expected support. Scores range from 0 to 10, representing the average of item scores. a The Persian versions of RHDS for both patients and nurses have been validated for reliability and validity by Mehraeen et al.^{38 39}

Secondary outcomes

Al training, The secondary outcomes will include (1) number of hospital readmissions, (2) duration of intubation, (3) length of ICU stay, (4) occurrence of atelectasis, (5) timing of the first mobilisation, (6) total length of <u>0</u> hospital stay and (7) levels of anxiety and depression during hospitalisation.

The number of hospital readmissions is defined as emergency and unexpected readmissions due to symptoms suggesting myocardial infarction, pneumonia or atelectasis. Participants will report the number of **G** readmissions within 30 days postdischarge, confirmed 🖁 through the hospital information system or medical records from other health centres if applicable.

Anxiety and depression during the hospital stay will be measured using the HADS, a 14-item tool developed by Sigmon and Snaith in 1983. HADS assesses anxiety and depression symptoms, scored on a 4-point scale (0-3), with a cut-off score of 11 indicating significant symptoms. The Persian version of HADS has been validated by Kaviani et al.^{40 41}

Bul

Other outcomes, including the duration of intubation, the occurrence of atelectasis (determined by chest X-ray and physician diagnosis), duration of ICU stay, length of hospitalisation and the initiation of mobility, will be measured using a researcher-made checklist.

Data collection

Data will be gathered using a demographic information questionnaire, HADS, RHDS-RN, RHDS-Pt (Readiness for Hospital Discharge Scale based on Registered Nurses' Score and patients' own score respectively), a researchermade checklist and a question on the number of readmissions. Demographic information will be collected through self-report and medical records at the study's onset. HADS will be completed at the study's start, on the night before the surgery and on the day of discharge. The RHDS forms will be filled out approximately 4 hours before discharge. 30 days postdischarge, the outcome assessor will inquire about emergency readmissions via phone. The data about the duration of intubation, the occurrence of atelectasis, duration of ICU stay, length of hospitalisation and the initiation of mobility will be extracted by the researcher by daily observation of patients' progress and data on the patients' documents.

Data displayed in table 1 provide the present study's enrolment, interventions and assessment schedule.

Sample size

Protected Based on the primary outcome and assuming a 5% type I error rate and 80% power, the sample size is ŝ calculated to be 26 patients per group, following the copyright, including for uses related to text and data mining, Al training, and similar technologies study by Nurhayati et al. Allowing for a 15% attrition rate, the estimated number of participants is 30 per

Table 1 Schedule for enrolment, interventions and assessments								
	Study period							
	Enrolment	Allocation	Intervention	Follow-up				
Time point	Baseline	0	Clinic meeting to day of the surgery	Day of the surgery to day of discharge	4 weeks after discharge			
Enrolment:								
Eligibility screen	*							
Informed consent	*							
Randomisation		*						
Intervention group:								
The pre-CABG programme			*					
Usual care			*	*	*			
Assessments:								
Demographic information	*							
Marital status	*							
Level of education	*							
Occupation status	*							
Income	*							
Medical history	*							
Smoking	*							
Alcohol	*							
Primary outcomes:								
Discharge readiness				*				
Secondary outcomes:								
Duration of intubation				*				
Duration of ICU stay				*				
Occurrence of atelectasis				*				
First day of starting mobility	,			*				
Duration of hospital stay				*				
Anxiety and depression during the hospital stay	*		*	*	*			
Hospital readmissions					*			
CABG, coronary artery bypass gra	afting.							

used for sample size calculation are: Mean readiness for discharge score: 7.11±0.59 Effect size (d): 0.71 Significance Level (α): 0.05 Power $(1 - \beta)$: 80% (β =0.2) Attrition rate: 15% Allocation ratio: 1

Recruitment

Participants will be recruited at the IKHC cardiac surgery clinic. Eligible patients will be screened by a nurse interventionist and a cardiac surgeon, and those meeting the criteria will receive detailed information about the study. Patients will be randomly assigned to the control or intervention group by nurse interventionists after providing written informed consent. The intervention will be

conducted in clusters of 6-8 participants, continuing until the sample size is reached (table 2).

Random allocation

Participants will be randomised into control or intervention groups using balanced block randomisation with a block size of six, stratified by gender, artery blockage percentage and symptom severity. An independent researcher will generate the allocation sequence, ensuring concealment with sealed opaque envelopes.

Blinding

Protected In this research, one-way blinding will be used. Patients will be admitted to separate rooms to prevent interaction ŝ between control and intervention groups. Ideally, patients will be placed in private, single-occupancy rooms to minimise the risk of interaction between the intervention

Table 2 Schedule for enrolm	ent, interventions	s and assessme	ents				
	Study period						
	Enrolment	Allocation	Intervention	Follow-up			
Time point	Baseline	0	Clinic meeting to day of the surgery	Day of the surgery to day of discharge	4 weeks after discharge		
Enrolment:							
Eligibility screen	*						
Informed consent	*						
Randomisation		*					
Intervention group:							
The pre-CABG programme			*				
Usual Care			*	*	*		
Assessments:							
Demographic information	*						
Marital status	*						
Level of education	*						
Occupation status	*						
Income	*						
Medical history	*						
Smoking	*						
Alcohol	*						
Primary outcomes:							
Discharge readiness				*			
Secondary outcomes:							
Duration of intubation				*			
Duration of ICU stay				*			
Occurrence of atelectasis				*			
First day of starting mobility				*			
Duration of hospital stay				*			
Anxiety and depression during the hospital stay	*		*	*	*		
Hospital readmissions					*		
CABG, coronary artery bypass gra	afting.						

groups and other patients. However, this arrangement may be subject to changes based on hospital policies and the availability of beds within the ward. In cases where private rooms are not available, the researcher will make every effort to admit study participants into the same room, ensuring that intervention groups are housed separately. If this is not feasible, the researcher will document and acknowledge this limitation in the final study results. The outcome assessor and the statistician will be kept uninformed and directed not to ask about the group allocations of the participants.

Data management

Data will be gathered through hard-copy questionnaires and lists, except for the 30-day follow-up, which will be conducted over the phone. Patients will be urged to complete the questionnaires to reduce missing data, and the person assessing the outcomes will receive training to ensure precise data collection. The data entered into SPSS software V.22 will undergo double verification to ensure accuracy. The data will be stored using anonymised ID codes to safeguard confidentiality, and the analysis will be carried out by a statistician specialised in epidemiology.

Statistical analysis

Frequency and percentage will describe qualitative data, while mean and SD (or median and first-third quartile) will describe quantitative variables. The normality of the data will be checked using the Shapiro-Wilk test and the P-P plot diagram, and the Levene test will check the homogeneity of the variance.

Independent t-tests will be used to compare means in two independent groups, and paired t-tests will be used in two dependent groups. A comparison of qualitative variables in two groups will be done using χ^2 or Fisher's exact test. The correlation of two quantitative variables will be investigated using the Pearson or Spearman correlation coefficient. In this study, analysis of covariance or linear regression will be used to eliminate the effect of confounders in examining the impact of the intervention. If the normality of the data is not met, the non-parametric equivalent of statistical tests will be used. All analyses will be done in SPSS22 software at a significance level of 0.05.

Progression criteria and evaluation

To ensure methodological rigour and consistency throughout the study, we have outlined specific progression criteria that will guide the execution of this research. These criteria will be evaluated at each stage of the study, from the initial planning phase to data analysis.

- 1. Clarity of research question: the research hypothesis, evaluating the effectiveness of a prehabilitation programme for improving discharge readiness and surgery outcomes in CABG patients, will be refined as necessary throughout the study to maintain alignment with the evolving research context.
- 2. Methodological rigour: the study design follows an RCT approach, adhering to the SPIRIT guidelines to

ensure methodological consistency. Randomisation procedures and blinding protocols will be closely monitored to minimise bias and maintain the integrity of the trial.

- 3. Intervention delivery: the pre-CABG intervention will be delivered as planned, with periodic monitoring to ensure that all elements of the programme, such as stress management techniques, educational sessions and preoperative health guidance, are consistently applied to all participants.
- 4. Outcome measurement: primary and secondary outcomes will be measured at predefined intervals using validated instruments such as the HADS and RHDS. Any missing data will be handled using appropriate methods, such as imputation or exclusion of incomplete cases, depending on the nature of the missing data.
- Statistical analysis: statistical analyses will be conducted in SPSS software to assess the impact of the intervention on the primary and secondary outcomes. Specific attention will be paid to potential confounders, and appropriate statistical tests will be employed to account for these variables.
 Discussion and study limitations: the findings of the secondary outcomes.
- 6. Discussion and study limitations: the findings of the study will be discussed in relation to existing literature, and limitations related to the study design, such as the inability to blind patients to the intervention, will be critically evaluated.

DISCUSSION

id data proposed RCT investigates the pre-CABG The programme's potential impact on discharge readiness and surgery outcomes for patients undergoing elective CABG. Coronary artery bypass graft surgery is a standard procedure for patients with severe coronary artery disease.⁷ However, the recovery process after CABG surgery can be \geq challenging and may involve complications such as infections, prolonged hospital stays and delayed discharge.¹⁰¹¹ This study aims to evaluate the impact of a pre-CABG programme on improving discharge readiness and surgery outcomes in patients undergoing elective CABG surgery. The pre-CABG programme provides patient education, encourages respiratory muscle training, offers preoperative nutritional guidance and introduces stress management techniques to optimise patient readiness for surgery and postoperative recovery. The pre-CABG programme prepares patients both physically and mentally for their & CABG surgery. Previous studies support the inclusion 2 of patient education in the pre-CABG programme.43-45 Prehabilitation before elective CABG, which includes patient education, exercise training and social support, has improved patients' physical and psychological readiness for surgery and reduced postoperative complications.^{46–48} The importance of patient education in promoting positive outcomes for CABG surgery patients has been highlighted by various studies. Varaei et al found that educational interventions effectively promote cardiac

lated

to text

an

Open access

self-efficacy in patients undergoing CABG surgery.⁴⁹ Additionally, Akbari and Çelik conducted a study that applied discharge training and counselling to patients undergoing CABG surgery, decreasing their problems.⁵⁰ Rief *et al* found that optimising patients' expectations presurgery improved outcomes, particularly regarding disability and quality of life, 6 months after treatment.⁵¹

Furthermore, the research article highlights the potential benefits of preoperative rehabilitation and prehabilitation. Patients who are waiting for CABG surgery often experience fear and anxiety, which can hinder their engagement in rehabilitation postoperatively.⁵² However, this waiting period offers an opportunity for preoperative rehabilitation or prehabilitation to improve the surgical intervention's safety and outcome and encourage ongoing postoperative engagement in rehabilitation.⁴⁷ Prehabilitation has gained recognition in cardiac surgery, with the idea that interventions before surgery can enhance patients' physical and psychological well-being and improve surgical outcomes. Furthermore, the emphasis on preoperative preparation in the pre-CABG programme may lead to shorter hospital stays and earlier patient discharge. This can improve patient satisfaction and alleviate the burden on healthcare facilities by freeing up resources for other patients in need of care. In conclusion, implementing a pre-CABG programme shows promise in improving discharge readiness and surgery outcomes for patients undergoing elective CABG surgery. Assessing the impact of such a programme through a randomised control trial will provide valuable insights into its effectiveness and potential for enhancing the overall care and recovery of patients undergoing this standard cardiac procedure.

Strengths and limitations

As with any study, this research has its limitations. Despite efforts to address them, certain challenges may persist. For instance, accurately completing the RHDS-RN and RHDS-Pt questionnaires precisely 4 hours before discharge may be difficult due to paperwork complications and unpredictable discharge times, which can sometimes occur at inconvenient hours like midnight. To manage this issue, the questionnaires will be filled out as soon as the discharge decision is made. Additionally, the nature of the intervention prevents blinding for participants and interventionists, although outcome assessors and the statistician will remain blinded. Remote interventions also carry risks of attrition and inadequate adherence, but we plan to motivate participants through valuable information, proper guidance and continuous follow-up.

We hypothesise that the comprehensive prehabilitation programme will enhance discharge readiness, improve surgery outcomes and reduce emergency readmissions. Positive results from this study could benefit patients undergoing elective CABG and be applicable in both Iran and other countries.

Patient and public involvement

There was no patient or public involvement in the development of this protocol.

Ethics and dissemination

The study received approval from the School of Nursing and Midwifery Research Ethics Committee on 10 October 2023, with the approval number IR.TUMS. IKHC.REC.1402.287. The study protocol has been registered with the ID IRCT20231019059768N1 in the Iranian Registry of Clinical Trials. All participants must provide written consent and be fully informed of their rights to voluntary participation and withdrawal. Data will be confidentially and anonymously stored and handled. Participants will be given contact information so the nurse researcher can address any queries. The findings will be shared with representative groups and published in peer-reviewed journals.

Trial status

The recruitment for this study began in April 2024 and is estimated to end in January 2025.

Author affiliations

¹Department of Medical-Surgical Nursing, Tehran University of Medical Sciences, Tehran, Iran (the Islamic Republic of)

²Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran (Islamic Republic of)

³Department of Community Medicine, School of Medicine, Cardiovascular Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran (the Islamic Republic of)

Acknowledgements This study protocol is a component of PD's Master of Science thesis in Medical-Surgical Nursing at TUMS. The achievement of this study has been facilitated by the invaluable assistance of the Education and Research Deputy and the Medical-Surgical Nursing Department at the School of Nursing and Midwifery, TUMS. We extend our deep appreciation for their endorsement of the thesis and their consistent support, which has played a crucial role in our research journey.

Contributors PDA leads the trial, and MZ supervises the project. The project was conceived by PDA, ZADA, MR, SMR, AK-K and MZ. PDA and MR will carry out the implementation. MR, who is a cardiothoracic surgeon at IKHC, will also assess the educational content and confirm patient eligibility. PDA and MZ will work together to create the educational material. SMR will provide statistical expertise for the trial design and conduct the statistical analyses. All authors were involved in drafting this paper and have given approval for the final manuscript. PDA is responsible for the overall content as guarantor. Chat GPT 3.5 was used to improve my writing as I am not a native English speaker.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Pouya Dolat Abadi http://orcid.org/0000-0002-7281-2863

Masoumeh Zakerimoghadam http://orcid.org/0000-0002-9762-1887

REFERENCES

- Amini M, Zayeri F, Salehi M. Trend analysis of cardiovascular disease mortality, incidence, and mortality-to-incidence ratio: results from global burden of disease study 2017. *BMC Public Health* 2021;21:401.
- 2 Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet* 2012;380:2095–128.
- 3 Sokhteh ASD, Mofrad ZP, Yaghoubinia F. The effectiveness of rehabilitation-education program on risk factors in patients with coronary artery disease. 2020.
- 4 Gaziano TA, Bitton A, Anand S, et al. Growing epidemic of coronary heart disease in low- and middle-income countries. Curr Probl Cardiol 2010;35:72–115.
- 5 Roth GA, Mensah GA, Johnson CO, et al. Global Burden of Cardiovascular Diseases and Risk Factors, 1990-2019: Update From the GBD 2019 Study. J Am Coll Cardiol 2020;76:2982–3021.
- 6 Sarrafzadegan N, Mohammmadifard N. Cardiovascular Disease in Iran in the Last 40 Years: Prevalence, Mortality, Morbidity, Challenges and Strategies for Cardiovascular Prevention. *Arch Iran Med* 2019;22:204–10.
- 7 Rodriguez AE, Pavlovsky H, Del Pozo JF. Understanding the Outcome of Randomized Trials with Drug-Eluting Stents and Coronary Artery Bypass Graft in Patients with Multivessel Disease: A Review of a 25-Year Journey. *Clin Med Insights Cardiol* 2016;10:195–9.
- 8 Serruys PW, Morice M-C, Kappetein AP, et al. Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. N Engl J Med 2009;360:961–72.
- 9 Noor Hanita Z, Khatijah LA, Kamaruzzaman S, et al. A pilot study on development and feasibility of the "MyEducation: CABG application" for patients undergoing coronary artery bypass graft (CABG) surgery. BMC Nurs 2022;21:40.
- Hussain SMA, Harky A. Complications of Coronary Artery Bypass Grafting. Int J Med Rev 2019;6:1–5.
- 11 AbuRuz ME, Al-Dweik G, Al-Akash HY. Checking the moderating effect of perceived control on the relationship between anxiety and postoperative hospital length of stay among coronary artery bypass graft patients. *Int J Gen Med* 2019;12:79–85.
- 12 Cebeci F, Celik SS. Discharge training and counselling increase selfcare ability and reduce postdischarge problems in CABG patients. J *Clin Nurs* 2008;17:412–20.
- 13 Tully PJ, Baker RA. Depression, anxiety, and cardiac morbidity outcomes after coronary artery bypass surgery: a contemporary and practical review. *J Geriatr Cardiol* 2012;9:197–208.
- 14 Yilmaz M, Sezer H, Gürler H, et al. Predictors of preoperative anxiety in surgical inpatients. J Clin Nurs 2012;21:956–64.
- Nemeroff C. Prevalence of depression and anxiety in CABG patients. J Watch Psychiatry 2003;18:12–9.
- 16 Middel B, El Baz N, Pedersen SS, et al. Decline in health-related quality of life 6 months after coronary artery bypass graft surgery: the influence of anxiety, depression, and personality traits. J Cardiovasc Nurs 2014;29:544–54.
- 17 AbuRuz ME, Masa'Deh R. Gender Differences in Anxiety and Complications Early After Acute Myocardial Infarction. J Cardiovasc Nurs 2017;32:538–43.
- 18 Nunes JKVRS, Figueiredo Neto JAd, Sousa RMLd, et al. Depression after CABG: a prospective study. *Rev Bras Cir Cardiovasc* 2013;28:491–7.
- 19 Blumenthal JA, Lett HS, Babyak MA, et al. Depression as a risk factor for mortality after coronary artery bypass surgery. The Lancet 2003;362:604–9.
- 20 Lespérance F, Frasure-Smith N. Depression in patients with cardiac disease: a practical review. J Psychosom Res 2000;48:379–91.
- 21 Williamson GM, Schulz R. Pain, activity restriction, and symptoms of depression among community-residing elderly adults. *J Gerontol* 1992;47:P367–72.
- 22 Laszlo G, Archer GG, Darrell JH, et al. The diagnosis and prophylaxis of pulmonary complications of surgical operation. Br J Surg 1973;60:129–34.
- 23 Shahood H, Pakai A, Kiss R, et al. Effectiveness of Preoperative Chest Physiotherapy in Patients Undergoing Elective Cardiac Surgery, a Systematic Review and Meta-Analysis. *Medicina (Kaunas)* 2022;58:911.
- 24 Nardi P, Pellegrino A, Pisano C, *et al.* The effect of preoperative respiratory physiotherapy and motor exercise in patients undergoing elective cardiac surgery: short-term results. *kitp* 2019;16:81–7.

25 Dąbek J, Pyka E, Piotrkowicz J, et al. Impact of post-hospital cardiac rehabilitation on the quality of life of patients after surgical treatment for coronary artery disease. *Kardiochir Torakochirurgia Pol* 2017;14:120–6.

Open access

- 26 Dayan V, Ricca R. Cardiac rehabilitation after coronary artery bypass surgery. Arch Cardiol Mex 2014;84:286–92.
- 27 Lee J-Y, Han S, Ahn J-M, et al. Impact of participation in phase I and phase II cardiac rehabilitation on long-term survival after coronary artery bypass graft surgery. Int J Cardiol 2014;176:1429–32.
- 28 Pack QR, Goel K, Lahr BD, et al. Participation in cardiac rehabilitation and survival after coronary artery bypass graft surgery: a communitybased study. Circulation 2013;128:590–7.
- 29 Dabek J, Pyka E, Piotrkowicz J, et al. Impact of post-hospital cardiac rehabilitation on the quality of life of patients after surgical treatment for coronary artery disease. kitp 2017;2:120–6.
- 30 Argunova Y, Belik E, Gruzdeva O, et al. Effects of Physical Prehabilitation on the Dynamics of the Markers of Endothelial Function in Patients Undergoing Elective Coronary Bypass Surgery. J Pers Med 2022;12:471.
- 31 Herdy AH, Marcchi PLB, Vila A, et al. Pre- and Postoperative Cardiopulmonary Rehabilitation in Hospitalized Patients Undergoing Coronary Artery Bypass Surgery. Am J Phys Med Rehabil 2008;87:714–9.
- 32 Jenkins CD, Stanton B-A, Savageau JA, et al. Coronary artery bypass surgery. Physical, psychological, social, and economic outcomes six months later. JAMA 1983;250:782–8.
- 33 Dressler D. Management of patients with coronary vascular disorders. In: *Brunner and suddarth's textbook of medical-surgical nursing.* 12th edn. 2010. Philadelphia: Wolters Kluwer Health, 2010: 775–9.
- 34 Pieper B, Sieggreen M, Freeland B, et al. Discharge Information Needs of Patients After Surgery. J Wound Ostomy Continence Nurs 2006;33:281–90.
- 35 Rushton M, Howarth M, Grant MJ, et al. Person-centred discharge education following coronary artery bypass graft: A critical review. J Clin Nurs 2017;26:5206–15.
- 36 Anthony MK, Hudson-Barr D. A patient-centered model of care for hospital discharge. *Clin Nurs Res* 2004;13:117–36.
- 37 Chan A-W, Tetzlaff JM, Gøtzsche PC, *et al.* SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ* 2013;346:e7586.
- 38 Mehraeen P, Jafaraghaee F, Paryad E, et al. Comparison of Nurses' and Patients' Readiness for Hospital Discharge: A Multicenter Study. J Patient Exp 2022;9.
- 39 Weiss ME, Costa LL, Yakusheva O, *et al.* Validation of patient and nurse short forms of the Readiness for Hospital Discharge Scale and their relationship to return to the hospital. *Health Serv Res* 2014;49:304–17.
- 40 Zigmond AS, Snaith RP. The hospital anxiety and depression scale. Acta Psychiatr Scand 1983;67:361–70.
- 41 Kaviani H, Seyfourian H, Sharifi V, et al. Reliability and validity of anxiety and depression hospital scales (HADS): Iranian patients with anxiety and depression disorders. *TUMJ* 2009;67:379–85.
- 42 Nurhayati N, Songwathana P, Vachprasit R. Surgical patients' experiences of readiness for hospital discharge and perceived quality of discharge teaching in acute care hospitals. *J Clin Nurs* 2019;28:1728–36.
- 43 Guo P. Preoperative education interventions to reduce anxiety and improve recovery among cardiac surgery patients: a review of randomised controlled trials. *J Clin Nurs* 2015;24:34–46.
- 44 Özdemir B, Önler E. The effect of a structured patient education intervention on the quality of life for coronary artery bypass grafting patients: A prospective randomised controlled study. *J Perioper Pract* 2021;31:124–31.
- 45 Shi W, Ghisi GLM, Zhang L, et al. Systematic review, metaanalysis and meta-regression to determine the effects of patient education on health behaviour change in adults diagnosed with coronary heart disease. J Clin Nurs 2023;32:5300–27.
- 46 Marmelo F, Rocha V, Moreira-Gonçalves D. The impact of prehabilitation on post-surgical complications in patients undergoing non-urgent cardiovascular surgical intervention: Systematic review and meta-analysis. *Eur J Prev Cardiol* 2018;25:404–17.
- 47 Sawatzky J-AV, Kehler DS, Ready AE, et al. Prehabilitation program for elective coronary artery bypass graft surgery patients: a pilot randomized controlled study. *Clin Rehabil* 2014;28:648–57.
- 48 Arora RC, Brown CH 4th, Sanjanwala RM, et al. 'NEW' Prehabilitation: A 3-Way Approach to Improve Postoperative Survival and Health-Related Quality of Life in Cardiac Surgery Patients. Can J Cardiol 2018;34:839–49.

Open access

- 49 Varaei S, Shamsizadeh M, Cheraghi MA, et al. Effects of a peer education on cardiac self-efficacy and readmissions in patients undergoing coronary artery bypass graft surgery: a randomizedcontrolled trial. *Nurs Crit Care* 2017;22:19–28.
 50 Akbari M, Celik SS. The effects of discharge training and
- 50 Akbari M, Celik SS. The effects of discharge training and postdischarge counseling on quality of life after coronary artery bypass graft surgery. *Iran J Nurs Midwifery Res* 2015;20:442–9.
- 51 Rief W, Shedden-Mora MC, Laferton JAC, *et al.* Preoperative optimization of patient expectations improves long-term outcome in heart surgery patients: results of the randomized controlled PSY-HEART trial. *BMC Med* 2017;15:4.
- 52 Rao A, Zecchin R, Newton PJ, et al. The prevalence and impact of depression and anxiety in cardiac rehabilitation: A longitudinal cohort study. Eur J Prev Cardiol 2020;27:478–89.