


BMJ Open Study protocol: diagnosis of atrial fibrillation in postoperative thoracic surgery using a smartwatch, an open-label randomised controlled study (THOFAWATCH trial)

Pierre Huette ^{1,2}, Christophe Beyls,² Momar Diouf,³ Azrat Ibrahima,² Guillaume Haye,² Mathieu Guilbart,² Thomas Lefebvre,² Guillaume Bayart,² Franck Lhotellier,¹ Michael Radji,¹ Katy-Anne Walczak,¹ Matthieu Caboche,¹ Florence De Dominicus,⁴ Olivier Georges,⁴ Pascal Berna,⁵ Geonie Merlusca,^{4,5} Alexis Hermida,⁶ Sarah Traullé,⁷ Herve Dupont,² Yazine Mahjoub,² Osama Abou-Arab,² On behalf of THOFAWATCH Study Group

To cite: Huette P, Beyls C, Diouf M, *et al.* Study protocol: diagnosis of atrial fibrillation in postoperative thoracic surgery using a smartwatch, an open-label randomised controlled study (THOFAWATCH trial). *BMJ Open* 2025;**15**:e097765. doi:10.1136/bmjopen-2024-097765

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-097765>).

Received 09 December 2024
Accepted 27 March 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Pierre Huette;
huette.pierre@chu-amiens.fr

ABSTRACT

Introduction Postoperative atrial fibrillation (POAF) affects approximately 20% of patients undergoing thoracic surgery and is associated with severe complications such as stroke, myocardial infarction, heart failure, and increased mortality. Early diagnosis is critical to mitigate these risks, but conventional monitoring is limited in detecting asymptomatic episodes. Smartwatches equipped with single-lead ECG and atrial fibrillation (AF) detection algorithms offer a novel approach for early POAF detection. This study aims to evaluate the effectiveness of smartwatch-based monitoring compared with standard care in identifying POAF following thoracic surgery.

Methods and analysis The THOFAWATCH trial is a randomised, bicentric open-label study enrolling 302 adult patients undergoing major thoracic surgery (pneumonectomy or lobectomy) with one-lung ventilation. Eligible patients will be randomised into two groups: (1) the 'Smartwatch Monitoring' group, where participants will undergo rhythm monitoring using a smartwatch and (2) the 'Conventional Monitoring' group, receiving standard care without smartwatch monitoring. In the intervention group, any smartwatch-detected POAF episodes will be confirmed by 12-lead ECG. The primary outcome is the incidence of POAF within 7-day postsurgery. Secondary outcomes include the rate of asymptomatic POAF, cardiovascular prognosis evaluated at 2 and 6 months (composite major adverse cardiovascular events outcome), feasibility of smartwatch usage (device usage time and success rate of single-lead ECGs) and recurrence or management of AF at follow-up. Inclusion criteria include adults (>18 years) undergoing scheduled thoracic surgery and able to use the smartwatch device. Exclusion criteria encompass patients with prior AF, those requiring telemetry, or undergoing reoperations. Statistical analysis will assess the primary outcome using χ^2 or Fisher's exact test ($\alpha=5\%$), while secondary outcomes will include descriptive and inferential statistics, with analysis conducted using SAS V.9.4.

Ethics and dissemination Ethical approval for this bicentric study has been granted by the institutional review board (IRB)

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The trial will be a prospective, bicentric, randomised trial and will include adult patients undergoing major thoracic surgery (pneumonectomy or lobectomy).
- ⇒ The intervention that is being investigated is the use of a smartwatch-based monitoring to detect the occurrence of postoperative atrial fibrillation (POAF) following thoracic surgery.
- ⇒ No study has yet evaluated the effectiveness of smartwatches in detecting POAF after thoracic surgery.
- ⇒ The primary endpoint will be the incidence of POAF within 7 days postsurgery.
- ⇒ The secondary endpoints will include the rate of asymptomatic POAF, cardiovascular prognosis evaluated at 2 and 6 months (composite major adverse cardiovascular events outcome), feasibility of smartwatch usage (device usage time and success rate of single-lead ECGs) and recurrence or management of atrial fibrillation at follow-up.

of the University Hospital of Amiens (Comité de Protection des Personnes sud-ouest et outre-mer 1, 21050 Toulouse, France, registration number ID RDB: 2022-A02028-27 in November 2024). The trial is registered under ClinicalTrials.gov (ID: (NCT06724718)). Results will be disseminated through peer-reviewed publications and scientific conferences to inform clinical practice regarding POAF detection and management following thoracic surgery.

Trial registration number NCT06724718; clinical trial.

INTRODUCTION

In patients undergoing major non-cardiac surgery, postoperative atrial fibrillation (POAF) is a common and significant complication associated with increased risks of heart

failure, stroke, myocardial infarction and death, leading to higher morbidity, prolonged hospital stays and elevated healthcare costs.^{1–4} POAF episodes are often paroxysmal and asymptomatic, raising the likelihood of developing permanent atrial fibrillation (AF) within 5 years by 4–5 times.² Most episodes occur between 2 and 5 days post-surgery, primarily in conventional surgical units without continuous cardiac monitoring, unlike critical care units where rhythm monitoring is routine.² The absence of monitoring in these settings leaves asymptomatic cases undiagnosed, exposing patients to greater risks of adverse outcomes.⁵ Treatment often involves bradycardic agents and, in cases exceeding 48 hours, anticoagulation therapy based on thromboembolic and bleeding risk scores.²

Smartwatches equipped with single-lead ECG capabilities and algorithms for AF detection provide a promising solution for identifying POAF with high accuracy.^{6,7} These devices use photoplethysmography (PPG), a non-invasive optical technique to analyse blood volume changes in superficial tissues, enabling heart rhythm monitoring and oxygen saturation (SpO₂) measurement.^{8,9} The European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery recommend their use to detect AF, aiming to reduce its economic impact.¹⁰ However, no studies have yet evaluated the effectiveness of smartwatches in detecting POAF after thoracic surgery. Their use could address the unmet need for continuous monitoring in conventional units, enabling early detection and management of POAF.

We hypothesise that the use of smartwatches in patients undergoing elective major thoracic surgery with lung exclusion could significantly enhance the early detection of POAF compared with conventional monitoring and improve diagnostic performance during the hospital stay.

METHODS AND ANALYSIS

Trial design

This is a randomised, prospective, multicentre (two centres), open-label study.

Study population

The inclusion criteria are as follows:

- ▶ Patients more than 18 years old.
- ▶ Patients undergoing major elective thoracic resection surgery with pulmonary exclusion (one-lung ventilation) within the past 48 hours.
- ▶ Elective pneumonectomy or lobectomy surgery.
- ▶ Postsurgery transfer to a conventional care unit.
- ▶ Patients with motor and cognitive ability to perform single-lead ECGs using the smartwatch.
- ▶ Social security coverage.
- ▶ Written informed consent from patient or next of kin.

The exclusion criteria are as follows:

- ▶ History of AF.
- ▶ Need for telemetry monitoring for atrioventricular block and/or rapid supraventricular or ventricular arrhythmias.

- ▶ Unstable condition requiring prolonged hospitalisation in a critical care unit with continuous telemetry.
- ▶ Dependence on atrial or ventricular pacing from a pacemaker.
- ▶ Inclusion in another interventional clinical trial affecting POAF incidence.
- ▶ Mediastinal surgery (eg, mediastinal mass resection or mediastinoscopy).
- ▶ Chest wall surgery (eg, lymph node biopsy or wall repair).
- ▶ Pleural surgery (eg, pneumothorax or pleurisy management).
- ▶ Reoperation of the surgical site (early or delayed).
- ▶ Surgery conducted >48 hours prior.
- ▶ Pregnancy.
- ▶ Coexisting illness with a high probability of death (inferior to 6 months).

Study protocol

Randomisation

The randomisation will be performed using a list with fixed-size random blocks. The randomisation will be stratified, and the stratification criteria will include age >65 years, diabetes, surgical approach via thoracotomy (as opposed to a minimally invasive video thoracoscopy approach). The study is open label, with an intention-to-treat analysis. Randomisation will be conducted using the Ennov Clinical software implemented by a data manager. Randomisation will occur within 48 hours postsurgery (up to the 48th hour) while the patient is hospitalised, provided there are no prior rhythm disturbances or indications for intensive care with telemetry monitoring. The result of randomisation will be displayed as 'Conventional Monitoring' or 'Smartwatch Monitoring' (figure 1). The diagram of the study process is shown in figure 2.

Intervention

In the 'Smartwatch Monitoring' group, POAF will be identified using a smartwatch (ScanWatch, Withings Move ECG, Withings, France), defined as an episode of AF lasting >20 s recorded by a single-lead ECG or POAF detected via PPG signals and/or a 12-lead ECG. More specifically, the smartwatch detects episodes of POAF using its artificial intelligence algorithm. Confirmation is then required, either through a 1-lead ECG recorded by the smartwatch or a 12-lead ECG.

During the second follow-up visit (V2), patients will return the smartwatch on day 7 after surgery or the day of hospital discharge. The main difference in this group is that, in addition to conventional monitoring, the patient can perform a 1-lead ECG independently, either on demand or in the presence of symptoms.

The smartwatch used in this study will be the ScanWatch 42 mm, which is CE-marked (CE 1282) for the diagnosis of AF. The smartwatch allows for heart rate monitoring (continuous monitoring via PPG using three integrated LEDs), single-lead ECG recordings (the smartwatch allows patients to perform 20 s ECG recordings by placing




Figure 1 Consort diagram.

one hand on the watch's case) and data integration (the device is equipped to record and transmit data and ECG tracings via Bluetooth to the Health Mate app). Data from the smartwatch will be retrieved via the Health Mate app by the investigator during follow-up visits (V1 and V2). The smartwatch will be collected from the patient during the second follow-up visit (V2, day 7 or hospital discharge).

'Conventional Monitoring' group postoperative care

In the 'Conventional Monitoring' group, patients from both centres (Amiens University Hospital and Victor Pauchet Clinic) will be screened for POAF using a 12-lead ECG, according to standard monitoring protocols. Specifically, POAF will be diagnosed using a 12-lead ECG, which will only be performed if clinical symptoms and/or significant heart rate variations are observed by medical or paramedical staff. No routine daily ECG will be conducted in the absence of such symptoms. Additionally, all patients (from both groups) will undergo standardised postoperative monitoring, which includes regular assessments every 4 hours throughout the hospital stay. These assessments will include the following parameters: non-invasive blood pressure, heart rate and SpO₂.

In both groups, the management of POAF will involve several essential steps:

- Assessment of haemodynamic stability: evaluate haemodynamic tolerance and perform cardioversion in cases of poor tolerance.
- Correction of coexisting disorders: address underlying issues such as hypoxemia, postoperative complications, sepsis or electrolyte imbalances.
- Treatment of AF: implement rhythm control strategies and initiate antiarrhythmic therapy.
- Decision on curative anticoagulation: in accordance with guidelines, anticoagulation decisions should be based on several factors, primarily the patient's thromboembolic risk as assessed by the CHA₂DS₂-VASc score and the absence of significant haemorrhagic risk.
- Follow-up consultation: provide a prescription and schedule a cardiology follow-up appointment after discharge.

Endpoints

Primary endpoint

The endpoints and definitions are presented in [table 1](#).

The primary outcome measure will be the occurrence of POAF after major elective thoracic surgery within seven postoperative days.

POAF is defined according to the 2020 ESC recommendations.²




Figure 2 Diagram of the study process. MACE, major adverse cardiovascular events; POAF, postoperative atrial fibrillation.

- POAF diagnosed by a smartwatch (ScanWatch, Withings Move ECG, Withings, France), defined as an episode of AF lasting >20s recorded by a single-lead ECG or POAF detected via PPG signals.
- POAF diagnosed in the standard care group will be confirmed using a 12-lead ECG.
- POAF diagnosis will be confirmed through interpretation of single-lead or 12-lead ECGs by a cardiologist.

Secondary endpoint

The secondary endpoints will be

- Asymptomatic POAF prevalence, assessed by the number of patients presenting with postoperative POAF without symptoms. Symptoms will be evaluated using the EHRA score (online supplemental appendix).
- Cardiovascular prognosis, assessed using a composite endpoint of postoperative cardiovascular complications (major adverse cardiovascular events (MACE) criteria) as defined by European Society of Anaesthesiology (ESA) and European Society of Intensive Care Medicine (ESCIM) standards.¹¹ MACE criteria will be analysed at 2 months and 6 months and validated in the presence of one of the followings:
 - *Stroke*: haemorrhagic or ischaemic stroke, defined by objective criteria including clinical assessment (persistent sensory, motor or cognitive deficits), neuroimaging and standardised scales (National

Institutes of Health Stroke Scale (NIHSS) or mRS).^{12 13}

- *Myocardial infarction*: elevated cardiac markers (eg, myoglobin, troponin I) above the 99th percentile per lab standards; ischaemic symptoms; ST-segment changes or left bundle branch block on ECG; Q waves or evidence of coronary thrombus via angiography or autopsy.
- *Congestive acute heart failure*: clinical syndrome characterised by symptoms and/or signs resulting from a structural and/or functional cardiac abnormality, accompanied by elevated natriuretic peptide levels and objective evidence of pulmonary or systemic congestion.^{14 15}
- *Mesenteric ischaemia*: mesenteric ischaemia confirmed by imaging or exploratory laparotomy, and/or ischaemic colitis confirmed by digestive endoscopy or laparotomy.
- *Resuscitated cardiac arrest*: loss of mechanical cardiac activity confirmed by the absence of clinical signs of circulation.
- *Cardiovascular death*.
- Feasibility of monitoring POAF using a smartwatch after thoracic surgery: Evaluated based on device wear time and the completion rate of patient-performed single-lead ECGs in case of POAF detection.
- Recurrence and management of POAF: rhythm status (AF yes/no) at 2 and 6 months, treatment for

Table 1 Endpoints and definitions

Endpoints	Definitions
Primary endpoint	
Occurrence of POAF after major elective thoracic surgery within seven postoperative days.	<p>POAF is defined according to the 2020 ESC recommendations.</p> <ul style="list-style-type: none"> ▶ POAF diagnosed by a smartwatch (ScanWatch, Withings Move ECG, Withings, France), defined as an episode of AF lasting >30s recorded by a single-lead ECG or POAF detected via PPG signals and confirmed by a 12-lead ECG ▶ POAF diagnosed in the standard care group will be confirmed using a 12-lead ECG ▶ POAF diagnosis will be confirmed through interpretation of single-lead or 12-lead ECGs by a cardiologist
Secondary endpoints	
Asymptomatic POAF prevalence	<ul style="list-style-type: none"> ▶ Number of patients presenting with postoperative POAF without symptoms. Symptoms will be evaluated using the EHRA score (online supplemental appendix).
Major cardiovascular and cerebral event (MACE)	<p>One of the following criteria (definitions above):</p> <ul style="list-style-type: none"> ▶ Stroke ▶ Myocardial infarction ▶ Acute kidney injury ▶ Mesenteric ischaemia ▶ Successful resuscitated cardiac arrest
Stroke	An embolic, thrombotic or haemorrhagic cerebral event with persistent residual motor, sensory or cognitive dysfunction (eg, hemiplegia, hemiparesis, aphasia, sensory deficit, impaired memory) diagnosed on a neuroimaging
Myocardial infarction	Myocardial infarction was diagnosed by the characteristics presentation, serial changes on 12-lead electrocardiographic suggesting infarction, and arise in cardiac markers, preferably cardiac troponins, with at least one value above the 99th percentile of the upper reference limit
Congestive acute heart failure	Clinical syndrome characterised by symptoms and/or signs resulting from a structural and/or functional cardiac abnormality, accompanied by elevated natriuretic peptide levels and objective evidence of pulmonary or systemic congestion Symptoms can be dyspnoea or fatigue, orthopnoea, paroxysmal nocturnal dyspnoea, increased jugular venous pressure or pulmonary rales on physical examination
Mesenteric ischaemia	Mesenteric ischaemia confirmed by imaging or exploratory laparotomy and/or ischaemic colitis confirmed by gastrointestinal endoscopy or exploratory laparotomy
Resuscitated cardiac arrest	Cessation of mechanical cardiac activity confirmed by the absence of clinical signs of blood flow
Feasibility of monitoring POAF using a smartwatch after thoracic surgery	Evaluated based on device wear time and the completion rate of patient-performed single-lead ECGs in case of POAF detection
Recurrence and management of POAF	Rhythm status (AF yes/no) at 3 and 6 months, treatment for cardioversion (eg, electrical cardioversion, ablation or medication)
AF, atrial fibrillation; EHRA, European Heart Rythm Association; POAF, postoperative atrial fibrillation.	

cardioversion (eg, electrical cardioversion, ablation or medication).

Data collection and outcome definitions

All collected variables are summarised in [table 2](#). The following data will be collected: age (years), gender, body mass index (kg/m), usual medication, medical history (coronary disease, peripheral vascular disease, stroke, smoking, diabetes, dyslipidaemia, chronic obstructive pulmonary disease, sleep apnoea syndrome, hypertension, chronic kidney disease, creatinine clearance, blood creatinine levels, surgery type, neurological conditions (stroke, transient ischaemic attack (TIA), dementia), deep vein thrombosis/pulmonary embolism, heart failure, ECG, CHAD₂-VASC₂ score and HASBLED score.

Also, the following perioperative data will be collected: date of surgery, nature of the surgical procedure performed, surgical approach: minimally invasive surgery with video thoracoscopy, robot-assisted surgery, conversion to thoracotomy, duration of anaesthesia, type of regional anaesthesia performed, total dose of phenylephrine, ephedrine, norepinephrine, nicardipine, dobutamine, epinephrine administered, administration of a beta-blocker or amiodarone during surgery. Fluid balance will be considered by taking into account intraoperative fluid management (crystalloids, colloids and volume in mL), administration of blood products (number and volume in mL), and administration of blood product derivatives (type and volume).

Table 2 Collected variables

Collected variables	Details of collected variables
Preoperative variables	
Demographic data	Age (years), gender, body mass index (kg m^{-2})
Usual medication	Calcium channel blockers, ACE inhibitors, aldosterone antagonists, beta-blockers, diuretics, statins, sacubitril/valsartan, bronchodilators, oral antidiabetic agents, insulin, antiplatelet agents, anticoagulants, immunosuppressants, antidepressants, antipsychotics, benzodiazepines.
Medical history	Coronary disease, peripheral vascular disease, stroke, smoking, diabetes, dyslipidaemia, chronic obstructive pulmonary disease, sleep apnoea syndrome, hypertension, chronic kidney disease, surgery type, neurological conditions (stroke, transient ischaemic attack, dementia), deep vein thrombosis/pulmonary embolism, heart failure. Baseline ECG, CHAD ₂ -VASC ₂ score and HASBLED score.
Biological data	Blood creatinine levels, creatinine clearance
Perioperative data	
Surgical variables	Date of surgery, nature of the surgical procedure performed (lobectomy or pneumectomy), surgical approach: minimally invasive surgery with video thoracoscopy, robot-assisted surgery, conversion to thoracotomy.
Medical variables	Duration of anaesthesia and type of regional anaesthesia performed. Total dose of phenylephrine, ephedrine, norepinephrine, nicardipine, dobutamine or epinephrine administered. Administration of a beta-blocker, amiodarone, magnesium sulfate or dexamethasone during surgery. Intraoperative fluid administration (type of fluid and volume in mL), administration of blood products (type of blood product, number and volume in mL), and administration of blood product derivatives (type and volume).
Postoperative data	
Medical variables	Duration of orotracheal intubation and reintubation. Duration, type and total dosage of catecholamines and postoperative VIS score. Thoracic drain output and urine output. Reoperation. As defined in table 1 , MACE criterion will be recorded. Also, we will assess respiratory, infectious, neurological and renal complications. Duration of stay in continuous care and duration of hospital stay.
Biological variables	Plasma haemoglobin, serum creatinine, urea, sodium, potassium, calcium, phosphorus, troponin US and BNP
General haemodynamic variables	Heart rate, systolic, diastolic, and mean blood pressure, oxygen saturation and oxygen therapy administered.
Follow-up visit	
Smartwatch Monitoring group	Number of single-lead ECGs performed, presence of POAF, duration of smartwatch usage, clinical symptoms (EHRA as presented in online supplemental appendix), duration of POAF and treatment provided for POAF.
Conventional Monitoring group	Number of 12-lead ECGs performed, presence of POAF, clinical symptoms (EHRA in online supplemental appendix), duration of POAF and treatment provided for POAF
EHRA, European Heart Rhythm Association; MACE, major adverse cardiovascular events; POAF, postoperative atrial fibrillation; VIS, vasoactive inotropic score.	

Additionally, thoracic drain output and urine output will be recorded.

Postoperative data include duration of orotracheal intubation, duration, type and total dosage of catecholamines, postoperative vasoactive inotropic score, reoperation, reintubation, MACE criterion, respiratory, infectious, neurological and renal complications, duration of stay in continuous care, and duration of hospital stay. Biological data will be recorded (plasma haemoglobin, serum creatinine, urea, sodium, potassium, calcium, phosphorus,

troponin US, leukocytes, CRP, TSH, BNP). During hospitalisation, the following clinical parameters will be recorded: general haemodynamic parameters (heart rate, systolic, diastolic and mean blood pressure, SpO₂, oxygen therapy administered) and the MACE criterion.

For the 'Smartwatch Monitoring' group: number of single-lead ECGs performed, presence of POAF, duration of smartwatch usage, clinical symptoms (EHRA, in online supplemental appendix), duration of POAF and treatment provided for POAF. For the 'Conventional

Monitoring' group: number of 12-lead ECGs performed, presence of POAF, clinical symptoms (EHRA, in online supplemental appendix), duration of POAF, and treatment provided for POAF.

Two months after surgery and 6 months after surgery, the following data will be collected: MACE criterion occurrence, the recurrence of POAF and management of POAF: rhythm status (AF yes/no), treatment for cardioversion (external electrical shock, ablation, pharmacological treatment). The recurrence of AF will be recorded in the hospital by performing an ECG if the patient is hospitalised. If the patient is not hospitalised, the data will be collected through a phone call to the patient and by reviewing the medical record (consultation for cardioversion, cardiology follow-up, follow-up for other conditions).

Endpoints will be assessed after thoracic surgery. Adverse events will be declared and notified in the electronic case report forms (eCRFs).

Standard definitions of postoperative outcomes established by the ESA will be used.¹¹ Cardiac arrest is defined as the cessation of cardiac mechanical activity, as confirmed by the absence of circulation signs. Stroke is defined as an embolic, thrombotic or haemorrhagic cerebral event with persistent residual motor, sensory or cognitive dysfunction (eg, hemiplegia, hemiparesis, aphasia, sensory deficit, impaired memory) diagnosed on a neuroimaging. Acute kidney injury is defined according to Kidney Disease Improving Global Outcomes criteria as an increase in serum creatinine of over 27 µmol/L within 48 hours or diuresis lower than 0.5 mL/kg/h.¹⁶ Myocardial injury is diagnosed by the characteristic presentation, serial changes on 12-lead electrocardiographic suggesting infarction, and arise in cardiac troponin, with at least one value above the 99th percentile of the upper reference limit.¹⁷ Mesenteric ischaemia will be confirmed by imaging or exploratory laparotomy, and ischaemic colitis will be confirmed by gastrointestinal endoscopy or exploratory laparotomy.

Intention-to-treat analysis

Patients with serious adverse events will be analysed according to their assigned group following the intention-to-treat principle.

Statistical method and sample size calculation

In recent studies, the incidence of POAF in the studied population was 20%, with AF predominantly occurring within the first 4 postoperative days.¹⁸ The percentage of POAF at day 7 is not precisely known, but since most cases occur relatively early, we estimate this percentage to be around 20%. Based on these assumptions, we would need 286 evaluable patients (142 per group) to detect an absolute difference of 12% (20% vs 8%) in the incidence of AF with a two-sided alpha risk of 5% and a power of 80%. Anticipating 5% of non-evaluable patients, we will randomise 302 patients (151 per group).

Primary endpoint (incidence of POAF) will be compared between the two groups using either a χ^2 test or Fisher's exact test, with a significance level set at 5%.

Regarding secondary endpoints, the percentage of asymptomatic POAF in the 'Smartwatch Monitoring' group will be calculated with a 95% CI. The percentage of MACE at 6 months among patients diagnosed with asymptomatic POAF in the 'Smartwatch Monitoring' group will also be calculated with a 95% CI. In the 'Smartwatch Monitoring' group, the duration of MI use will be expressed as mean±SD or median (range). The percentage of patients performing a single-lead ECG in the case of POAF will be calculated with a 95% CI. A p-value of 0.05 will be considered as significant. No intermediate analysis is planned in the trial. Statistical analyses will be performed using SAS software V.9.4.

Data management and monitoring

Registration

Data will be collected and registered using eCRFs by a dedicated local research technician. A research coordinator will centralise and verify the data.

Record keeping

Consent forms and eCRFs will be retained for 15 years at the University Hospital of Amiens in accordance with French law.

Study organisation

The study promotion is performed by the University Hospital of Amiens, France. The Victor Pauchet Clinic is an associated investigator centre.

Duration and timeline

Patients from Amiens University Hospital and Victor Pauchet Clinic can be included during a 2-year period. We plan to begin the study in April 2025 and complete it by 2027. The processes of developing the protocol, obtaining approval from the ethical committee, obtaining financial support and developing the eCRFs occurred in 2024. The database should be closed after all participants have been included, followed by data analysis, manuscript writing and submission for publication.

ETHICS AND DISSEMINATION

Ethical approval

The institutional review board (IRB) of the University Hospital of Amiens (*Comité de Protection des Personnes Sud-Ouest et outre-mer 1, 21050 Toulouse, France*) approved this bicentric study (registration number ID RDB: 2022-A02028-27 in November 2024). The THOFAWATCH study will be conducted in accordance with the Declaration of Helsinki and French law on clinical research¹⁹ and was registered on the 5 December 2024 on the ClinicalTrials.gov website with the trial identification number NCT06724718. THOFAWATCH trial follows Consolidated Standards of Reporting Trials (CONSORT) Statement—CONSORT diagram is given in figure 1.²⁰

Consent to participate

Written informed consent will be obtained from all participants or next of kin (the participant consent form is provided as an online supplemental file).

Patients or public involvement

Patients or the public will not be involved in the design, or conduct, or reporting or dissemination plans of our research. Authors will be involved in disseminating research findings (through attending conferences and co-authoring results 'papers').

Access to data

Data sharing is not applicable to this article because no datasets were generated or analysed during the current study. However, the investigators will provide authorised personnel with access to the necessary documents and individual data for study monitoring, quality control and auditing in compliance with applicable regulations.

DISCUSSION

The hypothesis suggests that utilising smartwatches for patients undergoing planned major thoracic surgery may notably enhance the detection of postoperative POAF compared with traditional monitoring methods.

POAF after thoracic surgery arises due to a combination of autonomic imbalance, systemic inflammation and atrial remodelling. The surgical stress response activates the sympathetic nervous system, which, along with inflammation from tissue injury and oxidative stress, contributes to electrophysiological instability.^{21–23} These factors interact dynamically during the perioperative period, leading to the development of transient or sustained AF episodes.²⁴

POAF is a recurrent issue following major surgery, particularly in thoracic surgery.² Smartwatches have demonstrated good diagnostic performance, with high sensitivity and specificity, in non-surgical settings.¹⁰ Several factors influence signal accuracy, including advanced arterial disease or hypothermia. The key challenge lies in detecting arrhythmias and diagnosing asymptomatic or highly transient AF episodes postoperatively. Most arrhythmias are paroxysmal, and a standard 12-lead ECG often fails to identify these episodes. Thus, we designed this study to evaluate the value of continuous smartwatch monitoring to overcome this limitation.

A recent study by Monteiro *et al* on 108 patients showed that smartwatches, compared with conventional monitoring, provided improved monitoring, early arrhythmia detection and better patient outcomes.²⁵

In non-surgical populations, smartwatches using PPG are effective for AF monitoring, with high detection accuracy. However, heart rate measurements may underestimate elevated rates. Also, improving diagnostic reliability in older adults may require training for participants and cardiologists, along with sufficient ECG recordings to ensure accuracy.^{7,26}

However, few studies to date have focused on this topic after thoracic surgery. We hypothesise that continuous smartwatch use can enhance AF detection following thoracic surgery. To limit the scope, smartwatch monitoring is restricted to a maximum of 7 days, as POAF predominantly occurs early in the postoperative period.

This study holds several key strengths. It is the first study, to our knowledge, that explores AF detection in the specific context of postoperative thoracic surgery, filling a crucial gap in the current literature. Furthermore, its multicentre design ensures diverse patient representation and robust data collection, while the inclusion of a significant sample size enhances the reliability and generalisability of its findings.

This study aims to enable early diagnosis of AF, reduce economic and hospitalisation burdens and improve patient outcomes through tailored care and remote monitoring protocols.

Author affiliations

¹Anesthesiology and critical care, Pauchet santé, Victor Pauchet Clinic, Amiens, France

²Anesthesiology and critical care, Amiens University Hospital, Amiens, Hauts-de-France, France

³Department of Statistics, Amiens Hospital University, Amiens, France

⁴Department of Thoracic surgery, Amiens Hospital University, Amiens, France

⁵Department of Thoracic Surgery, Pauchet santé, Victor Pauchet Clinic, Amiens, France

⁶Department of Cardiology, Amiens University hospital, Amiens, France

⁷Department of Cardiology, Victor Pauchet Clinic, Amiens, France

Acknowledgements We would like to sincerely thank Annabelle Boussault, project manager in the Clinical Research Department of the Amiens University Hospital, for her assistance in designing this research project.

Collaborators THOFAWATCH study group: Michaël Bernasinski, Adrien Coupez, Patricia Besserve, Alice Henry, Camille Daumin, Ammar BenAmmar, Florent Levie, Pierre-Yves Macq, Beatris Alina Labont, Gilles David, Julien Epailly, Paul-Emmanuel Esmard, Agathe Vernier, Pauline Bartoli, Jérôme Delmas, Sophie Romby, Simon Marx, Mohammed Fikri, Hélène Benoit-Fallet, Eric Nam, Jérôme Zakine, Amandine Zaatar, Thomas Flet.

Contributors PH and CB participated in the design of the study and helped to write the manuscript. OA, AI, GH, MG, TL, HD and YM participated in the design of the study. MD will perform the statistical analysis. All authors read and approved the final manuscript. PH acted as guarantor. Roles and responsibilities: Coordinating center: Boussault Annabelle, Amiens Hospital University. Data manager team: Esteban Soares, Amiens Hospital University. Monitoring committee: Momar Diouf, Amiens Hospital University. Monitoring of adverse events: Directorate of Clinical Research, Amiens Hospital University.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patients.

Provenance and peer review Not commissioned; externally peer-reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible

for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Pierre Huette <http://orcid.org/0000-0003-1563-4869>

REFERENCES

- 1 Conen D, Ke Wang M, Popova E, *et al*. Effect of colchicine on perioperative atrial fibrillation and myocardial injury after non-cardiac surgery in patients undergoing major thoracic surgery (COP-AF): an international randomised trial. *Lancet* 2023;402:1627–35.
- 2 Hindricks G, Potpara T, Dagres N, *et al*. Corrigendum to: 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J* 2021;42:4194.
- 3 Frendl G, Sodickson AC, Chung MK, *et al*. 2014 AATS guidelines for the prevention and management of perioperative atrial fibrillation and flutter for thoracic surgical procedures. *J Thorac Cardiovasc Surg* 2014;148:e153–93.
- 4 Caldonazo T, Kirov H, Rahouma M, *et al*. Atrial fibrillation after cardiac surgery: A systematic review and meta-analysis. *J Thorac Cardiovasc Surg* 2023;165:94–103.
- 5 Karakasis P, Pamporis K, Siontis KC, *et al*. Major clinical outcomes in symptomatic vs. asymptomatic atrial fibrillation: a meta-analysis. *Eur Heart J* 2025;46:1189–202.
- 6 Turakhia MP, Desai M, Hedlin H, *et al*. Rationale and design of a large-scale, app-based study to identify cardiac arrhythmias using a smartwatch: The Apple Heart Study. *Am Heart J* 2019;207:66–75.
- 7 Gruwez H, Ezzat D, Van Puyvelde T, *et al*. Real-world validation of smartphone-based photoplethysmography for rate and rhythm monitoring in atrial fibrillation. *Europace* 2024;26:euae065.
- 8 Guo Y, Wang H, Zhang H, *et al*. Mobile Photoplethysmographic Technology to Detect Atrial Fibrillation. *J Am Coll Cardiol* 2019;74:2365–75.
- 9 Pereira T, Tran N, Gadhoumi K, *et al*. Photoplethysmography based atrial fibrillation detection: a review. *NPJ Digit Med* 2020;3:3.
- 10 Van Gelder IC, Rienstra M, Bunting KV, *et al*. 2024 ESC Guidelines for the management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2024;45:3314–414.
- 11 Jammer I, Wickboldt N, Sander M, *et al*. Standards for definitions and use of outcome measures for clinical effectiveness research in perioperative medicine: European Perioperative Clinical Outcome (EPCO) definitions: a statement from the ESA-ESICM joint taskforce on perioperative outcome measures. *Eur J Anaesthesiol* 2015;32:88–105.
- 12 Ko CY, Hall BL, Hart AJ, *et al*. The American College of Surgeons National Surgical Quality Improvement Program: achieving better and safer surgery. *Jt Comm J Qual Patient Saf* 2015;41:199–204.
- 13 Hicks KA, Mahaffey KW, Mehran R, *et al*. 2017 Cardiovascular and Stroke Endpoint Definitions for Clinical Trials. *Circulation* 2018;137:961–72.
- 14 McDonagh TA, Metra M, Adamo M, *et al*. 2023 Focused Update of the 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2023;44:3627–39.
- 15 Ostrominski JW, DeFilippis EM, Bansal K, *et al*. Contemporary American and European Guidelines for Heart Failure Management: JACC: Heart Failure Guideline Comparison. *JACC Heart Fail* 2024;12:810–25.
- 16 Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. *Nephron Clin Pract* 2012;120:c179–84.
- 17 Thygesen K, Alpert JS, Jaffe AS, *et al*. Fourth universal definition of myocardial infarction (2018). *Eur Heart J* 2019;40:237–69.
- 18 Diallo E-H, Brouillard P, Raymond J-M, *et al*. Predictors and impact of postoperative atrial fibrillation following thoracic surgery: a state-of-the-art review. *Anaesthesia* 2023;78:491–500.
- 19 Toulouse E, Masseguin C, Lafont B, *et al*. French legal approach to clinical research. *Anaesth Crit Care Pain Med* 2018;37:607–14.
- 20 Schulz KF, Altman DG, Moher D, *et al*. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *Int J Surg* 2011;9:672–7.
- 21 Bessissow A, Agzarian J, Shargall Y, *et al*. Colchicine for Prevention of Perioperative Atrial Fibrillation in patients undergoing lung resection surgery: a pilot randomized controlled study. *Eur J Cardiothorac Surg* 2018;53:945–51.
- 22 Hunt TEF, Traaen GM, Aakerøy L, *et al*. Cardiac remodelling in patients with atrial fibrillation and obstructive sleep apnoea. *Open Heart* 2024;11:e002718.
- 23 Shahian DM, Paone G, Habib RH, *et al*. The Society of Thoracic Surgeons Preoperative Beta Blocker Working Group Interim Report. *Ann Thorac Surg* 2025;119:476–84.
- 24 Hayashi T, Sano Y, Tanaka K, *et al*. Predictors of postoperative atrial fibrillation after lung resection. *Curr Probl Surg* 2024;61:S0011-3840(24)00064-9.
- 25 Monteiro R, Rabello GCM, Moreno CR, *et al*. Enhancing cardiac postoperative care: a smartwatch-integrated remote telemonitoring platform for health screening with ECG analysis. *Front Cardiovasc Med* 2024;11:1443998.
- 26 Hibbitt K, Brimicombe J, Cowie MR, *et al*. Reliability of single-lead electrocardiogram interpretation to detect atrial fibrillation: insights from the SAFER feasibility study. *Europace* 2024;26:euae181.