

BMJ Open Barriers in implementing antibiotic stewardship programmes at paediatric units in academic hospitals in Thailand: a qualitative study

Suvaporn Anugulruengkitt ^{1,2}, Thidarat Jupimai,² Prissana Wongharn,² Thanyawee Puthanakit^{1,2}

To cite: Anugulruengkitt S, Jupimai T, Wongharn P, *et al.* Barriers in implementing antibiotic stewardship programmes at paediatric units in academic hospitals in Thailand: a qualitative study. *BMJ Open* 2025;**15**:e092509. doi:10.1136/bmjopen-2024-092509

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-092509>).

Received 16 August 2024
Accepted 12 May 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

¹Department of Pediatrics, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

²Center of Excellence for Pediatric Infectious Diseases and Vaccines, Chulalongkorn University, Bangkok, Thailand

Correspondence to

Dr Suvaporn Anugulruengkitt; suvaporn.a@chula.ac.th

ABSTRACT

Objective To explore the barriers that hinder and the facilitators that strengthen the implementation of the antimicrobial stewardship (AMS) programme at paediatric units in academic hospitals in Thailand.

Design A qualitative study using thematic analysis of interviews with healthcare staff.

Setting Five paediatric units in academic hospitals in Thailand.

Participants 20 healthcare workers and 10 AMS service providers who actively participated in the AMS programme in the sampled hospitals were included from purposive criterion.

Primary outcome measures Qualitative, interpretive description with semistructured individual interviews were digitally recorded and transcribed. The MAXQDA software was used to facilitate content analysis.

Results In total, 4 themes emerged from the data: (1) organisational hierarchical culture and individual behaviours influence the acceptance and adherence to AMS implementation, (2) changing healthcare workers' mindset to improve stewardship is crucial, (3) effective communication and collaboration among healthcare teams are the key to implementing the AMS programme and (4) dedication to antimicrobial stewardship despite resource limitations is important to improve AMS programme implementation.

Conclusions To implement antimicrobial stewardship in a paediatric setting, there are many issues to overcome. The key barriers to focus were organisational hierarchical culture and perception of healthcare workers. Support from hospital policy and effective communication with contextualised strategies should be considered to improve AMS programme implementation plans.

INTRODUCTION

Antimicrobial stewardship (AMS) programmes are a strategy to promote the judicious use of antibiotics, optimise patient outcomes and reduce the risk of developing resistance.¹ AMS has multifaceted strategies such as prospective audit and feedback, prior authorisation, education, clinical decision support and rapid diagnostics.² Strategies

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The study focuses on academic tertiary care hospitals, combined with specific inclusion and exclusion criteria.
- ⇒ Data were collected through interviews with both stewards and healthcare providers involved in antimicrobial stewardship.
- ⇒ A combined inductive and deductive approach was used for coding and data analysis.
- ⇒ The findings may have limited generalisability to other healthcare settings.

selection should be based on the availability of facility-specific resources for consistent implementation and tailored to local settings.¹ Implementing antibiotic stewardship can be challenging because it involves changing knowledge, deeply held attitudes, cultural norms and the emotionally influenced behaviours of clinicians and patients towards prescribing antibiotics and their use.³ Implementation science can help address this challenge. Implementation science principles can inform local antibiotic stewardship efforts and identify the gap between evidence-based practice and routine practice in real-world settings.⁴ Implementation frameworks should be identified and employed in research on implementing the AMS programme. Determinant frameworks such as the Consolidated Framework for Implementation Research (CFIR) can be used to understand different factors that enable or hinder the implementation process.^{5 6} The CFIR model has been used to study the perceptions of antibiotic stewardship personnel regarding why their programmes were successful by conducting a qualitative study across different hospital settings to examine contextual factors that facilitate the success of their programmes.⁷

Prescribing the correct antibiotics for children requires careful decision-making, as

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

BMJ Open: first published as 10.1136/bmjopen-2024-092509 on 23 May 2025. Downloaded from <http://bmjopen.bmj.com/> on June 10, 2025 at Agence Bibliographique de l'Enseignement Supérieur (ABES).

Table 1 Baseline characteristics of the 30 interview participants

Hospital levels	Number of interviewees	Profession	Gender	Mean age (range) (years)
Healthcare workers	20	Physician 19 Pharmacist 1	Male 3 Female 7	39.2 (29–57)
AMS service providers	10	Physician 5 Pharmacist 5	Male 8 Female 12	36.3 (26–56)
Overall	30	Physician 24 (80%) Pharmacist 6 (20%)	Male 11 (37%) Female 19 (63%)	37.0 (26–57)

AMS, antimicrobial stewardship.

errors in dosing can have serious consequences. However, in resource-limited settings, healthcare providers (HCPs) treating children often lack specialised training in paediatric infectious diseases or antibiotic stewardship, further affecting their ability to make informed prescribing decisions. In low- and middle-income settings, there are unique difficulties encountered in paediatric care regarding AMS such as limited access to diagnostics, unavailable antibiotic formulations, lack of education and limited resources of healthcare (staffing and equipment).⁸ A published study highlighted that the most commonly perceived barriers to paediatric AMS were a lack of education about antibiotics and a lack of support from hospital management. Furthermore, an additional barrier in low- and middle-income settings was lack of recognition of AMS importance by senior doctors.⁸ In addition, prescribing behaviour among colleagues and hierarchical structure within the medical field were found as factors under the social influences domain.⁹

In Thailand, AMS in the paediatric setting is not widely implemented, and the intervention is not highly accepted. Based on a previous study at a tertiary care centre, overall acceptance of AMS programme recommendations for carbapenem de-escalation was only 57.8%.¹⁰ Due to this low acceptance, implementation research is needed to explore the underlying reasons such as barriers and facilitators and to inform how interventions should be adapted. In this study, researchers conducted implementation research on AMS across five paediatric units in academic hospitals in Thailand to identify barriers and facilitators of implementing AMS across different settings.

METHODS

A qualitative study with semistructured interviews was conducted in 2022 to gather data from healthcare providers. The Ethics Committee of the Faculty of Medicine, Chulalongkorn University has approved the study (IRB 236/65).

Study design

Semistructured interviews were conducted with HCPs. The qualitative study was conducted to identify: (1) barriers and facilitators of current AMS intervention at King Chulalongkorn Memorial Hospital (KCMH) where AMS implementation has been launched in paediatric setting since 2017 and (2) barriers and facilitators to implement AMS at another four hospitals where AMS has not been established in paediatric setting. These include organisational context such as information technology support, staffing resources, organisational climate and culture and leadership support. The interviews also assessed local context and barriers to implementation: stakeholder perceptions and attitudes, current antibiotic prescribing problems and preferences for potential methods of implementation strategies.

Study setting and participants

The study was conducted by the Faculty of Medicine, Chulalongkorn University. We collected data from five academic tertiary care hospitals in Thailand. The site selection was recruited using convenient sampling, focusing on a centre where paediatric infectious diseases specialists had <10 years of work experience. Interviews were conducted with both healthcare providers involved in antibiotic prescribing and steward teams who were AMS services providers.

Target population

Healthcare providers

Inclusion criteria: HCPs who provide care at paediatric units at the Department of Paediatrics, KCMH and another four hospitals (academic tertiary care hospitals: Charoenkrung Pracharak Hospital, Vachira Phuket Hospital, Trang Hospital and Hat Yai Hospital)

Exclusion criteria: HCPs who provided <3 months of service at paediatric units.

Service providers of AMS teams

Inclusion criteria: Current HCPs who are steward teams at the Paediatrics Department, KCMH, such as physicians, nurses and pharmacists.

Exclusion criteria: HCPs who had been involved with AMS services for <3 months.

Sample size calculation for semistructured interviews

The initial target of the number of participants was to interview 20 healthcare workers and 10 AMS service providers (SPs). Interviews continued until thematic saturation was reached.¹¹ Ultimately, thematic saturation was achieved, and no new barriers or enablers were identified.

Interview guide and data collection

An interview guide was created by the research team with guidance from a comprehensive literature review.^{11–14} We used the CFIR model to inform interviews which were conducted among the steward team and HCPs to identify barriers and facilitators of AMS implementation. Open-ended questions were asked about current prescribing

practices, prior education about antibiotic stewardship and views about the stewardship programme at the hospital. Participants also answered a questionnaire regarding AMS implementation and interventions (online supplemental material). The interview guide was piloted on two clinicians and revised based on feedback from the study team. Co-investigators (PW and TJ) conducted the interviews. Each semistructured interview lasted 40–60 min. All participants provided informed consent to participate. The participant information sheet outlining the study was provided to all potential participants. Participants were informed of their rights, and confidentiality was assured. Their written consent was then obtained before each interview started. The interviews were audio recorded. The interviews began on 24 August 2022 and were completed on 6 Oct 2022. All interviews were conducted in Thai. Illustrative quotes were translated into English by one translator, then back-translated and reviewed by a second to ensure accuracy and maintain data integrity. The codebook was initially developed based on a review of the literature and the interview guide. It was then iteratively refined through a process of pilot coding and discussion among the research team.

Data analysis

Transcripts were analysed using MAXQDA (VERBI Software, 2022).¹⁵ Transcripts were coded inductively in an iterative manner by three independent reviewers (SA, PW and TJ). We employed intercoder reliability checks using Cohen's Kappa >0.8. The coding discrepancies have been further discussed and were resolved by consensus. The research team used a combined inductive and deductive approach to code and analyse data. Main themes were identified through an inductive process while CFIR domains were used to deductively code for subthemes.

Patient and public involvement

Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

RESULTS

In a total of 30 interviewees, there were 24 physicians and six pharmacists. Regarding gender, 63% of the participants were women. The mean age was 37 years (range 26–57 years). There were 14 (47%) healthcare workers who worked directly related to infectious diseases. Demographic characteristics of the participants were shown in [table 1](#).

Four themes were identified characterising the implementation of AMS in five paediatric units within academic hospitals in Thailand. These themes and subthemes are summarised in [table 2](#). Themes and subthemes linked to CFIR are presented in [figure 1](#).

The impact of organisation and individual factors

Hierarchical influence on prescribing

In Thailand, the context of seniority refers to the traditional cultural value placed on respecting and honouring

Table 2 Summary of themes and subthemes

Themes	Subthemes	Related CFIR domains
The impact of organisation and individual factors	Hierarchical influence on prescribing	Inner setting – culture
	Resource limitations and systemic barriers	Inner setting – available resources
	Knowledge and beliefs impacting prescribing	Characteristics of individuals – implementation team members and capability
Changing healthcare workers' mindset to improve stewardship	Addressing misconceptions and building trust	Characteristics of individuals – implementation team members
	Navigating authority and autonomy	Inner setting – relational connections and communications
Effective communication and collaboration among healthcare teams	Overcoming perceptual barriers and fostering mutual understanding	Characteristics of individuals – implementation team members
	Establishing communication channels for stewardship integration	Inner setting – communications
Dedication to antimicrobial stewardship despite resource limitations	Intrinsic motivation and professional fulfilment	Characteristics of individuals – motivation and need
	The reality of workload	Intervention characteristics – complexity and inner setting – implementation climate

CFIR, Consolidated Framework for Implementation Research.

elders or those who hold higher positions in an organisation. This concept is deeply rooted in Thai culture and is an essential aspect of social interactions and relationships. In a hospital setting, participants reported that the concept of seniority was significant. This affects the AMS when the steward team suggests an intervention to paediatric residents, but the final decision depends on the senior attending physician. Participants noted that the existing hospital system and culture did not readily support direct communication from the stewardship team to senior attending physician, potentially creating a barrier to implementing AMS recommendations.

Well, sometimes you know the senior attendings might prescribe antibiotic, and you think, 'maybe

Implementation of AMS in paediatric units in 5 academic hospitals in Thailand




Figure 1 Themes and subthemes linked to the Consolidated Framework for Implementation Research.

that's not the best choice,' but you don't really want to say anything. They have so much more experience. (HCP10)

Among medical staff, including attending physicians, there can be differing opinions about the appropriate use of antibiotics. This can lead to inconsistent empirical antibiotic therapy. Senior staff members often have the final say in antibiotic selection, which can override recommendations made based on ASP guidelines. (HCP02)

Resource limitations and systemic barriers

The participants reported the resource limitations such as lack of key personnel, competing priorities, lack of access to resources, problems with data and information systems, inadequate supply of laboratory diagnostic tests and limited available antibiotic options.

We as an ASP pharmacist had obstacles regarding the knowledge and number of key personnel. The expertise in antibiotics may still be limited to a small number of people. A challenge for us is the limited knowledge and number of staff with expertise, especially for complex antibiotic use. (HCP06)

It's put the effort to track and report antibiotic use because our electronic medical record system is not fully supported. This made really hard to monitor trends and identify areas for improvement. (SP02)

Knowledge and beliefs impacting prescribing

AMS stewards have to work to enhance the knowledge and understanding of HCPs regarding appropriate antibiotic use, including the latest guidelines and evidence-based practices. Individual behaviours and resistance to change are other factors, potentially due to established prescribing routines. The participants reported concerns of the frontline physicians about the benefits of the programme or fear of adverse outcomes and established prescribing habits which can act as barriers to change.

A significant barrier is the prescribing behavior of healthcare providers and their fear of adverse outcomes. Due to the complexity of the cases in the tertiary care setting, frontline physicians may be hesitant

to use narrower-spectrum antibiotics for fear of treatment failure or harm to the patient. While this is understandable, it can hinder the implementation of an ASP. (HCP04)

To address this, participants suggested the importance of integrating AMS principles and rational antimicrobial use into medical education at the medical student level. This early exposure, they argued, would foster a culture of teamwork and promote improved patient care within the future healthcare system.

We should start educating medical students about the stewardship program while they are still in medical school. This will prepare them to understand the importance of rational antibiotic use. When they graduate and start working, they will be ready to apply this knowledge immediately (SP01)

Changing healthcare workers' mindset to improve stewardship

Addressing misconceptions and building trust

Changing healthcare workers' mindset to improve stewardship and identify and address concerns or barriers that healthcare workers may have regarding AMS. This could include concerns about patient satisfaction, fear of under-treatment or lack of awareness. The participant from the AMS team suggested adjusting the mindset of the physician team. Different healthcare professionals may have varying perspectives and experiences regarding antibiotic stewardship. In some cases, physicians might have experienced conflicts with the stewardship team over specific patient cases, leading to a perception of friction between the two parties. Also, they might have concerns about the time it takes to consult or the potential delays in initiating antibiotics. The steward participants discussed correcting the AMS team is not the 'antibiotic police'; instead, they play a vital and supportive role in promoting responsible and effective antibiotic use in healthcare settings. They help HCPs make informed decisions about antibiotic prescriptions tailored to each patient's condition.

ASP team must understand that frontline clinicians may have concerns, anxieties or be dealing with

complex or unstable patients. Therefore, ASP team is there to assist in patient care, not to stop antibiotic therapy. (SP01)

Navigating authority and autonomy

The steward team reported lacking the authority to directly intervene in prescribing practices. Consequently, their recommendations were ignored by individual clinicians. This emphasises the need for effective communication and collaboration strategies. Social influences such as resistance from medical staff, lack of leadership and perceived unhelpful attitudes of clinicians also presented challenges. There is a tension between the steward team and clinicians who feel their autonomy is being undermined, leading to resistance and a lack of buy-in from the individual clinicians.

We can make suggestions, but at the end of the day, it's up to the primary physician to decide what to do. We don't have any real power to enforce. (SP06)

Certainly, a major challenge is gaining the understanding of physicians from other specialties or the primary care team. Many healthcare providers perceive the ASP as a team that simply stops antibiotic use. How can we change this perception and make primary teams see the ASP as a supportive specialty team focused on optimizing antimicrobial therapy? This is a significant challenge. (SP09)

Effective communication and collaboration for the success of AMS

Establishing communication channels for stewardship integration

Effective communication and collaboration among healthcare teams are crucial for the success of AMS. Issues related to communication gaps, interdisciplinary collaboration and hierarchies within healthcare settings were reported. The SPs suggested that effective strategies should be developed and established for a sustainable communication channel for the steward team, ensuring that all medical professionals are aware of the scope and work of the steward team.

Gaining the understanding of physicians from other specialties or the primary care team is crucial. Many healthcare providers perceive the ASP as a team that simply stops antibiotic use. How can we make the primary care team understand that the ASP is not a team that stops treatment, but rather a team that assists in patient care? We are essentially a specialty team focused on antimicrobial management. (SP01)

I would like to find a more sustainable way to inform the prescribers about the stewardship program, then it will be easier to work together. (SP08)

Overcoming perceptual barriers and fostering mutual understanding

This subtheme focuses on the need to address negative perceptions and misconceptions about AMS teams and

their role and to build a foundation of mutual understanding and respect between AMS teams and other healthcare professionals. The participants reported that they might have experienced conflicts with the stewardship team over specific patient cases. The participants also discussed the importance of engaging other specialties with high antibiotic prescribing rates, such as haematology-oncology, critical care and immunology, in the AMS programme.

Antimicrobial steward team often have to negotiate with the prescribing physician. It can be difficult to immediately change these prescribing practices. However, what we can do is make adjustments after the initial antibiotic choice has been made, after a few days of monitoring, discuss the possibility of making adjustments. (HCP01)

ASP team should start suggestion by fostering mutual understanding. We are all working together to provide the best care for the patients. (SP02)

Dedication to AMS despite resource limitations

Intrinsic motivation and professional fulfilment

Despite a lack of resources and staffing, the AMS steward team reported a sense of professional fulfilment experienced. The participants discussed working as the AMS team to prioritise patient safety by suggesting the appropriate use of antimicrobials. Furthermore, they are willing to continuously learn and stay updated on best practices in AMS, as well as to educate other healthcare professionals and patients about the importance of responsible antimicrobial use.

I feel like this work can improve patient care, especially in terms of reducing antibiotic resistance. We can provide healthcare providers involved in the ASP with valuable experience that they can apply in their future roles. If, overall, things improve, that would be a rewarding outcome. (HCP01)

I feel like it's a big job. It requires a lot of collaboration to manage, and I think it will take a long time to see any changes. We need to continuously educate involving medical staff, however, seeing those results takes time. (HCP04)

I enjoy doing this because I'm constantly learning new things. Additionally, it can be seen as a strategy to encourage pharmacists to develop their skills. (HCP06)

The reality of workload

A reported drawback was the participants' feeling of resistance due to additional workload, and some expressed concern about how to effectively communicate with physicians.

This task adds to the workload. I also feel it's a difficult and challenging task because it might involve dealing with many people and many groups like this.

And they might have both those who agree and those who disagree. (HCP04)

Concerning communication, generally speaking, even if we have a good reason, if our position is pharmacist, it's hard to speak up to doctors. (SP03)

Regarding the distributed questionnaire regarding AMS implementation and interventions, the top three rankings of the interest level among the participants to support AMS implementation at their hospital were: (1) a hospital policy and a formal organisational structure responsible for AMS (eg, a multidisciplinary committee focused on appropriate antibiotic use, pharmacy committee, patient safety committee or other relevant structure) (2) a formal AMS programme accountable for ensuring appropriate antibiotic use; and (3) available multidisciplinary AMS team, for example, greater than one staff member supporting clinical decisions and implementing a comprehensive programme to ensure appropriate antibiotic use. The top five rankings of the interest level in AMS programme interventions to be most effective in their hospital were: (1) de-escalation therapy (2) guideline implementation, (3) antibiogram, (4) prospective audit and feedback and (5) pharmacokinetics and pharmacodynamics application in antibiotic dosing.

DISCUSSION

The barriers and facilitators of implementing the AMS programme at paediatric units in academic hospitals in Thailand were identified. The major themes from this qualitative study were organisational hierarchical culture and individual behaviours, social influences aimed at improving healthcare workers' mindset, effective communication and collaboration and the positive mindset of the AMS programme team.

Our study identified organisational hierarchical culture and individual behaviours as key barriers to AMS implementation. These factors influence acceptance and adherence to AMS implementation. Understanding the specific cultural context of a healthcare setting is important for tailoring effective interventions. Studies from various settings consistently revealed inadequate knowledge and awareness of rational antibiotic use among physicians.¹⁶ To address knowledge gaps, enhancing education on appropriate antimicrobial prescribing while in medical school is a key target for improvement.¹⁷

Antibiotic prescribing practices vary across countries. While many countries involve a wide range of HCPs in antibiotic prescriptions,^{16 18} the prescribing practices in our study were distinct where specialists or senior physicians held primary decision-making authority. Consistent with a published study, hierarchical structures and a lack of recognition of AMS importance by senior doctors emerged as barriers in low- and middle-income settings.^{9 19} The findings demonstrated that the long-standing culture of seniority hindered AMS adoption.

Despite the existence of AMS policies in the studied hospitals, they were not consistently endorsed or applied in practice.

Many studies documented complex behavioural and social influences on antimicrobial prescribing.²⁰ Our findings align with other studies that have evaluated barriers in relation to clinical decision making. Prior studies reported physician insecurity in patients with severe or non-definitive diagnoses.^{21–23} Similar to the paediatric setting, most paediatricians stated that non-prescription of an antibiotic to potentially severely ill children generated more fear and insecurity than the consequences of an unnecessary antibiotic prescription.²³ Given the high rates of antimicrobial resistance in our study setting, paediatricians also experienced anxiety about not prescribing broad-spectrum antibiotics. To resolve these challenges, adaptable, evidence-based interventions should be developed and implemented across various healthcare settings. An effective feedback tool also needs to be developed and implemented to engage the HCP and impact on the patients' clinical outcomes.²⁴ Our study reported that HCP engagement was crucial for successful AMS implementation. ASP recommendations should be communicated collaboratively, focusing on improving patient outcomes rather than criticising the practices or being perceived as the 'antibiotic police'.^{12 25} From the previous study, formalised antibiotic stewardship meetings and ward rounds on a continuous and regular basis are one of the effective AMS interventions.²⁶ The AMS team can facilitate interdisciplinary rounds to address complex cases, bringing together expertise from specialists, infectious disease physicians, pharmacists and other healthcare professionals.²⁷ This collaborative approach encourages shared decision-making and can lead to more rational antibiotic use.

A recent national survey in Thailand indicated that nearly 90% of hospitals had an ASP in 2021, often with a multidisciplinary team.²⁸ However, there is a shortage of ASP team members nationwide. Most team members are general practitioners or pharmacists lacking formal infectious disease training.²⁹ Understaffing and underprioritising of ASP are also global challenges.³⁰ Despite resource constraints and heavy patient workloads in the region,¹⁶ our study found that the stewardship team maintained a positive attitude toward combating antimicrobial resistance and was committed to professional development, aligning with findings from the previous study.³¹ To further enhance their efforts, robust support systems, including hospital administrators, adequate laboratory capabilities and improved information technology infrastructure, are essential.²⁹

For the implication, antibiotic stewardship involves organisation and hospital policy, hospital staff and antibiotic prescribers, local context in each setting and individual behaviour change should be considered. Overall, the success of AMS programmes relies on understanding and addressing the perspectives and concerns of healthcare workers. Creating a positive and supportive

environment, providing ongoing education and recognising the contributions of those involved are essential components of a successful stewardship initiative. It requires a combination of educational, organisational and cultural interventions. Tailoring strategies to the specific challenges and culture of the healthcare environment is crucial for success. The Behaviour Change Wheel is a systematic framework designed to guide the development of effective interventions.³² It works by first requiring a thorough understanding of the target behaviour using the capability, opportunity and motivation model. For example, if a key barrier is someone's beliefs about the behaviour, potential interventions might focus on persuasion, modelling or education to change those beliefs. Similarly, if a lack of resources is identified as a barrier, an enablement intervention, such as endorsing a supportive hospital policy or providing staff and equipment, could be implemented.

Our findings highlight specific challenges and opportunities for AMS implementation within the context of inpatient paediatric care. Key strengths of the study include identifying institutional and cultural impediments to AMS adoption, especially in low- and middle-income situations. Furthermore, it identifies the dedication and professional fulfilment of AMS teams, which can be leveraged to improve AMS initiatives. However, it is important to acknowledge several methodological limitations. First, the findings may not be directly transferable to all settings due to differences in organisational structure, resource availability and patient populations. Second, conducting qualitative research in a hierarchical corporate culture presents challenges, potentially leading to power dynamics influencing participants' responses and a reluctance to express opinions. Finally, there is the possibility of social desirability bias influencing participants' responses during the interviews. Although we ensured anonymity and confidentiality to minimise this bias, we acknowledge that it may not have been entirely eliminated.

CONCLUSIONS

The implementation of AMS in paediatric settings is challenging. The key barriers to focus on were organisation hierarchical culture and the perception of healthcare workers. The next steps for implementing AMS programmes in Thai paediatric hospitals could involve integrating AMS principles into medical education, fostering an approach to promote shared decision-making regarding rational antibiotic use and improving healthcare workers' mindset towards stewardship.

Contributors SA and TJ conceived and designed the study. PW and TJ conducted and interviewed the participants. SA, PW and TJ analysed the data and interpreted the themes. TP supervised the study. SA wrote the first draft of the manuscript. All authors thoroughly reviewed and approved the manuscript. SA is responsible for the overall content as guarantor. AI-powered grammar checking software was used to assist in the identification and correction of grammatical errors within this manuscript. The authors ultimately bear responsibility for the accuracy and clarity of the final text.

Funding This work was supported by the Ratchadapiseksomphot Fund, Faculty of Medicine, Chulalongkorn University, Grant number RA66/013. The funder did not influence the results of the study despite author affiliation with the funders.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by The Ethics Committee of Faculty of Medicine, Chulalongkorn University has approved the study (IRB. 236/65). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Suvaporn Anugulruengkitt <http://orcid.org/0000-0002-6965-5537>

REFERENCES

- 1 Dellit TH, Owens RC, McGowan JE Jr, *et al*. Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America guidelines for developing an institutional program to enhance antimicrobial stewardship. *Clin Infect Dis* 2007;44:159–77.
- 2 Barlam TF, Cosgrove SE, Abbo LM, *et al*. Executive Summary: Implementing an Antibiotic Stewardship Program: Guidelines by the Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America. *Clin Infect Dis* 2016;62:1197–202.
- 3 Livorsi DJ, Drainoni M-L, Reisinger HS, *et al*. Leveraging implementation science to advance antibiotic stewardship practice and research. *Infect Control Hosp Epidemiol* 2022;43:139–46.
- 4 Proctor E. Dissemination and Implementation Research in Health: Translating Science to Practice. New York: Oxford University Press, 2018.
- 5 Damschroder LJ, Aron DC, Keith RE, *et al*. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implementation Sci* 2009;4:50.
- 6 Damschroder LJ, Reardon CM, Widerquist MAO, *et al*. The updated Consolidated Framework for Implementation Research based on user feedback. *Implement Sci* 2022;17:75.
- 7 Barlam TF, Childs E, Zieminski SA, *et al*. Perspectives of Physician and Pharmacist Stewards on Successful Antibiotic Stewardship Program Implementation: A Qualitative Study. *Open Forum Infect Dis* 2020;7:ofaa229.
- 8 Villanueva P, Coffin SE, Mekasha A, *et al*. Comparison of Antimicrobial Stewardship and Infection Prevention and Control Activities and Resources Between Low-/Middle- and High-income Countries. *Pediatr Infect Dis J* 2022;41:S3–9.
- 9 Abo YN, Freyne B, Kululanga D, *et al*. The Impact of Antimicrobial Stewardship in Children in Low- and Middle-income Countries: A Systematic Review. *Pediatr Infect Dis J* 2022;41:S10–7.
- 10 Rungsitsathian K, Wacharachaisurapol N, Nakaranurack C, *et al*. Acceptance and outcome of interventions in a meropenem de-escalation antimicrobial stewardship program in pediatrics. *Pediatr Int* 2021;63:1458–65.

- 11 Francis JJ, Johnston M, Robertson C, *et al*. What is an adequate sample size? Operationalising data saturation for theory-based interview studies. *Psychol Health* 2010;25:1229–45.
- 12 Szymczak JE, Kitt E, Hayes M, *et al*. Threatened efficiency not autonomy: Prescriber perceptions of an established pediatric antimicrobial stewardship program. *Infect Control Hosp Epidemiol* 2019;40:522–7.
- 13 Malone S, McKay VR, Krucylak C, *et al*. A cluster randomized stepped-wedge trial to de-implement unnecessary post-operative antibiotics in children: the optimizing perioperative antibiotic in children (OPerAtiC) trial. *Implement Sci* 2021;16:29.
- 14 Sayood SJ, Venkatram C, Newland JG, *et al*. Experiences from the Missouri Antimicrobial Stewardship Collaborative: A mixed methods study. *Infect Control Hosp Epidemiol* 2020;41:1455–7.
- 15 VERBI Software. MAXQDA 2022. VERBI Software Berlin, Germany; 2021. Available: <https://www.maxqda.com/>
- 16 Kakkar AK, Shafiq N, Singh G, *et al*. Antimicrobial Stewardship Programs in Resource Constrained Environments: Understanding and Addressing the Need of the Systems. *Front Public Health* 2020;8:140.
- 17 Pierce J, Apisarnthanarak A, Schellack N, *et al*. Global Antimicrobial Stewardship with a Focus on Low- and Middle-Income Countries. *Int J Infect Dis* 2020;96:621–9.
- 18 Barker AK, Brown K, Ahsan M, *et al*. What drives inappropriate antibiotic dispensing? A mixed-methods study of pharmacy employee perspectives in Haryana, India. *BMJ Open* 2017;7:e013190.
- 19 Tarrant C, Colman AM, Jenkins DR, *et al*. Drivers of Broad-Spectrum Antibiotic Overuse across Diverse Hospital Contexts-A Qualitative Study of Prescribers in the UK, Sri Lanka and South Africa. *Antibiotics (Basel)* 2021;10:94.
- 20 Bassetti M, Giacobbe DR, Vena A, *et al*. Challenges and research priorities to progress the impact of antimicrobial stewardship. *Drugs Context* 2019;8:212600.
- 21 Cabral C, Lucas PJ, Ingram J, *et al*. “It’s safer to ...” parent consulting and clinician antibiotic prescribing decisions for children with respiratory tract infections: An analysis across four qualitative studies. *Soc Sci Med* 2015;136–137:S0277-9536(15)00309-3:156–64:.
- 22 Marti D, Hamdy RF, Broniatowski DA. Gist Representations and Decision-Making Processes Affecting Antibiotic Prescribing for Children with Acute Otitis Media. *MDM Policy & Practice* 2022;7:23814683221115416.
- 23 Arnau-Sánchez J, Jiménez-Guillén C, Alcaraz-Quinonero M, *et al*. Factors Influencing Inappropriate Use of Antibiotics in Infants under 3 Years of Age in Primary Care: A Qualitative Study of the Paediatricians’ Perceptions. *Antibiotics (Basel)* 2023;12:727.
- 24 Cherian JP, Helsel TN, Jones GF, *et al*. Understanding the role of antibiotic-associated adverse events in influencing antibiotic decision-making. *ASHE* 2024;4:e13.
- 25 Szymczak JE, Newland JG. The social determinants of antibiotic prescribing. In: Barlam TF, Neuhauser MM, Tamma PD, *et al*, eds. *Practical Implementation of an Antibiotic Stewardship Program*. Cambridge, UK: Cambridge University Press, 2018: 45–62.
- 26 Chetty S, Swe-Han KS, Mahabeer Y, *et al*. Interprofessional education in antimicrobial stewardship, a collaborative effort. *JAC Antimicrob Resist* 2024;6:dlae054.
- 27 WHO. Antimicrobial Stewardship Programmes in Health-Care Facilities in Low- and Middle-Income Countries: a WHO Practical Toolkit, 2019. Available: <https://www.who.int/publications/i/item/9789241515481>
- 28 Patel PK, Watari T, Greene MT, *et al*. The current state of antimicrobial and urine culture stewardship in Thailand: Results from a national survey. *Am J Infect Control* 2024;52:191–4.
- 29 Rattanaumpawan P, Samanloh S, Thamlikitkul V. Feasibility of implementing antimicrobial stewardship programs in acute-care hospitals: A nationwide survey in Thailand. *Infect Control Hosp Epidemiol* 2022;43:1070–4.
- 30 Apisarnthanarak A, Kwa AL-H, Chiu C-H, *et al*. Antimicrobial stewardship for acute-care hospitals: An Asian perspective. *Infect Control Hosp Epidemiol* 2018;39:1237–45.
- 31 Khan MU, Hassali MAA, Ahmad A, *et al*. Perceptions and Practices of Community Pharmacists towards Antimicrobial Stewardship in the State of Selangor, Malaysia. *PLoS One* 2016;11:e0149623.
- 32 Michie S, Atkins L, West R. The Behaviour Change Wheel: A Guide to Designing Interventions. Silverback Publishing, 2014.