



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Outcomes associated with older patients who present or develop delirium in the emergency department: a protocol of a systematic review and meta-analysis

Journal:	BMJ Open
Manuscript ID	bmjopen-2024-095495
Article Type:	Protocol
Date Submitted by the Author:	23-Oct-2024
Complete List of Authors:	King, Sarah; Norwegian University of Science and Technology, Department of Circulation and Medical Imaging Uleberg, Oddvar; St. Olavs Hospital, Dept. of Emergency Medicine and Pre-hospital Services Pedersen, Sindre A; The Medicine and Health Library, Library Section for Research Support, Data and Analysis, Norwegian University of Science and Technology (NTNU), Trondheim, Norway Bjørnsen, Lars Petter; Norwegian University of Science and Technology
Keywords:	Delirium, Systematic Review, Emergency Departments < Emergency Service, Hospital, Adverse events < THERAPEUTICS, Aged

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Outcomes associated with older patients who present or develop delirium in the emergency department: a protocol of a systematic review and meta-analysis

Sarah King¹, Oddvar Uleberg^{1,2,3}, Sindre Andre Pedersen⁴, Lars Petter Bjørnsen^{1,3}

1 Trondheim Emergency Department Research Group (TEDRG), Department of Circulation and Medical Imaging, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

2 Department of Research and Development, Norwegian Air Ambulance Foundation, Oslo, Norway

3 Emergency Department, Clinic of Emergency Medicine and Prehospital Care, St. Olav's Hospital, Trondheim, Norway

4 Library Section for Research Support, Data and Analysis, The Medicine and Health Library, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

Corresponding author: sarah.king@ntnu.no

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

303 words

Abstract

Introduction Delirium is commonly observed in older patients who are admitted to the emergency department (ED). Previous systematic reviews have identified poor outcomes associated with delirium in surgical, intensive care and other hospital settings, yet none have specifically considered the ED. This systematic review aims to examine associations between older patients who present or develop delirium in the ED and adverse outcomes within the hospital and after discharge.

Methods and analysis Searches will be conducted in October 2024 in MEDLINE, Embase, Web of Science, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and the Cochrane Library. There will be no date or language restrictions. Key terms will include concepts related to delirium, the ED, and older adults. Observational studies or non-intervention clinical studies will be included that compare outcomes in older patients (i.e., ≥ 65 years) with and without delirium. Outcomes of interest will include length of hospital stay, non-home discharge (e.g., nursing home/residential aged care facility), cognitive impairment, decreased physical function, mortality, readmission to hospital, and quality of life measures. Two reviewers will independently screen the studies. Data extraction and quality assessment will be extracted by one reviewer and checked by a second reviewer with any disagreements resolved by discussion or by a third reviewer. Where appropriate (i.e., if the studies are clinically and statistically homogenous), data will be combined in a meta-analysis and a GRADE assessment will be made for each outcome. All methods will be guided by the Cochrane Handbook, the Centre for Reviews and Dissemination (CRD) and reported following the PRISMA statement as well as the recommendations set out by the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group.

Ethics and dissemination As this systematic review will use published data, ethical approval is not required. The results will be disseminated through a peer-reviewed publication and conference presentations.

PROSPERO registration number CRD42024594975

Keywords: Delirium, emergency department, adverse outcomes, systematic review

1,435 words

Introduction

Delirium is a syndrome that is characterized by an acute decline in cognitive functioning and can present as hypoactive (e.g., drowsy, lethargic, etc.), hyperactive (e.g., agitated, anxious, etc.), or a fluctuating mix of these psychomotor subtypes. It has been estimated that almost one in four older acute medical hospital patients present or develop delirium¹. Delirium is also commonly observed in the emergency department (ED)², with a recent systematic review reporting a prevalence rate of 15.2% in this setting (95% CI: 12.5% to 18.0% [pooled crude rate based on 30 studies])³. Given that delirium is often unrecognized (e.g., see ⁴⁻⁷), it is likely that some reported prevalence rates could represent underestimations.

Delirium is of concern because it has been associated with poorer outcomes in hospital settings, including mortality⁸, long-term cognitive decline^{9, 10}, as well as longer hospital stays and costs¹¹. Previous systematic reviews (SRs) have been conducted that assess outcomes associated with delirium in surgical patients (e.g., ¹²), intensive care units (e.g., ¹³⁻¹⁵) and hospitalized patients (e.g., ^{16, 17}), but none have specifically addressed the older ED patient, despite several new primary studies on this topic. A SR will be conducted by the Trondheim Emergency Department Research Group (TEDRG) to examine associations between delirium in older patients admitted to the ED and short and long-term adverse outcomes. By evaluating effect sizes of several outcomes derived from comparisons of patients with and without delirium, it will be possible to assess the magnitude and clinical significance of the problem in the ED context. The findings will be of relevance to clinicians and researchers who aim to prevent, detect and treat delirium in patients who present to the ED.

Methods and analysis

The primary research question of this SR is: What short and long-term adverse outcomes are associated with older patients (i.e., ≥ 65 years) who present or develop delirium in the ED compared to those who do not? To address this question, a SR will be undertaken according to guidance presented in the Cochrane Handbook¹⁸, the Centre for Reviews and Dissemination (CRD)¹⁹ and reported following the PRISMA statement²⁰ as well as the recommendations set out by the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group²¹. This protocol was developed following the guideline of the Preferred Reporting Items for Systematic Review and Meta-analysis Protocols (PRISMA-P)²² and is registered on PROSPERO at the University of York (CRD42024594975).

Inclusion and exclusion criteria

To be eligible for inclusion in the review, studies will have to meet all the following criteria in terms of population, intervention and comparator, outcomes, and study designs (PICOS):

Studies of older patients (i.e., ≥ 65 years) who visit the ED will be eligible for inclusion. Delirium assessments must be made using a diagnostic tool or defined criteria (e.g., Confusion assessment method [CAM], brief confusion assessment method [bCAM], 3-minute diagnostic interview for confusion assessment method [3D-CAM], 4AT, Diagnostic and statistical manual of mental disorder criteria [DSM] criteria etc.). Patients with self-

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

diagnosed delirium as well as patients with delirium tremens, drug-induced delirium or hepatic encephalopathy will be excluded. Studies examining older patients undergoing emergency surgery, and older patients with COVID-19 will not be included as other SRs have been conducted in these population groups/settings.

Studies reporting one or more of the following short or long-term (i.e., up to one year) outcomes will be eligible for inclusion: length of hospital stay (or a measure of prolonged hospital stay), non-home discharge (e.g., nursing home/residential aged care facility), cognitive impairment, decreased physical function, mortality (in hospital and out of hospital), readmission to hospital, and other quality of life (QoL) measures or an overall QoL assessment.

Only prospective observational studies or non-intervention clinical studies that compare outcomes in patients with and without delirium will be included. If there are less than five prospective studies, retrospective studies will also be considered. Controlled clinical trials or other types of intervention studies will be excluded as well as case series, commentaries and letters to the editor.

Conference abstracts will only be included if there is no associated full publication and if adequate data are reported (e.g., detailed information on the population, diagnostic criteria, and clear results with numerical data reported for those with and without delirium).

Search strategy

The literature search will be conducted in October 2024 in MEDLINE, Embase, Web of Science, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and the Cochrane Library. Key terms will include concepts related to delirium, the emergency department, and older adults (see supplementary file 1). There will be no language or date limitations. Grey literature will be searched in OAster (OpenGrey, NYAM Grey Literature Report, and the British Library EThOS are not currently available for searching). Handsearching will also be conducted to check references of relevant papers and reviews identified by the search. In addition, a citation search will be carried out to identify subsequent publications which have cited any of the included studies.

Study selection and data extraction

The records retrieved from the searches will de-duplicated in EndNote²³ and exported into Rayyan²⁴ for screening. During the first stage of study selection, titles and abstracts will be independently screened by two researchers against the inclusion/exclusion criteria. Full papers of all potentially relevant studies identified in the first stage will be screened by two reviewers working independently with any discrepancies resolved by discussion between the reviewers or with the assistance of a third reviewer.

The number of studies identified by the search and excluded at various stages will be recorded and reported in a PRISMA study flow diagram²⁰. After the second stage of screening, a table of excluded studies with detailed reasons for exclusion will be created.

Data to be extracted from each study will include information on the study (e.g., study type, setting, country, objective, inclusion criteria, study methods including delirium assessment tool) and patient characteristics (e.g., age, sex, main reason for ED visit, and for patients with delirium: subtype, severity), as well the results and any potential confounding factors identified by the study authors. A data extraction form will be developed using Microsoft Excel²⁵ and piloted by two independent reviewers using a minimum of three studies. Data will be extracted by one reviewer and checked by a second reviewer. Any discrepancies will be resolved through discussion. If any relevant information is missing from the studies or if any data is unclear, the reviewers will attempt to contact the authors.

Critical appraisal

The observational studies will be assessed using the Joanna Briggs Institute (JBI) critical appraisal tools²⁶. One reviewer will conduct the quality assessment, and a second reviewer will check the assessment with any discrepancies resolved through discussion and with a third reviewer.

Synthesis and analysis

Outcome comparisons between older patients with and without delirium will be presented. Where possible and appropriate (i.e., if the studies are clinically and statistically homogenous), data from the studies will be combined in a meta-analysis using RevMan²⁷. For continuous outcomes, means and standard deviations will be collected and used to estimate study-specific and pooled mean differences with 95% confidence intervals (CIs). For dichotomous outcomes, numerators and denominators will be collected, and Mantel-Haenszel (M-H) risk ratios (RRs) or odds ratios (ORs) and 95% CIs used to summarise effect sizes. If the scales/tools used to assess the outcomes differ between the studies, we will compute study-level standardised mean differences (SMD) between comparison groups with 95% CIs. The results will be statistically pooled (where possible) using both fixed-effect and random-effects models and the results from these different models will be compared. Statistical heterogeneity will be assessed using the *Chi*² test and the *I*² statistic, and by examining the random effects between study variance (*Tau*²). If possible, sensitivity analyses will be performed to assess the robustness of results by excluding studies deemed to have a high risk of bias. Funnel plots will be visually inspected to check for publication bias. For each outcome summarised, the GRADE system²⁸ will be used to provide an assessment of the quality of a body of evidence. One reviewer will rate the evidence using GRADEpro GDT²⁹ and a second reviewer will check the assessment, with any discrepancies resolved through discussion.

Exploratory subgroup analyses will be performed where possible by date of publication, delirium assessment method used by the study authors, and age of the study participants (if outcomes are categorized by age groups). If there are any studies conducted in specific patient groups (e.g., older patients with fractures), subgroup analysis may also be conducted by patient type.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

A narrative synthesis of the studies will be performed if the studies are too diverse to perform a meta-analysis, or if only one study reported on an outcome and pooling cannot be undertaken.

Author Contributions

SK conceived the idea for the SR. All authors read and contributed to writing different aspects of the protocol.

Amendments

Any amendments made to the protocol will be dated along with a rationale and a clear description of the change.

Funding Statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing Interests

The authors declare no competing interests.

Patient and Public Involvement

There was no patient or public involvement in the development of the protocol.

References

1. Gibb K, Seeley A, Quinn T, et al. The consistent burden in published estimates of delirium occurrence in medical inpatients over four decades: a systematic review and meta-analysis study. *Age Ageing* 2020; 49: 352-360. DOI: 10.1093/ageing/afaa040.
2. Han JH, Wilson A and Ely EW. Delirium in the older emergency department patient: a quiet epidemic. *Emerg Med Clin North Am* 2010; 28: 611-631. DOI: 10.1016/j.emc.2010.03.005.
3. Chen F, Liu L, Wang Y, et al. Delirium prevalence in geriatric emergency department patients: A systematic review and meta-analysis. *Am J Emerg Med* 2022; 59: 121-128. 20220701. DOI: 10.1016/j.ajem.2022.05.058.
4. El Hussein MT, Hirst S and Stares R. Delirium in Emergency Departments: Is it Recognized? *J Emerg Nurs* 2021; 47: 809-817. 20210311. DOI: 10.1016/j.jen.2021.01.009.
5. Lee JS, Tong T, Chignell M, et al. Prevalence, management and outcomes of unrecognized delirium in a National Sample of 1,493 older emergency department patients: how many were sent home and what happened to them? *Age Ageing* 2022; 51. DOI: 10.1093/ageing/afab214.
6. Meged-Book T, Frenkel R, Nikonov A, et al. Delirium screening in the emergency department: evaluation and intervention. *Isr J Health Policy Res* 2024; 13: 16. 20240402. DOI: 10.1186/s13584-024-00603-1.
7. Barron EA and Holmes J. Delirium within the emergency care setting, occurrence and detection: a systematic review. *Emerg Med J* 2013; 30: 263-268. 20120725. DOI: 10.1136/emered-2011-200586.
8. Aung Thein MZ, Pereira JV, Nitchingham A, et al. A call to action for delirium research: Meta-analysis and regression of delirium associated mortality. *BMC Geriatr* 2020; 20: 325. 20200907. DOI: 10.1186/s12877-020-01723-4.

9. Goldberg TE, Chen C, Wang Y, et al. Association of Delirium With Long-term Cognitive Decline: A Meta-analysis. *JAMA Neurol* 2020; 77: 1373-1381. DOI: 10.1001/jamaneurol.2020.2273.
10. Wilcox ME, Girard TD and Hough CL. Delirium and long term cognition in critically ill patients. *Bmj* 2021; 373: n1007. 20210608. DOI: 10.1136/bmj.n1007.
11. Dziegielewska C, Skead C, Canturk T, et al. Delirium and Associated Length of Stay and Costs in Critically Ill Patients. *Crit Care Res Pract* 2021: 6612187. DOI: 10.1155/2021/6612187.
12. Yan E, Veitch M, Saripella A, et al. Association between postoperative delirium and adverse outcomes in older surgical patients: A systematic review and meta-analysis. *J Clin Anesth* 2023; 90: 111221. 20230727. DOI: 10.1016/j.jclinane.2023.111221.
13. Krewulak KD, Stelfox HT, Ely EW, et al. Risk factors and outcomes among delirium subtypes in adult ICUs: A systematic review. *Journal of Critical Care* 2020; 56: 257-264. DOI: <https://doi.org/10.1016/j.jcrc.2020.01.017>.
14. Salluh JI, Wang H, Schneider EB, et al. Outcome of delirium in critically ill patients: systematic review and meta-analysis. *Bmj* 2015; 350: h2538. 20150603. DOI: 10.1136/bmj.h2538.
15. Zhang Z, Pan L and Ni H. Impact of delirium on clinical outcome in critically ill patients: a meta-analysis. *General Hospital Psychiatry* 2013; 35: 105-111. DOI: <https://doi.org/10.1016/j.genhosppsych.2012.11.003>.
16. Siddiqi N, House AO and Holmes JD. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. *Age and Ageing* 2006; 35: 350-364. DOI: 10.1093/ageing/afl005.
17. Witlox J, Eurelings LSM, de Jonghe JFM, et al. Delirium in Elderly Patients and the Risk of Postdischarge Mortality, Institutionalization, and Dementia: A Meta-analysis. *JAMA* 2010; 304: 443-451. DOI: 10.1001/jama.2010.1013.
18. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions version 6.5 (updated August 2024)* 2024.
19. Khan KB TRG, Glanville J, Sowden AJ, Kleijnen J. *Undertaking systematic reviews of research on effectiveness: CRD's guidance for those carrying out or commissioning reviews*. . York: NHS Centre for Reviews and Dissemination, University of York, 2001.
20. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372: n71. 20210329. DOI: 10.1136/bmj.n71.
21. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *Jama* 2000; 283: 2008-2012. DOI: 10.1001/jama.283.15.2008.
22. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015; 4: 1. 20150101. DOI: 10.1186/2046-4053-4-1.
23. The EndNote Team. EndNote. EndNote 21 ed. Philadelphia, PA: Clarivate, 2013.
24. Ouzzani M, Hammady H, Zbys Fedorowicz Z, et al. Rayyan — a web and mobile app for systematic reviews. *Systematic Reviews* 2016; 5. DOI: DOI: 10.1186/s13643-016-0384-4.
25. Microsoft Excel. Microsoft Corporation, 2019.
26. Joanna Briggs Institute. Checklist for systematic reviews and research syntheses. 2017.
27. Review Manager (RevMan). The Cochrane Collaboration, 2024.

- 1
2
3 28. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the
4 quality of evidence. *J Clin Epidemiol* 2011; 64: 401-406. 20110105. DOI:
5 10.1016/j.jclinepi.2010.07.015.
6
7 29. GRADEpro GDT. Hamilton (ON): McMaster University (developed by Evidence
8 Prime), 2023.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Example search strategy for Embase:

No.	Query
#1	'delirium'/exp
#2	'acute confusion'/exp
#3	deliri*:ti,ab,kw
#4	'acute confusion*':ti,ab,kw
#5	'acute organic psychosyndrome':ti,ab,kw
#6	'acute brain syndrome':ti,ab,kw
#7	'metabolic encephalopathy':ti,ab,kw
#8	'acute psycho-organic syndrome':ti,ab,kw
#9	'clouded state':ti,ab,kw
#10	'clouding of consciousness':ti,ab,kw
#11	'exogenous psychosis':ti,ab,kw
#12	'toxic psychosis':ti,ab,kw
#13	'toxic confusion':ti,ab,kw
#14	obnubilat*:ti,ab,kw
#15	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14
#16	'emergency ward'/exp
#17	'emergency care'/exp
#18	'emergency health service'/exp
#19	'emergency patient'/exp
#20	'emergency physician'/exp
#21	'emergency medical dispatcher'/exp
#22	'hospital emergency service'/exp
#23	(emergency NEAR/1 (care OR department? OR unit? OR center? OR ward? OR service?)):ti,ab,kw
#24	#16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23
#25	'aged'/exp
#26	'aged':ti,ab,kw
#27	'centenarian*':ti,ab,kw

Enseignement Supérieur (ABES) .
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

- #28 'elder*':ti,ab,kw
- #29 ((geriatric OR old* OR senior*) NEAR/2 (citizen* OR patient*)):ti,ab,kw
- #30 'senium':ti,ab,kw
- #31 'senile':ti,ab,kw
- #32 'senility':ti,ab,kw
- #33 'septuagenarian*':ti,ab,kw
- #34 'octogenarian*':ti,ab,kw
- #35 #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34
- #36 #15 AND #24 AND #35

BMJ Open

Outcomes associated with older patients who present or develop delirium in the emergency department: protocol for a systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2024-095495.R1
Article Type:	Protocol
Date Submitted by the Author:	26-Apr-2025
Complete List of Authors:	King, Sarah; Norwegian University of Science and Technology, Department of Circulation and Medical Imaging Uleberg, Oddvar; St. Olavs Hospital, Dept. of Emergency Medicine and Pre-hospital Services Pedersen, Sindre A; The Medicine and Health Library, Library Section for Research Support, Data and Analysis, Norwegian University of Science and Technology (NTNU), Trondheim, Norway Bjørnsen, Lars Petter; Norwegian University of Science and Technology
Primary Subject Heading:	Emergency medicine
Secondary Subject Heading:	Geriatric medicine, Public health
Keywords:	Delirium, Systematic Review, Emergency Departments < Emergency Service, Hospital, Adverse events < THERAPEUTICS, Aged

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Outcomes associated with older patients who present or develop delirium in the emergency department: protocol for a systematic review and meta-analysis

Sarah King¹, Oddvar Uleberg^{1,2,3}, Sindre Andre Pedersen⁴, Lars Petter Bjørnsen^{1,3}

1 Trondheim Emergency Department Research Group (TEDRG), Department of Circulation and Medical Imaging, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

2 Department of Research and Development, Norwegian Air Ambulance Foundation, Oslo, Norway

3 Emergency Department, Clinic of Emergency Medicine and Prehospital Care, St. Olav's Hospital, Trondheim, Norway

4 Library Section for Research Support, Data and Analysis, The Medicine and Health Library, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

Correspondence to:

Sarah King

sarah.king@ntnu.no

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

Abstract

Introduction: Delirium is commonly observed in older patients who are admitted to the emergency department (ED). Previous systematic reviews have identified poor outcomes associated with delirium in surgical, intensive care and other hospital settings, yet none have specifically considered the ED. This systematic review aims to examine associations between older patients who present or develop delirium in the ED and adverse outcomes within the hospital and after discharge.

Methods and analysis: Searches will be conducted in MEDLINE, Embase, Web of Science, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and the Cochrane Library. There will be no date or language restrictions. Key terms will include concepts related to delirium, the ED, and older adults. Observational studies or non-intervention clinical studies will be included that compare outcomes in older patients (i.e., ≥ 65 years) with and without delirium. Outcomes of interest will include length of hospital stay, non-home discharge (e.g., nursing home/residential aged care facility), cognitive impairment, decreased physical function, mortality, readmission to hospital, and quality of life measures. Two reviewers will independently screen the studies. Data extraction and quality assessment will be extracted by one reviewer and checked by a second reviewer with any disagreements resolved by discussion or by a third reviewer. Where appropriate, data will be combined in a meta-analysis and a GRADE assessment will be made for each outcome. All methods will be guided by the Cochrane Handbook, the Centre for Reviews and Dissemination (CRD) and reported following the PRISMA statement as well as the recommendations set out by the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group.

Ethics and dissemination: As this systematic review will use published data, ethical approval is not required. The results will be disseminated through a peer-reviewed publication and conference presentations.

Study registration: PROSPERO, CRD42024594975.

Keywords: Delirium, emergency department, adverse outcomes, systematic review

Strengths and limitations of this study

- A rigorous systematic review methodology will be used to address the research question.
- There will be no language or date restrictions to the search strategy.
- The review includes observational studies, which are subject to bias and confounding.
- Heterogeneity among the studies is expected and careful consideration of discrepancies will be needed to mitigate spurious conclusions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Delirium is a syndrome that is characterized by an acute decline in cognitive functioning and can present as hypoactive (e.g., drowsy, lethargic, etc.), hyperactive (e.g., agitated, anxious, etc.), or a fluctuating mix of these psychomotor subtypes. It has been estimated that almost one in four older acute medical hospital patients present or develop delirium¹. Delirium is also commonly observed in the emergency department (ED)², with a recent systematic review reporting a prevalence rate of 15.2% in this setting (95% CI: 12.5% to 18.0% [pooled crude rate based on 30 studies])³. Given that delirium is often unrecognized (e.g., see ⁴⁻⁷), it is likely that some reported prevalence rates could represent underestimations.

Delirium is of concern because it has been associated with poorer outcomes in hospital settings, including mortality⁸, long-term cognitive decline^{9, 10}, as well as longer hospital stays and costs¹¹. Previous systematic reviews (SRs) have been conducted that assess outcomes associated with delirium in surgical patients¹², intensive care units¹³⁻¹⁵ and hospitalized patients^{16, 17}, but none have specifically addressed the older ED patient, despite several new primary studies on this topic. A SR will be conducted by the Trondheim Emergency Department Research Group (TEDRG) to examine associations between older patients who have a positive delirium diagnosis in the ED and short- and long-term adverse outcomes. This SR specifically focuses on ED patients as they represent a distinct and vulnerable population. They often present with more acute medical or environmental stressors (e.g., sepsis, polypharmacy) than those who develop delirium postoperatively or during a hospital stay. Moreover, the ED is a critical entry point for early delirium recognition - which could potentially improve prognosis and thus hospital length of stay. The ED thus presents specific challenges and opportunities for delirium research and interventions. By evaluating effect sizes of several outcomes derived from comparisons of older patients with and without delirium, it will be possible to assess the magnitude of the current problem in the ED context. This SR will be of relevance to clinicians and researchers who aim to prevent, detect, and manage delirium in acute care patients.

METHODS AND ANALYSIS

The primary research question of this SR is: What short- and long-term adverse outcomes are associated with older patients (i.e., ≥ 65 years) who present or develop delirium in the ED compared to those who do not? To address this question, a SR will be undertaken according to guidance presented in the Cochrane Handbook¹⁸, the Centre for Reviews and Dissemination (CRD)¹⁹ and reported following the PRISMA statement²⁰ as well as the recommendations set out by the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group²¹. This protocol was developed following the guideline of the Preferred Reporting Items for Systematic Review and Meta-analysis Protocols (PRISMA-P)²² and is registered with PROSPERO (CRD42024594975).

Inclusion and exclusion criteria

To be eligible for inclusion in the review, studies will have to meet all the following criteria in terms of population, exposure/intervention and comparator, outcomes, and study designs (PECOS).

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

Studies that compare older patients (i.e., ≥ 65 years) who are delirium positive (at ED arrival or during their visit) with those who are delirium negative will be eligible for inclusion. Delirium assessments must be made in the ED using a diagnostic tool or defined criteria (e.g., Confusion assessment method [CAM], brief confusion assessment method [bCAM], 3-minute diagnostic interview for confusion assessment method [3D-CAM], 4AT, Diagnostic and statistical manual of mental disorder criteria [DSM] criteria etc.). Patients with delirium tremens, drug-induced delirium or hepatic encephalopathy will be excluded. Studies that specifically assess delirium after an ED visit will not be included. Studies examining older patients undergoing emergency surgery, and older patients with COVID-19 will not be included as other SRs have been conducted in these population groups/settings.

Studies reporting one or more of the following short- (i.e., within 30 days) or long-term (i.e., up to one year) outcomes will be eligible for inclusion: length of hospital stay (or a measure of prolonged hospital stay), non-home discharge (e.g., nursing home/residential aged care facility), cognitive impairment, decreased physical function, mortality (in hospital and out of hospital), readmission to hospital, and other quality of life (QoL) measures or an overall QoL assessment.

Prospective and retrospective observational studies or non-intervention clinical studies that compare outcomes between patients with and without delirium will be included. Controlled clinical trials or other types of intervention studies will be excluded as well as case series, commentaries and letters to the editor.

Conference abstracts will only be included if there is no associated full publication and if adequate data are reported (e.g., detailed information on the population, diagnostic criteria, and clear results with numerical data reported for those with and without delirium).

Search strategy

The literature search will be conducted in MEDLINE, Embase, Web of Science, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and the Cochrane Library. Key terms will include concepts related to delirium, the emergency department, and older adults (**supplementary file 1**). The search will not be limited by language or date. German, French and Scandinavian language papers will be assessed for inclusion and translated to English by members of the review team while Google Translate or DeepL will be used for all other languages. Grey literature will be searched in OAIster (OpenGrey, NYAM Grey Literature Report, and the British Library EThOS are not currently available for searching). Handsearching will also be conducted to check references of relevant papers and reviews identified by the search. In addition, a citation search will be carried out to identify subsequent publications which have cited any of the included studies.

Study selection and data extraction

The records retrieved from the searches will de-duplicated in EndNote²³ and exported into Rayyan²⁴ for screening. During the first stage of study selection, titles and abstracts will be independently screened by two researchers against the inclusion/exclusion criteria. Full papers of all potentially relevant studies identified in the first stage will be screened by two

reviewers working independently with any discrepancies resolved by discussion between the reviewers or with the assistance of a third reviewer.

The number of studies identified by the search and excluded at various stages will be recorded and reported in a PRISMA study flow diagram²⁰. After the second stage of screening, a table of excluded studies with detailed reasons for exclusion will be created.

Data to be extracted from each study will include information on the study (e.g., study type, setting, country, objective, inclusion criteria, study methods including the delirium assessment tool used as well as who did the assessment and any information on their experience or training) and patient characteristics (e.g., age, sex, main reason for ED visit, and for patients with delirium: subtype, severity), as well the results and any potential confounding factors identified by the study authors. A data extraction form will be developed using Microsoft Excel²⁵ and piloted by two independent reviewers using a minimum of four studies. This process will also be used to check consistency in data extraction between the reviewers. The remaining studies will then be extracted by one reviewer and checked by a second reviewer. Any discrepancies will be resolved through discussion. If any relevant information is missing from the studies or if any data is unclear, the reviewers will attempt to contact the authors.

Critical appraisal

The observational studies will be assessed using the Joanna Briggs Institute (JBI) critical appraisal tools²⁶. One reviewer will conduct the quality assessment, and a second reviewer will check the assessment with any discrepancies resolved through discussion and with a third reviewer.

Synthesis and analysis

Outcome comparisons between older patients with and without delirium will be presented. Where possible and appropriate (i.e., if the studies are clinically and statistically homogenous), data from the studies will be combined in a meta-analysis using RevMan²⁷. For continuous outcomes, means and standard deviations will be collected and used to estimate study-specific and pooled mean differences with 95% confidence intervals (CIs). For dichotomous outcomes, numerators and denominators will be collected, and Mantel-Haenszel (M-H) risk ratios (RRs) or odds ratios (ORs) and 95% CIs used to summarise effect sizes. If the scales/tools used to assess the outcomes differ between the studies, we will compute study-level standardised mean differences (SMD) between comparison groups with 95% CIs. The results will be statistically pooled (where possible) using both fixed-effect and random-effects models and the results from these different models will be compared. Statistical heterogeneity will be assessed using the *Chi*² test and the *I*² statistic, and by examining the random effects between study variance (*Tau*²). If possible, sensitivity analyses will be performed to assess the robustness of results by excluding studies deemed to have a greater risk of bias. Comparisons between crude and adjusted effect estimates will also be made. Funnel plots will be visually inspected to check for publication bias. For each outcome summarised, the GRADE system²⁸ will be used to provide an assessment of the quality of a

body of evidence. One reviewer will rate the evidence using GRADEpro GDT²⁹ and a second reviewer will check the assessment, with any discrepancies resolved through discussion.

Exploratory subgroup analyses will be performed where possible by date of publication, study type, delirium assessment method used in the study, assessor and training (if reported), and age of the study participants (if outcomes are categorized by age groups in the included studies). If there are any studies conducted in specific patient groups (e.g., older patients with fractures), subgroup analysis may also be conducted by patient type.

A narrative synthesis of the studies will be performed if the studies are too diverse to perform a meta-analysis, or if only one study reported on an outcome and pooling cannot be undertaken.

Patient and public involvement

None.

ETHICS AND DISSEMINATION

As this systematic review will use published data, ethical approval is not required. The results will be disseminated through a peer-reviewed publication and conference presentations. If any minor amendments are made to this protocol during the review process, they will be reported in the final results paper.

Contributors

SK conceived the idea for the SR. All authors read and contributed to writing different aspects of the protocol. Sarah King is responsible for the overall content as guarantor.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests

The authors declare no competing interests.

References

1. Gibb K, Seeley A, Quinn T, et al. The consistent burden in published estimates of delirium occurrence in medical inpatients over four decades: a systematic review and meta-analysis study. *Age Ageing* 2020; 49: 352-360. DOI: 10.1093/ageing/afaa040.
2. Han JH, Wilson A and Ely EW. Delirium in the older emergency department patient: a quiet epidemic. *Emerg Med Clin North Am* 2010; 28: 611-631. DOI: 10.1016/j.emc.2010.03.005.
3. Chen F, Liu L, Wang Y, et al. Delirium prevalence in geriatric emergency department patients: A systematic review and meta-analysis. *Am J Emerg Med* 2022; 59: 121-128. 20220701. DOI: 10.1016/j.ajem.2022.05.058.
4. El Hussein MT, Hirst S and Stares R. Delirium in Emergency Departments: Is it Recognized? *J Emerg Nurs* 2021; 47: 809-817. 20210311. DOI: 10.1016/j.jen.2021.01.009.
5. Lee JS, Tong T, Chignell M, et al. Prevalence, management and outcomes of unrecognized delirium in a National Sample of 1,493 older emergency department patients: how many were sent home and what happened to them? *Age Ageing* 2022; 51. DOI: 10.1093/ageing/afab214.
6. Meged-Book T, Frenkel R, Nikonov A, et al. Delirium screening in the emergency department: evaluation and intervention. *Isr J Health Policy Res* 2024; 13: 16. 20240402. DOI: 10.1186/s13584-024-00603-1.
7. Barron EA and Holmes J. Delirium within the emergency care setting, occurrence and detection: a systematic review. *Emerg Med J* 2013; 30: 263-268. 20120725. DOI: 10.1136/emered-2011-200586.
8. Aung Thein MZ, Pereira JV, Nitchingham A, et al. A call to action for delirium research: Meta-analysis and regression of delirium associated mortality. *BMC Geriatr* 2020; 20: 325. 20200907. DOI: 10.1186/s12877-020-01723-4.
9. Goldberg TE, Chen C, Wang Y, et al. Association of Delirium With Long-term Cognitive Decline: A Meta-analysis. *JAMA Neurol* 2020; 77: 1373-1381. DOI: 10.1001/jamaneurol.2020.2273.
10. Wilcox ME, Girard TD and Hough CL. Delirium and long term cognition in critically ill patients. *Bmj* 2021; 373: n1007. 20210608. DOI: 10.1136/bmj.n1007.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

11. Dziegielewska C, Skead C, Canturk T, et al. Delirium and Associated Length of Stay and Costs in Critically Ill Patients. *Crit Care Res Pract* 2021; 6612187. DOI: 10.1155/2021/6612187.
12. Yan E, Veitch M, Saripella A, et al. Association between postoperative delirium and adverse outcomes in older surgical patients: A systematic review and meta-analysis. *J Clin Anesth* 2023; 90: 111221. 20230727. DOI: 10.1016/j.jclinane.2023.111221.
13. Krewulak KD, Stelfox HT, Ely EW, et al. Risk factors and outcomes among delirium subtypes in adult ICUs: A systematic review. *Journal of Critical Care* 2020; 56: 257-264. DOI: <https://doi.org/10.1016/j.jcrc.2020.01.017>.
14. Salluh JI, Wang H, Schneider EB, et al. Outcome of delirium in critically ill patients: systematic review and meta-analysis. *Bmj* 2015; 350: h2538. 20150603. DOI: 10.1136/bmj.h2538.
15. Zhang Z, Pan L and Ni H. Impact of delirium on clinical outcome in critically ill patients: a meta-analysis. *General Hospital Psychiatry* 2013; 35: 105-111. DOI: <https://doi.org/10.1016/j.genhosppsych.2012.11.003>.
16. Siddiqi N, House AO and Holmes JD. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. *Age and Ageing* 2006; 35: 350-364. DOI: 10.1093/ageing/afl005.
17. Witlox J, Eurelings LSM, de Jonghe JFM, et al. Delirium in Elderly Patients and the Risk of Postdischarge Mortality, Institutionalization, and Dementia: A Meta-analysis. *JAMA* 2010; 304: 443-451. DOI: 10.1001/jama.2010.1013.
18. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions version 6.5 (updated August 2024)* 2024.
19. Khan KB TRG, Glanville J, Sowden AJ, Kleijnen J. *Undertaking systematic reviews of research on effectiveness: CRD's guidance for those carrying out or commissioning reviews*. . York: NHS Centre for Reviews and Dissemination, University of York, 2001.
20. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372: n71. 20210329. DOI: 10.1136/bmj.n71.
21. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *Jama* 2000; 283: 2008-2012. DOI: 10.1001/jama.283.15.2008.
22. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015; 4: 1. 20150101. DOI: 10.1186/2046-4053-4-1.
23. The EndNote Team. EndNote. EndNote 21 ed. Philadelphia, PA: Clarivate, 2013.
24. Ouzzani M, Hammady H, Zbys Fedorowicz Z, et al. Rayyan — a web and mobile app for systematic reviews. *Systematic Reviews* 2016; 5. DOI: DOI: 10.1186/s13643-016-0384-4.
25. Microsoft Excel. Microsoft Corporation, 2019.
26. Joanna Briggs Institute. Checklist for systematic reviews and research syntheses. 2017.
27. Review Manager (RevMan). The Cochrane Collaboration, 2024.
28. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011; 64: 401-406. 20110105. DOI: 10.1016/j.jclinepi.2010.07.015.
29. GRADEpro GDT. Hamilton (ON): McMaster University (developed by Evidence Prime), 2023.

Example search strategy for Embase:

No.	Query
#1	'delirium'/exp
#2	'acute confusion'/exp
#3	deliri*:ti,ab,kw
#4	'acute confusion*':ti,ab,kw
#5	'acute organic psychosyndrome':ti,ab,kw
#6	'acute brain syndrome':ti,ab,kw
#7	'metabolic encephalopathy':ti,ab,kw
#8	'acute psycho-organic syndrome':ti,ab,kw
#9	'clouded state':ti,ab,kw
#10	'clouding of consciousness':ti,ab,kw
#11	'exogenous psychosis':ti,ab,kw
#12	'toxic psychosis':ti,ab,kw
#13	'toxic confusion':ti,ab,kw
#14	obnubilat*:ti,ab,kw
#15	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14
#16	'emergency ward'/exp
#17	'emergency care'/exp
#18	'emergency health service'/exp
#19	'emergency patient'/exp
#20	'emergency physician'/exp
#21	'emergency medical dispatcher'/exp
#22	'hospital emergency service'/exp
#23	(emergency NEAR/1 (care OR department? OR unit? OR center? OR ward? OR service?)):ti,ab,kw
#24	#16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23
#25	'aged'/exp
#26	'aged':ti,ab,kw
#27	'centenarian*':ti,ab,kw

Enseignement Supérieur (ABES) .
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

- #28 'elder*':ti,ab,kw
- #29 ((geriatric OR old* OR senior*) NEAR/2 (citizen* OR patient*)):ti,ab,kw
- #30 'senium':ti,ab,kw
- #31 'senile':ti,ab,kw
- #32 'senility':ti,ab,kw
- #33 'septuagenarian*':ti,ab,kw
- #34 'octogenarian*':ti,ab,kw
- #35 #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34
- #36 #15 AND #24 AND #35