To cite: Lee JA. Park S.

treating primary

systematic review

Jung J, et al. Herbal medicine

dysmenorrhoea: a protocol for a

(Gyejibongneyong-hwan) for

of randomised controlled

trials. BMJ Open 2016;6:

e011071. doi:10.1136/

bmiopen-2016-011071

additional material is

Prepublication history and

available. To view please visit

the journal (http://dx.doi.org/

10.1136/bmjopen-2016-

Received 8 January 2016

Accepted 30 August 2016

Revised 20 June 2016

011071).

BMJ Open Herbal medicine (Gyejibongneyonghwan) for treating primary dysmenorrhoea: a protocol for a systematic review of randomised controlled trials

Ju Ah Lee,¹ Sunju Park,² Jeeyoun Jung,¹ Ji Hee Jun,³ Jiae Choi,³ Mveona Soo Lee³

ABSTRACT

Introduction: Gyejibongneyong-hwan (GBH), also known as Guizhi Fuling formula, and is widely used for uterine fibroids in East Asian countries. Many clinical trials assessing the efficacy and safety of GBH formula for the treatment of dysmenorrhoea have been reported. This review will assess the clinical evidence for and against the use of GBH formula as a treatment for dysmenorrhoea. It will also discuss the proposed mechanism(s) that could link herbal medicine to improvements in dysmenorrhoea.

Methods and analysis: Fourteen databases will be searched until September 2016. We will include randomised controlled trials (RCTs) examining GBH decoctions for any type of dysmenorrhoea. All RCTs of decoctions or modified decoctions will be included. The methodological gualities of the RCTs will be assessed using the Cochrane Collaboration tool for assessing risk of bias.

Ethics and dissemination: This systematic review will be published in a peer-reviewed journal. The review will also be disseminated electronically and in print. It will be updated to inform and guide healthcare practices.

Trial registration number: CRD42015023419.



¹KM Fundamental Research Division, Korea Institute of Oriental Medicine, Daejeon, South Korea ²Department of Preventive Medicine, College of Korean Medicine, Daejeon University, Daejeon, South Korea ³Clinical Research Division. Korea Institute of Oriental Medicine, Daejeon, South Korea

Correspondence to Dr Myeong Soo Lee; drmslee@gmail.com

INTRODUCTION

Dysmenorrhoea can be primary or secondary and is among the most painful menstrual problems of women of reproductive age.¹ This study focuses on primary dysmenorrhoea (PD), which is defined as painful menses in women without any uterine abnormalities.²

The prevalence of PD varies from 20% to 90% depending on the pain measurements employed and ethnicities examined.³ PD interferes with women's daily living and schoolgirls' activities. In Korea, 78.3% of adolescent girls have been reported to have PD during their menstrual periods, and this is the main cause of short-term school absences.⁴ The main symptoms of PD are

Strengths and limitations of this study

- The strength of this systematic review is unbiased search of various databases without any language restriction.
- The trial screening and data extraction will be conducted independently by two authors.
- We will use the GRADE approach to assess confidence in estimates of effect.
- In the field of complementary and alternative medicine, there are insufficient randomised controlled trials.

abdominal and lower back pain before and during menstruation. Fifteen per cent of girls require analgesics to relieve these symptoms.^b

Current thinking on the aetiology of PD is ĝ that the hormone prostaglandin triggers uterine muscle contractions which result in ≥ cramping and pain.⁶ ⁷ In Korean medicine, PD is thought to be caused by the stagnation of the blood or Qi in the lower abdomen **9** and during menstruation.⁸ Non-steroidal antiinflammatory drugs are considered the firstline treatment for PD.⁹ Other treatments, such as oral contraceptives and β-blockers, are also prescribed.¹⁰ However, these are ineffective and can even elicit adverse events, such as digestive disorders, which have occurred in up to 25% of women.¹¹ Thus, a large number of patients with PD seek complementary and alternative approaches, such as acupuncture, 12 moxibustion 13 14 herbal medicine. 15 16 and

In this review, we will investigate evidence related to the effectiveness of Gyejibongneyonghwan (GBH) formula, which is widely used in traditional Korean medicine (TKM) and traditional Chinese medicine (TCM) for treating the symptoms of PD.

and data

simi

METHODS Study register

Study registration

The protocol for this systematic review has been registered on PROSPERO 2015 under the number CRD42015023419.

Data sources

The following databases will be searched from inception to the current date: Medline, EMBASE, the Cochrane Central Register of Controlled Trials (CENTRAL), AMED and CINAHL. We will also search six Korean medical databases (OASIS, Korean Traditional Knowledge Portal (KTKP), Korean Studies Information Service System (KISS), KoreaMed, Korean Medical Database (KMBase) and DBpia) and three Chinese databases including CNKI China Academic Journal, China Doctoral (ie. Dissertations and Master's Theses Full-text Database, China Proceedings of Conference Full Text Database and Century Journal Project), Wanfang and VIP. Furthermore, we will conduct non-electronic searches of conference proceedings, our own article files, and nine

BMJ Open: first published as 10.1136/bmjopen-2016-011071 on 28 September 2016. Downloaded from http://bmjopen.bmj.com/ on June 10, 2025 at Agence Bibliographique de Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

Korean traditional medical journals (Journal of Korean Medicine, Journal of the Korean Acupuncture and Moxibustion Society, Korean Journal of Acupuncture, Journal of Acupuncture and Meridian Studies, Journal of Pharmacopuncture, Journal of Oriental Rehabilitation Medicine, Journal of Korea Chuna Manual Medicine for Spine and Nerves, Korean Journal of Oriental Physiology and Pathology and Journal of Korean Oriental Internal Medicine). The search strategies that will be applied to the Medline database and CNKI are presented in online supplements 1 and 2. Similar search strategies will be applied to the other databases.

Study selection will be documented and summarised in a PRISMA-compliant flow chart (http://www.prisma-statement.org) (figure 1).

Types of study

All prospective randomised controlled trials (RCTs) will be included if they are randomised studies of GBH formula as the sole treatment or as an adjunct to other treatments, as well as if the control group received the



Figure 1 PRISMA diagram for the included studies. NRS, non-randomised studies.

same treatment as the intervention group. Trials comparing GBH formula with any type of control intervention will also be included. No language restrictions will be imposed. Hard copies of all articles will be obtained and read in full.

Data extraction and guality assessment

Hard copies of all articles will be obtained and read in full. Two authors (SIP and IAL) will perform the data extraction and quality assessment using a predefined data extraction form (see online supplements 3-5). Any disagreement among the authors will be resolved by discussion between all of the authors. When the data are insufficient or ambiguous, MSL will contact the corresponding authors by e-mail or telephone to request additional information or clarification. The data screening and selection process will be performed independently by an author who is fluent in Chinese (IHI). The risk of bias will be assessed using the assessment tool for the risk of bias from the Cochrane Handbook V.5.1.0, which includes random sequence generation, allocation concealment, blinding of the participants and personnel, blinding of the outcome assessments, incomplete outcome data, selective reporting and other sources of bias.¹⁷ Our review will use 'L', 'U' and 'H' to indicate the results of the assessments: 'L' indicates a low risk of bias, 'U' indicates that the risk of bias was unclear, and 'H' indicates a high risk of bias. Disagreements will be resolved by discussion between all of the authors. When disagreements regarding selection cannot be resolved through discussion, the arbiter (MSL) will make the final decision.

Data collection and synthesis

Outcome measures

Primary outcomes

- ▶ Change in symptoms as indicated on a 100 mm visual analogue scale
- Pain duration (hours) according to the McGill questionnaire
- Improved effectiveness including total treatment efficacy-that is, the number of patients whose PD symptoms improve

Secondary outcomes

- Quality of life as measured using a validated questionnaire
- Impact of symptoms as measured with validated questionnaires
- Adverse events

Information related to GBH usage

- ▶ Pattern type in response based on TKM or TCM theory
- Range of dosage of GBH in each study
- Duration of treatment
- Details of formula compositions

Assessment of bias in the included studies

We will independently assess the bias of the included studies according to the criteria of the Cochrane Handbook, V.5.1.0: random sequence generation, allocation concealment, the blinding of participants and personnel, the blinding of outcome assessments, incomplete outcome data, selective reporting and other sources of bias.¹⁷

Data synthesis

The differences between the intervention and control groups will be assessed. For continuous data, we will use the mean differences (MDs) with 95% CIs to measure the treatment effects. We will convert other forms of data into MDs. In the case of outcome variables with different scales, we will use the standard mean difference (SMD) with 95% CIs. For dichotomous data, we will present the treatment effects as relative risks (RRs) with 95% CIs. We will convert other binary data into RR values.

We will use the GRADEpro software of Cochrane Systematic Reviews to create a Summary of Findings table. When disagreements regarding the selections ß cannot be resolved through discussion, the arbiter (MSL) will make the final decision.

uses All of the statistical analyses will be conducted using the Cochrane Collaboration's software programme Review re Manager (RevMan) V.5.3.5 for Windows. For studies with insufficient information, we will contact the corresponding authors to acquire and verify the data whenever possible. When appropriate, we will pool the data across studies for a meta-analysis using fixed or random effects.

Unit of analysis issues

For crossover trials, the data from the first treatment period will be used. For trials in which more than one control group is assessed, the primary analysis will combine the data from each control group. Subgroup analyses of the control groups will be performed. Each patient will be counted only once in the analyses.

Addressing missing data

Intention-to-treat analyses that include all of the randomised patients will be performed. For patients with missing outcome data, carry-forward of the last observed response procedure will be used. The individual patient data will be sought from the original source or the published trial reports when the individual patient data are initially unavailable.

We will use the random- or fixed-effects model for the set used to evaluate the heterogenetic and $I^2 > 50$ and $I^2 > 50$ will be considered indicative of high heterogeneity. When heterogeneity is observed, we will conduct subgroup analyses to explore the possible causes.¹⁸

Assessment of reporting biases

If a sufficient number of the included studies (at least 10 trials) are available, we will use funnel plots to detect

ited to

text

dat

>

training, and

<u>0</u>

tecl

reporting biases.¹⁹ However, funnel plot asymmetries are not identical with publication biases; therefore, we will attempt to determine the possible reasons for any asymmetries, such as small-study effects, poor methodological qualities, and true heterogeneities in the included studies.¹⁹

DISCUSSION

Among the alternative approaches, GBH, also known as Guizhi-Fuling-Wan (pill), is a widely used remedy for PD and has been employed since ancient times in East Asia.^{13 21 22} GBH was first prescribed in the Essential prescriptions from the golden cabinet (Jin Gui Yao Lue) by Zhang Zhongjing of the Han Dynasty (206BC-220ad).² The five ingredients of GBH are ramulus cinnamomi cassiae, sclerotium poriae cocos, cortex moutan radicis, semen pruni persicae and radix paeoniae lactiflorae, and these ingredients are combined in a ratio of 1:1:1:1:1. GBH is applied to remove blood stasis and masses in the abdominal region to promote blood circulation.^{15 24} There are various forms of GBH such as pills, capsules, tablets and decoctions. All types of GBH will be included in this study. Currently, no systematic reviews of the effects of GBH formula on PD have been published. This systematic review will provide a summary of the current evidence related to the effectiveness of GBH formula in the treatment of the symptoms of PD. In particular, we will consider special features of interventions in full reviews of GBH for PD. Although many systematic reviews have been conducted for herbal medicine, important information on its usage was not extracted and has been missed.²⁵ Therefore, we will identify subtypes that this remedy is particularly useful for (eg, certain pattern types based on TKM or TCM theory), identifying a range of dosages, modifications used to improve effectiveness, or comparing duration of treatment in full review of this protocol. The original GBH formula was composed of the five herbs mentioned above; however, primary studies show high heterogeneity in ingredients of GBH. Therefore, we will investigate the composition of each formula used in the primary studies. This evidence will be useful to practitioners, patients and health policy makers regarding the use of acupuncture in PD treatment.

Contributors JAL and SJP conceived the study, developed the criteria, searched the literature, performed the data analysis, and wrote the protocol. JAL, JJ and JC conducted the preliminary search. JJH and MSL assisted in searching the Chinese literature and extracting the data. All authors have read and approved the final manuscript.

Funding JAL, JJ, JHJ and JC were supported by a grant from the Korea Institute of Oriental Medicine (K16111). MSL was supported by Korea Institute of Oriental Medicine (K162921).

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement We will share any data needed. However, there are no additional unpublished data from this study.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license,

which permits others to distribute, remix, adapt, build upon this work noncommercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http:// creativecommons.org/licenses/by-nc/4.0/

REFERENCES

- 1. Kennedy S. Primary dysmenorrhoea. *Lancet* 1997;349:1116.
- Coco AS. Primary dysmenorrhea. Am Fam Physician 1999:60:489–96.
- Grandi G, Ferrari S, Xholli A, *et al.* Prevalence of menstrual pain in young women: what is dysmenorrhea? *J Pain Res* 2012;5:169.
- Kim HO, Lim SW, Woo HY, et al. Premenstrual syndrome and dysmenorrhea in Korean adolescent girls. Korean J Obstet Gynecol 2008;51:1322–9.
- Lee JC, Yu BK, Byeon JH, *et al.* A study on the menstruation of Korean adolescent girls in Seoul. *Korean J Pediatr* 2011;54:201–6.
- Dawood MY. Primary dysmenorrhea: advances in pathogenesis and management. Obstet Gynecol 2006;108:428–41.
- Novak E, REYNOLDS SR. The cause of primary dysmenorrhea: with special reference to hormonal factors. J AMA 1932;99:1466–72.
- Park JS, Park S, Cheon CH, et al. Effects and safety of gyejibongnyeong-hwan on dysmenorrhea caused by blood stagnation: a randomized controlled trial. Evid Based Complement Alternat Med 2013;2013:424730.
- 9. Lefebvre G, Pinsonneault O, Antao V, *et al.* Primary dysmenorrhea consensus guideline. *J Obstet Gynaecol Can* 2005;27:1117–46.
- Deligeorogiou E. Dysmenorrhea. Ann N Y Acad Sci 2000;900:237–44.
- Dawood MY. Nonsteroidal anti-inflammatory drugs and changing attitudes toward dysmenorrhea. *Am J Med* 1988;84:23–9.
- 12. Cho SH, Hwang ÉW. Acupuncture for primary dysmenorrhoea: a systematic review. *BJOG* 2010;117:509–21.
- 13. Yang J, Yu S, Lao L, *et al.* Use of moxibustion to treat primary dysmenorrhea at two interventional times: study protocol for a randomized controlled trial. *Trials* 2015;16:35.
- Yang J, Chen J, Lao L, *et al.* Effectiveness study of moxibustion on pain relief in primary dysmenorrhea: study protocol of a randomized controlled trial. *Evid Based Complement Alternat Med* 2014;2014:434978.
- Chen NN, Han M, Yang H, et al. Chinese herbal medicine Guizhi Fuling Formula for treatment of uterine fibroids: a systematic review of randomised clinical trials. BMC Complement Altern Med 2014;14:2.
- Chen HY, Lin YH, Su IH, *et al.* Investigation on Chinese herbal medicine for primary dysmenorrhea: implication from a nationwide prescription database in Taiwan. *Complement Ther Med* 2014;22:116–25.
- Higgins JPT, Altman DG, Sterne JAC. Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S, eds. Cochrane Handbook for Systematic Reviews of Interventions Version 510 (updated March 2011). The Cochrane Collaboration, 2011. http:// www.cochrane-handbook.org
- Deeks JJ, Higgins JPT, Altman DG. Chapter 9: Analysing data and undertaking meta-analyses. In: Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 510 (updated March 2011).* The Cochrane Collaboration, 2011. http://www.cochrane-handbook.org
- Sterne JAC, Egger M, Moher D. Chapter 10: Addressing reporting biases. In: Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 510 (updated March* 2011). The Cochrane Collaboration, 2011. http://www. cochrane-handbook.org
- Egger M, Davey Smith G, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. BMJ 1997;315:629–34.
- Zhu X, Proctor M, Bensoussan A, et al. Chinese herbal medicine for primary dysmenorrhoea. Cochrane Database Syst Rev 2007;(4): CD005288.
- Yu S, Yang J, Yang M, et al. Application of acupoints and meridians for the treatment of primary dysmenorrhea: a data mining-based literature study. *Evid Based Complement Alternat Med* 2015;2015:752194.
- 23. Chen J, Zhang Z. *Jin gui yao lue*. Beijing: People's Medical Publishing House, 2000.
- 24. Xinrong Y. *Encyclopedic reference of traditional Chinese medicine*. Springer Science & Business Media, 2003.
- Flower A, Lewith G, Liu JP, *et al.* Applying the principles of the Five Phase (Wu Xing) model to inform good practice for studies of Chinese herbal medicine. *Eur J Integr Med* 2016;8:191–7.