

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<u>http://bmjopen.bmj.com</u>).

If you have any questions on BMJ Open's open peer review process please email <u>info.bmjopen@bmj.com</u>

BMJ Open

Cutting care clusters: the creation of an inverse pharmacy care law? An area level analysis exploring the clustering of community pharmacies in England

Journal:	BMJ Open	
Manuscript ID	bmjopen-2018-022109	
Article Type:	Research	
Date Submitted by the Author:	01-Feb-2018	
Complete List of Authors:	Todd, Adam; Newcastle University, School of Pharmacy Thomson, Katie; Newcastle University, Institute of Health and Society Kasim, Adetayo; Durham University, Wolfson Research Institute for Health and Wellbeing Bambra, Clare; Newcastle University	
Keywords:	PUBLIC HEALTH, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, PRIMARY CARE	

SCHOLARONE[™] Manuscripts

BMJ Open

Cutting care clusters: the creation of an inverse pharmacy care law? An area level analysis exploring the clustering of community pharmacies in England

Adam Todd^{1,2,3}, Reader in Pharmaceutical Public Health, <u>adam.todd@newcastle.ac.uk</u> Katie Thomson^{2,3}, Post-doctoral Research Associate, <u>katie.thomson@ncl.ac.uk</u> Adetayo Kasim⁴, Associate Professor in Statistics, <u>a.s.kasim@durham.ac.uk</u> Clare Bambra² Professor of Public Health, <u>clare.bambra@newcastle.ac.uk</u>

1. School of Pharmacy, Newcastle University, Queen Victoria Road, Newcastle upon-Tyne, NE1 7RU, UK.

2. Institute of Health and Society, Newcastle University, Richardson Road, Newcastle upon Tyne, NE2 4AX, UK.

3. Fuse – the UKCRC Centre for Translational Research in Public Health, UK.

4. Wolfson Research Institute for Health and Wellbeing, Durham University, Stockton-on-Tees, TS17 6BH, UK.

Dr. Adam Todd is the corresponding author:

e-mail: adam.todd@newcastle.co.uk

Telephone: +44 (0) 191 208 2355

Abstract

Objectives: To (1) explore the clustering of community pharmacies in England; and (2) determine the relationship between community pharmacy clustering, urbanity and deprivation.

Design: An area level analysis spatial study.

Setting: England.

Primary and secondary outcome measures: Community pharmacy clustering determined as a community pharmacy located within 10 minutes walking distance to another community pharmacy.

Participants: Addresses and postcodes of each community pharmacy in England was used in the analysis. Each pharmacy postcode was assigned to a lower layer super output area (LSOA), which was then matched to urbanity (urban, town and fringe or village, hamlet and isolated dwellings) and deprivation decile (using the Index of Multiple Deprivation score).

Results:

Seventy five percent of community pharmacies in England were located in a 'cluster' (within 10 minutes walking distance of another pharmacy): 19% of community pharmacies were in a cluster of two, while 56% of community pharmacies were in clusters of three or more. There was a linear relationship between community pharmacy clustering and social deprivation – with clustering more prevalent in areas

BMJ Open

of higher deprivation: for community pharmacies located in areas of lowest deprivation (decile 1), there was a significantly lower risk of clustering compared to community pharmacies located in areas of highest deprivation (RR 0.12 [95% CI 0.10, 0.16]).

Conclusions:

Clustering of community pharmacies in England is common, although there is a positive trend between community pharmacy clustering and social deprivation, whereby clustering is more significant in areas of high deprivation. As such, arrangements for future community pharmacy funding should not solely focus upon distance from one pharmacy to another, as means of determining funding allocation, as this could penalise community pharmacies in our most deprived communities, and potentially have a negative effect on other healthcare providers, such as GP and A&E services by increasing their workload.

Keywords: community pharmacy; healthcare access; health inequalities; public health

Word of main manuscript: 2697

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES)

data mining, AI training, and similar technologies

Protected by copyright, including for uses related to text and

Strengths and limitations of this study

- This is the first study exploring community pharmacy clustering, and how this varies according to social deprivation, and urbanity
- We conceptualised a community pharmacy cluster using a 0.5 mile straight line, which is a limitation of the work
- We did not model what would happen for community pharmacy access if the clusters were removed due to community pharmacy closures

ore trick only

Introduction

In recent years, community pharmacies have emerged as strategically important settings, which play a crucial role in delivering healthcare and public health services.[1] In the UK, for example, community pharmacies offer a range of tiered services: the first level of service, the essential service, includes dispensing medication, and providing medication-related advice; the second, the advanced service, allows pharmacists, amongst other things, to provide flu vaccinations and medicines use reviews; the third, the locally commissioned services, are used to address the needs of the local population, and can include smoking cessation, and minor illness schemes.[2] Importantly, through these patient-focused roles, community pharmacy services are able to reduce pressure from other healthcare providers, such as GPs and Accident and Emergency departments.[3, 4]

Indeed, with the average waiting times for a non-urgent GP appointment around 13 days, and the number of people waiting longer than 4 hours in Accident and Emergency departments increasing, it is clear UK healthcare services are under unprecedented strain.[5] Furthermore, it is known that people living in areas of higher socio-economic deprivation are more likely to need primary and secondary health services but that provision is in fact lower in these high-need neighbourhoods: the *inverse care law*.[6] The problem of timely access to healthcare services is therefore adversely affecting people in the most deprived communities – potentially further widening health inequalities and thereby putting even more pressure on healthcare services in the longer term. Previous work has shown that, in contrast to the *inverse care law* for GPs and hospital services, a *positive pharmacy care law* exists – whereby access to community pharmacies is highest in areas of highest deprivation.[7] Community pharmacies are therefore able to reach people in the areas of greatest need thereby offering additional health care access in areas that are traditionally 'under doctored' – including rural areas.[8]

Despite the significant – and perhaps underutilised – potential of community pharmacies, austerity has seen the implementation of UK government funding cuts to the English community pharmacy sector – with some estimates suggesting one in four community pharmacies could close as a result of the reduced funding envelope.[9] It has been argued that because community pharmacies cluster together, some can be cut without impacting on service provision. In a letter to the Pharmaceutical Services Negotiating Committee (PSNC - the body that represents NHS pharmacy contractors in England), the Department of Health state that "40% of pharmacies are in a cluster where there are three or more pharmacies within ten minutes' walk" and that "in some parts of the country there are more pharmacies than are necessary to maintain good access".[10] Despite these claims, and the potential impact on funding allocations and therefore service provision, there are no published studies that explore the clustering of community pharmacies in England, and how such clustering is linked to socio-economic deprivation, and urbanity. The research, therefore, aimed to: (1) explore the clustering of community pharmacies in England by ten minutes walking distance; and (2) determine the relationship between community pharmacy clustering, urbanity and deprivation.

Methods

Design

This study used geographical information systems (GIS) to explore geographical access to community pharmacies, clustering and deprivation. For the

purposes of the study, a community pharmacy was defined as a premises registered with the General Pharmaceutical Council for the purposes of compounding, procurement, storage and distribution of medicines and appliances;[11] we excluded premises that were solely registered as Internet pharmacies in the analysis.

Data and Variables

Community pharmacy data were obtained from the Geo-healthcare access database.[12] This open access database contains data on the address and postcode of each community pharmacy premises in England (matched to their corresponding coordinates using the Office of National Statistics postcode directory, 2014). Community pharmacy locations were mapped and their corresponding 2011 lower layer super output areas (LSOA) were extracted. There are 32,844 LSOAs in England and these geographic areas comprise approximately 1 to 3000 people living in 400 to 1200 households, and were designed to improve the reporting of small area statistics. Urbanity is also a factor in access to healthcare services including community pharmacies so, using the Rural-Urban classification (2011) each community pharmacy LSOA was assigned to one of three categories: (1) urban, (2) town or fringe or village, and (3) hamlet and isolated dwellings.[8] These categories were aggregated from the original Rural-Urban classification which assigns areas to one of six rural or four urban settlement/context types. The 2015 Index of Multiple Deprivation (IMD) score was also matched to each community pharmacy's LSOA (from the Office of National Statistics). The IMD is an overall measure of multiple deprivation experienced by people living in an area, and comprises 37 separate indicators organized across seven domains of deprivation (income, employment,

health and disability, education, skills and training, crime, barriers to housing and services and living environment) which are combined, using appropriate weights.[13]

Data Analysis

Locations of community pharmacies in England were mapped using ArcMap (version 10.3) and 0.5-mile (straight-line) buffers were placed around the site to represent a 10-minute walk, using an average walking speed of 3 miles (4 km) per hour.[14] The number of community pharmacies which were isolated, or clustered in groups of two, or three or more were then extracted for all locations. A cluster was defined if the buffer around a single community pharmacy contained two or more unique sites. The number of community pharmacies within each individual buffer was then calculated and summarised. In addition to examining data for all community pharmacies, further breakdown of clustering took place depending on whether or not the LSOA of the pharmacy was 'urban', 'town or fringe' or 'village, hamlet and isolated dwellings'. This process was repeated for all deprivation deciles based on the IMD 2015 for England, whereby the most deprived decile (decile 1) equated to the most deprived 10% of LSOAs, while the least deprived decile (decile 10) represented the 10% least deprived LSOAs (Figure 1). The Relative Risk (RR) for community pharmacy clustering by deprivation were then calculated according to deprivation decile. For the deprivation analysis, a community pharmacy 'cluster' was considered as three or more community pharmacies within a 10-minute walking distance of each other.

BMJ Open

Results

Overall, our results show that, the percentage of community pharmacies within 10 minutes walking distance (0.5 mile) of one another is 75%: 19% of community pharmacies were in a cluster of two, while 56% of community pharmacies were in clusters of three or more. An example of community pharmacy clustering is shown visually in Figure 2.

Clustering of Community Pharmacies by Urban-Rural Classification

For community pharmacies located in urban areas (n=10,438), there was no clustering for 19% of community pharmacies, 19% were located in a cluster of two, while 62% were located in clusters of three or more. In town and fringe areas (n=1147), there was no clustering in 94% of community pharmacies, 5% were located in clusters of two, while 1% were located in clusters of three or more. In village areas (n=153), there was no clustering in 94% of community pharmacies, 4% were located in clusters of two, while 2% were located in clusters of three or more.

Clustering of Community Pharmacies by Index of Multiple Deprivation (IMD)

When stratifying by IMD, there was, overall, a linear relationship between community pharmacy clustering and social deprivation – with clustering more prevalent in areas of higher deprivation (Figure 3); the highest percentage of community pharmacy clustering was 62%, observed in deprivation decile ten, while the lowest percentage of community pharmacy clustering was 8%, observed in deprivation decile one (Figure 4). Community pharmacies located in the most deprived areas were significantly more likely to exist as clusters: for community pharmacy clusters of three or more, there was a significantly higher risk of clustering

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

BMJ Open

> in deprivation decile ten, compared to all the other deciles. When comparing community pharmacies in decile one (least deprived) to community pharmacies in decile ten (most deprived), there was an 88% lower risk of clustering for community pharmacies in the least deprived areas (Table 1).

Deprivation	No Clustering	Clustering of 2	Clustering of ≥ 3
Decile	RR (95% CI)	RR (95% CI)	RR (95% CI)
1	3.38 (3.06, 3.72)	1.20 (1.01, 1.43)	0.12 (0.10, 0.16)
2	3.08 (2.78, 3.39)	1.06 (0.89, 1.26)	0.27 (0.23, 0.31)
	<u> </u>		
3	2.78 (3.51, 3.08)	1.28 (1.09, 1.49)	0.31 (0.27, 0.35)
	R		
4	2.52 (2.27, 2.80)	1.49 (1.29, 1.72)	0.34 (0.30, 0.38)
5	2.48 (2.23, 2.74)	1.40 (1.21, 1.62)	0.38 (0.34, 0.42)
6	2.41 (2.17, 2.67)	1.50 (1.32, 1.73)	0.37 (0.33, 0.42)
7	1.93 (1.73, 2.15)	1.53 (1.34, 1.75)	0.53 (0.49, 0.56)
8	1.83 (1.65, 2.04)	1.39 (1.22, 1.58)	0.60 (0.56, 0.65)
9	1.47 (1.31, 1.64)	1.40 (1.23, 1.59)	0.73 (0.68, 0.78)
10	1	1	
(reference)		· · · · · · · · · · · · · · · · · · ·	

Table 1: Relative Risk (RR) of community pharmacy clustering according to deprivation decile (1 is the least deprived, while 10 is the most deprived).

Discussion

The key findings of the study show that clustering of community pharmacies in England is common – with around 56% of all community pharmacies existing in a

cluster of three or more. There is also a positive trend between deprivation, and community pharmacy clustering, whereby clustering is significantly more common in areas of high deprivation. Clustering of community pharmacies was also more common in urban areas, when compared to rural areas. This may well reflect both health need and population size in the respective neighbourhoods.

This is the first study that has empirically explored whether community pharmacies are clustered in England – and how such clustering varies according to deprivation, and urbanity. Previous literature has shown that geographical proximity of services can be an important consideration for healthcare utilization: for example, Turnbull and colleagues showed that call rates to primary care centres decreased with increasing distance. Furthermore, the same study also showed that higher call rates were associated with more deprived areas.[15]

In terms of study limitations, whilst we believe our results are robust, and have important policy implications for the way in which community pharmacies are funded, we acknowledge there are several: firstly, we recognize that a 10-minute walk (0.5 miles) from each community pharmacy was represented using a straight-line distance from the central point of each community pharmacy's postcode to create a buffer. This assumes people are able to walk in any direction from that postcode and always in a straight line, while, in reality people are often constrained to pathways that curve, or even be cut off by barriers (such as a lake, train tracks or busy roads). We also acknowledge that we did not consider community pharmacy utilization in our analysis; just because a community pharmacy is located in a particular area, does not necessarily mean people from that area choose to use it. In addition, we did not consider which types of service were being used from the community pharmacies; it is possible that service utilization will change according to health need of the

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES) Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

population (e.g. people living in areas with a high prevalence of smoking might utilise more community pharmacy smoking cessation services). We also acknowledge that we did not explore how community pharmacy clustering varied according to population type; for example, it is possible that some community pharmacies serve different types of populations, such as black and minority ethnic groups, or older populations. It would be prudent, therefore, that future work establishes the types of people that utilize community pharmacies services, and how this varies according to urbanity, deprivation, and local health need.

The policy implications of this work are clear: cutting the funding of community pharmacies by clustering will disproportionately affect community pharmacies located in the most deprived communities: potentially leading to an inverse pharmacy care law. Given that community pharmacies provide effective public health services, [16] this has the potential to further widen health inequalities – should community pharmacies in these areas close. Indeed, it would undermine the Department of Health's responsibilities to reduce inequalities in access to NHS services. There have been recent reports of some large community pharmacy organisations taking the decision to close some of their community pharmacies as they have become "commercially unviable", although it is not yet clear what areas will be affected.[17] We note that the Department of Health has recently introduced a Pharmacy Access Scheme (PhAS), whereby, according to specific criteria, certain community pharmacies will be protected from the reduction in funding.[18] Initially, community pharmacies would not qualify for the scheme if they were located within a one-mile radius of another community pharmacy, although we note that, in view of our preliminary work around community pharmacy clustering, the formula was modified to accommodate community pharmacies located in top 20 per cent of

deprived areas in England within a 0.8 mile radius of another community pharmacy.[19] At present, however, it is not clear if community pharmacy clustering is driven by health need of the local area, or, as many community pharmacies are located in urban areas, they benefit from the increased footfall by being located in these areas. Future work should seek to address this important issue. While we are encouraged by the concept of the PhAS recogising community pharmacies in deprived areas, we believe that future models of funding should be based upon quality metrics around the provision of healthcare and public health services, rather focus on distance or clustering of pharmacies. The Department of Health has also recently introduced a quality payment scheme whereby extra payments will be paid to community pharmacies if certain quality criteria are met. We believe this is significant progress, and it is important that this concept continue which will remove focus on the number of services a community pharmacy undertakes, to that of the quality of the service a community pharmacy undertakes. This will help direct funding to community pharmacies that are engaged in delivering the higher levels of service.

In terms of the impact on the wider healthcare community, previous work shows that if community pharmacy services were not present, people would use alternate – and more costly – branches of the NHS, such as GP and Accident and Emergency services.[3] It is also evident that people living in more deprived areas experience higher morbidity much earlier in life, compared to people living in affluent areas – as represented by lower healthy life expectancy is for example.[20] A study analyzing one million GP consultations in London showed that someone aged 50 years living in the most deprived quintile of English neighbourhoods consults with their GP at the same rate as someone aged 70 years in the most affluent quintile of

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES)

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

neighbourhoods.[21] So, community pharmacy closures – particularly in deprived areas – are likely to increase the workload of general practice. Community pharmacies are able to reach people that other services cannot. For example, a study has shown that owing to their wider accessibility (e.g. proximity; opening times, no appointment required), many people prefer to obtain healthcare from a community pharmacy setting [22] and a qualitative study revealed some people were more likely to use a community pharmacy for healthcare advice owing to their trusting relationship they develop with the pharmacist.[23]

Conclusion

Just over half of community pharmacies in England are located in clusters of three or more by ten minutes walking distance. There is also a positive trend between community pharmacy clustering and social deprivation, whereby clustering is significantly more common in areas of high deprivation. As such, arrangements for future community pharmacy funding should not solely focus upon distance from one pharmacy to another, as means of determining funding allocation, as this could penalise community pharmacies in our most deprived communities, and potentially have a negative effect on other healthcare providers, such as GP and A&E services by increasing their workload. Future funding models of community pharmacy should consider quality of healthcare and public health services provided, as well as health need of the local area.

Figure legends:

Figure 1: Map of England with LSOAs stratified according to deprivation.

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Figure 2: An example of community pharmacy distribution in England, showing: no clustering (blue sphere), clusters of 2 (yellow sphere) and clusters of 3 (green sphere).
Figure 3: Map of the North East of England showing; A: ordinance Survey Map; B: deprivation by LSOA; and, C: community pharmacy clustering.
Figure 4: Percentage clustering of community pharmacies in England according to

social deprivation (1 is the least deprived, while 10 is the most deprived).

References:

1. The Role of the Pharmacist in the Health Care System. The World Health Organization.

Available at: <u>http://apps.who.int/medicinedocs/en/d/Jh2995e/1.6.2.html</u> (accessed 23.01.18)

2. Community Pharmacy Contractual Framework. Pharmaceutical Services Negotiating Committee. Available at: <u>http://psnc.org.uk/contract-it/the-pharmacy-contract/</u> (accessed 23.01.18)

3. Baqir W, Learoyd T, Sim A, Todd A. Cost analysis of a community pharmacy 'minor ailment scheme' across three primary care trusts in the North East of England. *J Public Health (Oxf).* 2011;**33**(4):551-5.

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES)

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

4. Paudyal V, Watson MC, Sach T, Porteous T, Bond CM, Wright DJ, Cleland J, Barton G, Holland R. Are pharmacy-based minor ailment schemes a substitute for other service providers? A systematic review. *Br J Gen Pract.* 2013;**63**(612):e472-81.
5. Average waiting time for GP appointment increases 30% in a year. Pulse Today. Available at: <u>http://www.pulsetoday.co.uk/your-practice/access/average-waiting-time-for-gp-appointment-increases-30-in-a-year/20032025.article</u> (accessed 23.01.18)

6. Tudor Hart J. The inverse care law. The Lancet. 1971;297:405-412.

7. Todd A, Copeland A, Husband A, Kasim A, Bambra C. The positive pharmacy care law: an area-level analysis of the relationship between community pharmacy distribution, urbanity and social deprivation in England. *BMJ Open.* 2014;4(8):e005764.

8. Todd A, Copeland A, Husband A, Kasim A, Bambra C. Access all areas? An area level analysis of accessibility to general practice and community pharmacy services in England by urbanity and social deprivation. *BMJ Open.* 2015;**5**(5):e007328.

9. BBC News. Many high-street pharmacies in England face closure, says minister. Available at: http://www.bbc.co.uk/news/health-35418556 (accessed 23.01.18)

10. Letter to Sue Sharpe, Chief Executive of the PSNC, London. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/486941 /letter-psnc.pdf (accessed 23.01.18)

11. The General Pharmaceutical Council. Available at: https://www.pharmacyregulation.org (accessed 23.01.18)

12. Geo-healthcare access database. Available at: https://collections.durham.ac.uk/files/fq977t77k#.WJSlsRicZBw (accessed 23.01.18)

13. English Indices of Deprivation. Department of Health. Available at: https://www.gov.uk/government/statistics/english-indices-of-deprivation-2015 (accessed 23.01.18)

14. Towards an Urban Renaissance. The urban task force. London: Routledge, 1999.

15. Brown TJ, Todd A, O'Malley C, Moore HJ, Husband AK, Bambra C, Kasim A, Sniehotta FF, Steed L, Smith S, Nield L, Summerbell CD. Community pharmacydelivered interventions for public health priorities: a systematic review of interventions for alcohol reduction, smoking cessation and weight management, including meta-analysis for smoking cessation. *BMJ Open.* 2016;6(2):e009828.

16. Turnbull J, Martin D, Lattimer V, Pope C, Culliford D. Does distance matter? Geographical variation in GP out-of-hours service use: an observational study. *Br J Gen Pract.* 2008;**58**(552):471-7.

17. Lloydspharmacy to close or sell 190 branches across England. The Chemist and Druggist. Available at: <u>https://www.chemistanddruggist.co.uk/news/190-</u>lloydspharmacy-branches-close-across-england (accessed 23.01.18)

18. Community Pharmacy in 2016/17 and Beyond: the Pharmacy Access Scheme. The Department of Health. Available at: <u>https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/561497</u> /Pharmacy_access_scheme_A.pdf (accessed 23.01.18)

19. Community Pharmacy in 2016/17 and Beyond: Impact Assessment. The Department of Health. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/561496/Community_pharmacy_impact_assessment_A.pdf (accessed 23.01.18)

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) .

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

20. Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *The Lancet*. 2012: **380**;37-43.

21. Boomla K, Hull S, Robson J. GP funding formula masks major inequalities for practices in deprived areas. *BMJ*. 2014: **349**;g7648.

22. Anderson C, Thornley T. "It's easier in pharmacy": why some patients prefer to pay for flu jabs rather than use the National Health Service. *BMC Health Serv Res.* 2014; **14**: 35.

23. Lindsey L, Husband A, Steed L, Walton R, Todd A. Helpful advice and hidden expertize: pharmacy users' experiences of community pharmacy accessibility. *J Public Health (Oxf)*. 2017;**38**: 609–615.

Competing interests

The authors declare that they have no competing interests.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Contributions

CB and AT designed the study, led on data interpretation, and supervised all stages of the research. KT led on the GIS modelling, and AK conducted statistical analysis. AT led the drafting of the manuscript with input from CB, KT and AK. AT is guarantor of the work.

Acknowledgements

CB is a member of Fuse (funding reference MR/K02325X/1). Funding for Fuse comes from the British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, the National Institute for Health Research, under the auspices of the UK Clinical Research Collaboration, and is gratefully acknowledged (RF150334). The views expressed in this paper do not necessarily represent those of the funders or UKCRC. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Ethical approval and consent to participate

Ethical approval of this work was not required, as the study used non-patient identifiable secondary data; patients were not actively involved in this research.

Data sharing agreement

All of our raw data is available through the open access Geo-healthcare database:

https://collections.durham.ac.uk/files/fq977t77k#.WJSlsRicZBw



Figure 1: Map of England with LSOAs stratified according to deprivation. \bigcirc Crown Copyright and Database Right [2017]. Ordnance Survey (Digimap Licence)

183x234mm (600 x 600 DPI)

60

1 2 3 4 5 6 7 8 1°28'0"W 1°27'0"W 1°26'0"W 1°25'0"W 9 OUGHTO SPRING 10 F 11 12 13 14 15 54°50'0"N 16 54°50'0"N 17 18 19 Great Epple 20 HETTON-21 East A LE- Hetton Ly 22 23 aintor 24 54°49'0"N 54°49'0"N-25 26 obin 27 Mu M Low Moorsley 28 Easington Lane/ 29 High oorsley 30 31 32 -54°48'0"N 54°48'0"N etton le Hill Het 33 34 1°28'0"W 1°27'0"W 1°26'0"W 1°25'0"W 35 36 0.9 1.2 0 37 Miles 38 Legend 39 Pharmacy locations 40 Pharmacy clusters 3+ 41 2 42 43 44 45 Figure 2: An example of community pharmacy distribution in England, showing: no clustering (blue sphere), 46 clusters of 2 (yellow sphere) and clusters of 3 (green sphere). © Crown Copyright and Database Right 47 [2017]. Ordnance Survey (Digimap Licence) 48 247x331mm (300 x 300 DPI) 49 50 51 52 53 54 55 56 57 58





Figure 3: Map of the North East of England showing; A: ordinance Survey Map; B: deprivation by LSOA; and, C: community pharmacy clustering. © Crown Copyright and Database Right [2017]. Ordnance Survey (Digimap Licence)

434x225mm (300 x 300 DPI)





121x91mm (300 x 300 DPI)

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

BMJ Open

Cutting care clusters: the creation of an inverse pharmacy care law? An area level analysis exploring the clustering of community pharmacies in England

Journal:	BMJ Open
Manuscript ID	bmjopen-2018-022109.R1
Article Type:	Research
Date Submitted by the Author:	26-Apr-2018
Complete List of Authors:	Todd, Adam; Newcastle University, School of Pharmacy Thomson, Katie; Newcastle University, Institute of Health and Society Kasim, Adetayo; Durham University, Wolfson Research Institute for Health and Wellbeing Bambra, Clare; Newcastle University
Primary Subject Heading :	Public health
Secondary Subject Heading:	Health policy, Health services research
Keywords:	PUBLIC HEALTH, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, PRIMARY CARE



BMJ Open

2	
3	
4	
5	
6	
7	
8	
0	
9 10	
10	
11	
12	
13	
14	
15	
16	
17	
18	
10	
עו רכ	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
20	
20	
51	
32	
33	
34	
35	
36	
37	
38	
39	
40	
<u>⊿1</u>	
⊥⊤ 10	
+2 12	
45 44	
44	
45	
46	
47	
48	
49	
50	
51	
52	
52	
22	
54	
55	
56	
57	
58	
59	
60	

Cutting care clusters: the creation of an inverse pharmacy care law? An area level analysis exploring the clustering of community pharmacies in England

Adam Todd*^{1,2,3}, Reader in Pharmaceutical Public Health,

adam.todd@newcastle.ac.uk

Katie Thomson^{2,3}, Post-doctoral Research Associate, <u>katie.thomson@ncl.ac.uk</u> Adetayo Kasim⁴, Associate Professor in Statistics, <u>a.s.kasim@durham.ac.uk</u> Clare Bambra² Professor of Public Health, <u>clare.bambra@newcastle.ac.uk</u>

1. School of Pharmacy, Newcastle University, Queen Victoria Road, Newcastle upon-Tyne, NE1 7RU, UK.

2. Institute of Health and Society, Newcastle University, Richardson Road, Newcastle upon Tyne, NE2 4AX, UK.

3. Fuse – the UKCRC Centre for Translational Research in Public Health, UK.

4. Wolfson Research Institute for Health and Wellbeing, Durham University, Stockton-on-Tees, TS17 6BH, UK.

*Dr. Adam Todd is the corresponding author:

Corresponding author e-mail:

adam.todd@newcastle.co.uk

Telephone: +44 (0

+44 (0) 191 208 2355

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES)

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

Abstract

Objectives: To (1) explore the clustering of community pharmacies in England; and (2) determine the relationship between community pharmacy clustering, urbanity and deprivation.

Design: An area level analysis spatial study.

Setting: England.

Primary and secondary outcome measures: Community pharmacy clustering determined as a community pharmacy located within 10 minutes walking distance to another community pharmacy.

Participants: Addresses and postcodes of each community pharmacy in England was used in the analysis. Each pharmacy postcode was assigned to a lower layer super output area (LSOA), which was then matched to urbanity (urban, town and fringe or village, hamlet and isolated dwellings) and deprivation decile (using the Index of Multiple Deprivation score).

Results:

Seventy five percent of community pharmacies in England were located in a 'cluster' (within 10 minutes walking distance of another pharmacy): 19% of community pharmacies were in a cluster of two, while 56% of community pharmacies were in clusters of three or more. There was a linear relationship between community

pharmacy clustering and social deprivation – with clustering more prevalent in areas of higher deprivation: for community pharmacies located in areas of lowest deprivation (decile 1), there was a significantly lower risk of clustering compared to community pharmacies located in areas of highest deprivation (RR 0.12 [95% CI 0.10, 0.16]).

Conclusions:

Clustering of community pharmacies in England is common, although there is a positive trend between community pharmacy clustering and social deprivation, whereby clustering is more significant in areas of high deprivation. As such, arrangements for future community pharmacy funding should not solely focus upon distance from one pharmacy to another, as means of determining funding allocation, as this could penalise community pharmacies in our most deprived communities, and potentially have a negative effect on other healthcare providers, such as GP and A&E services by increasing their workload.

Keywords: community pharmacy; healthcare access; health inequalities; public health

Word of main manuscript: 3123

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) .

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES).

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

Strengths and limitations of this study

- This is the first study exploring community pharmacy clustering, and how this varies according to social deprivation, and urbanity
- We conceptualised a community pharmacy cluster using a 0.5 mile straight line, which is a limitation of the work
- We did not model what would happen for community pharmacy access if the clusters were removed due to community pharmacy closures

or review only

Introduction

In recent years, community pharmacies have emerged as strategically important settings, which play a crucial role in delivering healthcare and public health services.[1] In the UK, for example, community pharmacies offer a range of tiered services: the first level of service, the essential service, includes dispensing medication, and providing medication-related advice; the second, the advanced service, allows pharmacists, amongst other things, to provide flu vaccinations and medicines use reviews; the third, the locally commissioned services, are used to address the needs of the local population, and can include smoking cessation, and minor illness schemes.[2] Importantly, through these patient-focused roles, community pharmacy services are able to manage people with minor illnesses and some long-term conditions. This is advantageous, people with minor illness can be directed away from other healthcare providers, such as GPs and Accident and Emergency departments.[3, 4] This approach therefore has the potential to free up valuable healthcare resource, which allows other primary and secondary care services to mange people with more serious conditions where community pharmacy involvement would not be appropriate.

Indeed, with the average waiting times for a non-urgent GP appointment around 13 days, and the number of people waiting longer than 4 hours in Accident and Emergency departments increasing, it is clear UK healthcare services are under unprecedented strain.[5] Furthermore, it is known that people living in areas of higher socio-economic deprivation are more likely to need primary and secondary health services, but that provision is in fact lower in these high-need neighbourhoods: the *inverse care law*.[6] The problem of timely access to healthcare services is therefore

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

adversely affecting people in the most deprived communities – potentially further widening health inequalities and thereby putting even more pressure on healthcare services in the longer term. Previous work has shown that, in contrast to the *inverse care law* for GPs and hospital services, a *positive pharmacy care law* exists – whereby people living in areas of highest deprivation have better access to community pharmacies.[7] It is also known that the urbanity is an area is in important consideration for healthcare access, with people from urban areas living in greater proximity to GP services, when compared to people living in rural areas.[8] Community pharmacies therefore have the potential to reach people in the areas of greatest need thereby offering additional health care access in areas that are traditionally 'under doctored' – including rural areas.[8]

Despite the significant – and perhaps underutilised – potential of community pharmacies, austerity has seen the implementation of UK government funding cuts to the English community pharmacy sector – with some estimates suggesting one in four community pharmacies could close as a result of the reduced funding envelope.[9] It has been argued that because community pharmacies cluster together, some can be cut without impacting on service provision. In a letter to the Pharmaceutical Services Negotiating Committee (PSNC - the body that represents NHS pharmacy contractors in England), the Department of Health state that "40% of pharmacies are in a cluster where there are three or more pharmacies within ten minutes' walk" and that "in some parts of the country there are more pharmacies than are necessary to maintain good access".[10] Despite these claims, and the potential impact on funding allocations and therefore service provision, there are no published studies that explore the clustering of community pharmacies in England, and how such clustering is linked to socio-economic deprivation, and urbanity. The research, therefore, aimed to: (1)

BMJ Open

explore the clustering of community pharmacies in England by ten minutes walking distance; and (2) determine the relationship between community pharmacy clustering, urbanity and deprivation.

Methods

Design

This study used geographical information systems (GIS) to explore community pharmacy clustering according to urbanity and deprivation. For the purposes of the study, a community pharmacy was defined as a premises registered with the General Pharmaceutical Council for the purposes of compounding, procurement, storage and distribution of medicines and appliances;[11] we excluded premises that were solely registered as Internet pharmacies in the analysis.

Data and Variables

Community pharmacy data were obtained from the Geo-healthcare access database.[12] This open access database contains data on the address and postcode of each community pharmacy premises in England (matched to their corresponding coordinates using the Office of National Statistics postcode directory, 2014); the last update of the dataset was in 2016. Community pharmacy locations were mapped and their corresponding 2011 lower layer super output areas (LSOA) were extracted. There are 32,844 LSOAs in England and these geographic areas comprise approximately 1 to 3000 people living in 400 to 1200 households, and were designed to improve the reporting of small area statistics. Urbanity is also a factor in access to healthcare services including community pharmacy LSOA was assigned to one of three

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

categories: (1) urban, (2) town or fringe or village, and (3) hamlet and isolated dwellings. These categories were aggregated from the original Rural-Urban classification which assigns areas to one of six rural or four urban settlement/context types. The 2015 Index of Multiple Deprivation (IMD) score was also matched to each community pharmacy's LSOA (from the Office of National Statistics). The IMD is an overall measure of multiple deprivation experienced by people living in an area, and comprises 37 separate indicators organized across seven domains of deprivation (income, employment, health and disability, education, skills and training, crime, barriers to housing and services and living environment) which are combined, using appropriate weights.[13]

Data Analysis

Locations of community pharmacies in England were mapped using ArcMap (version 10.3) and 0.5-mile (straight-line) buffers were placed around the site to represent a 10-minute walk, using an average walking speed of 3 miles (4 km) per hour.[14] The number of community pharmacies which were isolated, or clustered in groups of two, or three or more were then extracted for all locations. A cluster was defined if the buffer around a single community pharmacy contained two or more unique sites. The number of community pharmacies within each individual buffer was then calculated and summarised. In addition to examining data for all community pharmacies, further breakdown of clustering took place depending on whether or not the LSOA of the pharmacy was 'urban', 'town or fringe' or 'village, hamlet and isolated dwellings'. This process was repeated for all deprivation deciles based on the IMD 2015 for England, whereby the most deprived decile (decile 1) equated to the most deprived 10% of LSOAs, while the least deprived decile (decile

BMJ Open

10) represented the 10% least deprived LSOAs (Figure 1). The Relative Risk (RR) for community pharmacy clustering by deprivation were then calculated according to deprivation decile. For the deprivation analysis, a community pharmacy 'cluster' was considered as three or more community pharmacies within a 10-minute walking distance of each other.

Patient and Public Involvement

As this study involved secondary data analysis from the Geo-healthcare access database, patients or the public were not involved in the design, or delivery of this research.

Ethical approval

Ethical approval of this work was not required, as the study used non-patient identifiable secondary data; patients were not actively involved in this research.

Results

Overall, our results show that, the percentage of community pharmacies within 10 minutes walking distance (0.5 mile) of one another is 75%: 19% of community pharmacies were in a cluster of two, while 56% of community pharmacies were in clusters of three or more. An example of community pharmacy clustering is shown visually in Figure 2.

Clustering of Community Pharmacies by Urban-Rural Classification

For community pharmacies located in urban areas (n=10,438), there was no clustering for 19% of community pharmacies, 19 % were located in a cluster of two,

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES)

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

while 62% were located in clusters of three or more. In town and fringe areas (n=1147), there was no clustering in 94% of community pharmacies, 5% were located in clusters of two, while 1% were located in clusters of three or more. In village areas (n=153), there was no clustering in 94% of community pharmacies, 4% were located in clusters of two, while 2% were located in clusters of three or more.

Clustering of Community Pharmacies by Index of Multiple Deprivation (IMD)

When stratifying by IMD, there was, overall, a linear relationship between community pharmacy clustering and social deprivation – with clustering more prevalent in areas of higher deprivation (Figure 3); the highest percentage of community pharmacy clustering was 62%, observed in deprivation decile ten, while the lowest percentage of community pharmacy clustering was 8%, observed in deprivation decile one (Figure 4). Community pharmacies located in the most deprived areas were significantly more likely to exist as clusters: for community pharmacy clustering in deprivation decile ten, compared to all the other deciles. When comparing community pharmacies in decile one (least deprived) to community pharmacies in decile ten (most deprived), there was an 88% lower risk of clustering for community pharmacies in the least deprived areas (Table 1).

Deprivation	No Clustering	Clustering of 2	Clustering of ≥ 3
Decile	RR (95% CI)	RR (95% CI)	RR (95% CI)
1	3.38 (3.06, 3.72)	1.20 (1.01, 1.43)	0.12 (0.10, 0.16)
2	3.08 (2.78, 3.39)	1.06 (0.89, 1.26)	0.27 (0.23, 0.31)
3	2.78 (3.51, 3.08)	1.28 (1.09, 1.49)	0.31 (0.27, 0.35)
4	2.52 (2.27, 2.80)	1.49 (1.29, 1.72)	0.34 (0.30, 0.38)
5	2.48 (2.23, 2.74)	1.40 (1.21, 1.62)	0.38 (0.34, 0.42)
6	2.41 (2.17, 2.67)	1.50 (1.32, 1.73)	0.37 (0.33, 0.42)
7	1.93 (1.73, 2.15)	1.53 (1.34, 1.75)	0.53 (0.49, 0.56)
8	1.83 (1.65, 2.04)	1.39 (1.22, 1.58)	0.60 (0.56, 0.65)
9	1.47 (1.31, 1.64)	1.40 (1.23, 1.59)	0.73 (0.68, 0.78)
10	1		1
(reference)			

Table 1: Relative Risk (RR) of community pharmacy clustering according to deprivation decile (1 is the least deprived, while 10 is the most deprived).

Discussion

The key findings of the study show that clustering of community pharmacies in England is common – with around 56% of all community pharmacies existing in a cluster of three or more. There is also a positive trend between deprivation, and community pharmacy clustering, whereby clustering is significantly more common in areas of high deprivation. Clustering of community pharmacies was also more

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

common in urban areas, when compared to rural areas. This may well reflect both health need and population size in the respective neighbourhoods.

This is the first study that has empirically explored whether community pharmacies are clustered in England – and how such clustering varies according to deprivation, and urbanity. Previous literature has shown that geographical proximity of services can be an important consideration for healthcare utilization: for example, Turnbull and colleagues showed that call rates to primary care centres decreased with increasing distance. Furthermore, the same study also showed that higher call rates were associated with more deprived areas.[15] Furthermore, Lin and colleagues examined travel distance to hospital and the associated effect on hospitalistions in Canada, and showed that admission rates were inversely proportional to hospital distance.[16] It has also been shown in the literature that proximity to healthcare services is not only an important consideration for healthcare utilization, but is an important factor for health outcome. For example, a study by Okwaraji and colleagues showed that, in a remote area of rural Ethiopia, the distance to the nearest health centre had a significant effect on child mortality – with children living greater than 1.5 hours walk from the health centre having a two to three greater risk of death compared to children who live within 1.5 hours walk.[17] Clearly the context of this work is different to healthcare settings in England, so limited comparisons can be drawn. However, given the well-established relationship between health and place, it would be prudent for future research to explore how community pharmacy distribution is associated with health of a particular area. The literature has also shown that healthcare access from a geographical perspective tends to be lower amongst rural communities when compared to more urban communities. For example, Lovatt and colleagues, who explore GP accessibility by car travel and public

transport, showed examples of rural areas in East Anglia (typically with low levels of personal mobility and high health need) where there was no daytime bus services or no community transport, which allowed travel/access to a GP.[18] These challenges have been summarised by Baird and Wright, who coined the term 'rural health deprivation', and argue that, to improve the health of the nation, more needs to be done to develop care pathways for rural communities.[19] Although we have not measured accessibility of community pharmacy by population, our findings that community pharmacy clustering is lower in rural areas lend support to the hypothesis that accessibility to healthcare is lower amongst rural communities.

In terms of study limitations, whilst we believe our results are robust, and have important policy implications for the way in which community pharmacies are funded, we acknowledge there are several: firstly, we recognize that a 10-minute walk (0.5 miles) from each community pharmacy was represented using a straight-line distance from the central point of each community pharmacy's postcode to create a buffer. This assumes people are able to walk in any direction from that postcode and always in a straight line, while, in reality people are often constrained to pathways that curve, or even be cut off by barriers (such as a lake, train tracks or busy roads). We also acknowledge that we did not consider community pharmacy utilization in our analysis; just because a community pharmacy is located in a particular area, does not necessarily mean people from that area choose to use it. In addition, we did not consider which types of service were being used from the community pharmacies; it is possible that service utilization will change according to health need of the population (e.g. people living in areas with a high prevalence of smoking might utilise more community pharmacy smoking cessation services). We also acknowledge that we did not explore how community pharmacy clustering varied according to

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

population type; for example, it is possible that some community pharmacies serve different types of populations, such as black and minority ethnic groups, or older populations. It would be prudent, therefore, that future work establishes the types of people that utilize community pharmacies services, and how this varies according to urbanity, deprivation, and local health need.

The policy implications of this work are clear: cutting the funding of community pharmacies by clustering will disproportionately affect community pharmacies located in the most deprived communities: potentially leading to an *inverse pharmacy care law.* Given that community pharmacies provide effective public health services, [20] this has the potential to further widen health inequalities should community pharmacies in these areas close. Indeed, it would undermine the Department of Health's responsibilities to reduce inequalities in access to NHS services. There have been recent reports of some large community pharmacy organisations taking the decision to close some of their community pharmacies as they have become "commercially unviable", although it is not yet clear what areas will be affected. [21] We note that the Department of Health has recently introduced a Pharmacy Access Scheme (PhAS), whereby, according to specific criteria, certain community pharmacies will be protected from the reduction in funding.[22] Initially, community pharmacies would not qualify for the scheme if they were located within a one-mile radius of another community pharmacy, although we note that, in view of our preliminary work around community pharmacy clustering, the formula was modified to accommodate community pharmacies located in top 20 per cent of deprived areas in England within a 0.8 mile radius of another community pharmacy.[23] At present, however, it is not clear if community pharmacy clustering is driven by health need of the local area, or, as many community pharmacies are

located in urban areas, they benefit from the increased footfall by being located in these areas. Future work should seek to address this important issue. While we are encouraged by the concept of the PhAS recogising community pharmacies in deprived areas, we believe that future models of funding should be based upon quality metrics around the provision of healthcare and public health services, rather focus on distance or clustering of pharmacies. The Department of Health has also recently introduced a quality payment scheme whereby extra payments will be paid to community pharmacies if certain quality criteria are met. We believe this is significant progress, and it is important that this concept continue which will remove focus on the *number* of services a community pharmacy undertakes, to that of the quality of the service a community pharmacy undertakes. This approach is critical, given that a recent study highlighted that community services are often not related to need of the local population, and can be influenced by pharmacy ownership type.[24] The development of a quality metric will help direct funding to community pharmacies that are engaged in delivering the higher levels of service.

In terms of the impact on the wider healthcare community, previous work shows that if community pharmacy services were not present, people would use alternate – and more costly – branches of the NHS, such as GP and Accident and Emergency services.[3] It is also evident that people living in more deprived areas experience higher morbidity much earlier in life, compared to people living in affluent areas – as represented by lower healthy life expectancy.[25] A study analyzing one million GP consultations in London showed that someone aged 50 years living in the most deprived quintile of English neighbourhoods consults with their GP at the same rate as someone aged 70 years in the most affluent quintile of neighbourhoods.[26] So, community pharmacy closures – particularly in deprived areas – are likely to BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

increase the workload of general practice. Community pharmacies are able to reach people that other services cannot. For example, a study has shown that owing to their wider accessibility (e.g. proximity; opening times, no appointment required), many people prefer to obtain healthcare from a community pharmacy setting [27] and a qualitative study revealed some people were more likely to use a community pharmacy for healthcare advice owing to their trusting relationship they develop with the pharmacist.[28]

Conclusion

Just over half of community pharmacies in England are located in clusters of three or more by ten minutes walking distance. There is also a positive trend between community pharmacy clustering and social deprivation, whereby clustering is significantly more common in areas of high deprivation. As such, arrangements for future community pharmacy funding should not solely focus upon distance from one pharmacy to another, as means of determining funding allocation, as this could penalise community pharmacies in our most deprived communities, and potentially have a negative effect on other healthcare providers, such as GP and A&E services by increasing their workload. Future funding models of community pharmacy should consider quality of healthcare and public health services provided, as well as health need of the local area.

Figure legends:

Figure 1: Map of England with LSOAs stratified according to deprivation.

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES)

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

BMJ Open

Figure 2: An example of community pharmacy distribution in England, showing: no clustering (blue sphere), clusters of 2 (yellow sphere) and clusters of 3 (green sphere).Figure 3: Map of the North East of England showing; A: ordinance Survey Map; B: deprivation by LSOA; and, C: community pharmacy clustering.Figure 4: Percentage clustering of community pharmacies in England according to

social deprivation (1 is the least deprived, while 10 is the most deprived).

to beet texter only

References:

1. The Role of the Pharmacist in the Health Care System. The World Health Organization.

Available at: <u>http://apps.who.int/medicinedocs/en/d/Jh2995e/1.6.2.html</u> (accessed 25.04.18)

2. Community Pharmacy Contractual Framework. Pharmaceutical Services Negotiating Committee. Available at: <u>http://psnc.org.uk/contract-it/the-pharmacy-contract/</u> (accessed 25.04.18)

3. Baqir W, Learoyd T, Sim A, Todd A. Cost analysis of a community pharmacy 'minor ailment scheme' across three primary care trusts in the North East of England. *J Public Health (Oxf)*. 2011;**33**(4):551-5.

4. Paudyal V, Watson MC, Sach T, Porteous T, Bond CM, Wright DJ, Cleland J, Barton G, Holland R. Are pharmacy-based minor ailment schemes a substitute for other service providers? A systematic review. *Br J Gen Pract.* 2013;**63**(612):e472-81.

5. Average waiting time for GP appointment increases 30% in a year. Pulse Today. Available at: <u>http://www.pulsetoday.co.uk/your-practice/access/average-waiting-time-for-gp-appointment-increases-30-in-a-year/20032025.article</u> (accessed 25.04.18)

6. Tudor Hart J. The inverse care law. *The Lancet*. 1971;**297**:405-412.

7. Todd A, Copeland A, Husband A, Kasim A, Bambra C. The positive pharmacy care law: an area-level analysis of the relationship between community pharmacy distribution, urbanity and social deprivation in England. *BMJ Open.* 2014;4(8):e005764.

BMJ Open

Protected by copyright, including fo		MJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 /
r uses related to text and data mining,	Enseignement Superieur (ABES) .	August 2018. Downloaded from http://b
Al training, and similar technologies.		mjopen.bmj.com/ on June 13, 2025 a
		t Agence Bibliographique de l

8. Todd A, Copeland A, Husband A, Kasim A	, Bambra C. Access all areas? An area
level analysis of accessibility to general practice	e and community pharmacy services
in England by urbanity and social deprivation.	BMJ Open. 2015; 5 (5):e007328.
9. BBC News. Many high-street pharmacies in	n England face closure, says minister.
Available at: <u>http://www.bbc.co.uk/news/health</u>	1-35418556 (accessed 25.04.18)
10. Letter to Sue Sharpe, Chief Executive o	of the PSNC, London. Available at:
https://www.gov.uk/government/uploads/system	n/uploads/attachment_data/file/486941
/letter-psnc.pdf (accessed 25.04.18)	
11. The General Pharmaceutical	l Council. Available at:
https://www.pharmacyregulation.org (accessed	25.04.18)
12. Geo-healthcare access dat	tabase. Available at:
https://collections.durham.ac.uk/files/fq977t77k	<u> (accessed 25.04.18)</u>
13. English Indices of Deprivation. De	partment of Health. Available at:
https://www.gov.uk/government/statistics/engli	sh-indices-of-deprivation-2015
(accessed 25.04.18)	
14. Towards an Urban Renaissance. The urban	task force. London: Routledge, 1999.
15. Turnbull J, Martin D, Lattimer V, Pope C	C, Culliford D. Does distance matter?
Geographical variation in GP out-of-hours ser	vice use: an observational study. Br J
Gen Pract. 2008; 58 (552):471-7.	
16. Lin G, Allan DE, Penning MJ. Examining	g Distance Effects on Hospitalizations
Using GIS: A Study of Three Health Reg	gions in British Columbia, Canada.
Environment and Planning A. 2002; 24 : 2037-20	053.
17. Okwaraji YB, Cousens S, Berhane Y, M	Mulholland K, Edmond K. Effect of
geographical access to health facilities on	child mortality in rural Ethiopia: a

18. Lovett A, Haynes R, Sünnenberg G, Gale S. Car travel time and accessibility by bus to general practitioner services: a study using patient registers and GIS. *Soc Sci Med.* 2002;**55**(1):97-111.

19. Baird AG, Wright N. Poor access to care: rural health deprivation? *Br J Gen Pract.* 2006;**56**(529):567-8.

20. Brown TJ, Todd A, O'Malley C, Moore HJ, Husband AK, Bambra C, Kasim A, Sniehotta FF, Steed L, Smith S, Nield L, Summerbell CD. Community pharmacydelivered interventions for public health priorities: a systematic review of interventions for alcohol reduction, smoking cessation and weight management, including meta-analysis for smoking cessation. *BMJ Open.* 2016;6(2):e009828.

21. Lloydspharmacy to close or sell 190 branches across England. The Chemist and Druggist. Available at: <u>https://www.chemistanddruggist.co.uk/news/190-</u>lloydspharmacy-branches-close-across-england (accessed 25.04.18)

22. Community Pharmacy in 2016/17 and Beyond: the Pharmacy Access Scheme. The Department of Health. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/561497 /Pharmacy access scheme A.pdf (accessed 25.04.18)

23. Community Pharmacy in 2016/17 and Beyond: Impact Assessment. The Department of Health. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/561496 /Community pharmacy impact assessment A.pdf (accessed 25.04.18)

24. Hann M, Schafheutle EI, Bradley F, Elvey R, Wagner A, Halsall D, Hassell K, Jacobs S. Organisational and extraorganisational determinants of volume of service delivery by English community pharmacies: a cross-sectional survey and secondary data analysis. *BMJ Open.* 2017;7(10):e017843.

BMJ Open

25. Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *The Lancet*. 2012: **380**;37-43.

26. Boomla K, Hull S, Robson J. GP funding formula masks major inequalities for practices in deprived areas. *BMJ*. 2014: **349**;g7648.

27. Anderson C, Thornley T. "It's easier in pharmacy": why some patients prefer to pay for flu jabs rather than use the National Health Service. *BMC Health Serv Res.*2014; 14: 35.

28. Lindsey L, Husband A, Steed L, Walton R, Todd A. Helpful advice and hidden expertize: pharmacy users' experiences of community pharmacy accessibility. *J Public Health (Oxf)*. 2017;**38**: 609–615.

Competing interests

The authors declare that they have no competing interests.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Contributions

CB and AT designed the study, led on data interpretation, and supervised all stages of the research. KT led on the GIS modelling, and AK conducted statistical analysis. AT led the drafting of the manuscript with input from CB, KT and AK. AT is guarantor of the work.

Acknowledgements

CB is a member of Fuse (funding reference MR/K02325X/1). Funding for Fuse comes from the British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, the National Institute for Health Research, under the auspices of the UK Clinical Research Collaboration, and is gratefully acknowledged (RF150334). The views expressed in this paper do not necessarily represent those of the funders or UKCRC. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Ethical approval and consent to participate

Ethical approval of this work was not required, as the study used non-patient identifiable secondary data; patients were not actively involved in this research.

Data sharing agreement

All of our raw data is available through the open access Geo-healthcare database: https://collections.durham.ac.uk/files/fq977t77k#.WJSlsRicZBw



Figure 1: Map of England with LSOAs stratified according to deprivation. © Crown Copyright and Database Right [2017]. Ordnance Survey (Digimap Licence)

183x234mm (600 x 600 DPI)





Figure 2: An example of community pharmacy distribution in England, showing: no clustering (blue sphere), clusters of 2 (yellow sphere) and clusters of 3 (green sphere). © Crown Copyright and Database Right [2017]. Ordnance Survey (Digimap Licence)

247x331mm (300 x 300 DPI)



Figure 3: Map of the North East of England showing; A: ordinance Survey Map; B: deprivation by LSOA; and, C: community pharmacy clustering. © Crown Copyright and Database Right [2017]. Ordnance Survey (Digimap Licence)

434x225mm (300 x 300 DPI)

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.





121x91mm (300 x 300 DPI)

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open: first published as 10.1136/bmjopen-2018-022109 on 1 August 2018. Downloaded from http://bmjopen.bmj.com/ on June 13, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.