


BMJ Open Scleroderma Patient-centered Intervention Network—Scleroderma Support group Leader EDucation (SPIN-SSLED) program: non-randomised feasibility trial

Brett D Thombs ^{1,2,3,4,5,6,7}, Laura Dyas,⁸ Mia Pépin,¹ Kylene Aguila,¹ Marie-Eve Carrier,¹ Lydia Tao,¹ Sami Harb,¹ Vanessa L Malcarne,^{9,10} Ghassan El-Baalbaki,¹¹ Sandra Peláez,^{1,6} Maureen Sauve,¹² Marie Hudson,^{1,4} Robert W Platt,^{1,3} SPIN-SSLED Patient Advisory Team

To cite: Thombs BD, Dyas L, Pépin M, *et al.* Scleroderma Patient-centered Intervention Network—Scleroderma Support group Leader EDucation (SPIN-SSLED) program: non-randomised feasibility trial. *BMJ Open* 2019;9:e029935. doi:10.1136/bmjopen-2019-029935

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-029935>).

Received 20 February 2019
Revised 27 September 2019
Accepted 16 October 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Brett D Thombs, Psychiatry, McGill University, Montreal, Quebec H3A 0G4, Canada; brett.thombs@mcgill.ca

ABSTRACT

Objectives The Scleroderma Patient-centered Intervention Network—Scleroderma Support group Leader EDucation (SPIN-SSLED) Programme was designed to improve confidence and self-efficacy and to reduce burden for support group leaders. Objectives were to (1) evaluate feasibility of programme delivery, including required resources, management issues and scientific aspects (eg, performance of outcome measures) and (2) assess user satisfaction and identify any modifications needed to improve programme content or delivery based on participant feedback.

Design Non-randomised feasibility trial.

Setting North American patient organisations.

Participants Current support group leaders or potential new leaders referred by patient organisations.

Intervention The programme included 13 modules delivered live via videoconference over 3 months (April to July 2018) in 60 to 90 min sessions.

Outcome measures (1) Elements of feasibility, including enrolment and consent procedures, percentage of referred group leaders who consented to participate, session attendance and technical support requirements; (2) programme usability, understandability, organisation and clarity; (3) leader satisfaction with the programme and (4) planned trial outcome measures, including support group leader self-efficacy, burnout, emotional distress and physical function.

Results All 12 referred potential participants consented to enrol, and 10 were included in two training groups of five participants each. Participants attended 95% of sessions. Required technical support was minimal, and videoconferencing technology functioned well. Overall programme satisfaction rating was 9.4/10. Mean item rating on the eight items of the Client Satisfaction Questionnaire-8 was 3.83 (1=low satisfaction; 4=high satisfaction). Pre-post scores on the Scleroderma Support Group Leader Self-efficacy Scale increased by 1.7 SDs (large effect); scores on burnout, emotional distress and physical function improved by 0.44, 0.38 and 0.45 SDs (moderate effects).

Strengths and limitations of this study

- This is the first trial to test the feasibility of delivering an education and training programme via videoconference to peer support group leaders.
- The education and training programme was developed in a partnership that included scleroderma peer support group leaders, patient organisation leaders and researchers and healthcare professionals.
- Trial outcomes included elements of feasibility (eg, management, resources, scientific aspects); programme usability, understandability, organisation and clarity; leader satisfaction with the programme; and planned trial outcome measures.
- This was a non-randomised feasibility trial that only included 10 participants in two training groups and did not include a control group; generalisability of results outside of scleroderma is not known.

Conclusion The SPIN-SSLED Programme was feasibly delivered, including management, resource and scientific aspects. Participant satisfaction was high. The programme is ready to be tested in a full-scale randomised controlled trial.

Trial registration number NCT03508661

INTRODUCTION

People with rare diseases face the same challenges as those with more common diseases plus unique challenges, including limited disease education and lack of specialised support options.^{1–12} Professionally organised support services for common diseases are often available through the healthcare system,^{13 14} but are not typically available for rare diseases.^{10 15} As a result, some people with rare diseases rely on peer-led support groups for disease-specific education and

support.^{16–20} Support group activities typically involve an educational or information-sharing component and the exchange of emotional and practical support.^{14 18–22}

Systemic sclerosis (SSc), or scleroderma, is a rare chronic, autoimmune connective tissue disease characterised by abnormal fibrotic processes and excessive collagen production.^{23–25} Support groups, most led by people with SSc, play an important role for many people with the disease.^{17 26–30} Many people with SSc, however, cannot access support groups because they are not available close to where they live, and many initiated support groups are not sustained due to challenges that could be addressed via leader training.^{18 19}

A systematic review of randomised controlled trials (RCTs) that evaluated the effects of training programme for patient leaders of illness-based support groups on the competency, self-efficacy, burden and emotional well-being of group leaders identified only one RCT that met inclusion criteria.³¹ That trial³² evaluated confidence and self-efficacy of cancer support group leaders randomised to either 4-month long high-resource (n=29; website, discussion forum, 2-day face-to-face training) or low-resource (n=23; website, discussion forum) interventions. The RCT did not find evidence that the high-resource programme was more effective. However, the trial was substantially underpowered, not enough information was provided to determine intervention content or how it was delivered, and the risk of bias was high due to methodological limitations.

The Scleroderma Patient-centered Intervention Network (SPIN) partnered with SSc patients and patient organisations to develop the Scleroderma Support group Leader EDucation (SPIN-SSLED) Programme. The programme is a 13-session group videoconference training programme designed to improve skills and self-efficacy, reduce burnout and improve emotional and physical function among support group leaders. The objectives of the SPIN-SSLED feasibility trial were to (1) evaluate the feasibility of steps needed to take place in a planned full-scale trial, including the required resources (eg, staffing, time and budget), management issues (eg, related to optimising performance of personnel and data systems) and scientific aspects (eg, recruitment rates of eligible leaders, acceptability of intervention to leaders, assessing performance of outcome measures) and (2) identify any modifications needed to improve the content or delivery of the SPIN-SSLED Programme based on participant feedback.

METHODS

The SPIN-SSLED feasibility trial was a non-randomised study. It was registered prior to enrolling participants (NCT03508661) and, although not a randomised study, is reported based on items from the Consolidated Standards of Reporting Trials (CONSORT) extension for randomised pilot and feasibility trials.³³ There were no

changes to the feasibility trial protocol and no changes to planned outcomes after commencement of the trial.

Participants

Eligible participants for the SPIN-SSLED feasibility trial were current SSc support group leaders or were identified by Scleroderma Canada or the Scleroderma Foundation (USA) as a new leader who will initiate a new support group, were able to use the internet to access and participate in training sessions and to complete study questionnaires online, were available to participate at times when sessions were scheduled and were English-speaking, since both groups in the feasibility trial were conducted in English. The full-scale trial will include groups conducted in French, but individuals who participated in the feasibility study will be excluded from the full-scale RCT. Thus, to ensure that there will be an adequate number of French-speaking participants in the full-scale RCT, only English-speaking leaders were included in the feasibility study.

Procedures

For the purpose of testing the feasibility of administering the SPIN-SSLED Programme, we sought 10 group leaders to participate in two separate training groups of five participants each. We asked Scleroderma Canada and the Scleroderma Foundation to generate an initial list of 12 interested potential participants and obtained permission for the SPIN team to send them an email with an invitation to participate in the feasibility trial and a copy of the consent form. Following the initial email, SPIN personnel contacted potential participants by phone within 48 hours to describe the study, assess their eligibility, review the consent form and answer questions they may have had about the study. Eligible leaders who verbally agreed to enrol in the study received a second email with the consent form again attached, and they were able to consent via email by replying, 'I have read the consent form and understand the terms of the feasibility study. I agree to participate in the study testing the feasibility of the SPIN-SSLED Program.' The first 10 people to respond and consent were enrolled, and the other 2 were put on a waiting list. All leaders who consented to participate and enrolled received an email invitation including a clickable link to the online data management platform where they were asked to complete baseline study measures. The email also included the date of their first training session, the topic of the first session and information on how to login to the videoconferencing system, as well as a link to the SPIN-SSLED online forum platform, where the programme manual and associated PowerPoint slides were available. Ongoing email and phone technical support was available to help leaders with the consent process, access to the data management platform to complete study measures and training sessions.

Intervention

The SPIN-SSLED Programme was developed by a team of researchers and healthcare professionals with expertise in SSc, patient organisation representatives and a Support Group Advisory Team comprised of people with SSc who are current SSc support group leaders. The programme content and design were based on results of our preliminary research on support groups in SSc, including individual interviews and surveys with leaders, members and non-attenders,^{17–20} and informed by instructional material for support group leaders in other diseases that we identified via the internet and by consultations with support group leaders. The programme uses a problem-based learning approach. Problem-based learning is a learner-centred approach that integrates theory and practice by providing the necessary knowledge and skills, presenting a complex, real-world problem, then working to identify an approach to solving the problem.^{34–37} To implement this, each module introduces a topic and provides an overview of key information. Then, there is a guided discussion among training group participants about possible approaches and solutions to problems.

The SPIN-SSLED Programme included 13 modules that are delivered live via videoconference over the course of 3 months. Each module is delivered in a 60 to 90 min session. Module topics include (1) the leader's role; (2) starting a support group; (3) structuring a support group meeting; (4) scleroderma 101; (5) successful support group culture; (6) managing support group dynamics I; (7) managing support group dynamics II; (8) grief and crisis in scleroderma; (9) marketing and recruitment; (10) the continuity of the group; (11) supporting yourself as a leader; (12) virtual support group meetings and (13) support group leader resources (see [table 1](#) for module content). The programme includes 11 filmed vignettes that demonstrate effective group facilitation techniques and ways to respond to problems that arise with the behaviours of specific group members or group interactions. In addition to the live modules, SPIN-SSLED participants receive a programme manual that summarises didactic material that is provided in the sessions.

Based on our previous experience with videoconferencing and consistent with previous trials of videoconference training, five group leaders were assigned to each training group to maximise effective interaction and participation.^{38–41} Training sessions were delivered using the GoToMeeting videoconferencing platform, a high-performance platform that has been used successfully for similar applications.^{42–44} In addition to the videoconference sessions, participants had access to a secure, monitored SPIN-SSLED online forum via the Slack communication tool to interact with other participants about programme content. The two training groups were held in the evening.

Feasibility outcomes

Outcomes related to process and resources were assessed throughout the duration of the feasibility trial, and leader

feedback was obtained on completion of the programme. The collected measures of feasibility included assessments of the (1) enrolment and consent procedure, (2) percentage of referred and eligible group leaders who consented to participate, (3) personnel requirements to assist participants with accessing the GoToMeeting videoconferencing platform for sessions and the online survey programme Qualtrics for online data collection pre-training and post-training, (4) technological performance of the videoconferencing system and (5) any challenges for study personnel.

Individual semistructured interviews were conducted with all participants via telephone on completion of the 13 modules using items based on the Patient Education Materials Assessment Tool for Audiovisual Materials (PEMAT)⁴⁵ and addressed topics related to usability, understandability, organisation and clarity of the SPIN-SSLED Programme, including its videoconference-based delivery. The PEMAT included a single rating of satisfaction from 0 (worst possible) to 10 (best possible).

SPIN-SSLED planned trial outcome measures

In the planned full-scale RCT, we will evaluate whether the SPIN-SSLED Programme is effective in improving SSc support group leaders' self-efficacy for carrying out their leader role (primary) and if the programme reduces burnout, improves emotional well-being and improves physical function among support group leaders (secondary). The SPIN-SSLED feasibility trial was not intended to test hypotheses and did not have adequate power for this, but we collected trial outcome measures at the time of consent to participate in the trial and following completion of the programme to evaluate the percentage of measures that were completed and to evaluate performance of the measures. Participants were emailed invitations to complete baseline and postintervention measures using the online survey programme Qualtrics.

Leader self-efficacy

The Scleroderma Support Group Leader Self-efficacy Scale (SSGLSS)⁴⁶ was developed by our research team, including the members of the SPIN Support Group Advisory Team, to measure support group leader self-efficacy for performing leader tasks. Initial items were obtained from the Group Leader Self-Efficacy Instrument, a 37-item self-report questionnaire that assesses self-efficacy for performing group leader skills.⁴⁷ The Group Leader Self-Efficacy Instrument is intended for use with group psychotherapy leaders, so many of its items are not relevant or appropriate for support group leaders. Items from this instrument were reviewed for relevancy, and relevant items were considered for inclusion, along with items from a questionnaire intended for leaders of cancer and multiple sclerosis support groups⁴⁸ and items that we generated from the results of a published study on the experiences of leaders of cancer support groups.⁴⁹ All items were then reviewed by members of our research team to remove items that were repetitive or not relevant



Table 1 Content of programme modules

Module title	Module description
1. The Support Group Leader's Role	This module discusses the benefits of being a support group leader, the expectations of what the role of leader involves (eg, facilitation of meetings and interactions but not giving medical advice) and tips for being an effective and supportive leader.
2. Starting a Support Group	This module discusses the purpose of a support group, what people with scleroderma hope to gain from support group, why some do not attend, establishing leadership (eg, one leader, co-leader), membership (eg, patients only, open to family and friends), logistics of starting a group (eg, time, place and meeting duration).
3. Structuring Support Group Meetings	This module discusses formatting group meeting and how to successfully integrate both educational activities with emotional and practical support for members, setting up a meeting agenda.
4. Scleroderma 101	This module shows a filmed conference by a physician specialised in scleroderma who explains the different types of scleroderma, symptoms, causes, treatments and alternative approaches. The module also includes tips to evaluate credibility of information sources on the Internet.
5. Successful Support Group Culture	This module discusses the importance of establishing expectations and guidelines for the support group with members, the importance of confidentiality, how to create and maintain positive and productive support group culture using (1) encouraging statements, (2) open-ended questions, (3) body language, (4) linking similar experiences between members and (5) summarising discussions. This module uses video vignettes to illustrate these techniques.
6. Managing Group Dynamics Part I	This module discusses managing difficult support group dynamics such as members who are 'quick fixers', overly talkative, how to maintain a positive group environment, conflict management and resolution for minor and larger issues. This module uses video vignettes to illustrate these techniques.
7. Managing Group Dynamics Part II	This module discussed how to identify and respond to members who are overly shy or who are chronically negative, and what to do when members bring unsubstantiated, inaccurate or potentially misleading medical information to the group. This module uses video vignettes to illustrate these techniques.
8. Grief and Crisis in Scleroderma	The module discusses different situations that may bring grief to members, stages of grief, supporting group members in grief, dealing with medical crises or death of group members.
9. Advertising and Recruiting for the Support Group	This module discusses how to advertise and promote a support group, how to recruit new members for support groups on ongoing basis, advertising through patient organisations and strategies to retain members.
10. The Continuity of the Group	This module discusses the importance of understanding and overcoming reluctance in seeking feedback, the importance of feedback in the support group experience, how to obtain and respond to feedback, how to identify reasons why members may stop attending meetings and strategies to help maintain membership, how to keep members engaged and move your support group forward by making changes.
11. Supporting Yourself as a Leader	This module discusses understanding what leader burnout is, understanding why it can happen and what the warning signs are, understanding the best way to address burnout including identifying methods of coping, understanding at what point it may be best for a leader to step down from his or her role, strategies to prevent experiencing leader burnout.
12. Remote support groups	This module discusses the benefits of an online support group, finding the right technology, scheduling and programming, advertising and reaching your target audience, tips for successful online meetings.
13. Resources	This module discusses strategies on how to obtain information and resources, how to communicate information effectively to members, the responsibilities of the national organisation, meeting the resource needs of the group, access to the SPIN-SSLED Online Resource Centre to download scleroderma talks, educational videos and research articles for leaders and members of support groups.

for SSc and to generate new items to reflect important SSc-specific content based on their own experiences or on qualitative interviews that we conducted with SSc support group leaders (n=10). Items were then reviewed iteratively by all members of the research team until a consensus was reached on included items. The resulting 32-item scale is scored on a 6-point Likert scale from 1 (strongly disagree) to 6 (strongly agree) with possible total scores from 32 to 192 and higher scores indicating greater self-efficacy. The SSGLSS was validated in two samples of SSc support group leaders (n=102, n=55) and found to have good internal consistency (Cronbach's alpha 0.96 and 0.95) and hypothesis-consistent convergent validity with the Oldenburg Burnout Inventory (OLBI). A strength of using the SSGLSS as the primary outcome measure is that

both the intervention and the SSGLSS were designed to reflect training needs of SSc support group leaders, and the items of the SSGLSS all reflect material covered in the programme.

Burnout

Leader burnout was assessed with the OLBI,^{50–52} which is a 16-item measure that assesses exhaustion and disengagement due to burnout. The OLBI was initially designed for work-related burnout, but has been adapted for numerous settings and in multiple countries and languages.⁵² Our research team revised the wording of each OLBI item to reflect the support group environment rather than a work environment (eg, 'I find my work to be a positive challenge' was revised to 'I find my role as a support

Table 2 Participant characteristics

Characteristics	No of participants=10
Female sex, n (%)	10 (100)
Age in years, mean (SD)	57.7 (11.1)
Country, n (%)	
Canada*	6 (60)
USA†	4 (40)
Race/ethnicity,‡ n (%)	
White	10 (100)
Aboriginal	1 (10)
Relationship status, n (%)	
Married or living as married	8 (80)
Separated or divorced	2 (20)
Education in years, mean (SD)	17.5 (2.7)
Occupational status, n (%)	
Homemaker	1 (10)
Part-time or full-time employment†	2 (20)
Disability	3 (30)
Retired	4 (40)
SSc diagnosis, n (%)	
Limited SSc	4 (40)
Diffuse SSc	5 (50)
Not diagnosed with SSc	1 (10)
Years since SSc diagnosis, mean (SD)	10.9 (7.4)
Current leader of SSc support group	10 (100)
Years as a SSc support group leader, mean (SD)	3.6 (3.7)

*Participants from British Columbia, Manitoba, Ontario, Quebec (2) and Saskatchewan.

†Participants from California (3) and Florida.

‡Participants could select more than one race/ethnicity. SSc, systemic sclerosis.

group leader to be a positive challenge'). The OLBI has a two-factor structure (exhaustion and disengagement) with good measurement properties.^{50–52} Items are scored on a 4-point scale; higher scores indicate higher levels of exhaustion and disengagement. Internal consistency reliability (Cronbach's alpha) in patients with SSc was 0.84 for exhaustion and 0.80 for disengagement.⁴⁶

Emotional distress

The Patient Health Questionnaire-8 (PHQ-8) was used to assess emotional distress.⁵³ The PHQ-8 items measure depressive symptoms over the last 2 weeks on a 4-point scale, ranging from 0 (not at all) to 3 (nearly every day) with higher scores indicating more depressive symptoms. The PHQ-8 performs equivalently to the PHQ-9,⁵³ which has been shown to be a valid measure of depressive symptoms in patients with SSc.⁵⁴

Physical function

Physical function was measured using the Physical Function subscale of the 29-item Patient-Reported Outcomes Measurement Information System (PROMIS-29) V.2.0. The PROMIS-29 measures eight domains of health status with four items for each of seven domains (physical function, anxiety, depression, fatigue, sleep disturbance, social roles and activities, pain interference) plus a single item for pain intensity. Items are scored on a 5-point scale (range 1–5), with different response options for different domains, and the single pain intensity item is measured on an 11-point rating scale. Higher scores represent more of the domain being measured; that is, better physical function. Total raw scores are obtained by summing item scores for each domain. The PROMIS-29 V.2.0 has been validated in SSc.⁵⁵

Participant satisfaction

Satisfaction with the SPIN-SSLED Programme was evaluated with the Client Satisfaction Questionnaire-8 (CSQ-8),⁵⁶ a standardised survey that is used to assess satisfaction with health services. Items are scored on a Likert scale from 1 (low satisfaction) to 4 (high satisfaction) with total scores ranging from 8 to 32. The CSQ-8 has been widely validated across a range of populations and health services programme.⁵⁷

Adverse events

Following each session, we emailed participants and requested that they report any concerns that they had about the sessions or their experience in the sessions.

Sample size

Guidance on appropriate sample size for feasibility trials varies substantially in the published literature. For the purposes of establishing feasibility of delivery of the SPIN-SSLED Programme, we determined that conducting two different training groups would allow us to evaluate intervention content and delivery aspects, and, thus, we sought to recruit a total of 10 participants to conduct two training groups.

Data analysis

Feasibility outcomes included leader eligibility and recruitment, leader enrolment and technological performance of the videoconferencing system. Qualitative information via interviews and weekly reports by participants was collected, and all suggestions for changes to the programme or trial methods that could be implemented prior to beginning a full-scale trial were recorded. Descriptive statistics were used to provide means and SD for SPIN-SSLED Programme outcome measures. Since the purpose of this feasibility trial was to evaluate feasibility and identify any modifications to the intervention or trial plan, the trial was not designed or powered to test hypotheses about outcomes. Thus, consistent with best practices,³³ hypothesis tests were not conducted, but effect sizes for pre-post differences are shown. Data were

Table 3 Summary of responses to the Patient Education Materials Assessment Tool for Audiovisual Materials (PEMAT) interviews

PEMAT item	Summary of responses
Are you currently a leader of a support group, a co-leader or planning on becoming a support group leader?	4 co-leaders, 5 leaders, 1 plan to become leader
PROCESS	
Did you find that the weekly frequency of the training sessions was adequate?	10 yes
Did you find the length of each training session appropriate?	10 yes; 3 added that grief module was not long enough, 1 said grief module was too long
Was it difficult to find the motivation to attend the training session every week?	9 no; 1 sometimes due to scheduling
PURPOSE	
Did you understand the objectives of the SPIN-SSLED training programme modules?	10 yes
Did you find the information provided in the SPIN-SSLED training programme relevant?	10 yes
WORDS AND LANGUAGE	
Did you find that the content of the programme manual was clear, concise and easy to follow?	10 yes
Did you find that the content delivered in the training sessions and the discussion about the content was easy to understand and useful?	10 yes
CONTENT AND ORGANISATION	
Did you find that the content of the SPIN-SSLED modules was presented logically and well-organised?	10 yes
Did you find that the order of the modules was logical and that linkages were clear?	9 yes; 1 said that order was 'staggered'
Did you find the discussion among other participants helpful?	10 yes; 1 mentioned that sometimes she had questions and challenges specific to herself that she did not have time to get addressed
VIDEO VIGNETTES	
Did the fact that the video vignette scenarios were performed by scleroderma patients make the programme more relatable?	9 yes; 1 had difficulty hearing the videos (participant with hearing impairment)
Were you able to clearly understand the people speaking in the videos?	9 yes; 2 no; 1 indicated there were small things they did not quite hear (participant with hearing impairment)
Did you develop an understanding of the challenges that could arise in a support group from watching the videos and obtain useful information or strategies to address them?	8 yes, 2 no (not hearing properly; 1 participant with hearing impairment)
TECHNOLOGY	
Did you use a computer, phone, tablet or all these devices to access the SPIN-SSLED Training Programme?	7 used computer, 4 used phone, 1 used tablet (>10 due to >1 method for some participants)
Did the initial invitation email provide you with the information you needed to understand how to log in for the training session?	10 yes
Did you experience any technical difficulties while using GoToMeeting?	8 no; 2 minor
Did you experience any technical difficulties while using Slack chatroom?	7 no; 3 did not use it. Use overall was minimal
Did you use the guide we provided to use GoToMeeting and the Slack chatroom?	5 yes; 5 no
OVERALL APPRECIATION	
Can you please tell us about your experience with the SPIN-SSLED Training Programme, including things that you liked about the programme and things that could be improved?	<p><i>Positive aspects of programme:</i> informative, organised, videos, 'Supporting Yourself as a Leader' and 'Grief and Crisis' modules were identified as very important.</p> <p><i>Positive aspects of SPIN-SSLED programme leader:</i> clear, conscientious, answered questions thoroughly, gave opportunities for feedback, was available between sessions.</p> <p><i>To improve:</i> discuss financial support for support group expenses, expand grief module over two sessions, do sessions in winter instead of summer so that individuals with scleroderma can only enjoy the outdoors in summer, adding more videos.</p>

Continued

Table 3 Continued

PEMAT item	Summary of responses
What grade (on a 0–10 scale, 0 being the worst and 10 being the best possible score) would you give the programme? 0 (worst) to 10 (best).	5 rated 10, 1 rated 9, 1 rated 9.5, 2 rated 8, 1 rated '9 or 10'. Mean score: 9.4/10
Would you recommend this programme to someone with scleroderma?	10 yes
Is there anything you want to give us feedback about that was not included in this interview?	The remote support group meetings module was less important to one participant. Suggestion to make recordings of support group discussions available online.

analysed using the statistics software programme, IBM SPSS.

Patient and public involvement

The SPIN Support Group Advisory Team has been involved in all stages of SPIN's research on support groups in SSc, including preliminary research on support groups in SSc, the development of the SPIN-SSLED programme and the design and implementation of the feasibility trial. Members of the team initially participated in the design of the Scleroderma Support Group Survey, which informed developing of the programme by collecting information on the experiences and training needs of SSc support group leaders, priorities of SSc support group members and reasons why people do not attend SSc support groups.^{17–20} Team members participated in the development of the SSGLSS,⁴⁶ which was administered in the feasibility trial and will be the primary outcome for the planned full-scale trial. Team members provided input into the development of the SPIN-SSLED Programme and its modules, filmed the vignettes used in the programme and were involved in decisions related to the conduct of the feasibility trial.

RESULTS

Participant characteristics

The trial was conducted between April and July 2018. Scleroderma Canada and the Scleroderma Foundation each provided our team with names of six potential participants. All agreed to participate in the programme. We initially enrolled 10 participants, but one was hospitalised prior to initiating the programme. Thus, prior to starting the trial, we added one participant who had been wait-listed.

All 10 participants were female. The mean age was 58 years (SD=11 years). There were six participants from Canada and four from the USA. All 10 described themselves as White, and one also described herself as Aboriginal. Of the 10 participants, 9 were people with SSc. Mean years since diagnosis among those with SSc was 11 years. Participant characteristics are shown in table 2.

Feasibility outcomes

All 10 participants completed all baseline and post-trial measures, including the PEMAT interview. Participant attendance at the weekly sessions was high (95%; 123 of

130 sessions). No sessions were missed or delayed due to technological difficulties, and time for technological support from our team was between 1 and 2 hours for the entire programme. Per the PEMAT interviews and per our observations, the GoToMeeting system worked fluidly and supported the training groups well.

A summary of responses to the PEMAT interviews is shown in table 3. As can be seen in the table, there were relatively minor suggestions for improving the programme. Overall, feedback was extremely positive. The overall mean grade given by participants for the SPIN-SSLED Programme was 9.4/10. No concerns related to adverse events were reported.

SPIN-SSLED planned trial outcome measures

Table 4 shows the responses to each of the 32 items of the SSGLSS, which will be the primary outcome measure in the full-scale trial. Pre-training, the mean (SD) was 124.4 (22.0), which was similar to the scores of our two international samples from the SSGLSS validation study (n=102, mean SSGLSS=122.9 (21.7); n=55, mean SSGLSS=123.9 (19.4)). Post-training, the mean total score increased to 159.2 (17.1). The standardised mean difference effect size was 1.70, which is considered a large effect size.⁵⁸ SSGLSS items are scored on a 1–6 scale, and the average item score increase pre-post training was 1.1 points.

Table 5 shows results for health outcomes, including burnout (OLBI), emotional distress (PHQ-8) and physical function (PROMIS-29). For all of these outcomes, the standardised mean difference effect size of post-trial score improvement was between 0.38 and 0.45, which are typically considered small to moderate effect sizes.⁵⁸

As shown in table 6, the mean post-training score on the CSQ-8 was 30.6 (2.2). On a per item basis, the mean item score (possible range 1–4) was 3.8, reflecting a very high level of satisfaction with the experience of trainees with the SPIN-SSLED Programme.

DISCUSSION

Feasibility of delivering the SPIN-SSLED Programme in the context of a trial, participant satisfaction and programme content was evaluated in the SPIN-SSLED feasibility trial. Results informed revisions to the content of the programme and provided confidence that the

Table 4 Pre-intervention and post-intervention item and total scores for the Scleroderma Support Group Leader Self-efficacy Scale: possible item scores range from 1 to 6 with higher scores reflecting greater self-efficacy

Items	Pre-trial N	Pre-trial mean (SD)	Post-trial N	Post-trial mean (SD)	Standardised mean difference effect size (95% CI—total only)
1. Obtain financial or other resources needed to run the group	10	3.5 (1.7)	10	4.8 (1.2)	0.61
2. Promote the group to health professionals as an important resource for patients	10	4.5 (0.7)	10	5.4 (0.7)	1.80
3. Share responsibilities, including administrative and practical tasks, with a co-facilitator or other group members	10	4.8 (1.6)	10	5.5 (0.5)	0.52
4. Manage group members who are overly talkative or monopolise the discussion	10	3.9 (1.1)	10	4.8 (0.6)	1.13
5. Manage group members who assume the role of the 'know-it-all'	10	3.9 (1.1)	10	4.6 (0.8)	0.73
6. Support members of the group who are grieving	10	3.3 (1.5)	10	4.7 (0.8)	0.97
7. Help overly shy group members feel comfortable interacting with the group	10	3.9 (1.1)	10	4.9 (0.7)	1.14
8. Help group members cope with difficult events, such as the death of a member	10	3.0 (1.8)	10	4.7 (0.8)	0.90
9. Effectively recruit new members	10	3.5 (1.4)	10	4.9 (0.7)	1.18
10. Address the different needs of group members at varying stages of the disease	10	3.7 (1.6)	10	4.8 (0.6)	0.77
11. Manage conflicts and disagreements between group members	10	3.3 (1.4)	10	4.5 (0.7)	0.95
12. Help the group establish appropriate group rules, such as maintaining confidentiality	10	4.3 (1.1)	10	5.8 (0.4)	2.31
13. Effectively publicise the group	10	3.5 (1.1)	10	5.1 (0.7)	1.86
14. Intervene effectively when group rules are not being followed	10	4.0 (1.1)	10	5.0 (0.7)	1.30
15. Obtain the support I need to cope with the emotional demands of leading the group	10	2.9 (1.2)	10	4.9 (1.0)	1.65
16. Respond constructively to feedback from group members	10	4.6 (0.8)	10	5.3 (0.8)	1.01
17. Help group members relate to other members of a different age	10	4.1 (1.0)	10	5.1 (0.7)	1.30
18. Provide the structure needed for successful meetings	10	5 (0.9)	10	5.6 (0.5)	1.03
19. Keep the group meetings interesting and relevant to both new and returning members	10	4.2 (0.9)	10	5.2 (0.8)	1.35
20. Manage group members who oversimplify or minimise the concerns of other members	10	3.5 (1.4)	10	4.9 (0.9)	0.99
21. Facilitate the group meetings so that all members have an opportunity to speak	10	4.6 (0.8)	10	5.1 (0.9)	0.68
22. Help the group stay focused on topics that are relevant to members	10	4.4 (0.7)	10	5.3 (0.7)	1.88
23. Obtain feedback from members about the group	10	4.3 (0.8)	10	5.0 (0.8)	1.04
24. Organise and plan activities for group members, such as having guest speakers	10	4.5 (1.4)	10	5.5 (0.7)	0.86
25. Help members feel comfortable in the group and relate to one another	10	4.4 (0.8)	10	5.2 (0.6)	1.43
26. Obtain feedback from members about my leadership	10	3.6 (0.7)	10	4.8 (0.9)	1.79
27. Help group members relate to other members of a different cultural background	10	3.9 (1.2)	10	4.5 (1.0)	0.50
28. Communicate reasonable boundaries about my availability outside of the group	10	3.7 (1.3)	10	4.5 (1.0)	0.64
29. Talk to a group member about her or his behaviour if it is disruptive to the group	10	2.7 (1.3)	10	4.5 (0.9)	1.58
30. Ask a member to leave the group due to her or his disruptive behaviour	10	1.6 (1.0)	10	4.2 (1.1)	2.32

Continued

Table 4 Continued

Items	Pre-trial N	Pre-trial mean (SD)	Post-trial N	Post-trial mean (SD)	Standardised mean difference effect size (95% CI—total only)
31. Help group members relate to other members of a different gender	10	4.4 (1.1)	10	5.0 (0.8)	0.65
32. Recruit a co-facilitator or other group members to help me with leadership responsibilities	10	4.9 (1.3)	10	5.1 (0.9)	0.16
Total score (possible range 32 to 192)	10	124.4 (22.0)	10	159.2 (17.0)	1.70 (0.67 to 2.72)

programme can be effectively and efficiently delivered in a full-scale trial.

With respect to overall experience with the programme and programme content, participants reported that the content was clear and well-organised. Overall satisfaction with their experience in the SPIN-SSLED Programme was rated as 9.4 out of 10 on average. Participant satisfaction was similarly high when evaluated with the eight items of the CSQ. Several participants encouraged the research team to expand on the single module related to grief. Based on comments and follow-up discussions with participants, the programme was revised to include two modules on grief, including one module on grief and loss that leaders have experienced because they or somebody close to them has been diagnosed with SSc and a second module on providing support to group members who are struggling with grief and loss. In order to add a second module on grief and loss, the two modules on managing group dynamics were reduced to a single module. To facilitate this, rather than viewing all of the short video vignettes included in those modules as part of the training sessions, participants suggested that they could view the vignettes prior to the sessions and then suggest specific modules for review and discussion in the training sessions.

With respect to programme delivery, participants indicated that they were able to access the sessions via the GoToMeeting platform and did not experience any technical difficulties that interrupted their training sessions. One difficulty that was reported involved a participant with a hearing impairment who was unable to hear all of the vignette videos well. This information will help us to

assess for reasons why participants may need additional support in the full-scale trial and will allow us to make accommodations. In the full-scale trial, we will assess for hearing and any other impairments that might limit participation, and we will seek appropriate assistance to be able to provide adaptations to meet participant needs. From a management standpoint, total time for technical support due to access difficulties across the trial period for the two groups was between 1 and 2 hours. Participants attended 95% of sessions, and all 10 participants completed all baseline and post-trial outcome assessments.

There were limitations that should be considered in evaluating the results of the SPIN-SSLED feasibility trial. First, we were provided with a small list of potential participants from our patient organisation partners, and it is possible that these support group leaders were more motivated or otherwise more likely to participate and engage than the leaders who will participate in the full-scale trial. Second, we did not randomise participants to the intervention and to a wait-list control group as we will do in the planned full-scale trial. Third, we only conducted two training groups and only included a total of 10 participants. The reason for not using a control group and limiting the feasibility trial to two groups is that there is a finite number of English-speaking and French-speaking SSc support group leaders, and we wanted to be able to assess feasibility aspects but maximise the number of participants eligible for the full-scale trial. It is possible that the pre-selection of potential participants and the lack of the possibility of randomised assignment to a non-intervention group may have resulted in overestimation of the percentage of participants who will enrol in the

Table 5 Pre-intervention and post-intervention total scores for secondary outcome measures

Measure	Pre-trial N	Pre-trial mean (SD)	Post-trial N	Post-trial mean (SD)	Standardised mean difference effect size (95% CI)
Oldenburg Burnout Inventory (higher scores=greater burnout)	10	33.2 (4.6)	10	31.0 (4.9)	0.44 (−0.44 to 1.33)
Patient Health Questionnaire-8 (higher scores=greater symptoms of depression)	10	10.8 (2.7)	10	9.8 (2.4)	0.38 (−0.50 to 1.27)
PROMIS-29 Physical Function (higher raw scores=greater function)	10	17.1 (2.2)	10	18.2 (2.4)	0.45 (−0.42 to 1.36)

PROMIS-29, 29-item Patient-Reported Outcomes Measurement Information System.



Table 6 Post-intervention items, frequencies and total scores for the Client Satisfaction Questionnaire-8: item response options vary across items, but all scored 1–4

Items	1 point (dissatisfied) N (%)	2 points (mildly satisfied) N (%)	3 points (mostly satisfied) N (%)	4 points (quite satisfied) N (%)	Item mean (SD)
1. How would you rate the quality of the SPIN-SSLED training?	0 (0)	0 (0)	1 (10)	9 (90)	3.9 (0.3)
2. Did the SPIN-SSLED programme provide you the kind of training you wanted?	0 (0)	0 (0)	2 (20)	8 (80)	3.8 (0.4)
3. To what extent has the SPIN-SSLED training met your needs?	0 (0)	0 (0)	4 (40)	6 (60)	3.6 (0.5)
4. If a friend were in need of similar training, would you recommend the SPIN-SSLED programme to him/her?	0 (0)	0 (0)	2 (20)	8 (80)	3.8 (0.4)
5. How satisfied are you with the amount of training you received from the SPIN-SSLED programme?	0 (0)	0 (0)	2 (20)	8 (80)	3.8 (0.4)
6. Has the SPIN-SSLED training helped you to deal more effectively with your support group leader role?	0 (0)	0 (0)	0 (10)	10 (100)	4.0 (0.0)
7. In an overall, general sense, how satisfied are you with the SPIN-SSLED training?	0 (0)	0 (0)	1 (20)	9 (90)	3.9 (0.3)
8. If you were to seek help again, would you come back to the SPIN-SSLED training?	0 (0)	0 (0)	2 (20)	8 (80)	3.8 (0.4)
Total score (possible range 8 to 32)					30.6 (2.2)

SPIN-SSLED, Scleroderma Patient-centered Intervention Network—Scleroderma Support group Leader Education.

full-scale trial and the degree to which they will actively participate. Given the small number of French-speaking leaders available for the full-scale trial, we did not include a French group in the feasibility trial. However, all measures have been used successfully previously with French-speaking research participants. Finally, the trial only included leaders of SSc support groups, and this may limit generalisability to other patient populations, but it will be useful to inform our planned full-scale trial of the SPIN-SSLED Programme.

There are no existing training programme for SSc support group leaders, and a systematic review did not identify any training or education programme that have been demonstrated to be effective for support group leaders in any medical condition.³¹ The planned full-scale SPIN-SSLED trial, which was recently funded by the Canadian Institutes of Health Research, is scheduled to begin in 2019 (http://webapps.cihr-irsc.gc.ca/decisions/p/project_details.html?applId=388187&lang=en). It will be a pragmatic RCT that will test whether providing the SPIN-SSLED Programme to leaders of SSc support groups will improve outcomes compared with leaders assigned to a wait-list control. Pragmatic RCTs differ from explanatory or mechanistic trials, in that they are intended to test the effectiveness of adding an intervention to routine practice in order to inform practice and policy decisions rather than explain intervention mechanisms.^{59 60} SSc support group leaders who are enrolled will be randomly allocated to the training programme or a wait-list control, and those allocated to training will be

clustered in training groups where they will interact with each other. To account for clustering in the training arm, but not the control arm, we will use a partially nested RCT (PN-RCT) trial design.^{61–63} The PN-RCT design is a hybrid between a conventional RCT, in which individual participants are randomised, and a cluster RCT, in which pre-existing clusters (eg, primary care practices, classrooms) are randomised to intervention or control arms. In the PN-RCT design, analyses account for dependence within intervention arm clusters, but treat leaders assigned to the control arm individually, as in a conventional RCT. Participants will be existing support group leaders or new leaders referred by Scleroderma Canada (English) or Sclérodermie Québec (French), the Scleroderma Foundation (USA), Scleroderma and Raynaud's UK, the Scleroderma Association of New South Wales (Australia) and Scleroderma New Zealand.

In sum, the SPIN-SSLED feasibility trial ensured that trial methodology was feasibly implemented and that the online intervention was user-friendly and acceptable to participants. Participants provided suggestions for adjustments to content that will be implemented before undertaking a full-scale RCT of the SPIN-SSLED programme to assess effectiveness.

Author affiliations

¹Lady Davis Institute for Medical Research, Jewish General Hospital, Montreal, Québec, Canada

²Department of Psychiatry, McGill University, Montreal, Québec, Canada

³Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, Quebec, Canada

⁴Department of Medicine, McGill University, Montreal, Quebec, Canada

⁵Department of Psychology, McGill University, Montreal, Quebec, Canada

⁶Department of Educational and Counselling Psychology, McGill University, Montreal, Quebec, Canada

⁷Biomedical Ethics Unit, McGill University, Montreal, Quebec, Canada

⁸Scleroderma Foundation Michigan Chapter, Southfield, Michigan, USA

⁹Department of Psychology, San Diego State University, San Diego, California, USA

¹⁰San Diego Joint Doctoral Program in Clinical Psychology, San Diego State University/University of California, San Diego, California, USA

¹¹Department of Psychology, Université du Québec à Montréal, Montreal, Quebec, Canada

¹²Scleroderma Society of Ontario, Scleroderma Canada, Hamilton, Ontario, Canada

Collaborators SPIN-SSLED Patient Advisory Team Members: Kerri Connolly, Director of Programs and Services of the Scleroderma Foundation, Danvers, Massachusetts, USA; Stephen Elrod, Southern California Patient Group, Los Angeles, California, USA; Catherine Fortuné, Ottawa Scleroderma Support Group, Ottawa, Ontario, Canada; Karen Gottesman, Director of Pharma & Biotech Engagement Scleroderma Foundation, Los Angeles, California, USA; Karen Nielsen, Scleroderma Society of Ontario, Hamilton, Ontario, Canada; Geneviève Guillot, Sclérodermie Québec, Longueuil, Quebec, Canada; Ken Rozee, Scleroderma Society of Nova Scotia, Halifax, Nova Scotia, Canada; Amy Gietzen, Scleroderma Foundation, Danvers, Massachusetts, USA; Michelle Richard, President of Scleroderma Canada, Halifax, Nova Scotia, Canada; Nancy Stephens, Michigan Patient Group, Detroit, Michigan, USA.

Contributors BDT, VLM, GE-B, SP, MS, MH, RWP and members of the SPIN-SSLED Patient Advisory Team were responsible for the study conception and design. BDT, LD, MP, KA, M-EC, LT and SH were responsible for implementation of the trial or acquisition, analysis and interpretation of trial data. BDT drafted the manuscript. All authors provided a critical review and approved the final manuscript. BDT is the guarantor.

Funding Funding from the Scleroderma Society of Ontario and the Canadian Institutes of Health Research (PJT-156429) was used for development of the SPIN-SSLED Program and preliminary research. The trial was funded by the Canadian Initiative for Outcomes in Rheumatology cAre. BDT was supported by a Fonds de Recherche Québec—Santé (FRQS) researcher award outside of the present work. RWP holds the Albert Boehringer I Chair in Pharmacoeconomics. No sponsor or funder was involved in the study design; in the collection, analysis and interpretation of the data; in the writing of the report or in the decision to submit the paper for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The trial was approved by the ethics committee of the Centre intégré universitaire de santé et de services sociaux du Centre-Ouest-de-l'Île-de-Montréal.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Brett D Thombs <http://orcid.org/0000-0002-5644-8432>

REFERENCES

- Kralik D. The quest for ordinariness: transition experienced by midlife women living with chronic illness. *J Adv Nurs* 2002;39:146–54.
- Karasz A, Ouellette SC. Role strain and psychological well-being in women with systemic lupus erythematosus. *Women Health* 1995;23:41–57.
- Kole A, Faurisson F. The voice of 12,000 patients: experiences and expectations of rare disease patients on diagnosis and care in Europe, 2009. Available: http://www.eurordis.org/IMG/pdf/voice_12000_patients/EURORDISCARE_FULLBOOKr.pdf [Accessed 10 Feb 2019].

- European Organisation for Rare Diseases. Rare diseases: understanding this public health priority, 2005. Available: http://www.eurordis.org/IMG/pdf/princeps_document-EN.pdf [Accessed 10 Feb 2019].
- Cohen JS, Biesecker BB. Quality of life in rare genetic conditions: a systematic review of the literature. *Am J Med Genet A* 2010;152A:1136–56.
- Huyard C. What, if anything, is specific about having a rare disorder? patients' judgements on being ill and being rare. *Health Expect* 2009;12:361–70.
- Nettleton S, Watt I, O'Malley L, et al. Understanding the narratives of people who live with medically unexplained illness. *Patient Educ Couns* 2005;56:205–10.
- van Walsem MR, Howe EI, Iversen K, et al. Unmet needs for healthcare and social support services in patients with Huntington's disease: a cross-sectional population-based study. *Orphanet J Rare Dis* 2015;10:124.
- Anderson M, Elliott EJ, Zurynski YA. Australian families living with rare disease: experiences of diagnosis, health services use and needs for psychosocial support. *Orphanet J Rare Dis* 2013;8:22.
- Dwyer AA, Quinton R, Morin D, et al. Identifying the unmet health needs of patients with congenital hypogonadotropic hypogonadism using a web-based needs assessment: implications for online interventions and peer-to-peer support. *Orphanet J Rare Dis* 2014;9:83.
- Kasprian NA, Rutstein A, Sansom-Daly UM, et al. Through the looking glass: an exploratory study of the lived experiences and unmet needs of families affected by von Hippel-Lindau disease. *Eur J Hum Genet* 2015;23:34–40.
- Holtzclaw Williams P. Policy framework for rare disease health disparities. *Policy Polit Nurs Pract* 2011;12:114–8.
- Newman S, Steed L, Mulligan K. Self-Management interventions for chronic illness. *The Lancet* 2004;364:1523–37.
- Davison KP, Pennebaker JW, Dickerson SS. Who talks? the social psychology of illness support groups. *Am Psychol* 2000;55:205–17.
- Kwakkenbos L, Jewett LR, Baron M, et al. The scleroderma patient-centered intervention network (spin) cohort: protocol for a cohort multiple randomised controlled trial (cmRCT) design to support trials of psychosocial and rehabilitation interventions in a rare disease context. *BMJ Open* 2013;3:e003563.
- Reimann A, Bend J, Dembski B. [Patient-centred care in rare diseases. A patient organisations' perspective]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2007;50:1484–93.
- Gumuchian ST, Delisle VC, Kwakkenbos L, et al. Reasons for attending support groups and organizational preferences: the European scleroderma support group members survey. *Disabil Rehabil* 2019;41:974–82.
- Delisle VC, Gumuchian ST, Peláez S, et al. Reasons for non-participation in scleroderma support groups. *Clin Exp Rheumatol* 2016;100:56–62.
- Gumuchian ST, Delisle VC, Peláez S, et al. Reasons for not participating in scleroderma patient support groups: a cross-sectional study. *Arthritis Care Res* 2018;70:275–83.
- Delisle VC, Gumuchian ST, El-Baalbaki G, et al. Training and support needs of scleroderma support group facilitators: the North American scleroderma support group facilitators survey. *Disabil Rehabil* 2018;25:1–6.
- Aymé S, Kole A, Groft S. Empowerment of patients: lessons from the rare diseases community. *The Lancet* 2008;371:2048–51.
- Barg FK, Gullatte MM. Cancer support groups: meeting the needs of African Americans with cancer. *Semin Oncol Nurs* 2001;17:171–8.
- Seibold J. Scleroderma. In: Harris ED, Budd RC, Firestein GS, eds. *Kelly's Textbook of Rheumatology*. Philadelphia: Elsevier, 2005: 1279–308.
- Mayes M. Systemic sclerosis: clinical features. In: Klippel JH, Stone JH, Crafford LJ, eds. *Primer on the rheumatic diseases*. 13th edn. New York: Springer and Arthritis Foundation, 2008: 343–50.
- Wigley FM, Hammers LK. Clinical features of systemic sclerosis. In: Hochberg MC, Silman AJ, Smolen JS, eds. *Rheumatology*. 3rd edn. Philadelphia: Mosby, 2003: 1463–80.
- Kwakkenbos L, Delisle VC, Fox RS, et al. Psychosocial aspects of scleroderma. *Rheum Dis Clin North Am* 2015;41:519–28.
- Scleroderma Canada. Find a support group. Available: www.scleroderma.ca/Support/Find-A-Support-Group.php [Accessed 10 Feb 2019].
- Scleroderma Foundation. Support groups. Available: www.scleroderma.org/site/PageServer?pagenam=patients_supportgroups#.Vbec7RZvdFw [Accessed 10 Feb 2019].

- 29 Scleroderma & Raynaud's UK. Find support. Available: www.sruk.co.uk/find-support/ [Accessed 10 Feb 2019].
- 30 Scleroderma Association of New South Wales. Regional support groups. Available: www.sclerodermansw.org/regional-support-groups [Accessed 10 Feb 2019].
- 31 Delisle VC, Gumuchian ST, Kloda LA, *et al.* Effect of support group peer facilitator training programmes on peer facilitator and support group member outcomes: a systematic review. *BMJ Open* 2016;6:e013325.
- 32 Zordan R, Butow P, Kirsten L, *et al.* Supporting the supporters: a randomized controlled trial of interventions to assist the leaders of cancer support groups. *J Community Psychol* 2015;43:261–77.
- 33 Eldridge SM, Chan CL, Campbell MJ, *et al.* Consort 2010 statement: extension to randomised pilot and feasibility trials. *BMJ* 2016;355:i5239.
- 34 Boud D, Feletti G. *The challenge of problem-based learning*. 2nd edn. London: Kogan Page, 1997.
- 35 Duch BJ, Groh SE, Allen DE. Why problem-based learning? A case study of institutional change in undergraduate education. In: Duch BJ, Groh SE, Allen DE, eds. *The power or problem-based learning*. Sterling, VA: Stylus, 2001: 3–11.
- 36 Hmelo-Silver CE. Problem-Based learning: what and how do students learn? *Educ Psychol Rev* 2004;16:235–66.
- 37 Torp L, Sage S. *Problems as possibilities: problem-based learning for K-16 education*. 2nd edn. Alexandria, VA: Association for Supervision and Curriculum Development, 2002.
- 38 Hay-Hansson AW, Eldevik S. Training discrete trials teaching skills using videoconference. *Res Autism Spectr Disord* 2013;7:1300–9.
- 39 Seibert DC, Guthrie JT, Adamo G. Improving learning outcomes: integration of standardized patients & telemedicine technology. *Nurs Educ Perspect* 2004;25:232–7.
- 40 Xavier K, Shepherd L, Goldstein D. Clinical supervision and education via videoconference: a feasibility project. *J Telemed Telecare* 2007;13:206–9.
- 41 Marziali E, Donahue P. Caring for others: Internet video-conferencing group intervention for family caregivers of older adults with neurodegenerative disease. *Gerontologist* 2006;46:398–403.
- 42 Fisher WW, Luczynski KC, Hood SA, *et al.* Preliminary findings of a randomized clinical trial of a virtual training program for applied behavior analysis technicians. *Res Autism Spectr Disord* 2014;8:1044–54.
- 43 Hommel KA, Gray WN, Hente E, *et al.* The telehealth enhancement of adherence to medication (team) in pediatric IBD trial: design and methodology. *Contemp Clin Trials* 2015;43:105–13.
- 44 Williams LK, McCarthy MC, Burke K, *et al.* Addressing behavioral impacts of childhood leukemia: a feasibility pilot randomized controlled trial of a group videoconferencing parenting intervention. *Eur J Oncol Nurs* 2016;24:61–9.
- 45 Shoemaker SJ, Wolf MS, Brach C. Development of the patient education materials assessment tool (PEMAT): a new measure of understandability and actionability for print and audiovisual patient information. *Patient Educ Couns* 2014;96:395–403.
- 46 Pal NE, Gumuchian ST, Delisle VC, *et al.* Development and preliminary validation of the scleroderma support group leader self-efficacy scale. *J Scleroderma Relat Disord* 2018;3:106–11.
- 47 Page BJ, Pietrzak DR, Lewis TF. Development of the group leader self-efficacy instrument. *JSGW* 2001;26:168–84.
- 48 Zordan RD, Juraskova I, Butow PN, *et al.* Exploring the impact of training on the experience of Australian support group leaders: current practices and implications for research. *Health Expect* 2010;13:427–40.
- 49 Butow PN, Ussher J, Kirsten L, *et al.* Sustaining leaders of cancer support groups: the role, needs, and difficulties of leaders. *Soc Work Health Care* 2005;42:39–55.
- 50 Demerouti E, Bakker AB, Vardakou I, *et al.* The convergent validity of two burnout instruments: a multitrait-multimethod analysis. *Eur J Psychol Assess* 2002;18:296–307.
- 51 Halbesleben JRB, Demerouti E. The construct validity of an alternative measure of burnout: investigating the English translation of the Oldenburg burnout inventory. *Work Stress* 2005;19:208–20.
- 52 Reis D, Xanthopoulos D, Tsaousis I. Measuring job and academic burnout with the Oldenburg burnout inventory (OLBI): factorial invariance across samples and countries. *Burn Res* 2015;2:8–18.
- 53 Kroenke K, Strine TW, Spitzer RL, *et al.* The PHQ-8 as a measure of current depression in the general population. *J Affect Disord* 2009;114:163–73.
- 54 Milette K, Hudson M, Baron M, *et al.* Comparison of the PHQ-9 and CES-D depression scales in systemic sclerosis: internal consistency reliability, convergent validity and clinical correlates. *Rheumatology* 2010;49:789–96.
- 55 Kwakkenbos L, Thombs BD, Khanna D, *et al.* Performance of the patient-reported outcomes measurement information System-29 in scleroderma: a scleroderma patient-centered intervention network cohort study. *Rheumatology* 2017;56:1302–11.
- 56 Larsen DL, Attkisson CC, Hargreaves WA, *et al.* Assessment of client/patient satisfaction: development of a general scale. *Eval Program Plann* 1979;2:197–207.
- 57 Kelly PJ, Kyngdon F, Ingram I, *et al.* The client satisfaction Questionnaire-8: psychometric properties in a cross-sectional survey of people attending residential substance abuse treatment. *Drug Alcohol Rev* 2018;37:79–86.
- 58 Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd edn. Hillsdale, NJ: Lawrence Erlbaum, 1988.
- 59 Zwarenstein M, Treweek S, Gagnier JJ, *et al.* Improving the reporting of pragmatic trials: an extension of the CONSORT statement. *BMJ* 2008;337:a2390.
- 60 Roland M, Torgerson DJ. Understanding controlled trials: what are pragmatic trials? *BMJ* 1998;316:285.
- 61 Lohr S, Schochet PZ, Sanders E. *Partially nested randomised controlled trials in educational research: a guide to design and analysis*. NCER 2014–2000. Washington, DC: National Center for Education Research, 2014.
- 62 Bauer DJ, Sterba SK, Hallfors DD. Evaluating group-based interventions when control participants are ungrouped. *Multivariate Behav Res* 2008;43:210–36.
- 63 Schweig JD, Pane JF. Intention-To-Treat analysis in partially nested randomized controlled trials with real-world complexity. *IJRES* 2016;39:268–86.