



# BMJ Open Severe varicose veins and the risk of mortality: a nationwide population-based cohort study

Nan-Chun Wu,<sup>1,2</sup> Zhih-Cherng Chen,<sup>3,4</sup> I-Jung Feng,<sup>5</sup> Chung-Han Ho <sup>5</sup>, Chun-Yen Chiang,<sup>3,4</sup> Jhi-Joung Wang,<sup>6</sup> Wei-Ting Chang <sup>3,7,8</sup>

**To cite:** Wu N-C, Chen Z-C, Feng I-J, *et al.* Severe varicose veins and the risk of mortality: a nationwide population-based cohort study. *BMJ Open* 2020;**10**:e034245. doi:10.1136/bmjopen-2019-034245

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-034245>).

N-CW and Z-CC contributed equally.

Received 12 September 2019  
Revised 28 February 2020  
Accepted 23 April 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Wei-Ting Chang;  
[cmcvecho2@gmail.com](mailto:cmcvecho2@gmail.com)

## ABSTRACT

**Objective** Varicose veins (VVs) are common and although considered benign may cause morbidity. However, the association between VV severity and cardiovascular and mortality risks remains unknown. The aim of this study was to investigate the factors associated with overall mortality in patients with VV.

**Methods** A total of 4644 patients with newly diagnosed VV between 1999 and 2013 were identified from Taiwan's National Health Insurance Database. VV severity was classified from grade 1 to 3 according to the presentation of ulcers or inflammation. Moreover, 9497, 2541 and 5722 age-matched, sex-matched and chronic cardiovascular risk factor-matched controls, as assessed based on propensity score, were separately selected for three grading VV groups. Enrolled patients were analysed using conditional Cox proportional hazards regression analysis to estimate risk of mortality and major adverse cardiovascular events (MACEs) in the VV and control groups.

**Results** Most patients with VV were free from systemic disease. However, compared with matched controls, patients with VV showed a 1.37 times increased risk of mortality (95% CI 1.19 to 1.57;  $p<0.0001$ ). Compared with matched controls, older (age  $\geq 65$  years) (adjusted HR: 1.38; 95% CI 1.17 to 1.62;  $p=0.0001$ ) and male patients with VV (adjusted HR 1.41; 95% CI 1.18 to 1.68;  $p=0.0001$ ) showed increased risk of mortality. Furthermore, compared with controls, patients with VV showed 2.05 times greater risk of MACE. Compared with matched controls, population at grade 3 increased 1.83 times risk of mortality and 2.04 to 38.42 times risk of heart failure, acute coronary syndrome, ischaemic stroke and venous thromboembolism.

**Conclusions** This nationwide cohort study demonstrated that patients with VV are at a risk of cardiovascular events and mortality. Our findings suggest that presence of VV warrants close attention in terms of prognosis and treatment.

## INTRODUCTION

Varicose veins (VVs) can be considered a common disease with prevalence ranging from 2% to 56% in the adult population.<sup>1</sup> Following clinical examination, VV diagnosis is primarily based on the presence of enlarged and twisted veins in the lower extremities.<sup>1 2</sup>

## Strengths and limitations of this study

- The strengths of this study are its population-based design with a large sample size including study and control cohorts.
- All insurance claims were reviewed by medical reimbursement specialists.
- However, some risk factors of varicose vein including smoking habits, lack of movement, overweight and glycated haemoglobin levels were not available in this database.
- Our novel findings indicated that patients at severe grades of varicose vein had higher risks of mortality and major adverse cardiovascular events.
- The presence of varicose vein should catch more awareness of potential coexisting risks of mortality and cardiovascular events.

Among people with VV, 1%–4% of individuals show higher severity grades (Clinical–Etiological–Anatomical–Pathophysiological (CEAP) classification, 5–6).<sup>1 2</sup> Although VVs lead to leg swelling, venous eczema and ulceration in some cases, they are regarded as a benign disease.<sup>3 4</sup> Moreover, the association between the severity of VV and risk of future adverse events remains unknown. In fact, the majority of the previous studies have focused on the importance of superficial venous thrombosis or deep vein thrombosis (DVT).<sup>5</sup> In a 30-year cohort study, mortality risk among patients with DVT and pulmonary embolism (PE) was markedly higher than that in age-matched and sex-matched patients, particularly within the first 30 days.<sup>6</sup> Similarly, another population-based case–control study demonstrated that having VV was a risk factor for venous thromboembolism, although the association of VV severity with survival and cardiovascular events remains unknown.<sup>7</sup> In addition, although age, family history and female sex are the known risk factors for VV, the effects of underlying diseases or sex on outcomes of VV remain unclear.<sup>1</sup> We hypothesised that presence of VV can be used as a

marker for cardiovascular risk. Therefore, the aim of this study was to investigate the association of VV with survival and cardiovascular outcomes.

## METHODS

### Data source

Taiwan launched a single-payer National Health Insurance (NHI) programme on 1 March 1995. This database contains details of almost every Taiwanese resident (coverage rate >98% in 2009), making it one of the world's largest and most complete population-based sources. The data used in this study were retrieved from the Longitudinal Health Insurance Database 2000 (LHID2000)—a subset of the NHI database containing all claims data from 1996 to 2013, covering 1 million beneficiaries randomly selected in 2000. At that time, there were no significant differences in age, sex and healthcare costs between patients with VV and matched controls. LHID2000 provided encrypted patient identification numbers; sex; date of birth; admission and discharge dates; International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes of diagnoses and procedures; prescription details; registry data in the Catastrophic Illness Patient Database; and costs covered and paid for by NHI. Details of the National Health Insurance Research Database (NHIRD) are described in previous studies.<sup>8,9</sup> Moreover, the accuracy of major disease diagnoses in the NHIRD, including stroke and acute coronary syndrome (ACS), has been validated.<sup>9</sup>

### Patient and public involvement

No patient involved.

### Study design

This nationwide population-based, retrospective cohort study was conducted to investigate the association between VV and subsequent mortality. Patients with at least three claims for outpatient VV diagnosis in 1 year or with one claim for inpatient VV diagnosis (ICD-9-CM codes 454, 454.0, 454.1, 454.2, 454.8 and 454.9) were considered as VV cases. Patients with a first-time diagnosis of VV from January 1999 to December 2012 were included in the cohort. Codes for VV were considered reliable for diagnosis based on clinical symptoms. The date of the first-time VV diagnosis was considered the index date in this study. To ensure accurate VV diagnosis, and to avoid potentially confounding effects, patients with DVT (ICD-9-CM codes 453.40, 459.1, 671.4, 671.3, 451.83, 459.3, 453.4, and 451.11) or PE (ICD-9-CM codes 415.1, 415.11, 673, 673.2, and 673.8) in an ambulatory setting before the index date were excluded. In addition, VV severity was categorised as grade 1 uncomplicated (ICD-9-CM code 454.9), grade 2 with ulcer (ICD-9-CM code 454.0) or inflammation (ICD-9-CM code 454.1) and grade 3 with both ulcer and inflammation (ICD-9-CM code 454.2).

Three control cohorts (n1=9497, n2=2541 and n3=5722; four control subjects for every enrolled patient

with VV), not diagnosed with VV from 1996 to 2013, were selected for three VV grade groups separately. To eliminate potential selection bias, the controls were matched using propensity score (PS) method at a 4:1 ratio for baseline characteristics of age, sex and chronic cardiovascular risk factors, including hypertension (ICD-9-CM codes 401–405, A260, A269, 4372), diabetes (ICD-9-CM codes 250, A181, A189, A229, A239, 3572, and 3620), hyperlipidaemia (ICD-9-CM code 272), and coronary artery disease (CAD; ICD-9-CM codes 410–414). The PS for identified VV cases and controls were estimated using the fitting logistic regression model. Based on greedy algorithm matching, eight control subjects (the nearest neighbour matching of VV) were selected as matched controls.<sup>10</sup> If a case failed to be assigned to the four matched controls, it was dropped from the set of matches. In addition, since the primary VV treatment was covered by insurance, it prevented VV from overdiagnosis. The matched controls were assigned the same index date as that of the corresponding VV patient.

### Outcomes

The primary outcome was mortality, and the secondary outcome was major adverse cardiovascular events (MACEs), including acute (ACS, ICD-9-CM codes 410, 410.7, 411.1, 411.81 and 414.8), congestive heart failure (CHF, ICD-9-CM codes 428, 428.0, 428.1, 428.2 and 428.9), ischaemic stroke (ICD-9-CM code 436), DVT and PE. Mortality was identified using the 'in-hospital death' or 'discharge under critical condition' codes at discharge. Enrolment in the NHI programme is mandatory for all people in Taiwan, and registration must be withdrawn within 30 days after death. Patients with the above-mentioned mortality-related codes and those withdrawn from the NHI programme within 30 days after discharge from the last hospitalisation were presumed to have died. All subjects were followed up from the index date to death (lost to follow-up) or until 31 December 2013, whichever was earlier.

### Validation of the accuracy of VV diagnosis and CEAP grading

To validate the accuracy of the VV diagnosis, we reviewed the charts of all patients (inpatients and outpatients) using ICD-9-CM diagnosis codes for VV who visited Chi-Mei Medical Center (Tainan, Taiwan) from 2010 to 2015. Our aim was to determine the accuracy and consistency of code usage. A vascular specialist reviewed patient discharge and clinical records. In addition to examining the accuracy of VV diagnosis, the reviewer compared CEAP stages with our ICD-9-CM-derived grades in inpatients. Subsequently, we further investigated the sensitivity, specificity and predictive value of the ICD codes for clinical diagnosis, as well as the applicability of our VV grading system. In particular, as ICD-9-CM coding and VV descriptions are associated with insurance payment, the accuracy of VV diagnosis and the reliability of VV severity grading increased. The consistency between CEAP and grading stages were evaluated by kappa score, whose

value between 0.8 and 1.0 was considered as an almost perfect agreement.

### Statistical analyses

Continuous and categorical baseline characteristics between the case and control groups were separately compared by standardised mean difference (SMD), an assessment approach for evaluating the balance between variables after PS matching. SMD greater than 0.1 is considered to denote a meaningful imbalance in variables.

Conditional Cox proportional hazards regression analysis was used to estimate the risk of mortality and MACE in the VV and control groups. Adjusted HRs were estimated by adjusting for chronic obstructive pulmonary disease (ICD-9-CM codes 490–496), cancer (ICD-9-CM codes 140–208), atrial fibrillation (ICD-9-CM codes 427.31), heart failure (ICD-9-CM codes 428), ischaemic heart disease (ICD-9-CM codes 410–414), chronic renal insufficiency (ICD-9-CM codes 403, 404, 582, 585–588). Moreover, the investigation was extended to stratified subgroup analysis. HRs between the VV and control groups were separately estimated in subgroups of population aged <65 years or ≥65 years; males or females; and subgroups with or without a diagnosis of hypertension, diabetes, hyperlipidaemia or CAD. The Kaplan-Meier method was used to separately estimate the 3-year 6-year and 9-year survival rates in the control and VV groups. Kaplan-Meier curves of mortality and MACE were plotted for controls and patients with three grades of VV severity. Differences in survival curves between the control and VV groups were examined using the log-rank test.

With respect to mortality, CHF, ACS, ischaemic stroke and DVT +PE endpoints, the risks for VV with three separate severity grades were further estimated by comparison against each matched controls.

Finally, sensitivity analyses were conducted to determine the influence from subjects with pregnancy history (ICD-9-CM codes V22, V23.2, 761.5), peripheral artery disease (PAD, ICD-9-CM codes 440.0, 440.2, 440.3, 440.8, 440.9, 443, 444.0, 444.22, 444.8, 447.8 and 447.9) medical history and patients treated with operations (ICD-9-CM procedure code 3859, 3889 and NHIRD order code 69013, 69014, 69015, 69016, 69017, 69019, 69020, 69021) including ligation and stripping procedures after VV diagnosis.

A two-tailed  $p < 0.05$  was considered statistically significant. All analyses were performed using the SAS software, V.9.4 (SAS) and Stata software V.15.0 (StataCorp)

## RESULTS

### Characteristics of the study population

A total of 4644 patients with newly diagnosed VV were identified during January 1999 to December 2012. Among them, 2467, 668 and 1509 VVs were separately classified into 1, 2 and 3 severity grade. For each VV

group, age-matched, sex-matched and chronic disease-matched patients without VV were separately included for comparison. The covariates between VV and matched groups are well balanced after PS matching. All patients were tracked from the index dates until achieving the primary outcomes or the end of the study. The mean age of patients with VV was  $55.70 \pm 16.03$  years, the majority of the patients were female (61.33%), and most of them did not present with chronic diseases such as hypertension, diabetes, hyperlipidaemia and CAD (table 1). Significantly different distribution of age, sex and diabetes among three severity VV groups were displayed ( $p < 0.05$ ) (online supplementary table 1). Interestingly, more female patients (68.67%) were diagnosed with a lower severity (grade 1). Also, the baseline characteristics and comorbid medical disorders for three grading VV groups and three separately matched controls were listed in online supplementary table 2.

### Long-term mortality risk

Compared with matched controls, the outcomes of patients with VV were worse. The estimated survival at 3, 6 and 9 years were 97.6%, 95.6%, and 93.5%, respectively, in patients with VV compared with 98.5%, 97.1%, and 95.6%, respectively, in controls (figure 1A). A log-rank test revealed a significant difference in survival curves of patients with VV and controls ( $p < 0.0001$ ). The survival curves of controls and patients with different severities of VV are presented in figure 1B. Lower survival rates over time were observed in patients with highest VV severity (grades 3) but not in those with grade 1–2. Significant difference between survival curves between VV grading 3 and corresponding controls were revealed by log rank test ( $p < 0.0001$ ). However, no significant differences were found between survival curves of patients with VV severity grades (1–2) and corresponding controls (grade 1:  $p = 0.3191$ ; grade 2:  $p = 0.3599$ ).

Overall, HR of all-cause mortality adjusted for chronic obstructive pulmonary disease, cancer, atrial fibrillation, heart failure, ischaemic heart disease, chronic renal insufficiency in patients with VV was 1.34 times higher (adjusted HR 1.37; 95% CI 1.19 to 1.57;  $p < 0.0001$ ) than that in controls (table 2). Stratified analysis revealed 1.38 and 1.41 times increased risks of mortality in older (age ≥65 years; adjusted HR 1.38; 95% CI 1.17 to 1.62;  $p = 0.0001$ ) and male patients with VV (adjusted HR 1.41; 95% CI 1.18 to 1.68;  $p = 0.0001$ ). Notably, despite no significant effect of VV on the survival of patients with hypertension, hyperlipidaemia or CAD was observed, patients with both VV and diabetes presented 1.50 times higher risk of mortality compared with those without VV (adjusted HR 1.50; 95% CI 1.05 to 2.15;  $p = 0.0254$ ). Furthermore, VV at grade 3 show 1.83 (95% CI 1.48, 2.27;  $p < 0.0001$ ) greater risk of mortality adjusted for chronic obstructive pulmonary disease, cancer, atrial fibrillation, heart failure, ischaemic heart disease, chronic renal insufficiency.





**Table 1** Baseline characteristics and comorbid medical disorders for the control cohort and patients with varicose vein


Characteristic, n (%)	Varicose vein, n=4644	Matched controls, n=17 742	Standardised difference
Age (years)			0.02275
<65	3164 (68.13)	12 275 (69.19)	
≥65	1480 (31.87)	5467 (30.81)	
Age (mean±SD)	55.70±16.03	56.10±16.04	0.02514
Gender			0.00944
Male	1796 (38.67)	6780 (38.21)	
Female	2848 (61.33)	10 962 (61.79)	
Hypertension			0.04519
No	3750 (80.75)	14 637 (82.50)	
Yes	894 (19.25)	3105 (17.50)	
Diabetes			0.05807
No	4247 (91.45)	16 501 (93.01)	
Yes	397 (8.55)	1241 (6.99)	
Hyperlipidaemia			0.08429
No	4413 (95.03)	17 157 (96.70)	
Yes	231 (4.97)	585 (3.30)	
Coronary artery disease			0.07832
No	4489 (96.66)	17 375 (97.93)	
Yes	155 (3.34)	367 (2.07)	

P value was calculated based on the two sample t-test and Pearson's  $\chi^2$  test.

### Long-term MACE risk

MACE risk significantly increased in patients with VV (HR 2.05; 95% CI 1.89 to 2.23;  $p<0.0001$ ), particularly in relatively younger (age, <65 years; adjusted HR 2.17; 95% CI 1.92 to 2.46;  $p<0.0001$ ) or male (adjusted HR 2.32; 95% CI 2.06 to 2.62;  $p<0.0001$ ) patients (table 3). In addition, patients with VV showing cardiovascular risk factors, including hypertension, diabetes, hyperlipidaemia

and CAD, were at a higher risk of MACE than were matched controls. In patients with VV, 3-year, 6-year and 9-year MACE-free rates were 91.17%, 84.99% and 79.27% (figure 2A). These rates dramatically declined further with disease severity (figure 2B). In terms of individual cardiovascular outcomes, patients with grade 3 VV were at a greater risk of CHF (adjusted HR 2.05; 95% CI 1.71 to 2.46;  $p<0.0001$ ), ACS (adjusted HR: 2.04;



**Figure 1** (A) Kaplan-Meier estimates of 12-year survival between patients with varicose veins (VV) and the matched control cohort. (B) Kaplan-Meier estimates of 12-year survival between patients with VV categorised by the disease severities and the matched control cohort.

**Table 2** Crude and adjusted HRs of all-cause mortality in patients with VV compared with the matched control cohort during the follow-up period

Cohort all (n=22 386)	Crude HR (95% CI)	P value	Adjusted HR (95% CI)†	P value
Overall analysis				
VV	1.43 (1.25 to 1.64)	<0.0001*	1.37 (1.19 to 1.57)	<0.0001*
Controls	1(reference)		1(reference)	
Stratified analysis				
Age (years)				
<65 (years)				
VV	1.49 (1.144 to 1.93)	0.0030*	1.2 (0.92 to 1.57)	0.1803
Controls	1(reference)		1(reference)	
≥65 (years)				
VV	1.41 (1.2 to 1.66)	<0.0001*	1.38 (1.17 to 1.62)	0.0001†
Controls	1(reference)		1(reference)	
Gender				
Male				
VV	1.46 (1.23 to 1.74)	<0.0001*	1.41 (1.181 to 1.68)	0.0001†
Controls	1(reference)		1(reference)	
Female				
VV	1.38 (1.1 to 1.74)	0.0058*	1.31 (1.04 to 1.65)	0.0227†
Controls	1(reference)		1(reference)	
Hypertension				
VV	1.14 (0.87 to 1.49)	0.3476	1.16 (0.88 to 1.52)	0.2957
Controls	1(reference)		1(reference)	
Diabetes				
VV	1.5 (1.06 to 2.14)	0.0226*	1.5 (1.05 to 2.15)	0.0254†
Controls	1(reference)		1(reference)	
Hyperlipidaemia				
VV	0.99 (0.39 to 2.54)	0.9865	1.01 (0.39 to 2.61)	0.9914
Controls	1(reference)		1(reference)	
Coronary artery disease				
VV	1.14 (0.64 to 2.03)	0.6565	1.05 (0.58 to 1.92)	0.8716
Controls	1(reference)		1(reference)	

\*P&lt;0.05.

†Adjusted for chronic obstructive pulmonary disease, cancer, atrial fibrillation, heart failure, ischaemic heart disease, chronic renal insufficiency.

VV, varicose veins.

95% CI: 1.58 to 2.63;  $p<0.0001$ ) and ischaemic stroke (adjusted HR 2.06; 95% CI 1.58 to 2.69;  $p<0.0001$ ) than were controls (table 4). In particular, with the highest VV severity there was an increasing risk of venous thrombotic events, including DVT and PE (grade 3: adjusted HR 38.4; 95% CI 16.4 to 90.1;  $p<0.0001$ ) (table 4).

#### Validation of the accuracy of VV diagnosis and ICD-9-CM-derived VV grading

During 2010–2015, a total of 2202 outpatients and 347 inpatients were reported to have VV in Chi-Mei Medical Center. Among the outpatients, 1188 were coded as

uncomplicated VV (ICD-9-CM code 454.9), 775 were coded as VV with inflammation (ICD-9-CM code 454.1), 152 were coded as VV with ulcers (ICD-9-CM code 454.0) and 87 were coded as VV with ulcer and inflammation (ICD-9-CM code 454.2) (online supplementary table 3). Notably, none were coded incorrectly. Compared with CEAP stage, as determined based on chart reviews, only a few inpatients were incorrectly or unclearly diagnosed using ICD-9-CM-derived VV codes (online supplementary table 4). For example, among patients with higher VV grades (CEAP stage 5–6), the positive and negative

**Table 3** Crude and adjusted HRs of MACE in patients with VV compared with the matched control cohort during the follow-up period

Cohort all (n=22386)	Crude HR (95% CI)	P value	Adjusted HR (95% CI)†	P value
Overall analysis				
VV	2.08 (1.91 to 2.25)	<0.0001*	2.05 (1.89 to 2.23)	<0.0001*
Controls	1(reference)		1(reference)	
Stratified analysis				
Age (years)				
<65 (years)				
VV	2.21 (1.95 to 2.5)	<0.0001*	2.17 (1.92 to 2.46)	<0.0001*
Controls	1(reference)		1(reference)	
≥65 (years)				
VV	1.98 (1.78 to 2.21)	<0.0001*	1.96 (1.76 to 2.18)	<0.0001*
Controls	1(reference)		1(reference)	
Gender				
Male				
VV	2.35 (2.09 to 2.65)	<0.0001*	2.32 (2.06 to 2.62)	<0.0001*
Controls	1(reference)		1(reference)	
Female				
VV	1.87 (1.67 to 2.09)	<0.0001*	1.85 (1.66 to 2.07)	<0.0001*
Controls	1(reference)		1(reference)	
Hypertension				
VV	1.65 (1.42 to 1.92)	<0.0001*	1.62 (1.39 to 1.89)	<0.0001*
Controls	1(reference)		1(reference)	
Diabetes				
VV	1.4 (1.11 to 1.76)	0.0042*	1.37 (1.08 to 1.72)	0.0081
Controls	1(reference)		1(reference)	
Hyperlipidaemia				
VV	1.5 (1.03 to 2.17)	0.0353*	1.56 (1.07 to 2.29)	0.0224
Controls	1(reference)		1(reference)	
Coronary artery disease				
VV	1.93 (1.38 to 2.7)	0.0001*	1.99 (1.41 to 2.82)	0.0001*
Controls	1(reference)		1(reference)	

\*P&lt;0.05.

†Adjusted for chronic obstructive pulmonary disease, cancer, atrial fibrillation, heart failure, ischaemic heart disease, chronic renal insufficiency.

.MACE, major adverse cardiovascular event; VV, varicose veins.


predictive values with ICD-9-CM-derived codes were 93% and 98.4%, respectively. Specifically, the sensitivity and specificity of ICD-9-CM-derived grading were up to 95.2% and 97.6%, respectively. The calculated kappa score between CEAP stages and grading severity is 0.92 (95% CI 0.88 to 0.96).

### Sensitivity analyses

VV and controls with pregnancy history were identified and examination the influence in sensitivity analysis (online supplementary table 5). After additionally adjustment for history of pregnancy, the results

remain showing great impacts on mortality and MACE (adjusted HR for death (1.37 (95% CI 1.19 to 1.57),  $p<0.0001$ ; adjusted HR for MACE 2.01 (95% CI 1.89 to 2.23),  $p<0.0001$ ).

After excluding 472 subjects with Myocardial infarction, stroke, coronary angioplasty or CABG, remaining VV and corresponding controls were included for sensitivity analysis. Comparing with corresponding matched controls, those conservatively treated VV patients were found 1.36 times risks of mortality (adjusted HR 1.36 (95% CI 1.18 to 1.57),  $p<0.0001$ ) and 1.95 times risks



**Figure 2** (A) Kaplan-Meier estimates of 12-year free from MACE between patients with varicose veins (VV) and the matched control cohort. (B) Kaplan-Meier estimates of 12-year free from MACE between patients with VV categorised by the disease severities and the matched control cohort. MACE, major adverse cardiovascular event.

of MACE (adjusted HR 1.95 (95% CI 1.80 to 2.12),  $p < 0.0001$ ).

## DISCUSSION

The primary findings of this study were that (1) patients with VV were at increasing risks of mortality and cardiovascular events, especially those with VV at grade 3

compared with matched controls; (2) having VV had a significant impact on the survival of male patients. To the best of our knowledge, this nationwide population-based study is the first to comprehensively describe the association of VV with patients' cardiovascular outcomes.

Although VV are common, their potential threat to health has not been well investigated previously.<sup>1 2</sup> Valve

**Table 4** The adjusted HRs of mortality and MACE in patients with VV compared with the matched control cohort during the follow-up period

	Grade 1 control	Grade 1	Grade 2 control	Grade 2	Grade 3 control	Grade 3
No of mortality, N (%)	343 (3.62)	99 (4.01)	147 (5.79)	44 (6.59)	266 (4.65)	136 (9.01)
Adjusted HR for mortality (95% CI)†	Referent	1.08 (0.86 to 1.36)	Referent	1.13 (0.8 to 1.6)	Referent	1.83 (1.48 to 2.27)*
No of CHF, N (%)	552 (5.82)	238 (9.65)	181 (7.12)	80 (11.98)	358 (6.26)	190 (12.59)
Adjusted HR for CHF (95% CI)†	Referent	1.68 (1.44 to 1.96)*	Referent	1.79 (1.37 to 2.34)†	Referent	2.05 (1.71 to 2.46)*
No of ACS, N (%)	291 (3.07)	125 (5.07)	72 (2.83)	24 (3.59)	174 (3.04)	95 (6.30)
Adjusted HR for ACS (95% CI)†	Referent	1.7 (1.37 to 2.11)*	Referent	1.25 (0.78 to 1.99)	Referent	2.04 (1.58 to 2.63)*
No of ischaemic stroke, N (%)	236 (2.49)	99 (4.01)	90 (3.54)	31 (4.64)	162 (2.83)	89 (5.90)
Adjusted HR for ischaemic stroke (95% CI)†	Referent	1.59 (1.25 to 2.01)*	Referent	1.4 (0.92 to 2.12)	Referent	2.06 (1.58 to 2.69)*
No of DVT+PE, N (%)	14 (0.15)	56 (2.27)	7 (0.28)	13 (1.95)	6 (0.10)	63 (4.17)
Adjusted HR for DVT+PE (95% CI)†	Referent	14.9 (8.26 to 26.86)*	Referent	6.27 (2.46 to 15.96)*	Referent	38.42 (16.38 to 90.13)*

\* $P < 0.05$ .

†Adjusted for chronic obstructive pulmonary disease, cancer, atrial fibrillation, heart failure, ischaemic heart disease, chronic renal insufficiency.

ACS, acute coronary syndrome; CHF, congestive heart failure; DVT, deep vein thrombosis; MACE, major adverse cardiovascular event; PE, pulmonary embolism; VV, varicose veins.







increased mortality with higher ICD-9-CM-derived grades indicated that our grading system specifically reflected the severity of VV but the cause of mortality was not available in this database.

## CONCLUSIONS

VVs are a common condition typically believed to be benign; however, our results suggest that they warrant close attention. Compared with matched controls, patients with VV were at increasing risks of mortality and cardiovascular events, especially those with VV at grade 3. Therefore, these findings should alert clinicians regarding the importance of detecting VV at an early stage.

## Author affiliations

- <sup>1</sup>Department of Cardiovascular Surgery, Chi Mei Medical Center, Tainan, Taiwan
- <sup>2</sup>Department of Hospital and Health Care Administration, Chia Nan University of Pharmacy and Science, Rende, Taiwan
- <sup>3</sup>Department of Cardiology, Chi Mei Medical Center, Tainan, Taiwan
- <sup>4</sup>Department of Pharmacy, Chia Nan University of Pharmacy and Science, Rende, Taiwan
- <sup>5</sup>Department of Healthcare Administration and Medical Informatics, Chi Mei Medical Center, Tainan, Taiwan
- <sup>6</sup>Medical Research, Chi Mei Medical Center, Tainan, Taiwan
- <sup>7</sup>Department of Biotechnology, Southern Taiwan University of Science and Technology, Tainan, Taiwan
- <sup>8</sup>Institute of Clinical Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan

**Contributors** All authors contributed to the revision of the manuscript and approved the final version. All agreed to be accountable. N-CW contributed to concept and design, critical revision. Z-CC contributed to data collection and critical revision. I-JF contributed to critical revision, data collection, statistical analysis and interpretation. C-HH contributed to critical revision, data collection, statistical analysis and interpretation. C-YC contributed to data collection and critical revision. J-JW contributed to data collection and critical revision. W-TC contributed to concept and design, data collection, interpretation, manuscript writing and critical revision.

**Funding** We received the research grant supported by Chi-Mei Medical Center

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Ethics approval** The present study was ethically approved by the Institutional Review Board of Chi-Mei Hospital (CV code: 10406-E01). All procedures followed the principles outlined in the Declaration of Helsinki.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open access repository. All the data are available in National Health Insurance Research Database (NHIRD) in Taiwan.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Chung-Han Ho <http://orcid.org/0000-0001-5925-8477>

Wei-Ting Chang <http://orcid.org/0000-0001-9525-2144>

## REFERENCES

- 1 Beebe-Dimmer JL, Pfeifer JR, Engle JS, *et al*. The epidemiology of chronic venous insufficiency and varicose veins. *Ann Epidemiol* 2005;15:175–84.
- 2 Cesarone MR, Belcaro G, Nicolaides AN, *et al*. 'Real' epidemiology of varicose veins and chronic venous diseases: the San Valentino vascular screening project. *Angiology* 2002;53:119–30.
- 3 Eberhardt RT, Raffetto JD. Chronic venous insufficiency. *Circulation* 2014;130:333–46.
- 4 Nicolaides AN. Investigation of chronic venous insufficiency: a consensus statement (France, March 5–9, 1997). *Circulation*;2000:E126–63.
- 5 Müller-Bühl U, Leutgeb R, Engeser P, *et al*. Varicose veins are a risk factor for deep venous thrombosis in general practice patients. *VASA* 2012;41:360–5.
- 6 Søgaard KK, Schmidt M, Pedersen L, *et al*. 30-Year mortality after venous thromboembolism: a population-based cohort study. *Circulation* 2014;130:829–36.
- 7 Heit JA, Silverstein MD, Mohr DN, *et al*. Risk factors for deep vein thrombosis and pulmonary embolism: a population-based case-control study. *Arch Intern Med* 2000;160:809–15.
- 8 Chang C-H, Lin C-H, Caffrey JL, *et al*. Risk of intracranial hemorrhage from statin use in Asians: a nationwide cohort study. *Circulation* 2015;131:2070–8.
- 9 Cheng C-L, Kao Y-HY, Lin S-J, *et al*. Validation of the National health insurance research database with ischemic stroke cases in Taiwan. *Pharmacoepidemiol Drug Saf* 2011;20:236–42.
- 10 Austin PC. An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivariate Behav Res* 2011;46:399–424.
- 11 Markovic JN, Shortell CK. Genomics of varicose veins and chronic venous insufficiency. *Semin Vasc Surg* 2013;26:2–13.
- 12 Komarow W, Hawro P, Lekston A, *et al*. Endothelial dysfunction in patients with chronic venous disease: an evaluation based on the flow-mediated dilatation test. *Int Angiol* 2015;34:36–42.
- 13 Jacob T, Hingorani A, Ascher E. Overexpression of transforming growth factor-beta1 correlates with increased synthesis of nitric oxide synthase in varicose veins. *J Vasc Surg* 2005;41:523–30.
- 14 Mäkiävaara LA, Ahti TM, Luukkaala T, *et al*. Persons with varicose veins have a high subsequent incidence of arterial disease: a population-based study in Tampere, Finland. *Angiology* 2007;58:704–9.
- 15 Piazza G, Goldhaber SZ. Venous thromboembolism and atherothrombosis: an integrated approach. *Circulation* 2010;121:2146–50.
- 16 Chang S-L, Huang Y-L, Lee M-C, *et al*. Association of varicose veins with incident venous thromboembolism and peripheral artery disease. *JAMA* 2018;319:807–17.
- 17 Sørensen HT, Horvath-Puho E, Lash TL, *et al*. Heart disease may be a risk factor for pulmonary embolism without peripheral deep venous thrombosis. *Circulation* 2011;124:1435–41.
- 18 Sørensen HT, Horvath-Puho E, Pedersen L, *et al*. Venous thromboembolism and subsequent hospitalisation due to acute arterial cardiovascular events: a 20-year cohort study. *Lancet* 2007;370:1773–9.
- 19 Lohr JM, Bush RL. Venous disease in women: epidemiology, manifestations, and treatment. *J Vasc Surg* 2013;57 (4 Suppl):37S–45S.
- 20 Fowkes FG, Lee AJ, Evans CJ, *et al*. Lifestyle risk factors for lower limb venous reflux in the general population: Edinburgh vein study. *Int J Epidemiol* 2001;30:846–52.
- 21 Hsieh C-Y, Su C-C, Shao S-C, *et al*. Taiwan's national health insurance research database: past and future. *Clin Epidemiol* 2019;11:349–58.