BMJ Open Comparing and determining factors associated with hypertension self-care profiles of patients in two multi-ethnic Asian countries: cross-sectional studies between two study populations

Sabrina Yi-Mei Wee,^{1,2} Hani Salim ^(b),³ Maliza Mawardi,³ Yi Ling Eileen Koh,¹ Hanifatiyah Ali,³ Sazlina Shariff Ghazali ^(b),^{3,4} Ping Yein Lee,⁵ Siew Mooi Ching ^(b) Nurainul Hana Shamsuddin,³ Ngiap Chuan Tan ^(b),^{1,2}

ABSTRACT

Objectives To compare the sociodemography, disease characteristics and hypertension self-care profiles and to determine the factors influencing Hypertension Self-Care Profiles (HTN-SCP) in two populations in primary care settings from Singapore and Malaysia.

Design Cross-sectional, cross national.

Setting Multi-centre, primary care clinics Malaysia and Singapore.

Participants 1123 adults with hypertension enrolled and analysed.

Primary and secondary outcome

measures Comparison between sociodemography, disease characteristics and the mean scores of HTN-SCP domains (behaviour, motivation and self-efficacy) and the factors influencing hypertension self-care.

Results 1123 adults with hypertension attending primary care clinics in Malaysia and Singapore were involved. The participants' mean age was 63.6 years (SD 9.7) in Singapore and 60.4 (SD 9.1) in Malaysia. Most of the participants in Singapore had tertiary education (22.3%) compared with Malaysia (13.0%), p<0.001. A higher proportion of participants from Singapore had controlled blood pressure (74.6%) compared with Malaysia (33.8%), p<0.001. The mean total score of HTN-SCP was significantly higher among Singapore participants compared with Malaysia participants 190 (SD 28) versus 184 (SD 23) (p<0.001). Similarly, the mean score for motivation domain 67 (SD 10) versus 65 (SD 9), followed by self-efficacy score 65 (SD 11) versus 62 (SD 9) and behaviour score (58 SD 9 vs 56 SD 9) were higher among Singapore participants. In both countries, the factors which influenced higher HTN-SCP mean scores across all domains were being Indian and had tertiary education.

Conclusions The study population in Singapore had a higher HTN-SCP mean score compared with Malaysia. The common factors influencing higher HTN-SCP mean scores at both study sites were ethnicity and level of education. Future intervention to improve self-care among people with hypertension may need to be tailored to their behaviour, motivation and self-efficacy levels.

- S-sectional studies opulations
 aliza Mawardi, ³ Yi Ling Eileen Koh,¹ ,^{3,4} Ping Yein Lee, ⁵ Siew Mooi Ching ⁽¹⁾, ³ Tan ⁽²⁾ ^{1,2}
 Stengths and limitations of this study in an ⁽²⁾ ^{1,2}
 The unprecedented cross-national comparison of patients with hypertension sharing similar sociocul-tural background but different economic and health financing environment contributes to this study's strength.
 The study examined the sociodemography and disease characteristics factors which influenced Hypertension Self-Care Profiles (HTN-SCP) domain scores in Malaysia and Singapore among people aged 40 years and above.
 Large overall sample size has enabled in-depth analyses of the individual domain of the HTN-SCP.
 Cross-sectional study design implies associations but not causation and does not allow conclusions about changes in behaviour during the course of illness.
 The sample size and recruitment methods differ between the countries; suggesting that the results may not be generalisable to the national population and the HTN-SCP domains mean scores comparison between the two countries must be interpreted with caution.
 Both Malaysia and Singapore inherited the British-style National Health Service and from its revenue.¹ Over the years, the dissim-ilarity in the rate of socioeconomic devel

from its revenue.¹ Over the years, the dissimilarity in the rate of socioeconomic development has significantly impacted the way resources being channelled to public services, particularly the health service. The Malaysian government funded medical and public health services through revenues derived from taxes, government revenues and income earned from government corporatised enterprises.² While in Singapore, it adopts a hybrid system to finance the health system whereby,

1

self-care profiles of patients in two multi-ethnic Asian countries: cross-sectional studies between two study populations. BMJ Open 2021;11:e044192. doi:10.1136/ bmjopen-2020-044192 Prepublication history and

To cite: Wee SY-M. Salim H.

Mawardi M, et al. Comparing

associated with hypertension

and determining factors

additional supplemental material for this paper are available online. To view these files, please visit the journal online (http://dx.doi.org/10.1136/ bmjopen-2020-044192).

SY-MW and HS contributed equally.

Received 26 August 2020 Accepted 27 May 2021



C Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BM.J.

For numbered affiliations see end of article.

Correspondence to Dr Hani Salim: hanisyahida@upm.edu.my



the cost of care is funded jointly by the government and the individual through insurance, revenue from taxes and personal medical saving accounts.³ The intent to cultivate personal responsibility towards taking charge of an individual's health underpins the shared healthcare financing concept. Such an approach is postulated to shape an individual's overall perceptions and attitudes towards self-efficacy and self-care,⁴ which is pivotal for the successful management of long term, non-communicable diseases such as hypertension.

Hypertension is a significant cause of morbidity and mortality arising from cardiovascular and kidney disease.⁵ The prevalence of adults with hypertension as 30% in Malaysia and 21.5% in Singapore.⁶⁷ However, a significant proportion of the affected population has yet to attain treatment goal. In Singapore, 49.7% of patients treated for hypertension in primary care were reported with good blood pressure (BP) control.⁷ While in Malaysia, the proportion of patients with good hypertension control is lower at 37.4%.⁸ The differences in the proportion of good BP control warrants a comparison of their health behaviour profiles in the two countries as they have similar multi-ethnic background and culture.

The management of hypertension does not only encompass pharmacological treatment prescribed from the attending physician, but patients themselves are required to perform self-care measures to improve their BP control.⁹ Self-care includes maintaining a healthy diet, performing regular physical activities, achieving ideal body weight and avoiding unhealthy lifestyle such as smoking.¹⁰ The success of these activities requires behaviour change, motivation and self-efficacy. Hypertension self-care profile among adult with hypertension aged 18 years and above were found to be moderate.¹¹ To date, such data in Singapore are relatively lacking. This study aimed to compare the sociodemography, disease characteristics and HTN-SCP between Malaysia and Singapore and to determine the factors influencing hypertension self-care among study populations in Malaysia and Singapore among people aged 40 years and above.

METHODOLOGY

The studies were cross-national surveys of self-care profiles of patients with hypertension conducted among two study populations in two countries between October 2016 and June 2017. In Malaysia, the study was conducted in three urban primary care clinics in Selangor, Malaysia and a polyclinic in Bukit Merah, Singapore.

The studies' inclusion criteria were adults aged 40 years and above, with underlying hypertension diagnosed by a physician. Pregnant women and those with underlying psychiatric illness or cognitive impairment were excluded from this study. The estimated sample size in Malaysia was 720 based on the mean score of self-care management among hypertensive aged <60 years and ≥ 60 years¹² using Lemeshow *et al* formula¹³ with 95% confidence level, 90% of power and 20% of ັດ

non-response rate. However, in comparison with our previous reporting, in this study, we only include people aged 40 years and above, which gave us a total of 702 participants in Malaysia.

As the Singapore study had no prior literature on the percentage of patients with high self-care, 50% was adopted to obtain the maximum sample size. With a 95%CI estimate and 5% precision, the sample size required was 385. The sample size was increased to 450 to account for 15% incomplete data and non-response rate.

Study instrument

Protected In this study, the Hypertension Self-Care Profiles (HTNby copyright SCP) was used to assess the hypertension self-care profile. In this tool's development, two underpinning theories were Orem's self-care model and motivational interviewing (MI).^{14 15} Orem's model described how people enabled self-care by performing specific actions to manage their illness.¹⁴ Understanding the reason behind these actions were crucial to self-care. MI facilitates the self-care process by promoting commitment and developing confidence for a behaviour change.¹⁵ Thus, using **o** HTN-SCP tool uses the domains of behaviour, motivation **g** and self-efficacy to assess self-care among patients with HTN-SCP tool uses the domains of behaviour, motivation hypertension.¹⁶

The HTN-SCP tool is a reliable tool with good internal consistency.¹⁶ It consists of 60-items and three domains which are behaviour, motivation and self-efficacy. It has been validated in Singapore,¹⁷⁻¹⁹ and the Cronbach's alpha for the subdomain's ranges from 0.857 to 0.948. For the Malay version, the Cronbach's alpha for the subdomains ranges from 0.851 to 0.945 whereas, for the Mandarin version, the Cronbach's alpha ranges from 0.838 to 0.929.^{18 19} There were 20-items in each domain, and the score of each domain ranges from 0 to 80 as each \mathbf{a} question using a 4-point Likert scale. Higher scores indi- ▶ cate a higher level of self-care behaviour, motivations and self-efficacy. A pretest of the questionnaire involving 30 participants was conducted in Malaysia to determine the B questionnaire's feasibility. Following the pretest, minor changes were made to the questionnaire.

The questionnaire was available in three languages which are English, Malay and Mandarin for participants to select their preferred version.^{17–19} It consists of three b select their preferred version.¹⁷⁻¹⁹ It consists of three tections: sociodemographic characteristics, medical rofiles on hypertension and the HTN-SCP tool. BP eading was taken from the patient's medical records. The definition of body mass index (BMI) was based sections: sociodemographic characteristics, medical profiles on hypertension and the HTN-SCP tool. BP reading was taken from the patient's medical records.

on the WHO recommendation for Asian population.²⁰ Underweight is defined if BMI $<18.5 \text{ kg/m}^2$, normal weight is defined if BMI is 18.5-22.9 kg/m², overweight is defined with BMI 23–27.4 kg/m² and obese is defined if BMI >27.5 kg/m². The definition of controlled BP was based on the Joint National Committee Eighth (JNC 8) guidelines.²¹ The BP of patients with underlying hypertension without diabetes was considered controlled if their BP <140/90 mm Hg, regardless of age.

Data collection Malaysia

In Malaysia, participants were recruited using a systematic random sampling method. A sampling interval of two was used as a constant difference between participants. The first patient (the reference point) was chosen using a draw lot method. Subsequently, every alternate patient was approached for study participation.

Singapore

In Singapore, potential participants were screened for eligibility at the waiting area outside the clinic consultation rooms and were invited to participate in the study.

Patients gave written informed consent to join the study. We obtained their sociodemographic data via self-administered proforma. The HTN-SCP questionnaire was administered through a face-to-face interview. We verified the patients' clinical information through their latest medical records.

Data analysis

We used SPSS V.22.0 in the data analysis. We used descriptive statistics to describe the demographic and disease profiles of the patients. We used percentages and frequencies for the categorical variables; mean and SD were used for the continuous variables if they were normally distributed. The normality of the continuous data was based on z-score of skewness and kurtosis, Kolmogorov-Smirnov, histogram and Q-Q plot. We used independent t-test or one-way ANOVA (analysis of variance) to determine the association for numerical data. We performed a χ^2 or Fisher's exact test for the categorical data. The significant level was set at p<0.05. multiple linear regression (MLR) model was performed to determine the predictors for hypertension self-care. Variables with p<0.25 from the univariate analysis were included in the MLR model. Level of significance was set at p<0.005. The MLR results were reported as beta coefficient, SE and 95% CI.

Patient and public involvement

This research was done with the involvement of the patients as research participants. They were not involved in the study design, recruitment, interpretation of the report's results and writing.

RESULTS

A total of 1123 adults with hypertension participated in this study, of which 702 and 421 were Malaysians and Singaporeans, respectively. The response rate in Malaysia was 93.8% (761/811). Non-participations were due to language barrier and time. The response rate was not reported in Singapore. The proportion of participants aged 60 years and above was higher in Singapore (63.4%) than Malaysia (54.6%). More female participants were recruited among the Singaporeans (50.6%) and Malaysians (49.4%). More Malays (42.9%) in the latter, whereas in the former, Chinese ethnicity constituted the highest proportion (69.6%). More participants were married in Malaysia (81.6%) than those in Singapore (67%). More patients had tertiary education in Singapore (22.3%), versus those in Malaysia (13.0%). Table 1 summarises the characteristics of the study participants.

A higher proportion of the participants from Malaysia were on three or more than four antihypertensive medications (20.9%) compared with Singapore (14.3%) (p=0.023). Regarding BP control to target, Malaysia (33.8%) had a significantly lower proportion of the treatment goal compared with Singapore (74.6%). Significantly more Malaysia participants had diabetes mellitus (65.8%) compared with Singapore participants (46.8%) (p<0.001) (table 2).

Table 3 illustrates the score of the HTN-SCP of the participants. The mean total score of HTN-SCP was significantly higher among Singapore participants (mean 189.9, SD 27.6) compared with Malaysia participants (mean 184.1, SD 22.8) (p<0.001). Similarly, for all the subdomains mean scores: motivation domain (p<0.001), self-efficacy domain (p<0.001) and behaviour score (p<0.001) were significantly higher among the Singapore participants compared with Malaysia participants.

uses rela Detail results of associations between HTN-SCP behaviour, motivation and self-efficacy scores and sociodemographic factors and disease characteristic in participants are in online supplemental files 1-3. As shown in table 4, in both countries, the factors that were significantly associated with HTN-SCP behaviour mean scores were e participants aged 60 years and above (Malaysia: adjusted beta=2.047, 95% CI 0.728 to 3.365, p=0.002) (Singapore: adjusted beta=2.473, 95% CI 0.671 to 4.275, p=0.007), of Indian ethnicity (Malaysia: adjusted beta=4.389, 95% CI 2.614 to 6.164, p<0.001) (Singapore: adjusted beta=3.271, ∃ 95% CI 1.09 to 5.452, p=0.003) and those with tertiary education (Malaysia: adjusted beta=4.274, 95% CI 2.175 ≥ to 6.373, p<0.001) (Singapore: adjusted beta=4.243, 95% CI 1.857 to 6.629, p<0.001). For Malaysia, Malay ethnicity (adjusted beta=3.192, 95% CI 1.719 to 4.665, p<0.001) ğ also was associated with higher HTN-SCP behaviour mean scores. For Singapore, other factors associated with HTN-SCP behaviour were women participants (adjusted beta=1.864, 95% CI 0.133 to 3.595, p<0.035), of other ethnicities (adjusted beta=9.25, 95% CI 2.714 to 15.786, p=0.006), and those with secondary education (adjusted beta=3.184, 95% CI 1.28 to 5.09, p=0.001).

Table 4 summarises the association between HTN-SCP motivation mean scores and sociodemographic and **G** disease characteristics among Malaysia and Singapore **B** participants. In both countries, the HTN-SCP motivation mean scores were significantly associated with Indian ethnicity (Malaysia: adjusted beta=5.099, 95% CI 3.359 to 6.838, p<0.001) (Singapore: adjusted beta=3.374, 95% CI 1.254 to 6.215, p=0.003), secondary education level (Malaysia: adjusted beta=3.125, 95% CI 1.693 to 4.557, p<0.001) (Singapore: adjusted beta=5.067, 95% CI 2.903 to 7.232, p<0.001), and tertiary education levels (Malaysia: adjusted beta=7.136, 95% CI 5.045 to 9.227,

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.	Enseignement Superieur (ABES) .	J Open: first published as 10.1136/bmjopen-2020-044192 on 14 June 2021. Downloaded from http://bmjopen.bmj.com/ on June 12, 2025 at Agence Bibliographique de
--	---------------------------------	---

Table 1 Sociodemographic cha	racteristics of the study participants by	country (N=1123)	
	Malaysia participants (n=702)	Singapore participants (n=421)	P value
	N (%)	N (%)	
Total	702 (62.5)	421 (37.5)	
Age, years (mean+SD)	60.4 (9.1)	63.6 (9.7)	<0.001*
≥60	383 (54.6)	267 (63.4)	0.004*
Gender			0.106
Women	208 (49.4)	213 (50.6)	
Ethnicity			<0.001*
Chinese	251 (35.8)	293 (69.6)	
Malay	301 (42.9)	40 (9.5)	
Indian	150 (21.4)	81 (19.2)	
Others	0 (0)	7 (1.7)	
Education level			<0.001*
Primary and below	252 (35.9)	157 (37.3)	
Secondary	359 (51.1)	170 (40.4)	
Tertiary	91 (13.0)	94 (22.3)	
Marital status			<0.001*
Married	572 (81.6)	282 (67)	
Single/separated/widowed	129 (18.4)	139 (33)	
Declined to answer	1 (0.1)	0	
Univariate analysis (χ^2 test or Fisher's	exact test).		

*P value <0.05.

Table 2 Clinical characteristics of the second	he study participants by country (N=112	3)	
Variable	Malaysia participants (n=702)	Singapore participants (n=421)	P value
	N (%)	N (%)	
Antihypertensive medications (n)			0.023*
0	7 (1)	7 (1.7)	
1	301 (42.9)	208 (49.6)	
2	247 (35.2)	144 (34.4)	
3	113 (16.1)	51 (12.2)	
>4	34 (4.8)	9 (2.1)	
Blood pressure well-controlled			<0.001*
Yes	237 (33.8)	314 (74.6)	
Body mass index, kg/m ²			0.191
Underweight (<18.5)	8 (1.2)	7 (1.7)	
Normal weight (18.5–22.9)	115 (16.7)	88 (21.3)	
Overweight (23–27.4)	236 (34.3)	140 (33.9)	
Obese (>27.5)	329 (47.8)	178 (43.1)	
Missing data	8 (1.1)	8 (1.9)	
Presence of diabetes mellitus			<0.001*
Yes	462 (65.8)	197 (46.8)	

Univariate analysis (χ^2 test or Fisher's exact test). *P value <0.05.

Table 3 Scores of Hypertension Self-C	are Profiles (HTN-	SCP)							
	Range of s	cores	Mean (SD)						
Domain	Malaysia	Singapore	Malaysia	Singapore	†P value	Adjusted beta	SE	95% CI	‡P value
Behaviour	31–79	27–80	56.5 (8.9)	58.4 (9.1)	0.001*	2.2	0.6	1 to 3.4.0	<0.001*
Motivation	23–80	27–80	65.2 (9.0)	67.1 (10.4)	0.001*	3.1	0.6	1.8 to 4.3	<0.001*
Self-efficacy	31–80	29–80	62.4 (8.8)	64.5 (10.6)	<0.001*	2.9	0.6	1.7 to 4.2	<0.001*
Total HTN-SCP score	102–238	92–240	184.1 (22.8)	189.9 (27.6)	<0.001*	8.2	1.6	5.0 to 11.4	<0.001*
*P value <0.05. †Univariate analysis (independent t-test).									

tMultivariate analysis assessing the difference between scores among two countries (controlling for age, gender, ethnicity, education, marital status, number of antihypertensive medications,

vell-controlled blood pressure, body mass index, presence of diabetes)

p<0.001) (Singapore: adjusted beta=5.528, 95% CI 2.863 to 8.193, p<0.001). For Malaysia, Malay ethnicity (adjusted beta=4.339, 95% CI 2.857 to 5.82, p<0.001) also was associated with higher HTN-SCP motivation mean scores.

Table 4 summarises the association between HTN-SCP self-efficacy mean scores and our participants' sociodemographic factors and disease characteristics. In both countries, the factors that were significantly associated with HTN-SCP self-efficacy mean scores were participants of Indian ethnicity (Malaysia: adjusted beta=6.174, 95% CI 4.433 to 7.914, p<0.001), (Singapore: adjusted beta=3.706, 95% CI 1.163 to 6.25, p=0.004) and those with tertiary education (Malaysia: adjusted beta=4.752, 95% CI 2.687 ŝ to 6.818, p<0.001) (Singapore: adjusted beta=4.179, 95%) CI 1.51 to 6.847, p<0.001). For Malaysia, Malay ethnicity pyright, (adjusted beta=4.003, 95% CI 2.537 to 5.468, p<0.001) and women (adjusted beta=1.747, 95% CI 0.475 to 3.02, p=0.007) were associated with higher HTN-SCP selfefficacy mean scores. For Singapore, other factors associated with HTN-SCP self-efficacy were of other ethnicities uding (adjusted beta=8.4, 95% CI 0.696 to 16.104, p=0.033) and those with secondary education (adjusted beta=3.921, for uses related to text 95% CI 1.698 to 6.145, p=0.001).

DISCUSSION Summary of findings

Compatible with the national population composition, more Malays and Chinese are present in the respective Malaysia and Singapore study populations. Demographically, the Singapore study population comprised higher proportions of those who are age 60 years (63.4% vs ta 54.6%) and older who were educated up to secondary level (87.0% vs 77.7%). Nearly half of the participants from both countries were treated with at least one antihy-≥ pertensive medication, with a significantly higher proportraining, tion of those from Malaysia's study population on three or more such medications (p=0.023). In terms of control, more than half of Singaporean participants attained BP and control goals based on JNC 8 guidelines, with fewer in Malaysia, attaining the mark.²¹

Singapore's participants in this study had significantly higher mean total HTN-SCP scores. In both countries, HTN-SCP behaviour, motivation and self-efficacy were technolog associated with Indian ethnicity and tertiary education. The HTN-SCP behaviour score was associated aged 60 years and above in both countries. The HTN-SCP motivation mean scores were associated with secondary education level in both countries. For Malaysia, Malay ethnicity was associated with higher HTN-SCP behaviour, motivation and self-efficacy scores. Other factors associated with HTN-SCP behaviour and mean self-efficacy scores were of other ethnicity and those with secondary education for Singapore. Women were associated with higher HTN-SCP behaviour mean scores in Singapore and HTN-SCP selfefficacy scores in Malaysia.

Table 4 Sum	imarv of factors in	fluencina Hv	vpertension Self-Care P	rofiles (HTN	N-SCP) behavic	our, motivation and	self-efficac	v scores		
	abayiour scores	ſ								
Countries	Malaysia					Singapore				
Variable	Behaviour domain score	P value	Adjusted beta (95% CI)	SE	P value	Behaviour domain score	P value	Adjusted beta (95% CI)	SE	P value
	Mean±SD					Mean±SD				
Age, years		0.052					0.068			
≥60	57.1 (9.1)		2.1 (0.7 to 3.4)	0.7	0.002*	59 (8.9)		2.5 (0.7 to 4.3)	0.9	0.007*
Gender		0.288					0.208			
Women	56.8 (8.8)		1		I	58.9 (9.0)		1.9 (0.1 to 3.6)	0.9	0.035*
Ethnicity		<0.001*					0.001*			
Chinese	55.6 (31.0)		Ref		I	57.9 (8.7)		Ref		I
Malay	58.3 (32.0)	0.751	3.2 (1.7 to 4.7)		<0.001*	55.8 (11.7)		to 0.9 (to 3.9 to 2.0)	1.5	0.534
Indian	59.7 (35.0)	0.906	4.4 (2.6 to 6.2)		<0.001*	60.7 (8.4)		3.3 (1.1 to 5.5)	1.1	0.003*
Others	0.0		1		I	68.0 (5.9)		9.3 (2.7 to 15.8)	3.3	0.006*
Education levi	0	<0.001*					0.004*			
Primary and below	56.8 (34.0)		Ref		I	56.5 (8.7)		Ref		I
Secondary	57.1 (31.0)		0.1 (to 1.3 to 1.5)	0.7	0.925	59.5 (9.1)		3.2 (1.3 to 5.1)	0.972	0.001*
Tertiary	62.1 (39.0)		4.3 (2.2 to 6.4)	1.1	<0.001*	59.5 (9.3)		4.2 (1.9 to 6.6)	1.217	<0.001*
HTN to SCP	notivation doma	in scores								
Countries	Malaysia					Singapore				
Variable	Motivation domain score	P value	Adjusted beta (95% CI)	SE	P value	Motivation domain score	P value	Adjusted beta (95% CI)	SE	P value
	Mean±SD					Mean±SD				
Gender		0.031*					0.208			
Women	65.8 (8.8)		2.1 (0.8 to 3.4)	0.7	0.002*	67.7 (10.5)		1.9 (to 0.1 to 3.8)	1.0	0.064
Ethnicity		<0.001*					0.013*			
Chinese	61.7 (9.6)		Ref		I	66.3 (10.2)		Ref		I
Malay	66.9 (7.9)		4.3 (2.9 to 5.8)	0.8	<0.001*	66 (13.7)	1.7	0.2 (to 3.1 to 3.5)		0.898
Indian	67.4 (8.5)		5.1 (3.4 to 6.8)	0.9	<0.001*	69.8 (8.7)	1.3	3.7 (1.3 to 6.2)		0.003*
Others	0.0					74 (7.6)	3.8	6.1 (to 1.4 to 13.6)	(0.109
Education level		<0.001*					<0.001*			
										Continued

6

BMJ Open: first published as 10.1136/bmjopen-2020-044192 on 14 June 2021. Downloaded from http://bmjopen.bmj.com/ on June 12, 2025 at Agence Bibliographique de I Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

Table 4 Con	tinued									
HTN to SCP b	ehaviour scores									
Countries	Malaysia					Singapore				
Variable	Behaviour domain score P va	ilue	Adjusted beta (95% CI)	SE	P value	Behaviour domain score	P value	Adjusted beta (95% CI)	SE	P value
Primary and below	62.6 (8.9)		Ref		1	64 (10.4)		Ref		1
Secondary	65.9 (8.7)		3.1 (1.7 to 4.6)	0.7	<0.001*	69.1 (10.1)		5.1 (2.9 to 7.2)	1.1	<0.001*
Tertiary	69.5 (8.4)		7.1 (5.0 to 9.2)	1.1	<0.001*	68.6 (9.8)		5.5 (2.9 to 8.2)	1.4	<0.001*
HTN to SCP :	self to efficacy domain	1 score	S							
Countries	Malaysia				Singapore					
Variable	Self to efficacy P va domain score	alue	Adjusted beta (95% CI)	SE	P value	Self to efficacy domain score	P value	Adjusted beta (95% CI)	SE	P value
	Mean±SD					Mean±SD				
Gender	0.01	2*					0.360	1		I
Women	63.2 (8.6)		1.7 (0.5 to 3.0)	0.649	0.007*	64.9 (10.3)				
Ethnicity	<0.0	101*					0.007*			
Chinese	59.2 (9.1)		Ref		I	63.7 (10.1)		Ref		I
Malay	63.5 (7.9)		4.0 (2.5 to 5.5)	0.748	<0.001*	63.2 (13.4)		0.1 (to 3.3 to 3.5)	1.733	0.975
Indian	65.6 (8.3)		6.2 (4.4 to 7.9)	0.888	<0.001*	67.1 (10.2)		3.7 (1.2 to 6.3)	1.298	0.004*
Others	0.0		I	I	I	73.6 (7.9)		8.4 (0.7 to 16.1)	3.931	0.033*
Education level	<0.0	01*					<0.001*			
Primary and below	60.6 (8.3)		Ref		I	61.9 (10.4)		Ref		I
Secondary	63 (8.7)		2.0 (0.6 to 3.4)	0.714	0.004	66 (10.8)		3.9 (1.7 to 6.1)	1.134	0.001*
Tertiary	65.3 (9.7)		4.8 (2.7 to 6.8)	1.054	<0.001*	66 (9.6)		4.2 (1.5 to 6.8)	1.361	0.002*
Univariate analy *P value <0.05.	/sis (t-test or ANOVA, anal) Variables from univariate a	ysis of <i>v</i> : analysis v	ariance). with p values of less than C).250 were in	cluded in the m	ultivariate analysis, ve	ariables which	I were not included we	re represented in	dashes (–).

6

BMJ Open: first published as 10.1136/bmjopen-2020-044192 on 14 June 2021. Downloaded from http://bmjopen.bmj.com/ on June 12, 2025 at Agence Bibliographique de I Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

7

Hypertension: the impact of self-care on health outcomes

In terms of control, three-quarter of the Singaporean participants in this study significantly attained BP control goals based on JNC 8 guidelines, with fewer in Malaysia reached the mark (73.6% vs 33.8%). The possible explanation could be due to the higher tertiary education background (22.3% vs 13% p=0.001) and lesser patients with diabetes (46.8% vs 65.8%, p<0.001) among study population from Singapore. In this study, Singapore participants attained significantly higher mean total HTN-SCP scores than their counterparts in Malaysia. These results apply to all the domains of the tool; behaviour, motivation and self-efficacy. Nearly two-thirds (62.2%) of Singaporean participants achieved BP goal versus one-third of those from Malaysia (34.5%). These findings were similar to the prevalence of BP controlled to target in population-based studies.¹⁷ Those with higher HPT-SCP scores reflects their higher levels of self-efficacy and self-care. Self-efficacy empowers them to take on daily self-care measures to control their BP and reduces cardiovascular risks.²²

The higher total HPT-SCP scores in Singapore patients may be partly due to its healthcare system and policy.²³ It is designed to enable the population to take on higher responsibility to manage their health through co-share healthcare financing, comprehensive individual and community empowerment and self-management programmes.^{24–26} Nevertheless, the implementation of these programmes^{27 28} remains challenging, with hindrance to reaching out to all patients, particularly those with lower health literacy and motivation.^{29 30}

Comparing self-care profiles

In this study, being of Indian ethnicity is associated with better self-care scores in all HPT-SCP domains for both countries than Chinese and Malay ethnicities. Despite good progress in healthcare accessibility, ethnic health disparity is still a challenge in both countries. While good self-care will result in better health outcomes, studies have shown that the incidence of metabolic syndromes, including raised BP, is high among Indian ethnicity with significant mortality risk in both Indian and Malay ethnicities.^{31–33} Although there may be a potential cultural influence of reporting desirable outcomes among patients of Indian ethnicity, further exploration may be of value to look at other factors, including the role of genetics on cardiovascular outcomes. We also found that Malay ethnicity in Malaysia's study population had better selfcare scores in all HPT-SCP domains than Singapore's study population. These findings may be related to ethnicity and medium of language used by healthcare staffs in the primary care setting. The majority of Malaysian healthcare staff are of Malay ethnicity with Malay language as the primary medium of communication. The similar medium of language used may have eased the access and understanding of health education across all aspect of self-care provided by the system.³⁴

Behaviour mean score was significantly associated with participants aged 60 years and above in both countries. ç

e

đ

Older patients with hypertension were reported to be more compliant in their BP monitoring and were more motivated to maintain weight.^{35 36} The longer duration of diagnosis increased their engagement with the health system over time. Thus, this might improve their knowledge about hypertension, and improved their coping skills to manage a chronic condition.

Studies have shown that higher education level is associated with adherence to self-care activities,^{30 37 38} as reflected in this study's results. Being more educated **u** allows people with hypertension to access and understand health information and resources to better manage their health better.^{11 39} They may be mindful of higher healthcare expenditure if they are hospitalised for 2 hypertension-related complications such as stroke and 8 maybe more conscious of the cost of maintaining their presence of the more conscious of the cost of maintaining their presence of the more gradient that the state of the more gradient of the more gr such complications. In this study, women had a significantly higher mean score for behaviour domain scores in Singapore and Malaysia's self-efficacy. Studies have found that women are likely to adopt the behaviour, leading to favourable lifestyle change and the self-efficacy to monitor their BP.^{38 40}

It has been shown that monitoring BP alone is not enough to improve cardiovascular outcomes.⁹ Self-care is not just about an individual responsibility to care for their health.⁴¹ Based on this study, self-care may be supported by the education policy and the healthcare system²³ Xt through better access to education and reducing gaps in health inequalities, that is, ethnicity and gender. Holistic management of hypertension is multi-faceted, including đ behaviour change approach and raising the motivation level. Enhancing self-efficacy to actualise self-care is one prerequisite for cost-effective and optimal long-term control of an individual's BP.

Strengths and limitations

mining, AI training, The unprecedented cross-national comparison of patients with hypertension that shares similar sociocultural background but different economic and health financing environment contributes to this study's strength. This is a study of comparison between two populations from two different countries, but not a comparison between two countries. It adds to the literature on the association between self-efficacy, self-care and BP treatment goal achievement. Large sample size has enabled in-depth analyses of the individual domain of the HTN-SCP. This study is not without limitations. The cross-sectional study **2** design does not allow causal effect relationship to be determined. The difference in assumptions (ie, conservative) resulted in different ways the sample size were calculated in both countries. Non-response rates adopted by the countries affected the sample size in each study. The small sample size in one country may inadequately power the study. As for recruitment, selection bias is inherent for convenience sampling in one study centre, suggesting that the results may not be generalisable to the national

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

population. Due to these reasons, the HTN-SCP domains mean scores comparison between the two countries must be interpreted with caution. We excluded an essential social variable, household income, due to the differences in how socio-economic categories being determined in each country. Malaysia has a national standard to categorise actual household income into three different levels (ie, low-income, middle-income and upper-income levels) and Singapore uses tax payment or housing type to ascertain this.

CONCLUSION

Patients with hypertension in Singapore's study population have a better overall self-care profile across behaviour, motivation and self-efficacy. In both study populations, being of Indian ethnicity and having tertiary education were predictors of higher self-care scores. Self-efficacy and skills in self-care are potentially modifiable. Future intervention to improve self-care among people with hypertension may need to be tailored to their behaviour, motivation and self-efficacy levels. This study's findings may be of interest for public health measures to tackle health inequality in multi-ethnic settings globally.

Author affiliations

¹SingHealth Polyclinics, Singapore

²Family Medicine Academic Clinical Programme, SingHealth Duke-NUS, Singapore
 ³Department of Family Medicine, Universiti Putra Malaysia, Serdang, Malaysia
 ⁴Malaysian Research Institute on Ageing, Universiti Putra Malaysia, Serdang, Malaysia

⁵UM eHealth Unit, University of Malaya, Kuala Lumpur, Malaysia

Acknowledgements We would like to thank all the participants of the study for their contribution. We also would like to thank Universiti Putra Malaysia for the support in publication under Dana Penerbitan Jurnal UPM (9001103).

Contributors HS and SY-MW contributed equally to this paper. HS, SY-MW, SSG, PYL, NCT, YLEK, SMC, MM, HA and NHS contributed to study design and planning, the data acquisition, the analysis and interpretation of data, drafted the first article and provided critical revision. HS and SY-MW had full access to the data and took responsibility for the study's overall conduct, the integrity of the data and accuracy of the analysis. All authors read and approved this article.

Funding This study was financially supported in part by the Malaysian Society of Hypertension (MSH RG/2017/HSS).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval We obtained ethical approval from the Medical Research & Ethics Committee of the Ministry of Health Malaysia (NMRR-17-1508-36071). The study was approved by the SingHealth Centralised Institutional Review Board (CIRB Reference number: 2017/2197).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The data used in this study are available from the Universiti Putra Malaysia and SingHealth Polyclinics. These data are not publicly available; therefore, restrictions apply as to its availability. However, the datasets during and/or analysed during the current study available from the corresponding author on reasonable request and with permission from Universiti Putra Malaysia and SingHealth Polyclinics.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content

includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Hani Salim http://orcid.org/0000-0002-7471-8678 Sazlina Shariff Ghazali http://orcid.org/0000-0002-5737-7226 Siew Mooi Ching http://orcid.org/0000-0003-0505-019X Ngiap Chuan Tan http://orcid.org/0000-0002-5946-1149

REFERENCES

- 1 Smullen A, Hong PK. Comparing the health care systems of highperforming Asian countries. *Asia Pac Policy Stud* 2015;2:347–55.
- 2 Kananatu K. Healthcare financing in Malaysia. *Asia Pac J Public Health* 2002;14:23–8.
- 3 von Eiff W, Massoro T, Voo YO, *et al*. Medical savings accounts: a core feature of Singapore's health care system. *Eur J Health Econ* 2002;3:188–95.
- 4 Bandura A. Social cognitive theory of self-regulation. Organ Behav Hum Decis Process 1991;50:248–87.
- 5 Roth GA, Abate D, Abate KH, et al. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the global burden of disease study 2017. *The Lancet* 2018;392:1736–88.
- 6 Institute For Public Health. National health and morbidity survey 2019, 2020. Available: http://iku.gov.my/index.php/nhms/nhms-2019
- 7 Koh KH, Goh CC, Goh SCP, et al. Blood pressure goal attainment in multi-ethnic Asian patients with hypertension and dyslipidaemia in primary care. Singapore Med J 2020;61:19.
- 8 Ab Majid NL, Omar MA, Khoo YY, et al. Prevalence, awareness, treatment and control of hypertension in the Malaysian population: findings from the National health and morbidity survey 2006–2015. J Hum Hypertens 2018;32:617–24.
- 9 Pearce G, Parke HL, Pinnock H, et al. The prisms taxonomy of self-management support: derivation of a novel taxonomy and initial testing of its utility. J Health Serv Res Policy 2016;21:73–82.
- 10 Fan AZ, Mallawaarachchi DSV, Gilbertz D, et al. Lifestyle behaviors and receipt of preventive health care services among hypertensive Americans aged 45 years or older in 2007. *Prev Med* 2010;50:138–42.
- 11 Salim H, Lee PY, Sazlina SG, et al. The self-care profiles and its determinants among adults with hypertension in primary health care clinics in Selangor, Malaysia. *PLoS One* 2019;14:e0224649.
- 12 Neminqani DM. Hypertensive patients: self-care management practices in Al-Taif, KSA. *Int J Sci Res* 2013;12:1705–14.
- 13 Lemeshow S, Hosmer DW, Klar J. Adequacy of sample size in health studies. Chichester: Wiley, 1990.
- 14 Orem DE, Taylor SG, Renpenning KML. *Nursing: concepts of practice*. Maryland Heights, Missouri: Mosby, 2001.
- 15 Miller WR, Rollnick S. Motivational interviewing: helping people change. New York: Guilford Publications, 2012.
- 16 Han H-R, Lee H, Commodore-Mensah Y, et al. Development and validation of the hypertension self-care profile: a practical tool to measure hypertension self-care. J Cardiovasc Nurs 2014;29:E11–20.
- 17 Koh YLE, Lua YHA, Hong L, *et al.* Using a web-based approach to assess test–retest reliability of the "Hypertension self-care profile" tool in an Asian population. *Medicine* 2016;95:e2955.
- 18 Seow KC, Mohamed Yusoff D, Koh YLE, et al. What is the test-retest reliability of the Malay version of the hypertension self-care profile self efficacy assessment tool? A validation study in primary care. BMJ Open 2017;7:e016152.
- 19 Ngoh SHA, Lim HWL, Koh YLE, et al. Test-retest reliability of the mandarin versions of the hypertension self-care profile instrument. *Medicine* 2017;96:e8568.
- 20 World Health Organization. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet* 2004;363:157–63.
- 21 James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the

Open access

panel members appointed to the eighth joint National Committee (JNC 8). *JAMA* 2014;311:507–20.

- 22 Tan FCJH, Oka P, Dambha-Miller H, et al. The association between self-efficacy and self-care in essential hypertension: a systematic review. BMC Fam Pract 2021;22:44.
- 23 Glynn LG, Murphy AW, Smith SM, et al. Interventions used to improve control of blood pressure in patients with hypertension. *Cochrane Database Syst Rev* 2010;7:CD005182.
- 24 Thumboo J, Yoon S, Wee S, *et al.* Developing population health research priorities in Asian City state: results from a multi-step participatory community engagement. *PLoS One* 2019;14:e0216303 –e03.
- 25 Nurjono M, Shrestha P, Ang IYH, *et al.* Implementation fidelity of a strategy to integrate service delivery: learnings from a transitional care program for individuals with complex needs in Singapore. *BMC Health Serv Res* 2019;19:177–77.
- 26 Tan ST, Quek RYC, Haldane V, et al. The social determinants of chronic disease management: perspectives of elderly patients with hypertension from low socio-economic background in Singapore. Int J Equity Health 2019;18:1.
- 27 Lee L-L, Arthur A, Avis M. Evaluating a community-based walking intervention for hypertensive older people in Taiwan: a randomized controlled trial. *Prev Med* 2007;44:160–6.
- 28 Kojuri J, Rahimi R. Effect of "no added salt diet" on blood pressure control and 24 hour urinary sodium excretion in mild to moderate hypertension. *BMC Cardiovasc Disord* 2007;7:34.
- 29 Lee J-E, Han H-R, Song H, *et al.* Correlates of self-care behaviors for managing hypertension among Korean Americans: a questionnaire survey. *Int J Nurs Stud* 2010;47:411–7.
- 30 Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/ American heart association Task force on clinical practice guidelines. J Am Coll Cardiol 2018;71:e127–248.

- 31 Hughes K, Yeo PP, Lun KC, et al. Cardiovascular diseases in Chinese, Malays, and Indians in Singapore. II. Differences in risk factor levels. J Epidemiol Community Health 1990;44:29–35.
- 32 Rampal S, Mahadeva S, Guallar E, et al. Ethnic differences in the prevalence of metabolic syndrome: results from a multi-ethnic population-based survey in Malaysia. PLoS One 2012;7:e46365.
- 33 de Hoog VC, Lim SH, Bank IEM, et al. Ethnic differences in clinical outcome of patients presenting to the emergency department with chest pain. Eur Heart J 2016;5:32–40.
- 34 Wittink H, Oosterhaven J. Patient education and health literacy. Musculoskelet Sci Pract 2018;38:120–7.
- 35 Hu H, Li G, Arao T. Prevalence rates of self-care behaviors and related factors in a rural hypertension population: a questionnaire survey. *Int J Hypertens* 2013;2013:1–8.
- 36 Niriayo YL, Ibrahim S, Kassa TD, et al. Practice and predictors of self-care behaviors among ambulatory patients with hypertension in Ethiopia. PLoS One 2019;14:e0218947.
- 37 Han H-R, Kim KB, Kang J, *et al.* Knowledge, beliefs, and behaviors about hypertension control among middle-aged Korean Americans with hypertension. *J Community Health* 2007;32:324–42.
- 38 Li W-W, Wallhagen MI, Froelicher ES. Hypertension control, predictors for medication adherence and gender differences in older Chinese immigrants. *J Adv Nurs* 2008;61:326–35.
- 39 Bacha D, Abera H. Knowledge, attitude and self-care practice towards control of hypertension among hypertensive patients on follow-up at St. Paul's Hospital, Addis Ababa. *Ethiop J Health Sci* 2019;29:421–30.
- 40 Martin MY, Person SD, Kratt P, et al. Relationship of health behavior theories with self-efficacy among insufficiently active hypertensive African–American women. *Patient Educ Couns* 2008;72:137–45.
- 41 Tucker KL, Sheppard JP, Stevens R, *et al.* Self-Monitoring of blood pressure in hypertension: a systematic review and individual patient data meta-analysis. *PLoS Med* 2017;14:e1002389.

ล