BMJ Open Global mapping of exercise interventions among COVID-19 survivors: protocol for a scoping review

Mukesh Kumar Sinha ,¹ Ajith Soman ,² Vaishali K ,¹ Rakesh Krishna Kovela,^{3,4} Monisha Lakshminarayanan,^{5,6} Revati Amin,¹ Ashwani Verma

ABSTRACT

survivors.

Introduction The COVID-19 pandemic has posed an

well-being. The Pan American Health Organization has

programmes for COVID-19 survivors. This scoping

acknowledged the crucial role of rehabilitation therapies

review outlines our strategy for systematically reviewing

volume of evidence for exercise interventions for COVID-19

published and unpublished literature to investigate the

Methods and analysis This scoping review will be

will be searched in August 2022 and articles will be

conducted based on the framework developed by Arksey

Web of Science, Cochrane Central, ProQuest and CINAHL)

considered for inclusion published till July 2022. Studies

will be screened by two independent reviewers at the title/abstract and full-text screening stages, as well as

data extraction and critical assessment. Data will be

extracted in a data extraction form and presented as

give a thorough understanding of the current literature

on exercise interventions for COVID-19 survivors, as

publication of this scoping review protocol.

Ethics and dissemination No human or animal

well as identify knowledge gaps that will guide future

figures and tables with narratives. This scoping review will

research. Preliminary searches will be conducted after the

participants were involved in this review. Therefore, ethical

disseminated through peer-review publications or through

committee approval is not required. Transparency will be

followed at every review stage. Review findings will be

and O'Malley. Six databases (PubMed/Medline, Scopus,

for COVID-19 and emphasised the importance of exercise

unparalleled threat to all dimensions of human health and

To cite: Sinha MK, Soman A, K V, *et al.* Global mapping of exercise interventions among COVID-19 survivors: protocol for a scoping review. *BMJ Open* 2022;**12**:e059987. doi:10.1136/ bmjopen-2021-059987

► Prepublication history for this paper is available online. To view these files, please visit the journal online (http://dx.doi. org/10.1136/bmjopen-2021-059987).

Received 08 December 2021 Accepted 18 October 2022



The

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to Dr Ashwani Verma; vamit1989@gmail.com INTRODUCTION The novel human COVID-19 is the fifth known pandemic since the 1918 influ-

conference presentations.

known pandemic since the 1918 influenza pandemic.¹ The WHO designated the COVID-19 outbreak as a public health emergency of international concern on 30 January 2020.² A recent report indicates that 26% of COVID-19 patients needed to be admitted to the hospital, with 14% requiring intensive care unit (ICU) admission and/or respiratory assistance.^{1 2} Individuals with a severe

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This is the first scoping review to map the existing exercise intervention literature among COVID-19 survivors.
- \Rightarrow This scoping review focused on effectiveness studies only.
- ⇒ Only English language studies will be considered for inclusion.

presentation of the illness are known to have underlying comorbidities and hospitalised patients require long durations of stay (around 4–53 days) or treatment in ICUs (around 4–19 days).^{3–5} Many patients are at risk of losing pulmonary functions and eventually having a poor health-related quality of life on discharge from hospitals.¹²

As the number of people recovering with COVID-19 rises, it is more important than ever to have a greater understanding of the healthcare challenges they confront. Previous training, research has shown that post-COVID-19 has an impact on multiple organs. Early data suggest that SARS-CoV-2 infection has left persistent effects such as fatigue, dyspnoea, myalgia, arthralgia, reduced cardio respira-<u>0</u> tory endurance, decreased functional exercise capacity and exercise tolerance, thereby limiting quality of life.^{6–15} All of these require long-term care and rehabilitation to regain functionality. Post-acute COVID-19 (Long COVID-19) is defined as persistent symptoms and/or delayed or long-term complications **g** beyond 4 weeks from the onset of symptoms.⁶

Prior research has indicated that COVID-19 survivors suffer long-term functional limitations following discharge.^{8–12} ¹⁶ A first case series undertaken by Carfi *et al* among the Italian population in 2020 reported the incidence of post-COVID-19 symptoms stands at 71.4% among post-COVID-19 survivors.¹⁷ Another cohort study conducted by Huang *et*

đ

e

data

Protected by copyright, including for uses related



alamong discharged COVID-19 patients in 2021 reported that 63% COVID-19 survivors were having musculoskeletal related symptoms with reduced 6 min walk distance.¹⁸ Impaired pulmonary diffusion capacities and abnormal radiographic features were evident among patients who were seriously ill during their hospital admissions. As a result, they were the major target group for long-term recovery interventions.¹⁸

COVID-19 rehabilitation research framework has been developed by the WHO Rehabilitation Programme and Cochrane Rehabilitation team to inform best practices and ensure that the health system offers its best to serve the COVID-19 survivors.¹⁹ The rehabilitation community has recently issued a 'call to action' to overcome multidomain impairments among COVID-19 survivors.^{8 20}

Exercise has been found to offer short and long-term health benefits in preventing, mitigating and even reversal of a wide range of metabolic, pulmonary, cardiovascular, neuropsychological, inflammatory and musculoskeletal illnesses. The benefits of exercise-induced immune response, including increased antipathogen activity, enhanced recirculation of anti-inflammatory cytokines and leucocytosis are relevant for fighting viral infections such as COVID-19.²¹

Rehabilitation professionals such as physiotherapists play a vital role in assessing and providing comprehensive pulmonary rehabilitation programme in long COVID-19 to restore functionality (cardiorespiratory, musculoskeletal).^{22 23} A preliminary report suggested the positive impact of pulmonary rehabilitation programme on dyspnoea and fatigue²⁴ among long COVID-19. However, the effectiveness of Pulmonary rehabilitation for long COVID-19 remains unclear.

Against this background, this scoping review will be conducted with an objective of mapping and collating the published literature on the exercise interventions among COVID-19 survivors.

METHODS

The present scoping review will be conducted in accordance with the Arksey and O'Malley's²⁵ framework which was further methodologically enhanced by the Levac et al²⁶ and Joanna Briggs Institute.²⁷ According to this framework, there are five main stages in undertaking a scoping review: (1) identification of research question; (2) identification of relevant studies; (3) study selection; (4) charting the data and (5) collate, summarise, and report the findings.

Stage 1: identify the research question

The research question for this scoping study was established by the team members after extensive brainstorming and further refinement of ideas. Based on the terminologies, preliminary literature search and current trends addressing long COVID-19 and exercises intervention, the research questions are presented in table 1. Also, narrative synthesis of change in outcome such as fatigue,

Table 1 Scoping review	research questions
Aspect	Research questions
Descriptive	How many studies published (year wise) on exercise intervention among COVID-19 survivors or long COVID? What is the region wise distribution (as per WHO classification) of published literature?
Methods	What type of study designs used in published literature? What are the research methods employed in published literature? What are the population characteristics available in published literature?
Exercise regimen	What is the type of exercise evaluated in published literature? What is the dose of exercise being reported in published literature?
Exercise and COVID-19 survivor or Long COVID	What is the impact of exercise on physical health among COVID-19 survivors as reported in published literature? What are the adverse events reported following exercise intervention (if any)?

dyspnoea, myalgia, arthralgia, reduced cardio respiratory endurance, decreased functional exercise capacity, exercise tolerance, health-related quality of life, quality of life, anxiety, stress, depression, self-esteem, mental health and mental well-being following intervention will be reported.

Stage 2: identifying relevant studies Based on Arksey and O'Malley's framework,²⁵ this stage entails the identification of the available literature. The **g** search will be based on the Medical Subject Heading (MeSH) terms (table 2) and the associated keywords identified through MeSH library.28 Drawing the help of an evidence specialist and a subject-matter expert, the search strategy (table 3) will be developed, which is likely to include a wide range of keywords and controlled vocabulary. Without using any time frames, a search of query will be developed in PubMed/Medline and this will **g** be reciprocated for additional specified databases such as $\overline{\mathbf{g}}$ Scopus, Web of Science, Cochrane Central, ProQuest and CINAHL. These databases have been included based on the authors' accessibility to them through their respective institutions and because they are comprehensive in terms of culminating information on exercise interventions. Studies will be screened for inclusion based on the Population, Concept and Context (table 4). Handsearch of the reference list of all included studies will be performed to ensure the inclusion of eligible and relevant studies.

Table 2 Search theme and terms		Table 3
Domain	Search terms	SI. no.
Population	Covid19 survivors, long COVID, post-acute COVID, COVID19, coronavirus survivors, COVID-19 patients, Coronavirus positive patients, Novel Coronavirus patients, 2019 Novel Coronavirus,	#1
Physical activity intervention Exercise, fitness, physical activity, physical education, physical fitness, aerobic exercise, resistance training, movement, physical excursion, exercise, mobility, walking, brisk walking, running, cycling, swimming, activity, basic activities, essential activities	#2	
	running, cycling, swimming, activity, basic activities, essential activities	#3
Outcomes Fatigue, dyspnea, myalgia, arthralgia, reduced cardio respiratory endurance, decreased functional exercise capacity, exercise tolerance, Health related Quality of Life, quality of life, anxiety, stress, depression, self-esteem, mental health and mental wollboing		
	Health related Quality of Life, quality of life, anxiety, stress, depression, self-esteem, mental health and mental wellbaing	#4
Study design	Randomized control trial, quasi experimental studies, before vs after studies, other matching designs	#5
		#6
		#7

Eligibility criteria for the inclusion of the studies are given below.

Inclusion criteria

All research articles eligible for selection must meet the following inclusion criteria:

- 1. Effectiveness studies (randomised control trials, quasiexperimental studies, other matching studies, before vs after study) were published in peer-reviewed journals.
- 2. Studies must be conducted among survivors of COVID-19 (Long COVID-19) after 4 weeks of recovery.
- 3. Must assess the exercise interventions (aerobic, resistance and concurrent exercise).
- 4. Studies conducted in the English language.

Exclusion criteria

- 1. Studies other than effectiveness studies will be excluded.
- 2. Reviews, reports, commentary and editorials will also be excluded.
- 3. Studies will be excluded where participants are cognitively not fit.

SI. no.	Search strategy
#1	"Covid19 patients" OR "Long Covid19" OR "post-acute Covid19" OR "Covid19 survivors" OR "Covid19" OR "coronavirus survivors" OR "coronavirus positive patients" OR "novel coronavirus patients" OR "2019 novel coronavirus" OR "2019-nCoV" OR "SARS- CoV-2"
#2	Exercise OR fitness OR "physical activity" OR "physical education" OR "physical fitness" OR "aerobic exercise" OR "resistance training" OR movement OR "physical excursion" OR exercise OR mobility OR walking OR "brisk walking" OR running OR cycling OR swimming OR activity OR "basic activities" OR "essential activities"
#3	"Health related Quality of Life" OR "quality of life" OR "physical health" OR tiredness OR fatigue OR pain OR myalgia OR "musculoskeletal pain" OR "exercise tolerance" OR rehabilitation OR "cardio respiratory endurance" OR "functional exercise capacity" OR anxiety OR stress OR depression OR "self-esteem" OR "mental health" OR "mental wellbeing"
#4	RCT OR Randomized control trial or quasi experimental study OR pre post study OR before vs after study OR experimental study OR effectiveness study
#5	#1 AND #2
#6	#4 AND #5
#7	#3 AND #5 AND #6

Search strategy for PubMed/Medline

Table 4 Eligibi	lity criteria for the studies
Population	COVID-19 survivors (Long COVID following 4 weeks recovery) aged above 18 years will be included.
Concept	Any exercise interventions (aerobic, resistance and concurrent exercise). Exercise dose must be reported as per frequency, intensity, type, time, volume and progression.
	Health-related quality of life, fatigue, musculoskeletal pain, exercise tolerance, cardiorespiratory fitness, mental health and well-being, emotional instability
Context	 All studies will be screened and reported based on: 1. WHO regional classification for countries. 2. At community level—both home based and centre based 3. Healthcare facilities

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Patient and public involvement

No patients were involved in this review.

Stage 3: study selection

This scoping review will follow a two-stage article selection process: (1) Title and abstract screening and (2) full-text screening using Rayyan.qcri.org software.²⁹ Inter-rater reliability will be calculated for the above two stages using Cohen's kappa statistics.³⁰ Searches will be conducted on the mentioned databases and duplicates will be removed making use of the Rayyan software. Both title and abstract screening, as well as full-text screening will be conducted independently by two reviewers (RA and RKK). Any discrepancy related to the inclusion or exclusion of the study among the reviewers will be resolved by discussion and mutual consensus. In the case of a disagreement among the reviewers, a third reviewer (MKS) will resolve it. Reporting guidelines and criteria set in Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews checklist will be used.³¹

Stage 4: charting the data

Arksey and O'Malley²⁵ demonstrated the fourth stage of the framework in charting the data from selected studies. To extract relevant data from the included studies, the study team (VK, MKS and AS) will design a data extraction tool in collaboration with subject knowledge experts with more than 7 years of experience in exercise prescription. These data collecting tool will be piloted for 5% of the studies to ensure that it captures the information needed for this scoping review by two independent authors (AV and MN). Two authors will extract data independently, and discrepancies will be solved by mutual consensus (RA and RKK). Data will be extracted on the following characteristics: (1) Bibliometric information (study title, authors, journal name, publication year, country, research setting, population characteristics like age, gender, body mass index, duration of hospitalisation if any); (2) Methodological characteristics (method of randomisation if applicable, sampling technique and sample size); (3) Intervention (type of exercise, method of delivery, the dose of exercise in terms of warm-up, coold own, frequency, intensity, type, time, volume and progression and type of comparator and (4) outcomes related to the quality of life, physical health, cardiorespiratory fitness and musculoskeletal pain.

Critical appraisal

All included studies will be critically appraised using Joanna Briggs's critical appraisal tool.³² Two authors (AV and MN) will independently assess the study quality and resolve disagreements through discussion and mutual consensus. The overall risk assessment will be presented in a tabular or graphical manner and will be described through systematic narrative synthesis.

Stage 5: collating, summarising and reporting the results

The descriptive and narrative summaries developed based on the findings of this scoping review will be documented in a report format that follows the Joanna Briggs Institute Reviewer's manual recommendation.²⁷ The key components of the domains in the included studies, such as study area, exercise dosage in the form of warm-up, cool down, frequency, intensity, type, time, volume and progression, sample size, and research outcomes, will be progression, sample size, and research outcomes, will be visualised using summary tables. Other relevant findings will be presented in a tabular manner with appropriate charts (bar and pie). DISCUSSION Residual changes in affected lungs and reduced quality of life (physical, health related) are evident among

of life (physical, health related) are evident among COVID-19 survivors.³³ Due to these clinical manifestations, exercise interventions play a significant role in the rehabilitation of COVID-19 survivors. This scoping review will map and collate the available literature on exercise interventions among COVID-19 survivors.

Author affiliations

¹Department of Physiotherapy, Manipal College of Health Professions, Manipal Academy of Higher Education, Manipal, Karnataka, India

²Department of Health Rehabilitation, College of Applied Medical Sciences, Shagra University, Shagra, Saudi Arabia

³Department of Neuro-physiotherapy, Datta Meghe Institute of Medical Sciences Sawangi (Meghe), Wardha, Maharashtra, India

⁴Nitte Institute of Physiotherapy, NITTE (Deemed to be University), Deralakatte, Mangaluru, Karnataka, India

⁵Delhi University, Delhi, India

⁶Campbell South Asia, Delhi, India

⁷School of Pharmaceutical and Populations Health Informatics, DIT University, Dehradun, Uttarakhand, India

Contributors All authors have made significant contribution in this scoping review manuscript. VK, MKS, RKK, MN, RA and AV have conceptualised this scoping review. AS provided the technical inputs for drafting the introduction and methodology section of the manuscript. All authors were involved in drafting and revising the manuscript

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work noncommercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/ licenses/by-nc/4.0/.

ORCID iDs

Mukesh Kumar Sinha http://orcid.org/0000-0002-7407-7153 Ajith Soman http://orcid.org/0000-0001-9890-2552 Vaishali K http://orcid.org/0000-0001-5746-1682

Open access

Lir ne J 2 nat Cou an j 3 ills for an err

- 16 Gentil P, de Lira CAB, Coswig V, *et al.* Practical recommendations relevant to the use of resistance training for COVID-19 survivors. *Front Physiol* 2021;12:637590.
- 17 Carfi A, Bernabei R, Landi F, et al. Persistent symptoms in patients after acute COVID-19. JAMA 2020;324:603.
- 18 Huang C, Huang L, Wang Y, *et al.* 6-Month consequences of COVID-19 in patients discharged from Hospital: a cohort study. *Lancet* 2021;397:220–32.
- 19 Negrini S, Mills J-A, Arienti C, *et al.* "Rehabilitation Research Framework for Patients With COVID-19" Defined by Cochrane Rehabilitation and the World Health Organization Rehabilitation Programme. *Arch Phys Med Rehabil* 2021;102:1424–30.
- 20 Stam HJ, Stucki G, Bickenbach J, *et al.* Covid-19 and post intensive care syndrome: a call for action. *J Rehabil Med* 2020;52:jrm00044.
- 21 Amatriain-Fernández S, Gronwald T, Murillo-Rodríguez É, et al. Physical exercise potentials against viral diseases like COVID-19 in the elderly. Front Med 2020;7:379.
- 22 Spruit MÁ, Holland AE, Singh SJ, et al. COVID-19: interim guidance on rehabilitation in the hospital and Post-Hospital phase from a European respiratory Society and American thoracic Societycoordinated international Task force. *Eur Respir J* 2020;56:2002197.
- Righetti RF, Onoue MA, Politi FVA, et al. Physiotherapy Care of Patients with Coronavirus Disease 2019 (COVID-19) - A Brazilian Experience. Clinics 2020;75:e2017.
- 24 Practice Update. ERS 2021: Pulmonary Rehabilitation Improves Long COVID Symptoms in Preliminary Study [Internet], 2021. Available: https://www.practiceupdate.com/content/ers-2021-pulmonaryrehabilitation-improves-long-covid-symptoms-in-preliminary-study/ 123701/8/20/2
- 25 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8:19–32.
- 26 Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010;5:69.
- 27 Peters MDJ, Marnie C, Tricco AC, et al. Updated methodological guidance for the conduct of scoping reviews. *JBI Evid Synth* 2020;18:2119–26.
- 28 U.S. National Library of Medicine. Medical subject Headings, 2021 2021. Available: https://meshb.nlm.nih.gov/ [Accessed 24 Mar 2022].
- 29 Ouzzani M, Hammady H, Fedorowicz Z, *et al*. Rayyan-a web and mobile APP for systematic reviews. *Syst Rev* 2016;5:210.
- 30 Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas* 1960;20:37–46.
- 31 Tricco AC, Lillie E, Zarin W, *et al.* PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018;169:467–73.
- 32 Joanna Briggs Institute. Critical Appraisal Tools [Internet]. Available: https://jbi.global/critical-appraisal-tools [Accessed 24 Mar 2022].
- 33 Strumiliene E, Zeleckiene I, Bliudzius R, et al. Follow-Up analysis of pulmonary function, exercise capacity, radiological changes, and quality of life two months after recovery from SARS-CoV-2 pneumonia. *Medicina* 2021;57:568.

REFERENCES

- 1 Liu Y-C, Kuo R-L, Shih S-R. COVID-19: the first documented coronavirus pandemic in history. *Biomed J* 2020;43:328–33.
- 2 World Health Organizaion. COVID-19 Public Health Emergency of International Concern (PHEIC) Global research and innovation forum [Internet], 2020. Available: https://www.who.int/publications/m/ item/COVID-19-public-health-emergency-of-international-concern-(pheic)-global-research-and-innovation-forum
- 3 Wang B, Li R, Lu Z, *et al.* Does comorbidity increase the risk of patients with COVID-19: evidence from meta-analysis. *Aging* 2020;12:6049–57.
- 4 Rees EM, Nightingale ES, Jafari Y, et al. COVID-19 length of hospital stay: a systematic review and data synthesis. BMC Med 2020;18:270.
- 5 Postigo-Martin P, Cantarero-Villanueva I, Lista-Paz A, *et al.* A COVID-19 rehabilitation prospective surveillance model for use by physiotherapists. *J Clin Med* 2021;10:1691.
- Nalbandian A, Sehgal K, Gupta A, et al. Post-acute COVID-19 syndrome. Nat Med 2021;27:601–15.
- 7 Zhu Y, Wang Z, Zhou Y, et al. Summary of respiratory rehabilitation and physical therapy guidelines for patients with COVID-19 based on recommendations of World Confederation for physical therapy and national association of physical therapy. J Phys Ther Sci 2020;32:545–9.
- Kemp HI, Corner E, Colvin LA. Chronic pain after COVID-19: implications for rehabilitation. *Br J Anaesth* 2020;125:436–40.
- 9 Zhao Y-M, Shang Y-M, Song W-B, et al. Follow-Up study of the pulmonary function and related physiological characteristics of COVID-19 survivors three months after recovery. *EClinicalMedicine* 2020;25:100463.
- 10 Moradi Y, Mollazadeh F, Karimi P, et al. Psychological disturbances of survivors throughout COVID-19 crisis: a qualitative study. BMC Psychiatry 2020;20:594.
- 11 Pant P, Joshi A, Basnet B, et al. Prevalence of functional limitation in COVID-19 recovered patients using the post COVID-19 functional status scale. JNMA J Nepal Med Assoc 2021;59:7–11.
- 12 Taboada M, Moreno E, Cariñena A, *et al.* Quality of life, functional status, and persistent symptoms after intensive care of COVID-19 patients. *Br J Anaesth* 2021;126:e110–3.
- Rudroff T, Fietsam AC, Deters JR, *et al.* Post-COVID-19 fatigue: potential contributing factors. *Brain Sci* 2020;10:E1012.
- 14 Phebe Tucker MD, Christopher S, Czapla MD. Post-COVID stress disorder: another emerging consequence of the global pandemic, 2021. Available: https://www.psychiatrictimes.com/view/post-covidstress-disorder-emerging-consequence-global-pandemic [Accessed 24 Mar 2022].
- 15 World Health Organizaion. Coronavirus disease Answers [Internet]. Available: https://www.who.int/emergencies/diseases/novelcoronavirus-2019/coronavirus-disease-answers [Accessed 24 Mar 2022].