BMJ Open Development and evaluation of a psychosocial adaptation questionnaire for women with breast cancer treated with adjuvant endocrine therapy: a singlecentre, cross-sectional study in China

Haoran Jiang ⁽¹⁾, ¹ Xiujie Zhang ⁽¹⁾, ² Yu Dong, ¹ Hui Xu, ³ Feng Jin¹

ABSTRACT

Objective To develop and evaluate a psychosocial adaptation (PSA) questionnaire of women with breast cancer treated with adjuvant endocrine therapy (AET). Design Cross-sectional study.

Setting Conducted study in a tertiary care centre in China from March to June 2021.

Participants Women with breast cancer who have received AET.

Methods Item development and extraction were based on literature, qualitative interviews (n=16) and assessment of content validity. 300 participants were recruited to perform item analysis, internal consistency reliability and exploratory factor analysis by cross-sectional studies. Finally, test-retest, criterion validity and confirmatory factor analysis of the questionnaire were performed with 96 respondents.

Results An initial questionnaire was devised. The exploratory factor analysis demonstrated four factors: emotional response (six items), self-worth (five items), physical impacts (three items) and social communication (two items). The cumulative contribution of the factors was 65.057%. Cronbach's alpha coefficient of the questionnaire was 0.876. The correlation coefficient between the questionnaire and General Self-Efficacy Scale was 0.565 (p<0.01). A 16-item guestionnaire and its factorial structure were confirmed.

Conclusions The 16-item questionnaire had good reliability and validity. This could be a useful tool to assess the PSA levels among women with breast cancer treated with AET and provide a basis for further research.

INTRODUCTION

Breast cancer (BC) is the most diagnosed cancer globally, with 2.3 million new cases in 2020, accounting for 11.7% of all cancer cases in that year .¹ BC accounts for one in four cancer cases and one in six cancer deaths among women. Approximately two-thirds of women with BC test positive for the oestrogen receptor (ER) and/or progesterone receptor (PR).² Due to early diagnosis and aggressive adjuvant therapy, the 5-year survival rate for

STRENGTHS AND LIMITATIONS OF THIS STUDY

- \Rightarrow The 16-item questionnaire can be answered quickly and would be easy to administer even in busy centres providing breast cancer treatment.
- \Rightarrow The questionnaire can be used as a comprehensive and cost-effective clinical assessment tool and to further study the mood, self-perception, self-image, social dimension and overall psychosocial adaptation level of patients with breast cancer receiving adjuvant endocrine therapy.
- \Rightarrow This study included only one centre in China. Therefore, it is not generalisable.
- \Rightarrow External validity needs further evaluation using a diversified sample.
- \Rightarrow The use of a self-report may have influenced the answers of the participants to show their intentions as more 'morally acceptable'.

ng, cancer is more than 91.2%, which is higher than that for other cancers.³⁴ Updated clinical practice guidelines recommend extending the use of adjuvant endocrine therapy (AET) from 5 to 10 years.²⁵

Breast cancer has gradually become a chronic disease with higher survival rates. <u>0</u> Breast cancer survivors (BCS) need to co-exist with cancer for a long time and learn to adapt to the disease.⁶ While physical functioning in quality of life improves over time after completion of aggressive cancer treatment, emotional and social functioning as well as $\mathbf{\hat{G}}$ some breast cancer-specific symptoms significantly worsen, with all symptoms of distress (except pain) being higher in breast cancer patients than in the general population.⁷ Compared with the general population, after AET, BCS experience physical symptoms and emotional issues as they enter remission.⁸ Long period of AET brings immense pain to the patient's body and mind. They often worry about disease recurrence and metastasis, and

≥

training

Protected by copyright, including for uses related to text and data mini

Dong Y, et al. Development and evaluation of a psychosocial adaptation questionnaire for women with breast cancer treated with adjuvant endocrine therapy: a singlecentre, cross-sectional study in China. BMJ Open 2022;12:e063082. doi:10.1136/ bmjopen-2022-063082

To cite: Jiang H. Zhang X.

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (http://dx.doi.org/10.1136/ bmjopen-2022-063082).

Received 24 March 2022 Accepted 17 September 2022

Check for updates

C Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Breast Surgery, First Affiliated Hospital of China Medical University, Shenyang, Liaoning, China ²Nursing department, First Affiliated Hospital of Dalian Medical University, Dalian, Liaoning China ³Breast Surgery, Cancer Hospital of China Medical University, Shenyang, Liaoning, China

Correspondence to Professor Feng Jin; jinfeng@cmu.edu.cn

Table 1	Sociodemographic and clinical characteristics of
the partic	pipants (n=396)

Characteristics	n (%)/mean±SD
Age(year)	50.628±9.024
BMI	24.112±2.903
Education level	
Junior high school or below	108 (27.272)
High school/college	190 (47.980)
Undergraduate college or above	98 (24.748)
Marital status	
Married	339 (85.606)
Unmarried	9 (2.273)
Divorced	31 (7.828)
Widowed	17 (4.293)
Professional	
Retired	166 (41.920)
Did not work	99 (25.000)
Temporary worked	13 (3.283)
Work as usual	118 (29.798)
Years taking endocrine medications	3.333±2.152
BMI, body mass index.	

most have experienced hot flashes, night sweats, insomnia, stiff joints, weight gain, anxiety, depression, and a series of side effects.¹⁰⁻¹⁴ In addition, BCS also suffer from a variety of psycho-social distress due to self-blame (negative perceptions of cancer) and social stigma.⁸ Adaptation on a multi-dimensional level, such as social and sexual, while tolerating the symptoms of diagnosis and treatment is called psycho-social adaptation.¹⁵ Psycho-social adaptation (PSA) in BCS is an important part of BCS evaluation.¹⁶

Londono and McMillan used literature analysis to form the intermediate domain theory of the concept of PSA, showing its multidisciplinary characteristics and properties of variability, process, persistence, interac-tivity and individuality.^{17 18} PSA involves the integration of illness or disability into an individual's life, identity, self-concept and self-image.¹⁹ Based on the perspective of positive psychology, PSA refers to the transition process of the person with disability from a state of disability to an enabling state, which is characterised by the transition from negative to positive happiness.²⁰ PSA has both relation and difference with self-efficacy and health-related quality of life. BCS with an optimistic disposition, and with a positive perception of illness have better health-related quality of life.²¹ Self-efficacy is important for better PSA.²² Based on previous theories and studies, when women have BC treated with AET, they will adopt either a positive or negative attitude and behaviour to face it, which results in the changes of emotional dimension, self-cognition, selfimage and social dimension. After an extensive literature

<page-header><page-header><text><text><text><text><text><text><text>

conducted through interviews and participatory observation.²⁷ The literature reviewed and the information **3** obtained from the participants (n=16) provided abundant data for the development of the item pool and the initial questionnaire. The PSA of women with BC treated with AET was closely related to emotional response, selfcognition, self-image and social situation.

Questionnaire development

Five experts (two BC specialists, two psychologists and one nursing specialist) assessed the questionnaire's 6

Table 2	Results of reliability and validity analysis for the PSA que	stionnaire am	iong BC women	with AET			
Factor	Item	Factor loading	Eigenvalue	Explanatory variation (%)	Cumulative explanatory variation (%)	Cronbach's alpha coefficient	Correlation coefficient
-	Q1. I am worried about metastasis or recurrence	0.845	5.920	36.997	36.997	0.899	0.857*
	Q2. Breast problems make me feel nervous	0.802					
	Q3. I am worried that breast disease will get worse	0.800					
	Q4. Breast problems make me feel scared	0.795					
	Q5. I am worried about the side effects of the drug	0.705					
	Q6. Breast problems make me very depressed	0.670					
2	#Q7. I still think of myself as an attractive person	0.784	1.954	12.211	49.208	0.790	0.685*
	#Q8. Overall, I am satisfied with myself	0.761					
	#Q9. I think I am still a valuable person	0.738					
	#Q10. I can share happiness and sadness with friends and colleagues	0.660					
	#Q11. I am still interested in things I was interested in before	0.649					
e	Q12. Breast problems affect my sex life	0.795	1.468	9.174	58.382	0.664	0.698*
	Q13. I can not face the weight gain	0.681					
	Q14. Disease affects my appearance and makes me unhappy	0.638					
4	Q15. I am not willing to discuss breast problems with others	0.861	1.068	6.675	65.057	0.730	0.573*
	Q16. I do not like others asking about my condition	0.839					
Confirm: *At level AET, adju	atory factor analysis. 0.01 (double tail), the correlation is significant. # is the reverse scoring uvant endocrine therapy; BC, breast cancer; PSA, psychosocial adapt.	j item. ation.					



Scree plot. Figure 1

content validity through two rounds of panel discussions. In preparation for the formal survey, participants who met the criteria were chosen prior to the survey, and the difficulty, suitability and complexity of the items were assessed.

Investigation

There are two parts to the investigation. The sample size was determined by the item number, which met the requirement of 5-10 times the item number. In the first part, the initial questionnaire contained 56 items (56×5=280). A total of 300 participants (>280) were included to perform item analysis, internal consistency reliability and exploratory factor analysis (EFA). In the second part, the final questionnaire contained 16 items (16×5=80). Test-retest, criterion validity and confirmatory factor analysis (CFA) of the questionnaire were performed with 96 respondents (>80).

Sociodemographic characteristics of the participants were obtained in the survey, which included populationrelated and disease-related variables. The PSA questionnaire was developed for women with BC who have received AET. The answers were scored on a five-point Likert scale: all the time, often, sometimes, rarely and never. The higher the score, the better was the PSA level. The 10-item General Self-Efficacy Scale (GSES), developed by Schwarzer in 1981, has been widely used to measure self-efficacy and has been shown to have good reliability and validity.

The items were analysed using the critical ratio, discriminative index (DI) and item-total correlation (ITC). Items with values below the standard value were deleted. Construct validity was assessed using EFA and CFA. The internal consistency of the questionnaire was evaluated using Cronbach's alpha coefficient. Test-retest reflects the stability and consistency of the test over time. After 10-18 days, 31 patients completed the questionnaire again, and the correlation coefficients of the two measures were used to assess the reliability of the retest. The Spearman correlation coefficient test was used to assess the validity of scale stability. The GSES was used to assess criterion validity.²⁸

Statistical analysis

If an item was not answered, the score was replaced with the average. If two or more questions were not answered, the questionnaire was excluded from analysis. In the qualitative part, all data were managed and analysed using NVivo V.11.0. In the quantitative stage, IBM SPSS Statistics 25.0 and AMOS 23.0, were used for statistical analysis.

Patient and public involvement

No patient or public involvement.

RESULTS

Participant characteristics

A total of 409women with BC treated with AET were selected between May and June 2021, of which 396 patients agreed to participate (response rate=96.82%). The sociodemographic and clinical characteristics of the study participants are presented in table 1.

Development and analysis of items

Item pool

The initial questionnaire included items related to the literature, qualitative research and group discussion.



Figure 2 The standardised estimate of each coefficient in the confirmatory factor analysis model.

പ്പ

Items			Estimate	SE	Z	P value	Estimate	CR	AVE
Q1	<	F1	1				0.781	0.887	0.577
Q2	<	F1	1.076	0.109	9.869	***	0.908		
Q3	<	F1	1	0.12	8.309	***	0.79		
Q4	<	F1	1.088	0.118	9.243	***	0.86		
Q5	<	F1	0.575	0.131	4.392	***	0.452		
Q6	<	F1	0.806	0.117	6.904	***	0.678		
Q7	<	F2	1				0.779	0.803	0.461
28	<	F2	0.781	0.126	6.216	***	0.669		
29	<	F2	0.91	0.123	7.42	***	0.819		
Q10	<	F2	0.494	0.13	3.791	***	0.416		
Q11	<	F2	0.78	0.132	5.921	***	0.639		
212	<	F3	1				0.371	0.632	0.390
213	<	F3	0.973	0.329	2.953	0.003	0.542		
214	<	F3	1.891	0.598	3.164	0.002	0.86		
Q15	<	F4	1				0.727	0.747	0.596
Q16	<	F4	1.308	0.378	3.463	***	0.815		
**p<0.00	1. Ivant endoci	rine therapy	/; BC, breast canc	er; CR, critical	ratio; PSA, ps	ychosocial adar	otation.		

Content validity was evaluated by five specialists (three breast specialists and two psychologists) through two rounds of panel discussions. The content validity index (CVI) of 76 questionnaires ranged from 0.769 to 0.893, and the average of the first round was 0.832. The content validity after deleting the 19 items was 0.930. A pre-experiment was conducted with 37 women with BC receiving AET to test the reliability of the questionnaire contents, of which 31 completed questionnaires were returned (recovery rate: 83.78%). Women with BC receiving AET judged the content of the questionnaire for appropriateness and accuracy. Cronbach's alpha coefficient of the questionnaire was 0.934 (p<0.05).

Item analysis

After an independent sample t-test, item 2, which proved that the discrimination was poor, was deleted. Items with DI and ITC <0.3 were deleted at the same time. Consequently, a total of 17 items were deleted.

Exploratory factor analysis

Twenty-three items were deleted in 21 rounds of EFA. From the perspective of factor load and content, the first subscale was 'emotional response' consisting of six items. The second subscale was 'self-worth', which consisted of five items. The third subscale was 'physical impacts', consisting of three items, and the fourth subscale was social communication', consisting of two items (table 2, online supplemental file 1). The Kaiser-Meyer-Olkin (KMO) index of the 16-item questionnaire was 0.880, and the result of Bartlett's spherical test was 2129.914 (p=0.000), which was suitable for factor analysis. When the eigenvalue was >1.0 (figure 1), four factors were \vec{a} extracted through principal factor analysis and varimax ⋗ rotation, and the cumulative contribution rate of the factors was 65.057%.

CFA was used to further evaluate the construct validity, ھ and the results are shown in figure 2. Except for Item 10, the load coefficient of the standardised factor obtained was >0.50. The path coefficients of the four subscales were highly correlated (p<0.001, table 3). The discrimination degree of the questionnaire and common fitting indices of the CFA model are listed in table 4, and the Pearson nologies correlation coefficient between subscales is 0.165-0.760 (p<0.001).

Table 4 Discrimination and model fit of confirmatory factor analysis for the questionnaire										
Items	Emotional response	Self- worth	Physical impacts	Social communication	CMIN/DF	RMSEA	CFI	PNFI	AIC	
Emotional response	0.760				1.366*	0.062	0.941	0.667	209.819	
Self-worth	0.290	0.679								
Physical impacts	0.655	0.478	0.625							
Social communication	0.399	0.165	0.405	0.772						
*p<0.05.										

ar

tra

Psychometric properties

Internal consistency

The Cronbach's alpha coefficient of the PSA questionnaire among women with BC treated with AET and the four subscales were 0.876, 0.899, 0.790, 0.664 and 0.730, respectively (table 2). Correlation analysis showed that the correlation coefficients between each dimension and the total questionnaire ranged from 0.573 to 0.857 (p<0.01; table 2).

Test-retest reliability

After 10–18 days, 30 patients answered the questionnaire again, with a correlation coefficient of 0.846 for both measures (p<0.01).

Convergent validity

The correlation coefficient between the PSA questionnaire among women with BC treated with AET and the GSES was 0.565 (p<0.01).

DISCUSSION

According to the adaptive model theory proposed by Roy, an American nursing theorist, individuals determine the best coping strategies based on objective analysis and thinking of various stress stimuli based on self-cognition and emotional state evaluation, so as to adopt adaptive response behaviours.²⁹ Based on Roy's adaptation theory and the PSA defined by Londono and McMillan, a literature review found that PSA is associated with psychological, social and other factors.^{17 30} Qualitative analysis found that the self-cognition of women with BC treated with AET played a crucial role in the process of PSA. Based on this, a questionnaire for women with BC treated with AET was developed, which consisted of sixteen items and four subscales (emotional response, self-worth, physical impacts and social communication). According to the indicators of the CFA value, the questionnaire was considered to have good compatibility from three aspects: Goodness of Fit Index (GFI), adjusted GFI and contracted GFI.³¹

The item pool was derived from reviews, qualitative research and two rounds of panel discussions. The DI and ITC among the 16 items of the PSA questionnaire among women with BC treated with AET was >0.30, which confirms that the entries are sufficiently differentiated and relevant.^{32 33}

In practice, the four-factor structure can display the connotation of PSA in patients with BC with AET more finely and intuitively. In short, these statistical correlations and structural relationships can be reasonably explained by logical relationships.³⁴ The KMO of the 16 question-naires was 0.880, which was considered suitable for factor analysis.³⁵ Four subscales were extracted, and the factor load of each item of the questionnaire was >0.60.³⁶

The correlation coefficient between each subscale was smaller than that between each subscale and the overall questionnaire, indicating that the correlation coefficient between each subscale and the overall question naire was significant. $^{\rm 37}$

Standard correlation validity reflects the level of consistency between the research tools and other measurement criteria. The 16-item questionnaire was positively correlated with GSES,³⁸ indicating that the lower the PSA level of psychosocial patients with BC assisted by endocrine therapy is, the lower is their self-efficacy. The correlation coefficient of standard correlation validity was 0.4–0.8, indicating a good level of correlation.^{39 40} The correlation coefficient between the 16-item questionnaire and the GSES met this.

By convention, an alpha of 0.60 or higher, is needed **b** to keep an item on a 'sufficient' scale.^{41 42} Therefore, the 16-item questionnaire was considered to have good pinternal consistency. Test–retest reliability could reflect the stability of the questionnaire, which was used to test the consistency of the same tool, and the same participants were evaluated twice or more.⁴³ The test–retest reliability of 16-item questionnaire was high.⁴¹ To sum up, this questionnaire had good reliability.

From a clinical perspective, the 16-item questionnaire **for** can be answered quickly; that is, it can effectively and pertinently evaluate the PSA of women with BC treated with AET, and is easy to administer in busy centres practicing treatment for BC. Women with BC treated with AET will be completed directly, and their PSA will be assessed by their healthcare provider based on the scoring method listed below. The questionnaire can be used as a comprehensive and cost-effective clinical assessment tool and overall PSA level of patients with BC with AET.

The use of a self-report may have influenced the answers of the participants to show their intentions as more 'morally acceptable'. The 16-item questionnaire mainly focused on the subjective feelings of patients, which can easily produce measurement bias, and there may be differences between subjective and objective indicators. Further evaluation of external validity is required. Therefore, we encourage action to acculturate, translate and validate the psychosocial adjustment questionnaire for women with BC receiving AET and to refine the normative component by adding descriptive and ethical norms.^{35 41}

The reliability and validity of the 16-item questionnaire, which can be used as a tool for clinical evaluation and further study, were verified in this study. This will help healthcare providers to evaluate PSA in women with AET for BC and provide an appropriate basis for women with AET to adapt to it and improve their quality of life.

Acknowledgements The authors thank the breast cancer specialists, psychologists and statisticians who were consulted during the development of the questionnaire and all the patients who participated in the study

Contributors HJ and FJ wrote the article; XZ, YD, HX and FJ designed the research; HJ performed the research; HJ, XZ and YD analysed the data. All authors approved the final version of the article, including the authorship list. FJ is guarantor for the work.

Open access

Funding Study on psychosocial factors influencing quality of life in breast cancer patients. This work was supported by the Science and Technology Department of Liaoning Province (2008225010-3)

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval This study involves human participants and was approved by This study was approved by the ethics committee at First Affiliated Hospital of China Medical University (Project No.2016-5-2). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Not applicable.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Haoran Jiang http://orcid.org/0000-0002-4792-6607 Xiujie Zhang http://orcid.org/0000-0002-0873-7977

REFERENCES

- Sung H, Ferlay J, Siegel RL, *et al*. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71:209–49.
- 2 Burstein HJ, Lacchetti C, Anderson H, et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: ASCO clinical practice guideline focused update. J Clin Oncol 2019;37:423–38.
- 3 Pagani O, Francis PA, Fleming GF, et al. Absolute improvements in freedom from distant recurrence to tailor adjuvant endocrine therapies for premenopausal women: results from text and soft. J Clin Oncol 2020;38:1293–303.
- 4 Ruhstaller T, Giobbie-Hurder A, Colleoni M, et al. Adjuvant letrozole and tamoxifen alone or sequentially for postmenopausal women with hormone receptor-positive breast cancer: long-term follow-up of the big 1-98 trial. J Clin Oncol 2019;37:105–14.
- 5 National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology. Breast cancer version 4, 2020. Available: https://www. nccn.org [Accessed 31 May 2020].
- 6 Hsiao F-H, Kuo W-H, Jow G-M, *et al*. The changes of quality of life and their correlations with psychosocial factors following surgery among women with breast cancer from the post-surgery to posttreatment survivorship. *Breast* 2019;44:59–65.
- 7 Finck C, Barradas S, Zenger M, *et al.* Quality of life in breast cancer patients: associations with optimism and social support. *Int J Clin Health Psychol* 2018;18:27–34.
- 8 van Vulpen JK, Peeters PHM, Velthuis MJ, *et al.* Effects of physical exercise during adjuvant breast cancer treatment on physical and psychosocial dimensions of cancer-related fatigue: a meta-analysis. *Maturitas* 2016;85:104–11.
- 9 Pitman A, Suleman S, Hyde N, *et al.* Depression and anxiety in patients with cancer. *BMJ* 2018;361:k1415.
- 10 Sebaoun P, Frederic M, Weryha G, et al. [Tolerance of Tamoxifen as an adjuvant therapy and long-term follow up of 55 premenopausal breast cancer women, cared for at the Institut de cancérologie de Lorraine, treated with Tamoxifen]. Bull Cancer 2019;106:S75–100.

- 11 Kuba S, Ishida M, Nakamura Y, *et al.* Persistence and discontinuation of adjuvant endocrine therapy in women with breast cancer. *Breast Cancer* 2016;23:128–33.
- 12 Toivonen KI, Williamson TM, Carlson LE, et al. Potentially modifiable factors associated with adherence to adjuvant endocrine therapy among breast cancer survivors: a systematic review. *Cancers* 2020;13. doi:10.3390/cancers13010107. [Epub ahead of print: 31 12 2020].
- 13 Choo SB, Saifulbahri A, Zullkifli SN, et al. Adjuvant endocrine therapy side-effects among postmenopausal breast cancer patients in Malaysia. *Climacteric* 2019;22:175–81.
- 14 Condorelli R, Vaz-Luis I. Managing side effects in adjuvant endocrine therapy for breast cancer. *Expert Rev Anticancer Ther* 2018;18:1101–12.
- 15 Stanton AL, Bower JE. Psychological adjustment in breast cancer survivors. Adv Exp Med Biol 2015;862:231–42.
- 16 Brandão T, Schulz MS, Matos PM. Psychological adjustment after breast cancer: a systematic review of longitudinal studies. *Psychooncology* 2017;26:917–26.
- 17 Londono Y, McMillan DE. Psychosocial adaptation: an evolutionary concept analysis exploring a common multidisciplinary language. *J Adv Nurs* 2015;71:2504–19.
- 18 AntonakRF, M. J.Leahy. Case management forrehabilitation health professionals. Osage Beach, MO: Aspen ProfessionalServices. In: Chan F, Saunders J, eds. *Psycho-Social aspectsof chronic illness* and disability. 2, 2005: 2. 3–43.
- 19 Chan F, da Silva Cardoso E, Chronister JTurner JR, ed. Encyclopedia of behavioral medicine. New York: Springer, 2013.
- 20 Livneh H, Antonak RF. Case management forrehabilitation health professionals. In: Chan F, Leahy MJ, Saunders J, eds. *Psycho-Social aspectsof chronic illness and disability*. 2. Osage Beach: Aspen ProfessionalServices, 2005.: 2. 3–43.
- 21 Ośmiałowska E, Staś J, Chabowski M, et al. Illness perception and quality of life in patients with breast cancer. *Cancers* 2022;14. doi:10.3390/cancers14051214. [Epub ahead of print: 25 02 2022].
- 22 Bakan G, Inci FH. Predictor of self-efficacy in individuals with chronic disease: stress-coping strategies. *J Clin Nurs* 2021;30:874–81.
- 23 Derogatis LR. The psychosocial adjustment to illness scale (PAIs). J Psychosom Res 1986;30:77–91.
- 24 Kolokotroni P, Anagnostopoulos F, Missitzis I. Psychosocial adjustment to illness scale: factor structure, reliability, and validity assessment in a sample of Greek breast cancer patients. *Women Health* 2017;57:705–22.
- 25 Schwab DP. Book review: research methods for organizational studies. *Organ Res Methods* 2006;9:572–4.
- 26 Ezzedine K, Bennani M, Shourick J, et al. A method for designing a patient burden questionnaire in dermatology. *Clin Cosmet Investig Dermatol* 2020;13:521–8.
- 27 Charmaz K. Constructing grounded theory. London: Sage, 2006.
- 28 Mystakidou K, Parpa E, Tsilika E, et al. General perceived selfefficacy: validation analysis in Greek cancer patients. Support Care Cancer 2008;16:1317–22.
- 29 Roy C. Extending the Roy adaptation model to meet changing global needs. Nurs Sci Q 2011;24:345–51.
- 30 Roy C. Pearson Education. In: *The Roy adaptation model, upper saddle river NJ*, 2009.
- Batista-Foguet JM, Coenders G, Alonso J. [Confirmatory factor analysis. Its role on the validation of health related questionnaires]. Med Clin 2004;122 Suppl 1:21–7.
- 32 Quadrelli S, Davoudi M, Galíndez F, et al. Reliability of a 25-item lowstakes multiple-choice assessment of bronchoscopic knowledge. Chest 2009;135:315–21.
- 33 University. NCS.Reliability analysis: key concepts and terms. Available: http://www2.chass.ncsu.edu/garson/pa765/reliab.htm [Accessed 18 Oct 2008].
- 34 Williams LS, Weinberger M, Harris LE, et al. Development of a Stroke-Specific quality of life scale. Stroke 1999;30:1362–9.
- 35 Tang W-K, Mak K-K, Kam PM-H, et al. Reliability and validity of the euthanasia attitude scale (EAS) for Hong Kong medical doctors. Am J Hosp Palliat Care 2010;27:320–4.
- 36 University. NCS. Validity analysis: key concepts and terms. Available: http://www2.chass. ncsu.edu/garson/pa765/reliab.htm. [Accessed 18 Oct 2008].
- 37 Grafton KV, Foster NE, Wright CC. Test-Retest reliability of the shortform McGill pain questionnaire: assessment of intraclass correlation coefficients and limits of agreement in patients with osteoarthritis. *Clin J Pain* 2005;21:73–82.
- 38 Wang X-ling, Zhao T-en, Zhang X-qin. [Assessment on the reliability and validity of the Dermatology Life Quality Index in Chinese version]. Zhonghua Liu Xing Bing Xue Za Zhi 2004;25:791–3.

Open access

- 39 Qiu H. Quantitative research and statistical analysis. In: . 4. Chongqing: Chongqing university press, 2013.
- 40 Burgess C, Cornelius V, Love S, *et al.* Depression and anxiety in women with early breast cancer: five year observational cohort study. *BMJ* 2005;330:702.
- 41 Cronbach LJ, Warrington WG. Time-limit tests: estimating their reliability and degree of speeding. *Psychometrika* 1951;16:167–88.
- 42 Heo M, Kim N, Faith MS. Statistical power as a function of Cronbach alpha of instrument questionnaire items. *BMC Med Res Methodol* 2015;15:86.
- 43 Wild D, Grove A, Martin M, *et al.* Principles of good practice for the translation and cultural adaptation process for patient-reported outcomes (pro) measures: report of the ISPOR Task force for translation and cultural adaptation. *Value Health* 2005;8:94–104.