


BMJ Open Evaluation of a mental health screening tool using cross-sectional surveys in a workplace setting

Joe Xu ¹, Alexander Willems,¹ Vincy Li,¹ Nick Glozier,² Philip J Batterham,³ Victoria Malone,¹ Richard W Morris ⁴, Chris Rissel⁵

To cite: Xu J, Willems A, Li V, *et al*. Evaluation of a mental health screening tool using cross-sectional surveys in a workplace setting. *BMJ Open* 2022;**12**:e052155. doi:10.1136/bmjopen-2021-052155

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-052155>).

Received 13 April 2021
Accepted 14 April 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹NSW Office of Preventive Health, NSW Health, Sydney, New South Wales, Australia

²Brain and Mind Research Institute, University of Sydney, Sydney, New South Wales, Australia

³Centre for Mental Health Research, Australian National University, Canberra, Australian Capital Territory, Australia

⁴Central Clinical School, University of Sydney, Sydney, New South Wales, Australia

⁵School of Public Health, University of Sydney, Sydney, New South Wales, Australia

Correspondence to

Dr Joe Xu; jx1158@gmail.com

ABSTRACT

Objectives The Brief Health Check (BHC) is a health screener used by the Get Healthy at Work programme, which identifies workers with chronic disease risk and provides them with advice and referrals to support services. The BHC was revised to include mental health to provide a holistic approach to workplace health. This study aimed to evaluate the acceptability and appropriateness of the revised BHC by comparing the results around psychological distress and future risk with previous research, and a participant feedback survey.

Method Data collection took place between October 2018 and May 2019. The study used data that were collected as part of programme delivery, as well as a participant feedback survey that was administered after the health check was completed.

Results BHCs were completed by n=912 workers, out of which, n=238 completed the feedback survey. The mean Distress Questionnaire 5 score was 10.5, and 10% of participants met the threshold for 'high' future risk. The feedback survey revealed that the majority of participants found the mental health advice to be useful (76%), agreed with their mental health distress and risk ratings (92%–94%) and most intended on using the referred services (62%–68%).

Conclusion The findings around mental health risk were comparable to previous findings in employed samples. The inclusion of mental health assessments, advice and referral pathways into the BHC was found to be acceptable and the subsequent referrals were appropriate, indicating that this approach could be scaled up and implemented to help address worker's mental ill health

INTRODUCTION

Mental health issues are very prevalent in the Australian population, with one in five adults (aged 18–85) having experienced mental disorders within the last 12 months, and 45.5% of the total population having experienced a mental disorder at some point in their lifetime.¹ Mental health issues in the working population can be costly to employers in terms of lost productivity and turnover, as well as to society at large in the form of health service use, where US\$9.9 billion was spent on mental health related services in Australia

Strengths and limitations of this study

- ⇒ The study is the first to use the Distress Questionnaire 5, a general population health screener tool, in a working population.
- ⇒ This is the first health screener in Australia to include both current psychological distress and future mental health risk.
- ⇒ The study did not employ a longitudinal design, and future research could follow-up with employees to assess the impact of the health check.
- ⇒ The study did not ask about demographics in the participant feedback survey, so it is uncertain how the participant feedback survey sample compared with the larger sample which completed the Brief Health Check.

from 2017 to 2018.^{2–5} For individual workers, mental health issues can impact negatively on workplace engagement as well as overall quality of life.^{6 7} In recent years, governing bodies in Australia have implemented strategies to facilitate the promotion of mental well-being in the workplace.^{8 9} One of the channels through which this strategy is implemented is through existing workplace health programmes, which have significant reach in the working population and present opportunities for promoting mental well-being (eg, the Mentally Healthy Workplaces programme from SafeWork New South Wales (NSW)).¹⁰

Workplace health programmes are health promotion and protection strategies implemented in the workplace,¹¹ with the goal of establishing organisational cultures that promote and provide healthy lifestyle choices. Systematic reviews of such programmes have found positive impacts on the health and well-being of workers as well as the productivity of the organisation.^{12–14} In NSW, the Get Healthy at Work programme was launched in 2014, along with a Brief Health Check (BHC) with the aim of reducing type 2 (T2) diabetes and cardiovascular disease risk among workers.

The supports workplaces to create health promoting structures and processes, along with a BHC designed to help individual workers to reduce their lifestyle risk factors (ie, waist circumference, diet, physical activity and smoking). The BHC identifies workers with high T2 diabetes and cardiovascular risk, refers them to external support services, and offers personalised advice.¹⁵

In late 2018, the Get Healthy at Work programme sought to include mental health into the BHC to provide a holistic assessment for employee health and well-being. The BHC was expanded to include mental health assessments, referral pathways to mental health support services, as well as personalised mental health advice. Similar to the development of the original BHC,¹⁶ the development of the mental health items followed a translational formative evaluation process,¹⁷ which began with synthesising the evidence, consulting with practitioners/academics, as well as stakeholders. From this process, programme managers decided to use the Distress Questionnaire 5 (DQ5), which is a short assessment of current psychological distress.¹⁸

The BHC also sought to prevent future incidence of psychological distress in participating workers. Therefore, the revised BHC includes a risk algorithm developed by Morris and Glozier (an unpublished internal report) to identify participants who are at risk of experiencing mental health issues within the next 12 months.

Based on advice from the clinical advisory panel, the revised BHC refers participants with high current distress (according to the DQ5 score) to the MindSpot free online supported mental health clinic,¹⁹ as well as to a general practitioner. Those found to have moderate current distress are referred to myCompass,²⁰ an online mental health programme that is self-guided. Both myCompass and MindSpot^{21–23} have demonstrated efficacy in improving mental health outcomes. Participants with high future risk scores are given advice to help manage their mental well-being. Further, because of the importance of positive lifestyle modification in promoting mental well-being,^{24–26} the BHC offers personalised advice around how individuals could improve their mental well-being by modifying their lifestyle through improved diet and physical activity.

Following the translational formative evaluation process,¹⁷ the current study aims to evaluate the revised BHC within workplaces to assess whether it can be scaled up for state-wide delivery and identify ways in which the tool can be improved. The key implementation research questions to be examined were: (1) Comparability: How do the findings around current psychological distress and high mental health risk in the applied setting compare with previous research? (2) Acceptability: Do workers find the new mental health questions easy to understand? Do participants agree with the results they received? Is there any potential harm in using these assessments? Do participants agree with the risk ratings they received? (3) Uptake and engagement: What is the uptake of referrals made? Do participants intend on using the services

to which they were referred? Do participants find the personalised mental health advice useful?

METHOD

The study used BHC cross sectional survey data that was collected as part of regular programme delivery to determine the comparability of results and uptake of referral pathways. A cross-sectional feedback survey was administered after completing the BHC. The feedback survey was included to help answer the research questions around acceptability, uptake of referrals and engagement with advice.

Sample

The revised BHC was first administered within two NSW government organisations that consented to using the revised BHC: the Department of Education, and a government insurance and workers compensation unit. Data collection for the current study ran from October 2017 to May 2018. The worksites for both organisations were in metro and regional/rural areas. Each participating organisation promoted the BHC at each worksite, and participants who completed the BHC were asked to complete the feedback survey immediately after completing the BHC. The participant feedback survey was administered at worksites that allowed the participant feedback survey to be administered (ie, 13 of the 35 worksites that were involved in the pilot). The study made use of all BHC data that was collected during the study period, as well as all participants who consented to provide feedback via the survey. The BHC sample was large enough to detect small effect sizes (Cohens's $d=0.2$ at 80% power) when comparing samples on the DQ5.

Participant involvement

Participants provided data for the study and were not involved in the design, reporting or dissemination for this project.

Measures

Distress Questionnaire 5

The DQ5 has greater sensitivity than other widely used measures (ie, Kessler 6 and 10) for identifying individuals currently at risk for specific anxiety disorders. The development of the DQ was described in detail in the paper by Batterham *et al.*¹⁸ The BHC uses the cut-points defined by Batterham *et al.*¹⁸ to classify participants into different levels of current distress. That is, participants with DQ5 scores equal to or greater than 11 were identified as having 'moderate' current distress, where a participant is likely to meet the criteria for a wide range of disorders, and those with DQ5 scores equal to or greater than 14 were identified as having 'high' current distress, where a participant is likely to meet the criteria for specific disorders with a lower rate of false positives compared with participants who are classified as having 'moderate' distress.

Table 1 Future risk model

Participant characteristics	Beta coefficient (log odds)
Aged 18–35	None*
Aged 35–39	0.030
Aged 40–44	–0.167
Aged 45–54	–0.040
Aged 55–64	–0.167
Aged 65 and over	–0.207
Origin Australia	None*
Origin Asia	0.104
Origin Middle East/N. Africa	–0.011
Origin Other	–0.080
Aboriginal or Torres Strait Islander	0.032
Male	–0.085
Recent history of mental illness (2 year)	0.672
Agree with ‘I am pushed around’	0.281
Satisfaction with health	–0.068
Agree with ‘I don’t have anyone to confide in’	0.151
Five or more standard alcoholic drinks in any single day (last 7 days)	0.047
Are you a current smoker?	0.158
Exercise less than once per week	0.056
Constant	–1.288

Future risk tool

The future risk tool used in the BHC was adapted from Fernandez *et al*²⁷ by Morris and Glozier, which is the first mental health risk algorithm to be created for the working population in Australia. Morris and Glozier updated the algorithm using 2015 and 2016 data from the Household and Income Labour Dynamics in Australia survey and obtained a comparable C-index (0.71) and positive predictive value (0.28) in validation (The formula for the future risk algorithm is: $Y_i = -1.288 + (0.03) \text{ age: } 35\text{--}39 + (-0.167) \text{ age: } 40\text{--}44 + (-0.04) \text{ age: } 45\text{--}54 + (-0.167) \text{ age: } 55\text{--}64 + (-0.207) \text{ age: } 65 \text{ and over} + (0.104) \text{ country of origin: Asia} + (-0.011) \text{ country of origin: Middle East/N. Africa} + (-0.080) \text{ country of origin: other} + (0.032) \text{ Aboriginal or Torres St. Islander} + (-0.085) \text{ sex: male} + (0.672) \text{ recent mental illness} + (0.281) \text{ bullied} + (-0.068) \text{ health satisfaction} + (0.151) \text{ loneliness} + (0.047) \text{ binge drink} + (0.158) \text{ Smoker} + (0.056) \text{ physically inactive}$). The coefficients for the future risk algorithm are presented in table 1. For future risk scores, the revised BHC uses thresholds defined by Morris and Glozier, in which participants who exceed the algorithm’s threshold for high risk are expected to have a 28% chance of experiencing psychological distress in the next 12 months. Participants who exceed the threshold for moderate risk are expected to have a 22% chance of experiencing psychological distress in the next 12 months.

Analysis

1. Comparability: The results around current distress and prevalence of future risk categories in the BHC were compared with previous research. The BHC sample was weighted for age and gender before the results were compared with previous data, which examined findings at the population level. The weight values were based on the 2016 Australian Census filtered for individuals who were employed.²⁸ A two-sample t-test was used to compare the mean DQ5 score from the current study with the results from Batterham *et al*,¹⁸ and the prevalence of future risk was compared with the models that informed the development of the future risk tool by descriptive statistics.
2. Acceptability: The feedback survey asked participants whether the questions were difficult to understand, and whether participants felt uncomfortable about answering any of the mental health questions. Both were examined using ‘Yes/No’ questions followed by open-ended questions to identify the items that were difficult or made participants feel uncomfortable. These questions aimed to assess any potential issues with comprehension and harm associated with the revised BHC.
3. Uptake and engagement: The uptake of the referred services was recorded in the BHC questionnaire, where participants have ‘accepted’ referrals if they agreed to be referred during the BHC session by the health professional, or indicated that they will register for the service after the BHC. Referral outcomes were stratified by current help seeking behaviour (ie, whether participants are currently seeing a mental health professional), as well as demographic characteristics to assess the rate of uptake in those who are not receiving help, and a range of population groups. Uptake of the referred services was also examined through the participant feedback survey, which asked participants whether they intend on using the service to which they were referred in the BHC (examined using multiple choice ‘Yes/No/Intend to use at a later time’). The feedback survey also asked participants whether they found the mental health advice useful on a five-point scale. The authors do not have visibility of the number of participants who access their referred service after the BHCs were conducted.

Participants who did not complete the DQ5 or future risk questionnaires were excluded from the analyses. Participants who did not answer a question in the feedback survey were removed from the analysis of that question.

Brief Health Check

The revised BHC was administered face to face within participating worksites by trained health professionals, such as accredited dietitians or exercise physiologists. Participants completed a questionnaire related to diet, physical activity, demographic characteristics, and physical and mental health risk profiles, distress (DQ5) and

health-related behaviours. The BHC questionnaires were completed on paper (n=198) or equivalent digital forms (n=714). Once the risk scores were calculated, the health practitioners provided feedback about the risk scores, and provided appropriate referrals and advice depending on the risk profile of the participant. Health professionals then recorded whether participants accepted referrals. A single BHC session took around 20 min to complete.

Participant feedback survey

Between 1 and 3 interviewers were present at each of the 13 worksites participating in the feedback survey. Once participants completed the BHC, they were asked to participate in a survey administered by the trained interviewers, who were blinded to the results from the BHC. The surveys included seven questions that were administered verbally and responses were collected on paper forms. The survey took no longer than 10 min to complete. The questions were a combination of closed and open-ended responses that were developed for this study. The open-ended responses about difficulties understanding questions and feeling uncomfortable about answering questions were analysed by two coauthors (JX and VM) using closed-coding to identify the specific BHC questions referenced in participant feedback. Open-ended responses around participant feedback, agreement with current distress and future risk ratings, and intention to use services were analysed using open coding. The authors coded the responses independently, and then met to reach a consensus on the assignment of the codes.

RESULTS

Profile of worksites

A total of 35 worksites participated in the study and 13 worksites allowed participant feedback surveys to be administered. The total number of completed BHCs was n=912, and a total of n=238 participants completed the feedback survey. The authors did not have visibility of the number of employees within each organisation that were invited to complete the BHC. Based on an estimate of the number of employees across the worksites (n=7200), and

the assumption that all employees at each worksite were invited, a conservative estimate of the response rate for the BHC (with n=912 completes) is 12.7%. A breakdown of completed BHCs, number of surveys within organisations and the location of the worksite is presented in [table 2](#). The urban/rural/remoteness of the worksite was based on postcode, using the Accessibility and Remoteness Index of Australia.²⁹

Comparability

The characteristics of participants, including the current distress results and prevalence of future risk categories, are presented in [table 3](#). Two participants did not complete the DQ5 and were excluded from the analyses. Future risk scores were only calculated for participants who did not have a high level of current distress (ie, those with DQ5 scores<14). In this study, the weighted mean DQ5 score was 10.5 (SD=4.2). This was significantly higher than the weighted mean scores from the study by Batterham *et al*¹⁸ (mean DQ5 score=9.28, SD=4.08), via an independent samples t-test: $t(4083)=7.8$, $p<0.001$, and the difference was small-medium in terms of effect size (Cohen's $d=0.29$). For the prevalence of future risk in the weighted sample, 9.6% of participants met the threshold to be in the 'high' future risk category, in which 28% of participants are expected to experience psychological distress within 12 months. This is consistent with the population proportion that was expected to meet this threshold according to the future risk algorithm (10% or 90th percentile).

Acceptability

Mental health questions

The participant feedback survey revealed that 17.2% (n=41) of respondents found the mental health questions difficult to understand. Participants reported that they found one (13.4%, n=32) or two (2.5%, n=6) questions difficult, and the remainder reported that their difficulties were due to general comprehension or recall (1.2%, n=3). The responses were back-coded to identify the specific questions that were difficult to understand, which showed that 10.9% (n=26) of participants found

Table 2 Breakdown of Brief Health Checks and surveys by organisation

Organisation name	No of worksites	Participants	
		Brief Health Checks n=912 (%)	Feedback surveys n=238 (%)
Department of Education			
Major cities	13	363 (39.8)	124 (52.1)
Regional/remote	16	264 (28.9)	51 (21.4)
Insurance and Workers Compensation Unit			
Major cities	6	285 (31.3)	63 (26.5)
Total major cities	19	648 (71.1)	187 (78.6)
Total regional/remote	16	264 (28.9)	51 (21.4)
Total	35	912	238

Table 3 Characteristics of participants who completed Brief Health Checks (BHC) (n=912)

	Unweighted N (%)	Weighted N (%)
Age group*		
18–34 years	256 (28.1)	326.9 (35.8)
35–39 years	148 (16.2)	101.2 (11.1)
40–44 years	115 (12.6)	104.1 (11.4)
45–54 years	238 (26.1)	200.9 (22.0)
55–64 years	142 (15.6)	139.8 (15.3)
65 years or over	13 (1.4)	39.1 (4.3)
Gender		
Male	253 (27.7)	480.2 (52.7)
Female	659 (72.3)	431.8 (47.3)
Current distress categories†		
High	208 (22.9)	188.3 (20.7)
Moderate	216 (23.7)	228.0 (25.0)
Low	486 (53.4)	494.0 (54.3)
Future risk categories‡		
High	77 (11.0)	69.3 (9.6)
Moderate	143 (20.3)	140.7 (19.4)
Low	482 (68.7)	513.7 (71.0)

*Age data were collected using the categories below. These groupings are used by the BHC to determine type two diabetes risk which is not a focus of the current study.

†n=2 participants did not complete the DQ5 questions.

‡Age data were collected using the categories below. These groupings are used by the BHC to determine type 2 diabetes risk which is not a focus of the current study.
DQ5, Distress Questionnaire 5.

the future risk questions to be difficult, and 6.7% (n=16) of participants found the DQ5 questions to be difficult. Of the participants who found the future risk questions to be difficult, themes emerged regarding whether the question around 'satisfaction with your health' referred to mental or physical health, and whether the question 'Have you had mental health problems in the past 2 years' referred to mental health issues that were diagnosed or included all mental health problems. Most of the participants who had trouble understanding the DQ5, linked their difficulties to the question 'I found social settings upsetting' and whether 'social settings' referred to all social settings or just those in the workplace. Many participants who had trouble with DQ5 or the future risk questions also reported that the health professionals conducting the BHC offered useful prompts which helped them answer these questions. A small proportion (7.6%, n=18) reported that they felt uncomfortable about answering one or more of the mental health questions. When probed further about the specific questions they had concerns about, most of these participants indicated that they felt uncomfortable about talking about mental health in general (n=12, 5%), while 1.3% (n=3) linked

their response to the DQ5, and 2.1% (n=5) linked their response to the future risk questions.

Agreement with risk ratings

From the participant feedback surveys, only 5.9% (n=14) of participants disagreed with their current distress scores, and 8.0% (n=19) disagreed with their future risk scores. Of the participants who disagreed with their either their current or future mental health risk, there was a mix of those who expected their scores to be higher (current: n=1, 0.4%; future: n=3, 1.3%) or lower (current: n=4, 1.7%; future: n=7, n=2.9%) than what they received. For those who disagreed with their current distress or future risk scores, some participants did not disagree with the rating per se but expressed scepticism that the questions could provide an accurate assessment of their mental health state or predict their future risk: 'Assessing risk for the future seems unrealistic—impossible to know what will happen in the future. Not sure how the assessment/questions work'.

Uptake and engagement

Uptake of referrals

The breakdown of participants who accepted referrals during the BHC session are presented in table 4. Questions around whether participants were currently seeing a mental health professional were introduced later in the pilot, and so the sample size for table 4 is smaller than the total number of completed BHCs. Of participants who had high current distress and were not currently receiving support from a mental health professional (n=139), the majority (n=95, 68.3%) accepted referrals to MindSpot, and most participants accepted referrals to their GP for mental health support (n=86, 61.9%). Referral outcomes were further examined by age, gender and cultural background to assess whether referral rates differ across population groups. For participants with high current distress, there were no significant differences between any demographic groups in accepting referrals to MindSpot or their GP (using χ^2 tests; p's>0.05). Females (n=97, 65.1%) were significantly more likely than males (n=33, 49.3%) to accept a referral to myCompass ($\chi^2=4.2$, p=0.04).

Based on the participant feedback surveys, the majority of participants indicated that they intended to access the mental health services to which they were referred (myCompass n=62/81, 76.5%; MindSpot n=31/43, 72.1%; n=21/29, GP 72%). Some participants who indicated that they did not plan on accessing MindSpot or myCompass suggested that they would prefer face-to-face mental health support: 'No, not likely to go online...I would rather see someone face-to-face'. However, a number of participants suggested that they might use these services in the future: 'I don't think I need [MindSpot] right now, but it is good to know about it if I need to access it later'.

Advice

Out of the participants who received mental health advice during the BHC, most reported that the advice

Table 4 Brief Health Checks referral outcomes for mental health

Referrals from Brief Health Checks	Accept/self-referral	Declined/not referred
MindSpot		
Currently seeing a mental health professional (n=41)	23 (56.1%)	18 (43.9%)
Not currently seeing a mental health professional (n=139)	95 (68.3%)	44 (31.7%)
Mental Health GP referral		
Currently seeing a mental health professional (n=41)	21 (51.2%)	20 (48.8%)
Not currently seeing a mental health professional (n=139)	86 (61.9%)	53 (38.1%)
myCompass		
Currently seeing a mental health professional (n=17)	14 (82.4%)	3 (17.6%)
Not currently seeing a mental health professional (n=149)	95 (63.8%)	54 (36.2%)

The question around whether participants were currently seeking support were introduced later in the pilot. For this table, the base for high current distress n=180; and the base for moderate risk n=166.
GP, general practitioner.

they received was useful (n=89, 76.1% reported that the advice was 'Very useful'/'Fairly useful'; n=26, 22.2% reported that the advice was 'A little useful'/'Not useful at all'; and n=2, 1.7% indicated that they 'Don't know'). When asked to provide further feedback about the advice they received, some participants suggested that the advice helped them learn more about their mental well-being: 'I knew much of the information on physical health, but mental health was all new to me. Surprised about the links between physical health and mental health...I didn't previously ever even consider my mental health'. Participants who suggested that the advice confirmed what they already know, saw this as a useful instance of reinforcing their understanding of healthy lifestyle behaviours: '[I] already know about own mental and physical states, but was good to get confirmation and reminder'. Participants who felt that the advice was 'A little useful'/'Not useful' indicated that the advice was not specific enough: 'I am [already] conscious of my physical and mental health, the check-up was very broad'.

DISCUSSION

The findings from his study suggest that the revised BHC is appropriate for assessing both current and future mental health risk in the workplace context. The mean DQ5 score from the current study is higher than that from Batterham *et al*, which is consistent with previous research. Specifically, Jarman *et al*³⁰ compared the psychological distress from a general population with the findings from an employee well-being survey among public servants in Tasmania. The authors found that the mean psychological distress (using the Kessler 10) scores from public service workers was higher than the general population, and suggested that the differences could be attributed to workplace specific stressors such as the rationalisation of the workforce, job insecurity and effort-reward imbalance.^{31 32} The lower levels of well-being among public sector employees has also been found in other jurisdictions.³³ A recent study of secondary school

teachers in NSW by Parker *et al*³⁴ found a mean DQ5 value (ie, mean=11.25, SD=3.8), a similar result to the current study, although this was from a small sample. The prevalence of high future mental health risk is comparable with the models that informed the development of the future risk tool, which used the same measures in a statewide sample across many different occupational groups. While different occupational groups commonly report very different levels of mental ill health,³⁵ the similarity in risk prevalence between this study and earlier work suggests that there are common drivers of mental ill health risk across industries (eg, prior ill health, discrimination).

The majority of those who were not receiving mental health support at the time of the BHC accepted referrals to mental health support services (ie, MindSpot, mental health GP referrals and myCompass) based on their risk scores. There were no differences in the demographic characteristics of participants with high current distress that accepted referrals compared with those who did not accept referrals. The findings from the participant survey suggest that only a small number of participants felt uncomfortable about answering the mental health questions, and most participants agreed with their mental health risk scores. Participants mostly reported that the advice that was offered as part of the BHC was useful and that they intended on using the mental health services to which they were referred. Overall, these results suggest that the revised BHC is suitable for use among workers.

The participant feedback survey revealed that around one out of five participants found the mental health questions difficult to understand, which would require the BHC to be refined to facilitate understanding. The findings from the survey also highlighted ways in which comprehension could be improved. Specifically, confusion around the DQ5 question 'I found social settings upsetting', and around whether the future risk question for whether participants have had 'mental health problems in the past 2 years' could be addressed by providing participants with suitable prompts. For the future risk

question around 'satisfaction with health', prompts could be offered to clarify that health refers to both mental and physical health, or reorder the question to a location where the participant would not be biased towards interpreting the question as referring to either physical or mental health. In terms of next steps, it is recommended that the prompts for the DQ5 and future risk tool are added to the revised BHC before it is implemented on a wider scale. The prompts will only be provided by the health professional if a worker has trouble with the instrument and are not expected to impact on the validity of those instruments.

For participants who reported that they prefer a more comprehensive health check or were sceptical that their future mental health risk can be accurately determined from a small number of questions, their experience could be improved by setting more realistic expectations about the programme. That is, the BHC should be introduced as a concise screener tool used to identify participants who are 'at risk' and refer them to clinical support services, as opposed to a definitive diagnostic test, consistent with the messaging from other online assessment tools such as the Black Dog Institute's Online Clinic assessment.³⁶ The information about how future risk is calculated (ie, a combination of physical and mental health questions, modifiable and non-modifiable factors) as well as noting that the future risk score is based on existing research, will help assure participants who are sceptical about the validity of the assessments. To improve workers' experience with the tool, it is recommended that these adjustments are incorporated into the standard BHC protocol.

As an adaptation, the BHC could be implemented as an online assessment (eg, with automated scoring, advice and referrals), which presents an opportunity to scale up the programme and extend the reach to a larger number of organisations and remote locations. Future research could explore whether participants would find an online BHC to be as useful as a face-to-face version, given that the participants have responded positively to the personalised advice delivered by health professionals. The ease of administering the revised BHC as an online tool presents opportunities for a mental health screener to be deployed at scale in the workplace, while offering relevant advice and referral pathways. The introduction of an accessible health screening tool aligns with the recommendation from public and mental health professionals to improve the mental health of workers.^{37 38} However, the BHC with feedback and advice might not be sufficient in isolation, as studies have suggested that improvements to some health outcomes are better achieved through a combination of health assessments and other health promotion activities (eg, health education, policy and environmental change),³⁹ which highlights the importance of implementing other workplace health initiatives prescribed by the Get Healthy at Work programme alongside the BHC.

A limitation of this study is that the current study did not collect demographic information in the

participant feedback survey, so the sample from the feedback survey cannot be compared with the BHC sample. Additionally, the current study does not provide any insight into the long-term benefits of the programme. Future research can also use the BHC to track the health of workers longitudinally and examine the relative impacts of the workplace health programme on the health outcomes of workers. The predictive accuracy of the future risk tool may also be a limitation of the current study. Although the tool has modest predictive accuracy, there are no established risk prediction tools that perform better in identifying the risk of future mental ill health. Predictive validity of such tools will be limited by a multitude of risk factors that influence distress and the relatively low base rate of distress in general population settings.

The current research suggests that the revised BHC with mental health assessments, referral pathways and advice are acceptable and suitable for the workplace setting, but also highlights ways in which the revised BHC could be improved. To our knowledge, this is first study to assess the acceptability and appropriateness of the DQ5, a population health screener, in a workplace setting. Additionally, the revised BHC is the first mental health assessment that tests for both current and future mental health risk in the workplace.

Acknowledgements The authors would like to thank Ms Jillian Green (SafeWork NSW) and Prof Samuel Harvey (University of New South Wales) for their support and advice.

Contributors JX drafted the manuscript and conducted data analyses. JX and VM conducted qualitative analyses on the survey responses. All other authors (AW, VL, NG, PJB, RWM and CR) contributed to revising the manuscript. JX is the guarantor for this work.

Funding This work was funded by the Centre for Population Health, NSW Ministry of Health.

Disclaimer The contents of this paper are solely the responsibility of the individual authors and do not reflect the view of NSW Ministry of Health.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s)

Ethics approval Ethical approval for the analysis of routine programme data and participant feedback was obtained from South Western Sydney Local Health District Human Ethics Committee (Ref: ETH12061). The ethics approval covered the routine analysis of programme data (BHCs) and the participant feedback survey, for which verbal consent was obtained from participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Data are available on reasonable request to the authors.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Joe Xu <http://orcid.org/0000-0002-6034-4118>

Richard W Morris <http://orcid.org/0000-0002-5018-1239>

REFERENCES

- 1 Department of Health. Prevalence of mental disorders in the Australian population [Internet]. Available: <https://www1.health.gov.au/internet/publications/publishing.nsf/Content/mental-pubs-m-mhaust2-toc~mental-pubs-m-mhaust2-hig~mental-pubs-m-mhaust2-hig-pre> [Accessed 15 Mar 2021].
- 2 LaMontagne AD, Sanderson K, Cocker F, Victorian Health Promotion Foundation. *Estimating the economic benefits of eliminating job stress as a risk factor for depression: full report*. Melbourne: VicHealth, 2010.
- 3 LaMontagne AD, Martin A, Page KM, et al. Workplace mental health: developing an integrated intervention approach. *BMC Psychiatry* 2014;14:131.
- 4 Yu S, Glozier N. Mentally healthy workplaces a return-on-investment study for Safework NSW, 2017. Available: https://www.safework.nsw.gov.au/__data/assets/pdf_file/0011/320132/Mentally-healthy-workplaces-A-return-on-investment-study-August-2017-SW08735.pdf
- 5 Australian Institute of Health and Welfare. Mental health services in Australia, Expenditure on mental health-related services [Internet]. Available: <https://www.aihw.gov.au/reports/mental-health-services/mental-health-services-in-australia/report-contents/expenditure-on-mental-health-related-services> [Accessed 15 Mar 2021].
- 6 Lehman AF, Postrado LT, Rachuba LT. Convergent validation of quality of life assessments for persons with severe mental illnesses. *Qual Life Res* 1993;2:327–33.
- 7 Slep GR, Vella-Brodick DA. Optimising employee mental health: the relationship between intrinsic need satisfaction, job crafting, and employee well-being. *J Happiness Stud* 2014;15:957–77.
- 8 National Mental Health Commission. National Workplace Initiative [Internet]. Available: <https://www.mentalhealthcommission.gov.au/Mental-health-Reform/National-Workplace-Initiative> [Accessed 31 Mar 2020].
- 9 NSW mentally healthy workplaces strategy 2018–22. Available: https://www.safework.nsw.gov.au/__data/assets/pdf_file/0006/362274/NSW_mentallyhealthyworkplacesstrategy_2018_22.pdf
- 10 Safework NSW. Mentally healthy workplaces [Internet]. Safework Mental Health, 2018. Available: <https://www.safework.nsw.gov.au/safety-starts-here/mental-health-at-work-the-basics/mental-health-at-work/mentally-healthy-workplaces> [Accessed 24 Apr 2020].
- 11 Matson Koffman DM, Lang JE, Chosewood LC. Cdc resources, tools, and programs for health promotion in the worksite. *Am J Health Promot* 2013;28:TAHP2–5.
- 12 Goetzel RZ, Henke RM, Tabrizi M, et al. Do workplace health promotion (wellness) programs work? *J Occup Environ Med* 2014;56:927–34.
- 13 Grossmeier J, Terry PE, Anderson DR, et al. Financial impact of population health management programs: Reevaluating the literature. *Popul Health Manag* 2012;15:129–34.
- 14 Bellew B. Managing health and wellbeing in the workplace: an Evidence Check rapid review brokered by the Sax Institute (www.saxinstitute.org.au) for SafeWork NSW [Internet], 2018. Available: <https://www.saxinstitute.org.au/wp-content/uploads/Managing-health-and-wellbeing-in-the-workplace.pdf> [Accessed 24 Apr 2020].
- 15 SWSLHD - Office of Preventive Health [Internet]. Available: <http://www.preventivehealth.net.au/GHaW.html> [Accessed 31 Mar 2020].
- 16 Lloyd B, Khanal S, Macoun E, et al. Development of a multiple risk factor brief health check for workplaces. *Public Health Res Pract* 2016;26:2641649.
- 17 O'Hara BJ, Phongsavan P, King L, et al. 'Translational formative evaluation': critical in up-scaling public health programmes. *Health Promot Int* 2014;29:38–46.
- 18 Batterham PJ, Sunderland M, Carragher N, et al. The distress Questionnaire-5: population screener for psychological distress was more accurate than the K6/K10. *J Clin Epidemiol* 2016;71:35–42.
- 19 About MindSpot – our team, vision and values - MindSpot Clinic [Internet], 2020. Available: <https://mindspot.org.au/about-mindspot> [Accessed 27 Mar 2020].
- 20 myCompass - MyCompass [Internet], 2020. Available: <https://www.mycompass.org.au/> [Accessed 27 Mar 2020].
- 21 Proudfoot J, Clarke J, Birch M-R, et al. Impact of a mobile phone and web program on symptom and functional outcomes for people with mild-to-moderate depression, anxiety and stress: a randomised controlled trial. *BMC Psychiatry* 2013;13:312.
- 22 Titov N, Dear BF, Staples LG, et al. MindSpot clinic: an accessible, efficient, and effective online treatment service for anxiety and depression. *Psychiatr Serv* 2015;66:1043–50.
- 23 Titov N, Dear BF, Staples LG, et al. The first 30 months of the MindSpot clinic: evaluation of a national e-mental health service against project objectives. *Aust N Z J Psychiatry* 2017;51:1227–39.
- 24 Meegan AP, Perry IJ, Phillips CM. The association between dietary quality and dietary guideline adherence with mental health outcomes in adults: a cross-sectional analysis. *Nutrients* 2017;9:238.
- 25 Rivenes AC, Harvey SB, Mykletun A. The relationship between abdominal fat, obesity, and common mental disorders: results from the HUNT study. *J Psychosom Res* 2009;66:269–75.
- 26 Teychenne M, Ball K, Salmon J. Physical activity and likelihood of depression in adults: a review. *Prev Med* 2008;46:397–411.
- 27 Fernandez A, Salvador-Carulla L, Choi I, et al. Development and validation of a prediction algorithm for the onset of common mental disorders in a working population. *Aust N Z J Psychiatry* 2018;52:47–58.
- 28 Australian Bureau of Statistics. Census [Internet], 2016. Available: <https://www.abs.gov.au/websitedbs/censushome.nsf/home/2016> [Accessed 29 May 2020].
- 29 Australian Bureau of Statistics [Internet]. The Australian Statistical Geography Standard (ASGS) remoteness structure [Internet]. Available: <https://www.abs.gov.au/websitedbs/D3310114.nsf/home/remoteness+structure> [Accessed 17 Apr 2020].
- 30 Jarman L, Martin A, Venn A, et al. Prevalence and correlates of psychological distress in a large and diverse public sector workforce: baseline results from Partnering Healthy@Work. *BMC Public Health* 2014;14:125.
- 31 McHugh M. Rationalization as a key stressor for public sector employees: an organizational case study. *Occup Med* 1998;48:103–12.
- 32 Murcia M, Chastang J-F, Niedhammer I. Psychosocial work factors, major depressive and generalised anxiety disorders: results from the French national SIP study. *J Affect Disord* 2013;146:319–27.
- 33 Trincheri E, Borgonovi E, Farr-Wharton B. Leader-member exchange, affective commitment, engagement, wellbeing, and intention to leave: public versus private sector Italian nurses. *Public Money & Management* 2014;34:381–8.
- 34 Parker BL, Anderson M, Batterham PJ, et al. Examining the preliminary effectiveness and acceptability of a web-based training program for Australian secondary school teachers: pilot study of the beam (building educators' skills in adolescent mental health) program. *JMIR Ment Health* 2021;8:e29989.
- 35 Bültmann U, Kant I, van Amelsvoort LG, et al. Differences in fatigue and psychological distress across occupations: results from the Maastricht cohort study of fatigue at work. *J Occup Environ Med* 2001;43:976–83.
- 36 Black Dog Institute's Online Clinic [Internet]. Sydney NSW, 2019. Available: <https://onlineclinic.blackdoginstitute.org.au/> [Accessed 05 May 2020].
- 37 Goetzel RZ, Roemer EC, Holingue C, et al. Mental health in the workplace: a call to action proceedings from the mental health in the Workplace-Public health Summit. *J Occup Environ Med* 2018;60:322–30.
- 38 The CDC worksite health scorecard; an assessment tool for employers to prevent heart disease, stroke, & related health conditions [Internet]. Available: <https://stacks.cdc.gov/view/cdc/12023> [Accessed 12 Dec 2021].
- 39 Hymel PA, Loeppke RR, Baase CM, et al. Workplace health protection and promotion: a new pathway for a healthier--and safer--workforce. *J Occup Environ Med* 2011;53:695–702.