To cite: Qu Y. Li C. Lv J.

et al. Joint trajectories of

body mass index and waist

circumference in early-life to

mid-life adulthood and incident

hypertension: the China Health

and Nutrition Survey. BMJ Open

2022;12:e059556. doi:10.1136/

Prepublication history and

for this paper are available

online. To view these files,

(http://dx.doi.org/10.1136/

bmjopen-2021-059556).

Accepted 12 April 2022

please visit the journal online

Received 24 November 2021

additional supplemental material

bmjopen-2021-059556

# **BMJ Open** Joint trajectories of body mass index and waist circumference in early-life to mid-life adulthood and incident hypertension: the China Health and Nutrition Survey

Yanlin Qu,<sup>1,2</sup> Chunxia Li,<sup>1,2</sup> Jiali Lv,<sup>1,2</sup> Bingbing Fan,<sup>1,2</sup> Ying Liu,<sup>1,2</sup> Chang Su,<sup>3</sup> Xiangjuan Zhao <sup>1</sup>/<sub>2</sub> <sup>4</sup>

# ABSTRACT

**Objective** This longitudinal study aims to identify distinct trajectories of body mass index (BMI) and waist circumference (WC) during 20–60 years old, and explore their joint effect on incident hypertension.

 $\textbf{Design} \ \ \textbf{A} \ \textbf{longitudinal cohort study}.$ 

**Setting** China Health and Nutrition Survey, 1993–2011. **Participants** The longitudinal cohort included 6571 participants (3063 men) who had BMI and WC repeatedly measured 3–7 times before incident hypertension or loss to follow-up.

**Outcomes** Hypertension was defined as systolic blood pressure/diastolic blood pressure>140/90 mm Hg or diagnosis by medical records or taking antihypertensive medication.

Results Two distinct trajectories were characterised for both BMI and WC: low-increasing and high-increasing. Jointly, subjects were divided into four groups: normal (n=4963), WC-increasing (n=620), BMI-increasing (n=309) and BMI&WC-increasing (n=679). Compared with the normal group, the adjusted HRs and 95% CIs for hypertension were 1.43 (1.19 to 1.74), 1.51 (1.19 to 1.92) and 1.76 (1.45 to 2.14) for WC-increasing, BMI-increasing and BMI&WC-increasing groups, respectively. The model-estimated levels and slopes of BMI and WC were calculated at each age point in 1-year interval according to the model parameters and their first derivatives, respectively. The associations between model-estimated levels and hypertension increased with age, with adjusted ORs and 95% Cls ranging from 0.92 (0.86 to 0.98) to 1.57 (1.47 to 1.67) for BMI and 0.98 (0.92 to 1.05) to 1.44 (1.35 to 1.53) for WC. Conversely, the ORs (95% Cls) of leveladjusted linear slopes decreased with age, ranging from 1.47 (1.38 to 1.57) to 0.97 (0.92 to 1.03) for BMI and 1.36 (1.28 to 1.45) to 0.99 (0.93 to 1.06) for WC. **Conclusions** Our study demonstrates that the joint trajectories of BMI and WC have significant effect on future hypertension risk, and the changing slopes of BMI and WC during young adulthood are independent risk factors. Both BMI and WC should be paid more attention to prevent hypertension, and young adulthood may be a crucial period for intervention.

# STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The community-based longitudinal study, with large sample size and repeated measurements, was used to explore the distinct trajectories of body mass index (BMI) and waist circumference (WC) in early-life to mid-life adulthood.
- ⇒ Distinct trajectories of BMI and WC were identified by the latent class growth mixed model, and their joint effect on incident hypertension was explored.
- ⇒ The trajectory parameters, model-estimated levels and linear slopes, were examined to identify the critical period for early prevention of hypertension.
- ⇒ Unmeasurable covariates may cause potential confounding bias.

# INTRODUCTION

As a major risk factor for cardiovascular **nig**, A training disease, hypertension is a severe problem the public health.<sup>1</sup> The prevalence of hypertension has increased rapidly over decades,<sup>2</sup> causing heavy burden to the health system in China.<sup>34</sup> Obesity is regarded as an important risk factor for incident hypertension,<sup>56</sup> and it could be evaluated by many indicators, such as body mass index (BMI) and waist circumference (WC).<sup>78</sup> The relationship between single obesity indicators and incident hypertension has been well established in observation studies.<sup>5–9</sup> However, the joint effect of those indicators on hypertension is largely unknown in a life course perspective.

Generally, obesity is divided into systemic obesity and abdominal obesity.<sup>8</sup> <sup>9</sup> Systemic obesity is mainly defined by BMI, while the abdominal obesity is often defined by WC.<sup>8</sup> The levels of BMI and WC reflect different aspects of obesity,<sup>5–12</sup> and their joint trajectories present the heterogeneous longitudinal changing patterns of body shape. Previous research has showed that BMI and WC were

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC

permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Check for updates

For numbered affiliations see end of article.

Correspondence to Dr Xiangjuan Zhao; xzhao6@hotmail.com



associated with hypertension in both cross-sectional<sup>5-12</sup> and longitudinal studies.<sup>13–15</sup> However, these studies mainly focused on the BMI and WC separately,<sup>10-15</sup> and the joint effect of BMI and WC trajectories on incident hypertension is still unclarified. We hypothesised that individuals in different joint trajectory group of BMI and WC may present different risk of hypertension, and a critical period may exist in early-life to mid-life adulthood for early prevention of hypertension.

Using data from the China Health and Nutrition Survey (CHNS), this study aims to identify BMI and WC trajectories during early-life to mid-life adulthood (20-60 years), explore the joint effect of BMI and WC trajectories on incident hypertension, and determine the potential critical period for the development of hypertension.

# **Subjects and methods**

### Study cohort

As an ongoing longitudinal cohort, the CHNS is implemented by national and local governments.<sup>16 17</sup> It is aimed at understanding how the economic and social transformation of Chinese society affects the health and nutritional status of Chinese population. Data from Beijing, Shanghai, Guizhou, Henan, Chongqing, Heilongjiang, Liaoning, Jiangsu, Zhejiang, Hubei, Hunan, Shaanxi, Shandong, Guangxi and Yunnan were collected by a multistage, random cluster process. During 1989-2011, a total of 9 cross-sectional surveys have been completed, covering 4400 households with 33 348 individuals.

In this current study, we excluded individuals younger than 20 years old or older than 60 years old (n=12 147), with BMI lower than 15 kg/m<sup>2</sup> or higher than 40 kg/m<sup>2</sup> (n=799), with WC greater than 120 cm (n=1632), and with less than three follow-up visits before loss to follow-up or incident hypertension (n=12 199). Finally, 6571 normotensive adult subjects, with 3-7 times visits, were included in this study. The mean follow-up time was 11.8 years (SD=4.2 years). BMI and WC measurements after the onset of hypertension (outcome) were excluded in our analyses. Online supplemental table S1 shows excluded respondents (n=26 777) were generally younger man, with lower baseline BMI, higher baseline systolic blood pressure (SBP) and diastolic blood pressure (DBP), lower proportion of smokers and alcohol consumers than those included.

### **Examinations**

Standing height without shoes was measured to the nearest 0.2 cm using a portable SECA stadiometer (SECA, Hamburg, Germany) at each follow-up.<sup>18</sup> Weight was measured to the nearest 0.1 kg in light clothing without shoes on a dedicated scale that was routinely calibrated.<sup>18</sup> We calculated BMI as weight in kilograms divided by height in metres squared. Smoker was defined as ever smoking cigarettes (including ex-smoker and current smoker). Alcohol drinker was defined as alcohol

consumption (including beer, liquor or wine)  $\geq 25$  g (for man) or  $\geq 15$  g (for woman) per week.

WC was defined as the midway between the lowest costal margin and the superior margin of the iliac crest, and was measured in centimetres using a SECA tapeline during 1993–2011.<sup>18</sup> After resting 5 min in a seated position, SBP and DBP were measured three times on the right upper arm using a standard mercury sphygmomanometer during 1991–2011.<sup>18</sup> The mean of the last two measurements was used for analyses.

## **Outcome**

Protected Hypertension was defined as SBP/DBP≥140/90 mm Hg and/or diagnosis by medical records and/or taking antihypertensive medication. Data for SBP, DBP, taking antihypertensive medication and incidence of diagnosis hypertension were first collected in 1991, and subse-quently in 1993, 1997, 2000, 2004, 2006, 2009 and 2011 survey. Statistical methods Hypertension was defined as SBP/DBP≥140/90 mm

The latent class growth mixed model (LCGMM) was used 2

Statistical methods The latent class growth mixed model (LCGMM) was used to identify different trajectory patterns of BMI and WC.<sup>19</sup> Structions of age (centred to 42 years, the mean age of the cohort). Multiple LCGMMs with different trajectory shapes, including linear and non-linear parameters were tested using the strategy we previously described.<sup>19</sup> We trajectory shapes, including the trajectories of BMI and WC trajectory and posterior probability. In the joint analyses, compre-hensively considering the trajectories of BMI and WC trajectory and nostrior probability. In the joint group members, buildent hypertension were explored using Cox, Artining and incident hypertension were explored using the ta-seline age, gender and baseline BMI, model 3 with further adjustment for baseline SBP and model 4 with a further adjustment for smoking and alcohol drinking. Statistical methods. We established four models for the trajectories of BMI and WC trajectory and locate the solution of the explored using Cox model 1 with no covariates, model 2 with adjustment for baseline WC was not adjusted due to its collinearity with truther adjustment for smoking and alcohol drinking. Multiple to avoid collinearity of levels and linear slopes of BMI and WC were calculated at each age point in ordels were used to examine the associations of model-estimated incident hypertension. Before logistic regression madyses, the model-estimated linear slope values of BMI and WC at each age point were adjusted for their corre-sponding levels to avoid collinearity of levels and linear slopes in the same model.<sup>20,21</sup> Standardised ORs of levels and level-adjusted slopes of BMI and WC for incident hypertension were estimated, adjusted for gender, base-ine SBP, smoking and alcohol drinking.

Table 1 Baseline characteristics by incident hypertension at follow-up							
Variable	Total	Normotensives	Incident hypertension	P value			
Ν	6571	5156	1415				
Age, years	35.4 (8.6)	34.8 (8.8)	37.8 (7.4)	<0.001			
Males, n (%)	3063 (46.6)	2329 (45.2)	734 (51.9)	<0.001			
BMI, kg/m <sup>2</sup>	22.0 (2.7)	21.8 (2.6)	22.9 (2.9)	<0.001			
WC, cm	75.8 (8.4)	75.2 (8.2)	78.0 (8.8)	<0.001			
SBP, mm Hg	110.0 (103.0, 120.0)	110.0 (101.0, 120.0)	116.0 (108.0, 120.0)	<0.001			
DBP, mm Hg	74.0 (70.0, 80.0)	72.5 (68.0, 80.0)	77.5 (70.0, 80.0)	<0.001			
Smoker, n (%)	2123 (32.5)	1578 (30.8)	545 (38.8)	<0.001			
Drinker, n (%)	2312 (35.5)	1712 (33.6)	600 (42.7)	<0.001			
Follow-up, years	11.8 (4.2)	12.0 (4.3)	11.3 (3.9)	<0.001			
Study variables are prese	ented as mean (SD), median (IQ )BP diastolic blood pressure: S	R) or n (%), appropriately.	waist circumference				

Variables were described using mean (SD), median (IQR) and n (%), as appropriate. All analyses were performed using R V.4.0.4. Hypothesis tests were two sided, and p<0.05 was considered statistically significant.

# Patient and public involvement

Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

# RESULTS

Table 1 presents baseline characteristics of study variables by incident hypertension groups during follow-up. Incident hypertension subjects (n=1415) were more likely to be older males, with higher BMI, WC, SBP and DBP, higher proportions of smokers and alcohol drinkers than normotensives.

Online supplemental tables S2 and S3 summarise LCGMM results of the BMI and WC trajectory model fitting process. We fitted models from one class to three classes of linear, quadratic and cubic curves. According to statistical criteria above, a model of quadratic parameters with two classes was chosen for both BMI and WC as the best fit. Online supplemental tables S4 and S5 present parameter estimates for the best-fitting two-class quadratic LCGMM of BMI and WC, respectively.

Figure 1 shows the longitudinal trajectories of BMI (figure 1A) and WC (figure 1B) during 20-60 years old for 6571 participants. Both BMI and WC had two distinct trajectories, labelled as low-increasing (n (%)=5583 (84.96%) for BMI and 5272 (80.53%) for WC) and high-increasing (n (%)=988 (15.04%) for BMI and 1299 (19.47%) for WC). Compared with the subjects in lowincreasing group, those in the high-increasing group had similar predicted levels at age 20 years but higher linear slopes from 20 to 60 years.

Online supplemental table S6 shows HRs and 95% CIs for the association between the BMI trajectory group

membership and incident hypertension. Compared with the BMI low-increasing group, HR (95% CI) for the BMI a high-increasing group was 1.49 (1.26 to 1.75), adjusted for **ਰ** baseline age, gender, baseline BMI, baseline WC, baseline SBP, smoking and alcohol drinking. Online supplemental table S7 presents HRs and 95% CIs for the relationship <u>e</u> between the WC trajectory classes and incident hypertension. Compared with the WC low-increasing group, to texi HR (95% CI) for the WC high-increasing group was 1.43 (1.22 to 1.68), adjusted for the same covariates above.

Jointly, subjects were divided into four joint groups according to their BMI and WC trajectories, named as normal (n=4963), WC-increasing (n=620), BMIincreasing (n=309) and BMI&WC-increasing (n=679). Individuals in the normal, WC-increasing, BMI-increasing and BMI&WC-increasing group had low-increasing WC and BMI, low-increasing BMI and high-increasing WC, high-increasing BMI and low-increasing WC, and highincreasing BMI and WC, respectively. Table 2 summarises the baseline characteristics of study variables by joint **a** classes. Compared with the normal group, individuals Dd in the BMI&WC-increasing group were more likely to be younger men with higher baseline BMI, WC, SBP and DBP, and higher proportions of smoking and alcohol drinking.

Table 3 presents HRs and 95% CIs for the association between joint group membership and incident hypertension. Compared with the normal group, the HRs (95% CIs) for WC-increasing, BMI-increasing and BMI&WCincreasing group were 1.55 (1.31 to 1.83), 1.69 (1.36 to 2.10) and 1.98 (1.71 to 2.28) in unadjusted model, respectively. After adjustment for baseline age, gender, baseline BMI, SBP, smoking and drinking status, these joint groups were still significantly associated with incident hypertension, whose HRs (95% CIs) were 1.43 (1.19 to 1.74), 1.51 (1.19 to 1.92) and 1.76 (1.45 to 2.14), respectively.

Online supplemental tables S8 and S9 show modelestimated levels and linear slopes of BMI and WC in means Α



Figure 1 (A) Predicted trajectories of BMI during 20–60 years old. The trajectories were shown in solid lines, and the 95% CIs were shown in shadow. The proportions in each trajectory were shown below solid lines. BMI, body mass index (see detailed information on the curve parameters in online supplemental table S4). (B) Predicted trajectories of WC during 20-60 years old. The trajectories were shown in solid lines, and the 95% CIs were shown in shadow. The proportions in each trajectory were shown below solid lines. WC, waist circumference (see detailed information on the curve parameters in online supplemental table S5).

(SD) by incident hypertension at follow-up. Figure 2 shows the change of model-estimated levels and linear slopes of BMI and WC during age 20-60 years in hypertension and normotension group. In figure 2A, the levels increased and linear slopes decreased with age during 20-60 in both hypertension and normotension groups. However, the hypertension group had higher levels and steeper rate of change than the normotension group. Similar results were found in WC (figure 2B). Noticeably, the difference in change rate of WC linear slopes was large between the hypertension group and normotension group.

Figure 3 presents ORs and 95% CIs of model-estimated levels and level-adjusted linear slopes of BMI and WC for incident hypertension, adjusted for gender, baseline SBP, smoking and alcohol drinking. The association between a model-estimated levels and hypertension increased during a 20–60 years old, with adjusted ORs and 95% CIs ranging  $\exists$  from 0.92 (0.86 to 0.98) to 1.57 (1.47 to 1.67) for BMI and from 0.92 (0.86 to 0.98) to 1.57 (1.47 to 1.67) for BMI and õ 0.98 (0.92 to 1.05) to 1.44 (1.35 to 1.53) for WC. This association became significant at age 25 or above for BMI and at ≥ age 23 or above for WC. Conversely, the standardised ORs (95% CIs) of level-adjusted linear slopes decreased with age,

Table 2 Characteristics at baseline and incident hypertension at follow-up by joint trajectory groups									
Variable	Normal	WC-increasing	<b>BMI-increasing</b>	BMI&WC-increasing	P value				
Ν	4963	620	309	679					
Age, years	36.3 (8.6)	32.8 (8.2)	34.3 (8.3)	31.8 (7.7)	< 0.001				
Males, n (%)	2019 (40.7)	557 (89.8)	33 (10.7)	454 (66.9)	< 0.001				
BMI, kg/m <sup>2</sup>	21.3 (2.2)	22.8 (2.3)	24.5 (3.1)	25.0 (3.4)	<0.001				
WC, cm	73.8 (7.0)	81.3 (8.7)	78.9 (8.2)	84.1 (9.9)	< 0.001				
SBP, mm Hg	110.0 (101.0, 120.0)	117.5 (110.0, 120.0)	110.0 (105.0, 120.0)	116.0 (108.0, 120.0)	<0.001				
DBP, mm Hg	72.5 (68.0, 80.0)	77.0 (70.0, 80.0)	73.0 (68.8, 80.0)	77.0 (70.0, 80.0)	<0.001				
Smoker, n (%)	1435 (29.1)	364 (59.4)	30 (9.8)	294 (43.5)	<0.001				
Drinker, n (%)	1563 (31.8)	370 (60.3)	58 (18.9)	321 (48.0)	<0.001				
Hypertension, n (%)	935 (18.8)	159 (25.6)	89 (28.8)	232 (34.2)	<0.001				

Study variables are presented as mean (SD), median (IQR) or n (%), appropriately.

BMI, body mass index; DBP, diastolic blood pressure; SBP, systolic blood pressure; WC, waist circumference.

Table 3 HRs and 95% CIs of joint trajectory groups for incident hypertension							
	Model 1	Model 2	Model 3	Model 4			
Joint trajectory groups							
Normal	Reference	Reference	Reference	Reference			
WC-increasing	1.55 (1.31 to 1.83)	1.51 (1.25 to 1.82)	1.41 (1.17 to 1.71)	1.43 (1.19 to 1.74)			
BM-increasing	1.69 (1.36 to 2.10)	1.47 (1.16 to 1.87)	1.48 (1.17 to 1.89)	1.51 (1.19 to 1.92)			
BMI&WC-increasing	1.98 (1.71 to 2.28)	1.74 (1.44 to 2.10)	1.73 (1.42 to 2.10)	1.76 (1.45 to 2.14)			
Covariates							
Baseline age		1.07 (1.06 to 1.08)	1.07 (1.06 to 1.08)	1.07 (1.06 to 1.08)			
Female		0.77 (0.69 to 0.87)	0.89 (0.79 to 1.00)	1.01 (0.85 to 1.18)			
Baseline BMI		1.10 (1.08 to 1.13)	1.07 (1.05 to 1.10)	1.07 (1.05 to 1.10)			
Baseline SBP			1.03 (1.02 to 1.04)	1.03 (1.02 to 1.04)			
Smoker				1.12 (0.96 to 1.30)			
Drinker				1.10 (0.96 to 1.26)			

BMI, body mass index; WC, waist circumference; SBP, systolic blood pressure.

ranging from 1.47 (1.38 to 1.57) to 0.97 (0.92 to 1.03) for BMI and 1.36 (1.28 to 1.45) to 0.99 (0.93 to 1.06) for WC. The associations of level-adjusted linear slopes with incident hypertension were no longer significant at age 55, indicating the importance of BMI and WC slopes in young adulthood.

# DISCUSSION

In this longitudinal study, we identified two distinct trajectories for both BMI and WC during 20-60 years old in Chinese population, and found a joint effect of BMI and WC trajectories on incident hypertension. The model-estimated levels and level adjusted slopes at each age point in a 1-year interval were further calculated for both BMI and WC, and their associations with hypertension were estimated. Though

Protected by copyright, including previous studies have demonstrated BMI or WC trajectories were associated with hypertension separately, the joint effect ġ of BMI and WC trajectories was unclarified. Our observauses rela tions support that BMI and WC trajectories have synergistic effect on future hypertension risk, and the changing slopes of BMI and WC during young adulthood are independent risk factors. Therefore, we suggest controlling both BMI and WC in young adulthood to prevent hypertension. 5

In the current study, we identified two trajectories for BMI. The low-increasing group, with baseline BMI around  $20 \text{ kg/m}^2$ , increased slowly during the 20–60 years period, while the high-increasing group with lower baseline BMI increased rapidly. Some studies, using data from China, the UK and Finland, explored BMI trajectories



(A) The mean of model-estimated levels and level-adjusted linear slopes of BMI during 20-60 years old by incident Figure 2 hypertension. (B) The mean of model-estimated levels and level-adjusted linear slopes of WC during 20-60 years old by incident hypertension. BMI, body mass index; WC, waist circumference.

text and



Figure 3 Standardised ORs and 95% Cls of modelestimated levels and level-adjusted linear slopes of BMI and WC during 20-60 years old by age for incident hypertension, adjusted for gender, baseline SBP, smoking and alcohol drinking, BMI, body mass index: SBP, systolic blood pressure; WC, waist circumference.

for childhood and early adulthood, showed that BMI may have 2–6 trajectories.<sup>13 14 20–25</sup> Though the numbers of trajectories were different among these studies, they all discovered a low-increasing or low-stable group and a high-increasing group,<sup>13 1420-25</sup> which was in line with our results. Two distinct trajectories of WC were also identified in this study. The low-increasing group, with baseline WC around 70 cm, persisted 70-80 cm during early-life to mid-life adulthood, while the high-increasing group with same baseline WC increased gradually from 70 to 95 cm. To our knowledge, the literatures on WC trajectories are still limited. Jeon *et al*<sup>15</sup> showed five trajectories for WC in individuals aged 40 years or older using data from Korea. Cheng *et al*<sup>26</sup> presented that WC may have four trajectories among Chinese population during 18-50 years old. However, these previous researches did not consider the subject-specific random effect, which was a strength in our study.<sup>15 26</sup>

Generally, obesity is divided into systemic obesity and abdominal obesity. Only studying the relationship between BMI or WC and hypertension is insufficient.<sup>13-15 20 21</sup> The joint effect of BMI and WC on hypertension should be explored. In this study, we divided participants into four subgroups according to their BMI and WC trajectories. Compared with the normal group, BMI&WC-increasing group had a higher risk of incident hypertension after adjustment for some covariates, indicating that the joint effect of BMI and WC trajectories was significant associated with hypertension. Previous researches showed BMI or WC trajectories were associated with hypertension.<sup>13–15</sup> <sup>20 26</sup> However, these studies mainly focused on the BMI and WC trajectories separately, ignoring the complementary effect between BMI and WC on incident hypertension.<sup>13–15 20 21 26</sup> Hu *et al*<sup> $\delta$ </sup> found a joint effect of BMI and WC on hypertension in a cross-sectional study, nevertheless, the longitudinal changing patterns of both BMI and

e

≥

S

WC were not considered in that study. To our knowledge, the joint effect of BMI and WC trajectories was largely unknown. Our study supports a synergistic effect between BMI and WC trajectories on incident hypertension, highlighted that we should take more attention to control both BMI and WC in early adulthood.

In this longitudinal study, the model-estimated levels increased with age and the model-estimated slopes decreased with age for both BMI and WC, which was in line with our previous studies.<sup>20 21</sup> The standardised ORs were calculated for both model-estimated levels and leveladjusted slopes, after adjusting for gender, baseline SBP, smoking and alcohol drinking. Our results showed that for both BMI and WC, level-adjusted slopes had higher 2 ORs than model-estimated levels before 30 years old. It 2 may indicate slopes of BMI and WC are more important than their levels in early adulthood and 20-30 may be a critical period for early prevention of hypertension, which is similar with previous studies.<sup>20 21</sup> Interestingly, our longitudinal study found that the linear slopes of BMI had higher ORs than those of WC during 20-25 years old. Similarly, in 43-60 years old, the levels of BMI had higher ORs than those of WC. These results may indicate that BMI plays a more important role than WC in incident with the adult-hypertension during both early-life and mid-life adulthood in Chinese population. ated

According to the life course epidemiology theory,<sup>27-29</sup> exposure at a particular period in the life span has a longterm effect on the physiological function and anatomical structure and may eventually lead to some diseases. In the current study, individuals with high-increasing trajectory patterns of BMI and WC may have some changes in physiological aspect during early adulthood, including **a** aorta root thickening, premature vascular ageing and  $\Xi$ endothelial dysfunction.<sup>30–32</sup> These changes may have accumulated effects on the way to incident hypertension. Previous research has reported meaningful high prevalence of premature vascular ageing in younger than 40 years people.<sup>31</sup> Endothelial dysfunction may also play an important role in developing hypertension<sup>32</sup> and it may impact early aorta root thickening.<sup>30</sup> Furthermore, aorta root size may play a causative role in the pathogenesis of systolic hypertension.<sup>30</sup>

There are some important strengths in our study. The current study is a community-based longitudinal study with large sample size and repeated measurements. It allows us to use LCGMM to explore trajectory of BMI 2 and WC. The LCGMM can identify distinct trajectory & for subgroup participants and permits us to analyse the **3** relationship between model-estimated levels and leveladjusted slopes and hypertension to reveal the critical period in life course. Moreover, the joint effect of BMI and WC trajectories on incident hypertension was explored, which was rarely reported in previous researches. On the other hand, some limitations should be acknowledged. First, the covariates we used may be insufficient. Some unavailable covariates, such as physically fitness, may impact the prevalence of incident hypertension. Second,

CHNS was a survey concentrated on Chinese population, suggesting our study may not be generalizable to other ethnic population. Finally, there are many indicators of obesity, besides BMI and WC. Further studies should consider more indicators such as body fat percentage and fat free mass to explore the relationship between those indicators and incident hypertension.

In conclusion, the current study identified two distinct trajectories of BMI and WC from early-life to mid-life adulthood, and found complementary trajectory patterns between BMI and WC (620 (9.4%) and 309 (4.7%) participants for WC-increasing only trajectory and BMIincreasing only trajectory). From a life course perspective, our study demonstrates the joint trajectories of BMI and WC have significant effect on future hypertension risk, and the changing slopes of BMI and WC during young adulthood are independent risk factors. These results emphasise that both BMI and WC should be paid more attention to prevent hypertension, and young adulthood may be a crucial period for intervention. More targeted strategies in the prevention of hypertension should be emphasised among early adulthood. Public health intervention for controlling modifiable risk factors during young adulthood may reduce the future prevalence and burden of hypertension among Chinese.

### **Author affiliations**

<sup>1</sup>Department of Biostatistics, School of Public Health, Cheeloo College of Medicine, Shandong University, Jinan, Shandong, China

<sup>2</sup>Institute for Medical Dataology, Cheeloo College of Medicine, Shandong University, Jinan, Shandong, China

<sup>3</sup>National Institute for Nutrition and Health, Chinese Center for Disease Control and Prevention, Beijing, China

<sup>4</sup>Department of gynaecology, Shandong Province Maternal and Child Health Care Hospital, Jinan, Shandong, China

**Contributors** YQ, CL, JL, BF, YL, CS and XZ generated the hypothesis and contributed to analytic strategy and wrote the manuscript. CS and XZ supervised the field activities and data collection and edited the manuscript. XZ was the guarantor of this study.

Funding This study was supported by grants 81973147 from National Natural Science Foundation of China and the Cheeloo Young Scholars Program of Shandong University.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. The protocol was approved by the Ethics Committee of the National Institute for Nutrition and Health, China CDC (no. 201524). Informed consent was obtained from all subjects before the investigation. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines,

terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

### ORCID iD

Xiangjuan Zhao http://orcid.org/0000-0001-5360-4324

### REFERENCES

- 1 Li Y, Yang L, Wang L, *et al.* Burden of hypertension in China: a nationally representative survey of 174,621 adults. *Int J Cardiol* 2017;227:516–23.
- 2 Gu D, Reynolds K, Wu X, et al. Prevalence, awareness, treatment, and control of hypertension in China. *Hypertension* 2002;40:920–7.
- 3 Ye R, Liu K, Zhang Z, *et al.* Health-Related quality of life of hypertension in China: a systematic review and meta-analysis. *J Cardiovasc Med* 2018;19:430–8.
- 4 Feng YJ, Wang HC, Li YC, et al. Hypertension screening and follow-up management by primary health care system among Chinese population aged 35 years and above. *Biomed Environ Sci* 2015;28:330–40.
- 5 Shihab HM, Meoni LA, Chu AY, *et al.* Body mass index and risk of incident hypertension over the life course: the Johns Hopkins precursors study. *Circulation* 2012;126:2983–9.
- 6 Robinson RF, Batisky DL, Hayes JR, et al. Body mass index in primary and secondary pediatric hypertension. *Pediatr Nephrol* 2004;19:1379–84.
- 7 Dong B, Wang Z, Yang Y, et al. Intensified association between waist circumference and hypertension in abdominally overweight children. Obes Res Clin Pract 2016;10:24–32.
- 8 Hu L, Huang X, You C, *et al.* Prevalence and risk factors of prehypertension and hypertension in southern China. *PLoS One* 2017;12:e0170238.
- 9 Feng R-N, Zhao C, Wang C, et al. Bmi is strongly associated with hypertension, and waist circumference is strongly associated with type 2 diabetes and dyslipidemia, in northern Chinese adults. J Epidemiol 2012;22:317–23.
- 10 Javed F, Aziz EF, Sabharwal MS, et al. Association of BMI and cardiovascular risk stratification in the elderly African-American females. *Obesity* 2011;19:1182–6.
- 11 Levine DA, Calhoun DA, Prineas RJ, et al. Moderate waist circumference and hypertension prevalence: the REGARDS study. Am J Hypertens 2011;24:482–8.
- 12 Soltani S, Shirani F, Chitsazi MJ, et al. The effect of dietary approaches to stop hypertension (DASH) diet on weight and body composition in adults: a systematic review and meta-analysis of randomized controlled clinical trials. Obes Rev 2016;17:442–54.
- 13 Buscot M-J, Thomson RJ, Juonala M, et al. Distinct child-to-adult body mass index trajectories are associated with different levels of adult cardiometabolic risk. *Eur Heart J* 2018;39:2263–70.
- 14 Ahanchi NS, Ramezankhani A, Munthali RJ, et al. Body mass index trajectories from adolescent to young adult for incident high blood pressure and high plasma glucose. PLoS One 2019;14:e0213828.
- 15 Jeon J, Jung KJ, Jee SH. Waist circumference trajectories and risk of type 2 diabetes mellitus in Korean population: the Korean genome and epidemiology study (KoGES). *BMC Public Health* 2019;19:741.
- 16 Popkin BM, Du Š, Zhai F, et al. Cohort Profile: The China Health and Nutrition Survey--monitoring and understanding socioeconomic and health change in China, 1989-2011. Int J Epidemiol 2010;39:1435–40.
- 17 Zhang B, Zhai FY, Du SF, et al. The China health and nutrition survey, 1989-2011. Obes Rev 2014;15 Suppl 1:2–7.
- 18 Huang L, Wang H, Wang Z, et al. Regional disparities in the association between cereal consumption and metabolic syndrome: results from the China health and nutrition survey. Nutrients 2019;11:764. doi:10.3390/nu11040764
- 19 Proust-Lima C, Philipps V, Liquet B. Estimation of extended mixed models using latent classes and latent processes: the R package lcmm. J Stat Softw 2015;78:1–56. doi:10.18637/jss.v078.i02
- 20 Fan B, Yang Y, Dayimu A, et al. Body mass index trajectories during young adulthood and incident hypertension: a longitudinal cohort in Chinese population. J Am Heart Assoc 2019;8:e011937.

# **Open access**

- 21 Lv J, Fan B, Wei M, *et al.* Trajectories of early to mid-life adulthood BMI and incident diabetes: the China health and nutrition survey. *BMJ Open Diabetes Res Care* 2020;8:e000972.
- 22 Kelly Y, Patalay P, Montgomery S, *et al.* Bmi development and early adolescent psychosocial well-being: UK millennium cohort study. *Pediatrics* 2016;138:e20160967.
- 23 Nano J, Dhana K, Asllanaj E, et al. Trajectories of BMI before diagnosis of type 2 diabetes: the Rotterdam study. Obesity 2020;28:1149–56.
- 24 Viner RM, Costa S, Johnson W. Patterns of BMI development between 10 and 42 years of age and their determinants in the 1970 British cohort study. *J Epidemiol Community Health* 2019;73:79–85.
- 25 Wang X, Dong B, Huang S, *et al.* Body mass index trajectory and incident hypertension: results from a longitudinal cohort of Chinese children and adolescents, 2006-2016. *Am J Public Health* 2020;110:1689–95. –.
- 26 Cheng C, Li Y, Ma W, et al. Trajectories of waist circumference during young adulthood and incident hypertension: the China health and nutrition survey. J Hum Hypertens 2021. doi:10.1038/s41371-021-00563-y. [Epub ahead of print: 18 Jun 2021].

- 27 Ben-Shlomo Y, Kuh D. A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. *Int J Epidemiol* 2002;31:285–93.
- 28 Lynch J, Smith GD. A life course approach to chronic disease epidemiology. *Annu Rev Public Health* 2005;26:1–35.
- 29 Mishra GD, Anderson D, Schoenaker DAJM, et al. InterLACE: a new international collaboration for a life course approach to women's reproductive health and chronic disease events. *Maturitas* 2013;74:235–40.
- 30 Farasat SM, Morrell CH, Scuteri A, et al. Pulse pressure is inversely related to aortic root diameter implications for the pathogenesis of systolic hypertension. *Hypertension* 2008;51:196–202.
- 31 Cunha PG, Cotter J, Oliveira P, et al. Pulse wave velocity distribution in a cohort study: from arterial stiffness to early vascular aging. J Hypertens 2015;33:1438–45.
- 32 Scuteri A, Tesauro M, Rizza S, et al. Endothelial function and arterial stiffness in normotensive normoglycemic first-degree relatives of diabetic patients are independent of the metabolic syndrome. Nutr Metab Cardiovasc Dis 2008;18:349–56.