

		Visit 1 (randomisation)	Fortnightly pre-op telephone call	Visit 2: Surgery	Visit 3 (6 months)*	Visit 4 (12 months)	Visit 5 (18 months)	Visit 6 (24 months)	Visit 7 (30 months)	Visit 8 (36 months)	Visit 9 (42 months)	Visit 10 (48 months; End of Intervention phase	60days after treatment end	Extended follow up phase (60, 72 & 84 months)
Clinical assessments and procedures	Data Collection from hospital records	x	x	x	x	x	x	x	x	x	x	x	x	x
	CT (thorax/abdomen/pelvis)			x*	x	x	x	x		x		x		
	Body Weight	x		x	x	x	x	x	x	x	x	x		
	Adverse event assessment		x	x	x	x	x	x	x	x	x	x	x	
	Concomitant medications	x	x	x	x	x	x	x	x	x	x	x		
	Compliance monitoring		x	x	x	x	x	x	x	x	x	x		
	Participant Questionnaire Pack	x		x	x	x	x	x	x	x	x	x		
	Pregnancy test	x												
	Blood sample for fatty acid levels	x		x	x									
	Biospecimen collection (optional)	x		x	x	x	x	x	x	x	x	x		
Trial interventions	Randomisation	x												
	IMP dispensing	x		x	x	x	x	x	x	x	x			

*In the event that macroscopic tumour remains following liver resection and curative treatment is not possible, a CT scan is carried out as soon as possible to measure the extent of existing disease.

Supplementary Figure 1