

BMJ Open Incidence of admission ionised hypocalcaemia in paediatric major trauma: protocol for a systematic review and meta-analysis

Owen Hibberd ^{1,2}, James Price ^{1,3}, Tim Harris,²
Ed Benjamin Graham Barnard ^{1,4}

To cite: Hibberd O, Price J, Harris T, *et al.* Incidence of admission ionised hypocalcaemia in paediatric major trauma: protocol for a systematic review and meta-analysis. *BMJ Open* 2023;**13**:e077429. doi:10.1136/bmjopen-2023-077429

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2023-077429>).

Received 04 July 2023

Accepted 20 October 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

¹Emergency and Urgent Care Research in Cambridge (EURECa), PACE Section, Department of Medicine, Cambridge University, Cambridge, UK

²Queen Mary University of London, Blizard Institute, London, UK

³Department of Research, Audit, Innovation, and Development (RAID), East Anglian Air Ambulance, Norwich, UK

⁴Academic Department of Military Emergency Medicine, Royal Centre for Defence Medicine, Birmingham, UK

Correspondence to

Dr Owen Hibberd;
oh296@cam.ac.uk

ABSTRACT

Introduction Hypocalcaemia forms part of the ‘diamond of death’ in major trauma, alongside hypothermia, acidosis and coagulopathy. In adults, admission hypocalcaemia prior to transfusion is associated with increased mortality, increased blood transfusion requirements and coagulopathy. Data on paediatric major trauma patients are limited. This systematic review and meta-analysis aims to describe and synthesise the available evidence relevant to paediatric trauma, admission hypocalcaemia and outcome.

Methods and analysis The Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols guidelines will be used to construct this review. A planned literature search for articles in the English language will be conducted from inception to the date of searches using MEDLINE on the EBSCO platform, CINAHL on the EBSCO platform and Embase on the Ovid platform. The grey literature will also be searched. Both title and abstract screening and full-text screening will be done by two reviewers, with an adjudicating third reviewer. Heterogeneity will be assessed using the I² test, and the risk of bias will be assessed using the ROBINS-I tool. A meta-analysis will be undertaken using ratio measures (OR) and mean differences for measures of effect. When possible, the estimate of effect will be presented along with a CI and a p value.

Ethical review and dissemination Ethical review is not required, as no original data will be collected. Results will be disseminated through peer-reviewed publications and at academic conferences.

PROSPERO registration number CRD42023425172.

BACKGROUND

Major trauma is one of the leading causes of death in children in the UK.^{1 2}

A key cause of potentially survivable death from trauma is haemorrhage.³ Uncontrollable haemorrhage may be related to the injury mechanism itself or as a result of trauma-induced coagulopathy (TIC).⁴ TIC is common, occurring in at least a quarter of haemorrhagic deaths, and has a number of proposed pathophysiological mechanisms

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The protocol follows the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols guidelines.
- ⇒ This is a novel review that addresses an area of uncertainty in the current evidence base surrounding paediatric major trauma through a systematic review and meta-analysis of published data and the grey literature.
- ⇒ The review methodology is at risk of limitation by publication bias. Where appropriate, this will be assessed using funnel plots.

that generally involve injury and shock, provoking an immunological, endothelial and platelet response.⁴ All forms of haemorrhage are further exacerbated by the ‘lethal triad’ of coagulopathy, hypothermia and acidosis.^{5–7} More recently, biochemical abnormalities such as hyperkalaemia and hypocalcaemia have been recognised to contribute to deaths from haemorrhage.^{6 8} In particular, calcium’s role is important for clot formation, vascular tone and cardiac contractility, with hypocalcaemia contributing to coagulopathy and cardiovascular decompensation.^{5 6} As such, the ‘lethal triad’ is now considered a ‘diamond of death’ with hypocalcaemia forming a key component of this deleterious combination.^{5 6} The early recognition and treatment of these components in the ‘diamond of death’ are essential for trauma resuscitation.^{5 7 9}

Rationale

The free form of calcium (ionised calcium (iCa)) is the physiologically relevant component of calcium in the blood.¹⁰ iCa is measured on blood gases, which are often taken on arrival for major trauma patients, and there is good agreement between arterial and venous measurements.¹¹ Blood

gas measurements will also record the pH and lactate, which can affect the availability of iCa.^{12 13} Ionised hypocalcaemia (iHypoCa) in major trauma patients is multifactorial.^{5–7 14} The infusion of citrated blood products is a recognised cause of hypocalcaemia in trauma due to calcium chelation with citrate.^{6 15} There is also emerging evidence in adults that early hypocalcaemia may occur in trauma patients prior to the receipt of blood products containing citrate.^{16–19} Potential pathophysiological mechanisms underpinning this include calcium binding by lactate, the intracellular influx of calcium due to ischaemia and reperfusion, impaired calcium homeostasis secondary to trauma and dilution by crystalloid fluid resuscitation.^{5–7} A systematic review and meta-analysis, which included a total of 1213 major trauma patients, 18 years or older, with a documented iCa level on admission, explored the incidence and outcomes associated with admission iHypoCa.²⁰ Studies that involved patients whose calcium concentration may have been confounded by prior blood transfusions were excluded.²⁰ Overall, the incidence of admission ionised hypocalcaemia (iHypoCa) was 56.2%, and iHypoCa was associated with increased mortality, increased blood transfusion requirements and coagulopathy.^{16–20} Evidence of admission iHypoCa and the association with adverse outcomes in adult trauma patients has led to the early measurement and replacement of calcium being recommended in adult trauma guidelines.^{21 22} Paediatric major trauma data are limited. Given the different physiology of children compared with adults, children may be more vulnerable to the effects of iHypoCa, and the results of studies involving adult major trauma patients may not be able to be extrapolated to a paediatric cohort.²³ A search of PROSPERO did not find any similar planned systematic reviews or meta-analyses. Moreover, a preliminary search of the literature has found a few heterogeneous studies, which indicate that admission iHypoCa may be less prevalent in children compared with adults.^{24–27}

Aims

The primary aim of this systematic review and meta-analysis is to explore the limited evidence related to the incidence of admission iHypoCa in paediatric major trauma patients. The review also aims to explore whether admission iHypoCa, compared with normocalcaemia, is associated with adverse clinical outcomes.

METHODS

Eligibility criteria

This proposed systematic review and meta-analysis will explore the incidence of iHypoCa in paediatric (<18 years old) major trauma patients (Injury Severity Score (ISS) >15) and explore whether admission iHypoCa (iCa <1.16 mmol/L), compared with normocalcaemia (iCa ≥1.16 mmol/L) is associated with a greater incidence of adverse outcomes.¹² An iCa of <1.16 mmol/L was chosen to reflect different levels of hypocalcaemia thresholds

across the literature and facilitate the inclusion of all relevant studies.^{24–26} The Population, Intervention, Comparison, Outcomes and Study Design (PICOS) eligibility criteria are detailed in [table 1](#).

Information sources

A planned literature search for articles in the English language will be conducted from inception to the search date using MEDLINE on the EBSCO platform, CINAHL on the EBSCO platform and Embase on the Ovid platform. The reference lists of all included studies and the grey literature will also be searched.

Search strategy

The search strategy can be found in online supplemental tables 1–3.

The search will also involve checking reference lists of retrieved articles, conference abstracts and online study results. If the data are incomplete, then the corresponding authors will be contacted for additional information.

Study records

The search strategy will be undertaken by a trained librarian and information specialist. The combined abstracts from the search strategy will be independently screened by two reviewers to identify studies meeting inclusion criteria; any duplications will be removed manually. For abstracts meeting inclusion criteria, full texts will be retrieved and again independently reviewed against the inclusion and exclusion criteria by two reviewers and an adjudicating third reviewer.

A standardised data sheet (Microsoft Excel for Mac, V.16.72, 2023) will be used to extract data from included studies to facilitate data synthesis and assessment of quality and risk of bias. The extracted data will be independently verified by the second reviewer, and any discrepancies again be adjudicated by the third reviewer.

The following data items will be extracted:

1. Hospital setting.
2. Study type.
3. Country of treatment.
4. Cohort size.
5. ISS.
6. Abbreviated Injury Scale score for injury regions.
7. Ionised hypocalcaemia definitions.
8. Incidence of admission iHypoCa (iCa <1.16 mmol/L).¹²
9. Definitions and presence of coagulopathy
10. The presence of hyperkalaemia (>5.5 mmol/L).²⁸
11. The presence of hyperlactataemia (>2.0 mmol/L).²⁹
12. Haemodynamic instability (hypotension (based on age-specific Advanced Paediatric Life Support (APLS) values) or elevated Shock Index Paediatric Age-Adjusted (SIPA)).^{30–32}
13. Administration of exogenous calcium.
14. Vasoactive medication requirements within the first 24 hours.
15. Total blood product transfusion requirement during the first 24 hours.

Table 1 PICOS strategy for inclusion and exclusion

PICOS strategy	Inclusion criteria	Exclusion criteria
P	Paediatric (<18 years) major trauma patients (injury severity score >15) with a documented iCa level on admission.	iCa level taken after the administration of blood products in the emergency department.
I	Hypocalcaemia on admission (iCa <1.16 mmol/L)	N/A
C	Normocalcaemia on admission (iCa ≥1.16 mmol/L)	N/A
O	Primary outcome include the incidence of admission ionised hypocalcaemia. Secondary outcomes include the association with physiological abnormalities: <ul style="list-style-type: none"> ▶ Haemodynamic instability ▶ Hyperkalaemia ▶ Hyperlactataemia ▶ pH abnormalities ▶ Coagulopathy and adverse outcomes: <ul style="list-style-type: none"> ▶ Vasopressor requirement within 24 hours ▶ Transfusion requirement within 24 hours ▶ Activation of the major haemorrhage protocol within 24 hours ▶ Requiring invasive (operative or interventional radiology) intervention within 24 hours ▶ Hospital length of stay ▶ Paediatric Intensive Care Unit length of stay ▶ Early mortality within 24 hours and medium mortality during episode of hospital admission (>24 hours) or within 30 days 	N/A
S	Clinical trials (randomised and non-randomised), observational studies (cohort and case-controlled) case reports, case series and literature reviews.	Systematic reviews. Opinion articles.

iCa, ionised calcium; PICOS, Population, Intervention, Comparison, Outcomes, and Study Design.

- Activation of the major haemorrhage protocol within the first 24 hours.
- Requirement for invasive (operative or interventional radiology) intervention within 24 hours.
- Hospital length of stay (LOS) (days).
- Paediatric Intensive Care Unit (PICU) LOS (days).
- Early mortality within 24 hours and medium mortality during an episode of hospital admission (>24 hours) or within 30 days.

Outcomes and prioritisation

The primary outcome of this systematic review and meta-analysis is the overall incidence of admission iHypoCa. Secondary outcomes are the associations with physiological abnormalities and adverse outcomes. Physiological abnormalities are classified dichotomously as the presence of hypotension (based on age-specific APLS values)³⁰ or elevated SIPA (0–6 years: >1.22, 7–12 years: >1.00 and 13–16 years: >0.90),^{31 32} hyperkalaemia (>5.5 mmol/L)²⁸ and hyperlactataemia (>2.0 mmol/L).²⁹ Adverse outcomes are classified dichotomously as the requirement for vasopressors, transfusion, activation of the major haemorrhage protocol or invasive (operative or interventional radiology) intervention in the first 24 hours and mortality within 30 days. Hospital LOS and PICU LOS in days are classified continuously.

Ratio measures (OR) and mean differences will be used for measures of effect. When possible, the estimate of effect will be presented along with a CI and a p value.

Risk of bias

The risk of bias will be assessed for all included studies. For any randomised controlled trials, the Grading of Recommendations Assessment, Development and Evaluation methodology will be used, and for observational studies, the Risk Of Bias In Non-randomized Studies of Interventions tool will be used.^{33 34}

The risk of publication bias will be assessed with funnel plots as appropriate.³⁵

Data synthesis

The data will be synthesised following Preferred Reporting Items for Systematic Review and Meta-Analysis guidelines. Studies will be assessed clinically (PICO) and methodologically (study design, comparability, outcome ascertainment and risk of bias). Given that current evidence is likely to be limited, the minimum number of studies is two. A preliminary search has identified four studies.^{24–27} The I² test will be conducted to determine if the data are suitable for quantitative synthesis.³⁶

Meta-analysis of effect estimates is intended and will be displayed using a forest plot. If there is limited evidence for a prespecified comparison, then the haemodynamic instability and vasopressor PICO groups may be combined. Definitions of hypocalcaemia will also be combined if required, providing values are iCa<1.16 mmol/L. Other elements are unlikely to be suitable as contingencies for a combination. If different effect measures are used, attempts will be made to transform the effect measures for meta-analysis.

A narrative synthesis and summary of effect measures (with the use of box-and-whisker plots) will be conducted if heterogeneity is deemed too substantial across studies to allow for meaningful meta-analysis or if there are major concerns about bias from the three reviewers.

Meta-analysis or narrative synthesis of elements will focus on the incidence of hypocalcaemia in paediatric trauma patients and the trend towards adverse outcomes. Subgroup analysis may be undertaken for severe iHypoCa (iCa<1.0 mmol/L).

Patient and public involvement

None.

ETHICS AND DISSEMINATION

Ethical review is not required, as no original data will be collected. Results will be disseminated through peer-reviewed publications and at academic conferences.

Acknowledgements The authors would like to acknowledge and thank Catherine Hancox and the Defence Medical Academic Library Team for their assistance with the search strategy.

Contributors OH conceptualised the protocol. OH, JP, TH and EBGB all contributed to the design, data interpretation, critical revision and final approval of the protocol.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Owen Hibberd <http://orcid.org/0000-0002-3839-1874>

James Price <http://orcid.org/0000-0002-9643-692X>

Ed Benjamin Graham Barnard <http://orcid.org/0000-0002-5187-1952>

REFERENCES

- Wolfe I, MacFarlane A, Donkin A, et al. *Why children die: death in infants, children and young people in the UK*. Royal College of Paediatrics and Child Health, 2014. Available: <https://www.rcpch.ac.uk/sites/default/files/2017-06/Why%20children%20die%20-%20Death%20in%20infants%20and%20children%20and%20young%20people%20in%20the%20UK%20-%20Part%20A%202014-05.pdf>
- Elbourne C, Cole E, Marsh S, et al. At risk child: a contemporary analysis of injured children in London and the South east of England: a prospective, Multicentre cohort study. *BMJ Paediatr Open* 2021;5:e001114.
- Davis JS, Satahoo SS, Butler FK, et al. An analysis of Prehospital deaths: who can we save *J Trauma Acute Care Surg* 2014;77:213–8.
- Moore EE, Moore HB, Kornblith LZ, et al. Trauma-induced Coagulopathy [published correction appears in *NAT Rev Dis Primers*. *Nat Rev Dis Primers* 2022;7:30.
- Wray JP, Bridwell RE, Schauer SG, et al. The Diamond of death: hypocalcemia in trauma and resuscitation. *Am J Emerg Med* 2021;41:104–9.
- DeBot M, Sauaia A, Schaid T, et al. Trauma-induced hypocalcemia. *Transfusion* 2022;62 Suppl 1:S274–80.
- Ditzel RM Jr, Anderson JL, Eisenhart WJ, et al. A review of Transfusion- and trauma-induced hypocalcemia: is it time to change the lethal Triad to the lethal diamond *J Trauma Acute Care Surg* 2020;88:434–9.
- Brohi K, Gruen RL, Holcomb JB. Why are bleeding trauma patients still dying *Intensive Care Med* 2019;45:709–11.
- Braasch MC, Turco LM, Cole EM, et al. The evolution of initial-Hemostatic resuscitation and the void of Posthemostatic resuscitation. *J Trauma Acute Care Surg* 2020;89:597–601.
- Hamroun A, Pekar J-D, Lionet A, et al. Ionized calcium: Analytical challenges and clinical relevance. *J Lab Precis Med* 2020;5:22.
- Bilkovski RN, Cannon CM, Adhikari S, et al. Arterial and venous ionized calcium measurements: is there a difference *Annals of Emergency Medicine* 2004;44:S56.
- Egi M, Kim I, Nichol A, et al. Ionized calcium concentration and outcome in critical illness. *Crit Care Med* 2011;39:314–21.
- Mirzazadeh M, Morovat A, James T, et al. Point-of-care testing of electrolytes and calcium using blood gas Analysers: it is time we trusted the results. *Emerg Med J* 2016;33:181–6.
- Kronstedt S, Roberts N, Ditzel R, et al. Hypocalcemia as a Predictor of mortality and transfusion. A Scoping review of hypocalcemia in trauma and Hemostatic resuscitation. *Transfusion* 2022;62:S158–66.
- Byerly S, Inaba K, Biswas S, et al. Transfusion-related hypocalcemia after trauma. *World J Surg* 2020;44:3743–50.
- Webster S, Todd S, Redhead J, et al. Ionised calcium levels in major trauma patients who received blood in the emergency Department. *Emerg Med J* 2016;33:569–72.
- Cherry RA, Bradburn E, Carney DE, et al. Do early ionized calcium levels really matter in trauma patients *J Trauma* 2006;61:774–9.
- Magnotti LJ, Bradburn EH, Webb DL, et al. Admission ionized calcium levels predict the need for multiple transfusions: a prospective study of 591 critically ill trauma patients. *J Trauma* 2011;70:391–5.
- Vasudeva M, Mathew JK, Fitzgerald MC, et al. Hypocalcaemia and traumatic Coagulopathy: an observational analysis. *Vox Sang* 2020;115:189–95.
- Vasudeva M, Mathew JK, Groombridge C, et al. Hypocalcemia in trauma patients: A systematic review. *J Trauma Acute Care Surg* 2021;90:396–402.
- Rossaint R, Afshari A, Bouillon B, et al. The European guideline on management of major bleeding and Coagulopathy following trauma. *Crit Care* 2023;27:80.
- Henry S. ATLS 10th edition offers new insights into managing trauma patients Bulletin of the American college of Surgeons. *Am Coll Surg* 2018;56. Available: https://bulletin.facs.org/2018/06/atls-10th-edition-offers-new-insights-into-managing-traumapatient/#Chapter_3_Shock
- Barcelona SL, Thompson AA, Coté CJ. Intraoperative pediatric blood transfusion therapy: a review of common issues. part I: hematologic and physiologic differences from adults; metabolic and infectious risks. *Paediatr Anaesth* 2005;15:716–26.
- Ciaraglia A, Lumbard D, Deschner B, et al. The effects of hypocalcemia in severely injured pediatric trauma patients. *J Trauma Acute Care Surg* 2023;95:313–8.
- Gimelraikh Y, Berant R, Stein M, et al. Early hypocalcemia in pediatric major trauma: A retrospective cohort study. *Pediatr Emerg Care* 2022;38:e1637–40.
- Epstein D, Ben Lulu H, Raz A, et al. Admission hypocalcemia in pediatric major trauma patients—an uncommon phenomenon associated with an increased need for urgent blood transfusion. *Transfusion* 2022;62:1341–6.
- Cornelius BG, Clark D, Williams B, et al. A retrospective analysis of calcium levels in pediatric trauma patients. *Int J Burns Trauma* 2021;11:267–74.
- Wolf J, Geneen LJ, Meli A, et al. Hyperkalemia following blood transfusion—a systematic review assessing evidence and risks. *Transfus Med Rev* 2022;36:133–42.
- Rishu AH, Khan R, Al-Dorzi HM, et al. Even mild Hyperlactatemia is associated with increased mortality in critically ill patients. *Crit Care* 2013;17:R197.

- 30 Samuels M, Wieteska S. Advanced Paediatric life support. In: *Advanced Paediatric Life Support: A Practical Approach to Emergencies*, 6th edn. Manchester, UK: Advanced Paediatric Life Support Group; New Jersey, USA: John Wiley & Sons Ltd, 2016.
- 31 Linnaus ME, Notrica DM, Langlais CS, *et al.* Prospective validation of the shock index pediatric-adjusted (SIPA) in blunt liver and spleen trauma: an ATOMAC+ study. *J Pediatr Surg* 2017;52:340–4.
- 32 Nordin A, Coleman A, Shi J, *et al.* Validation of the age-adjusted shock index using pediatric trauma quality improvement program data. *J Pediatr Surg* 2017:S0022-3468(17)30645-0.
- 33 Atkins D, Best D, Briss PA, *et al.* Grading quality of evidence and strength of recommendations. *BMJ* 2004;328:1490.
- 34 Sterne JA, Hernán MA, Reeves BC, *et al.* ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:i4919.
- 35 Sterne JAC, Sutton AJ, Ioannidis JPA, *et al.* Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011;343:bmj.d4002.
- 36 Migliavaca CB, Stein C, Colpani V, *et al.* Meta-Analyses of prevalence: I2 Statistic and how to deal with heterogeneity. *Res Synth Methods* 2022;13:363–7.

Online Supplementary Tables**Supplementary Table 1. CINAHL search strategy for a systematic review and meta-analysis exploring the incidence and associated outcomes with admission hypocalcaemia in paediatric major trauma**

Search ID#	Search Terms	Last Run Via	Results from 3 rd July 2023
S29	S25 AND S28	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	2
S28	S26 OR S27	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	95,090
S27	AB trauma	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	77,482
S26	TI trauma	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	39,417
S25	S23 OR S24	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	11
S24	AB paediatric hypocalcaemia or pediatric hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	2
S23	TI paediatric hypocalcaemia or pediatric hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	9
S22	S3 AND S17	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	351
S21	S3 AND S12	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	0
S20	S18 OR S19	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	19
S19	AB admission hypocalcaemia or admission hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	18

S18	TI admission hypocalcaemia or admission hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	1
S17	S15 OR S16	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	835,597
S16	AB outcomes	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	732,991
S15	TI outcomes	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	213,098
S14	S3 AND S9 AND S12	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	0
S13	S3 AND S6	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	2
S12	S10 OR S11	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	83
S11	AB normocalcaemia or normocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	78
S10	TI normocalcaemia or normocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	8
S9	S7 OR S8	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	1,406
S8	AB hospital arrival	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	1,349
S7	TI hospital arrival	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	107
S6	S4 OR S5	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	1,745
S5	AB hypocalcaemia	Interface - EBSCOhost Research Databases	1,451

	or hypocalcemia	Search Screen - Advanced Search Database - CINAHL	
S4	TI hypocalcaemia or hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	584
S3	S1 OR S2	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	1,003
S2	AB paediatric trauma patients or pediatric trauma patients	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	805
S1	TI paediatric trauma patients or pediatric trauma patients	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - CINAHL	364

Supplementary Table 2. MEDLINE search strategy for a systematic review and meta-analysis exploring the incidence and associated outcomes with admission hypocalcaemia in paediatric major trauma

Search ID#	Search Terms	Last Run Via	Results from 3 rd July 2023
S30	S26 AND S29	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	5
S29	S27 OR S28	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	276,599
S28	AB trauma	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	245,191
S27	TI trauma	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	87,100
S26	S24 OR S25	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	38
S25	AB paediatric hypocalcaemia or pediatric hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	22
S24	TI paediatric hypocalcaemia or pediatric hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	18

S23	S6 AND S22	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	4
S22	S3 AND S17	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	746
S21	S3 AND S12	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	0
S20	S18 OR S19	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	72
S19	AB admission hypocalcaemia or admission hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	71
S18	TI admission hypocalcaemia or admission hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	4
S17	S15 OR S16	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	2,179,197
S16	AB outcomes	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	2,053,450
S15	TI outcomes	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	450,999
S14	S3 AND S9 AND S12	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	0
S13	S3 AND S6	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	5
S12	S10 OR S11	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	870
S11	AB normocalcaemia or normocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	837
S10	TI normocalcaemia	Interface - EBSCOhost Research Databases	52

	a or normocalcemia	Search Screen - Advanced Search Database - MEDLINE	
S9	S7 OR S8	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	80,231
S8	AB hospital arrival or hospital admission	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	76,431
S7	TI hospital arrival or hospital admission	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	8,797
S6	S4 OR S5	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	11,987
S5	AB hypocalcaemia or hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	10,707
S4	TI hypocalcaemia or hypocalcemia	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	3,153
S3	S1 OR S2	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	2,051
S2	AB paediatric trauma patients or pediatric trauma patients	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	1,764
S1	TI paediatric trauma patients or pediatric trauma patients	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - MEDLINE	649

Supplementary Table 3. EMBASE search strategy for a systematic review and meta-analysis exploring the incidence and associated outcomes with admission hypocalcaemia in paediatric major trauma

#	Query	Results from 3rd July 2023
1	(paediatric trauma patients or pediatric trauma patients).m titl.	384
2	(paediatric trauma patients or pediatric trauma patients).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword heading word, floating subheading word, candidate term word]	1,172

3	limit 2 to abstracts	1,155
4	1 or 3	1,172
5	(hypocalcaemia or hypocalcemia).m_titl.	3,811
6	(hypocalcaemia or hypocalcemia).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword heading word, floating subheading word, candidate term word]	31,406
7	limit 6 to abstracts	26,928
8	5 or 7	27,890
9	(hospital arrival or hospital admission).m_titl.	4,381
10	(hospital arrival or hospital admission).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword heading word, floating subheading word, candidate term word]	292,976
11	limit 10 to abstracts	245,800
12	9 or 11	246,388
13	(normocalcaemia or normocalcemia).m_titl.	63
14	(normocalcaemia or normocalcemia).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword heading word, floating subheading word, candidate term word]	1,223
15	limit 14 to abstracts	1,213
16	13 or 15	1,223
17	4 and 8 and 12	1
18	4 and 8	6
19	(paediatric hypocalcaemia or pediatric hypocalcemia).m_titl.	2
20	(paediatric hypocalcaemia or pediatric hypocalcemia).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword heading word, floating subheading word, candidate term word]	4
21	limit 20 to abstracts	2
22	19 or 21	4
23	(admission hypocalcaemia or admission hypocalcemia).m_titl.	2
24	(admission hypocalcaemia or admission hypocalcemia).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword heading word, floating subheading word, candidate term word]	11
25	limit 24 to abstracts	11
26	23 or 25	11
27	4 and 16	0