# **BMJ Open** Preventive strategies for low anterior resection syndrome: a protocol for systematic review and evidence mapping

Xin-Yu Zhang,<sup>1,2</sup> Ke-Lu Yang,<sup>3</sup> Yang Li,<sup>4,5</sup> Rui-Shu Li,<sup>1</sup> Shi-Qi Wang,<sup>6</sup> Xiao-Nan Liu,<sup>1</sup> Quan Wang 💿 1

#### ABSTRACT

To cite: Zhang X-Y, Yang K-L, Li Y, et al. Preventive strategies for low anterior resection syndrome: a protocol for systematic review and evidence mapping. BMJ Open 2023:13:e077279. doi:10.1136/ bmjopen-2023-077279

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (http://dx.doi.org/10.1136/ bmjopen-2023-077279).

Received 30 June 2023 Accepted 31 October 2023

Check for updates

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

#### **Correspondence to**

Quan Wang; wangquan2013@126.com and Xiao-Nan Liu; liuxnxjh@163.com

Introduction Rectal cancer is one of the top 10 cancers worldwide. Up to 80% of patients with rectal tumours have had sphincter-saving surgery, mainly due to the large expectation of anal preservation. However, patients tend to experience low anterior resection syndrome (LARS) after rectal resection, which is disordered bowel function that includes faecal incontinence, urgency, frequent defecation, constipation and evacuation difficulties. LARS, with an estimated prevalence of 41%, has been reported to substantially decrease the quality of life of patients. However, no comprehensive preventive strategies are currently available for LARS. This systematic review aims to synthesise evidence on the current LARS preventive strategies.

Methods and analysis This protocol is reported according to the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) checklist. Literature in PubMed (via Medline), Embase and the Cochrane Library from inception to July 2023 will be searched to identify articles relevant to preventive effectiveness against LARS. The Cochrane Collaboration's risk of bias tool for randomised controlled trials and the Newcastle-Ottawa Scale for clinical controlled trials, cohort studies and case-control studies will be used to assess the risk of bias. We will group the included studies by the type of LARS prevention strategy and present an overview of the main findings in the form of evidence mapping. A meta-analysis is planned if there is no substantial clinical heterogeneity between the included studies. The Grading of Recommendations, Assessment, Development and Evaluation (GRADE) will be used to evaluate the quality of the evidence.

Ethics and dissemination Ethical approval is not needed for systematic review of published data. The findings will be published in a peer-reviewed journal and disseminated at scientific conferences.

PROSPERO registration number CRD42023402886.

#### **INTRODUCTION**

Rectal cancer is one of the top 10 cancers worldwide, with an estimated 732210 new cases (3.8% of the total cases) and 339022 new deaths (3.4% of the total deaths due to cancer) in 2020.<sup>1</sup> Radical rectectomy has remained the cardinal treatment for rectal tumours. Due to advancements in surgical

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- $\Rightarrow$  This study will comprehensively identify and summarise evidence relevant to low anterior resection syndrome (LARS) prevention strategies for perioperative management.
- $\Rightarrow$  The review will be conducted in accordance with the Cochrane Handbook for Systematic Reviews of Interventions, and the protocol was registered in PROSPERO, indicating the transparent process.
- $\Rightarrow$  Evidence mapping, with an overall and clear understanding of the current diagnosis and treatment of LARS, will be performed to contribute to clinical awareness and practical management.
- $\Rightarrow$  Large heterogeneity might exist among the included studies due to the usage of different scoring systems for evaluating bowel dysfunction.
- $\Rightarrow$  Our findings may be affected by potential publication bias because of the restriction to the English language.

and data mini techniques and patients' expectations for anal preservation, more and more patients ≥ prefer to avoid a permanent colostomy, and training, sphincter-saving surgery is performed in up to 80% of patients with rectal cancer.<sup>2</sup> However, the structural preservation of the anus does not assure the complete functional preservation of the anal sphincter complex. Ś Up to 90% of such patients were reported to experience major changes in bowel habits, including faecal incontinence, urgency, frequent defecation, diarrhoea, constipation and evacuation difficulties occurring alone or in combination.<sup>3</sup> This wide spectrum of  $\boldsymbol{\hat{G}}$ disturbed bowel movements occurring after **3** rectal resection is termed low anterior resection syndrome (LARS).<sup>4</sup>

LARS was pragmatically defined as 'disordered bowel function after rectal resection, leading to a detriment in quality of life' in 2012.<sup>3</sup> A meta-analysis found that the estimated prevalence of major LARS (regarded to have a significant impact on quality of life) was 41% (95% CI: 34% to 48%).<sup>5</sup> The

to text

Protected by copyright, including for uses related

rectum plays a role as the reservoir of stool.<sup>3</sup> Thus, resection of the rectum results in a partial loss of this function. In addition, radiotherapy, with accompanying surgical damage to the anal sphincter and anastomotic leakage, could also contribute to the deterioration of defecatory function.<sup>2-6</sup> Of more than 30 symptoms associated with LARS reported in 128 studies, the most frequently mentioned symptom was faecal incontinence (97% of the studies), followed by stool frequency (80%), flatus incontinence (70%), urgency (67%) and pad-wearing (66%).<sup>4</sup> Toileting dependence caused by such daily episodes of troublesome defecation has a strong impact not only on patients' social and daily activities but also on their mental and emotional well-being.<sup>7</sup> Furthermore, LARS tends to persist for a prolonged period, even up to 15 years.<sup>2 3</sup> And worse, a proportion of patients undergoing sphincter-saving surgery require a colostomy again to eliminate LARS.<sup>89</sup>

No current comprehensive preventive strategies are presently available for LARS. LARS is a series of multifactorial postoperative complications, and the causative factors have been identified to include radiotherapy, tumour height, anastomotic technique and diverting stoma.<sup>56</sup> Trials have been conducted to explore the modifiable variables for preventing LARS, such as preoperative consult,<sup>10</sup> colonic I-pouch or side-to-end anastomosis,<sup>1</sup> pelvic floor rehabilitation<sup>12</sup> and dietary modifications.<sup>13</sup> However, data were scattered and not systematic, and little evidence on preventive interventions for LARS has been reported in guidelines and expert consensuses. One management guideline for LARS covers its prevention and supportive care. However, most recommendations were based on expert opinions.<sup>14</sup> Therefore, this study aims to conduct a systematic review to synthesise evidence on current preventive strategies concerning LARS, which could contribute to clinical awareness and practical management. We also aim to demonstrate available evidence in the form of evidence mapping to depict a clear understanding of LARS for medical staff and patients.

#### **METHODS**

The protocol was designed in line with the Cochrane Handbook for Systematic Reviews of Interventions and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocol (PRIS-MA-P).<sup>15 16</sup>

#### Patient and public involvement

No patients will be involved in this study.

#### **Research questions**

In directing the systematic review, research questions were identified consistent with our research objectives, as follows:

1. What is the current evidence on preventive strategies for LARS before, during or after sphincter-preserving surgery?

uses rel

2. Would these interventions be effective in reducing the incidence or severity of LARS?

# **Eligibility criteria**

Studies will be included if (1) the study participants diagnosed with rectal cancer had or will have sphincterpreserving surgery; (2) the interventions of interest could avoid LARS or minimise the incidence and severity of LARS, such as faecal incontinence, urgency, frequent defecation, diarrhoea, constipation and evacuation difficulties, before, during or after surgery; (3) the outcomes of interest include at least one type of bowel dysfunction evaluated by validated scoring systems (eg, LARS scale, Jorge-Wexner scale, International Prostate Syndrome g Score or Cleveland Clinic Incontinence Score) or objective measurements (eg, bowel squeeze pressure, resting pressure, tolerated volume or the incidence of bowel dysfunction) and (4) the study type is a randomised controlled trial (RCT), clinical controlled trial, cohort study or case-control study, in which the baseline characteristics of the participants were comparable in all groups. ßu

We will exclude (1) conference abstracts, (2) ğ animal studies and (3) cross-sectional studies lacking comparisons.

#### Search strategy

lated We will develop search strategies combining LARSrelated terms used in systematic reviews and guidelines ç in this area.<sup>4 6 14</sup> A comprehensive literature search will be conducted in PubMed (via Medline), Embase and the Cochrane Library from inception until July 2023. A a manual search will also be performed by screening the references of the included studies and similar reviews to identify relevant reports missed in the search strategies. Only articles in English will be included, without restrictions on publication time. The search strategy to be used in Pubmed (via Medline) is presented in table 1. Similar ۷. strategies will be performed in Embase and the Cochrane training, Library (online supplemental file 1).

## **Study selection**

Two reviewers (X-YZ and K-LY) will independently screen the titles and abstracts of the articles identified from the S databases based on the eligibility criteria using Endnote and Rayvan software and review the full text of potentially eligible studies. Any discrepancies in the selection process will be resolved through discussion with the third reviewer (QW). After exclusion, we will report the literature selection and final totals in a PRISMA flow diagram.

#### **Data extraction**

Two reviewers (YL (Air Force Military Medical University) and RSL) will independently extract data from the included studies using a standardised data collection form designed for our study. The data charted will include but not be limited to the following items: first author, year of publication, country of origin, study design, purpose, sample size, diagnosis, tumour height, surgery, radiotherapy, anastomosis, diverting stoma,

Table 1	Search strategy in PubMed (via Medline) database
No.	Search term
#1	'Low Anterior Resection Syndrome' [MeSH] OR 'Low Anterior Resection Syndrome' [Title/Abstract] OR 'Anterior Resection Syndrome' [Title/Abstract]
#2	'Transanal Endoscopic Surgery' [MeSH] OR 'Transanal Endoscopic Microsurgery' [MeSH] OR 'Sphincter- Preserving Surgery' [Title/Abstract] OR 'Sphincter Saving Surgery' [Title/Abstract] OR 'Low Anterior Resection' [Title/Abstract] OR 'Intersphincteric Resection' [Title/Abstract] OR 'Transanal Local Resection' [Title/Abstract] OR 'Transanal Excision' [Title/Abstract] OR 'Transanal Endoscopic Microsurgery' [Title/Abstract] OR 'Transanal Minimally Invasive Surgery' [Title/Abstract] OR 'Natural Orifice Transluminal Endoscopic Surgery' [Title/Abstract] OR 'Anterior Perineal Plane for Ultra-Low Anterior Resection of the Rectum' [Title/Abstract] OR 'Transanal Total Mesorectal Excision' [Title/Abstract] OR 'Coloanal Anastomosis' [Title/Abstract]
#3	'Diarrhea' [MeSH] OR 'Fecal Incontinence' [MeSH] OR 'Constipation' [MeSH] OR Diarrhea*[Title/Abstract] OR Diarrhoea [Title/Abstract] OR 'Bowel Incontinence' [Title/Abstract] OR 'Fecal Incontinence' [Title/Abstract] OR 'Faecal Incontinence' [Title/Abstract] OR 'Faeces Incontinence' [Title/Abstract] OR 'Rectal Incontinence' [Title/ Abstract] OR Constipation [Title/Abstract] OR Obstipation [Title/Abstract] OR Dyschezia [Title/Abstract] OR 'Faecal Urgency' [Title/Abstract]
#4	#2 AND #3
#5	#1 OR #4

intervention characteristics, follow-up duration and incidence or severity of LARS. All extracted information will be cross-checked in Microsoft Excel by two reviewers for validation.

## **Risk of bias assessment**

Two reviewers (YL (Capital Medical University) and S-OW) will independently assess the risk of bias in the included studies. We will use the risk of bias tool recommended by the Cochrane Collaboration to assess RCTs.<sup>17</sup> The tool contains six domains, including selection bias, performance bias, detection bias, attrition bias, reporting bias and other bias, which can be assessed as low, unclear or high risk of bias. We will use the Newcastle-Ottawa Scale (NOS) to assess nonrandomised studies (NRS, including CCTs, cohort studies and case-control studies). The NOS contains eight items, with a total scale score of 9.<sup>18</sup> Higher scores suggest a lower risk of bias. Any disagreements regarding quality ratings will be resolved by consulting the third reviewer (QW).

#### Data synthesis and analysis

We will group the included studies by the type of LARS prevention strategies and summarise the population characteristics, intervention details and LARS measurements in each group, along with other main findings. We will calculate the relative risk with 95% CI for dichotomous data and mean differences with 95% CIs for continuous data. We will synthesise and present the results concerning the effectiveness of LARS prevention in the included studies in the form of forest plots using Review Manager (RevMan V.5.3) if there is no significant clinical heterogeneity. The results of each prevention strategy will be subtotaled and totaled simultaneously based on the different study types. However, we will not pool the data if there is significant clinical heterogeneity. The Chisquared  $(I^2)$  test will be used to evaluate the statistical heterogeneity of results between the included studies. We will use the random-effects model to synthesise the data regardless of statistical heterogeneity, as there are likely to be many sources

Protected by copyright, including of heterogeneity. We will also demonstrate the overview of ₫ our main findings in the form of evidence mapping, such as graphs and tables, for clear understanding.

We plan to conduct a subgroup analysis according to risk factors, such as radiotherapy, tumour height, anastomotic technique and diverting stoma if the relevant data are available. Tests of interaction terms will be performed to determine statistical differences in the effect estimates in different subgroups.

If more than 10 studies are included in the metaanalysis, we will assess publication bias using funnel plots.

#### **Quality of evidence**

data min Two reviewers (X-YZ and K-LY) will independently assess the quality of evidence produced in this systematic review using the Grading of Recommendations, Assessment, ⊳ Development and Evaluation (GRADE) approach.<sup>19</sup> Evidence in different study types will be assessed and graded separately using the corresponding evaluation criteria. The quality of evidence will be graded as high, moderate, low or very low. Five factors will be used to rate the quality of the evidence for risk of bias, inconsistency of results, indirectness of evidence, imprecision and publication bias. Any divergence will be resolved by r technologies discussion between two other reviewers (QW and X-NL).

Study selection is being performed and is anticipated to be completed in December 2023.

## DISCUSSION

In this review, we will systematically appraise and summarise LARS prevention strategies in the perioperative period of sphincter-preserving surgery. Sphincter-saving surgery was originally developed to preserve normal anal defecation and avoid impairing body image. However, its surgical outcomes have been disputed as a result of the extremely low life quality.<sup>20</sup> An international consensus definition of

ē

to texi

ar

## **Open access**

LARS identified eight symptoms and eight consequences, including physiology, psychology, emotion, social activities, relationships and responsibility, which covered almost all essential aspects.<sup>7</sup> Two scoping reviews reported intervention options for managing bowel symptoms after sphincter-saving rectal cancer surgery. However, they both only focused on postoperative interventions.<sup>21 22</sup> Prevention strategies presurgery and during surgery are lacking. Therefore, comprehensive prevention strategies for avoiding or minimising the incidence or severity of LARS with evidence mapping will be reported on the completion of this systematic review. According to our findings, future research recommendations will be suggested, and an evidence-based preventive strategy for preventing LARS may be conducted.

#### **Author affiliations**

<sup>1</sup>Ambulatory Surgery Center, Xijing Hospital, Air Force Military Medical University, Xi'an, China

<sup>2</sup>West China School of Nursing, Sichuan University, Chengdu, China

<sup>3</sup>Academic Center for Nursing and Midwifery, Department of Public Health and Primary Care, University of Leuven (KU Leuven), Leuven, Belgium

<sup>4</sup>Department of Nursing, Air Force Military Medical University, Xi'an, China
<sup>5</sup>Department of General Surgery, Beijing Friendship Hospital, Capital Medical University, Beijing, China

<sup>6</sup>Department of Gastrointestinal Surgery, Xijing Hospital of Digestive Diseases, Xijing Hospital, Air Force Military Medical University, Xi'an, China

**Contributors** QW, X-NL and YL (Capital Medical University) conceived the idea and designed the study protocol. X-YZ and K-LY developed the search strategy. YL (Air Force Military Medical University), R-SL, YL (Capital Medical University) and S-QW extracted data and assessed the risk of bias of included studies. X-YZ and QW drafted the manuscript. All authors read and approved the final version.

**Funding** This work is financially supported by Disciplinary Booster Programme of Xijing Hospital, China (No. XJZT21CM27, XJZT19X11 and XJZT18Z22) and Key Research and Development Project of Shaanxi Province (No. 2022ZDLSF04-04). The funders are not involved in study design, paper screening, data extraction, data synthesis, evidence evaluation and conclusions.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

#### **ORCID iD**

Quan Wang http://orcid.org/0000-0002-2821-5017

#### REFERENCES

- Sung H, Ferlay J, Siegel RL, *et al.* Global cancer Statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71:209–49.
- 2 Martellucci J. Low anterior resection syndrome: A treatment algorithm. *Dis Colon Rectum* 2016;59:79–82.
- 3 Bryant CLC, Lunniss PJ, Knowles CH, *et al*. Anterior resection syndrome. *Lancet Oncol* 2012;13:e403–8.
- 4 Keane C, Wells C, O'Grady G, et al. Defining low anterior resection syndrome: a systematic review of the literature. *Colorectal Dis* 2017;19:713–22.
- 5 Croese AD, Lonie JM, Trollope AF, et al. A meta-analysis of the prevalence of low anterior resection syndrome and systematic review of risk factors. Int J Surg 2018;56:234–41.
- 6 Sun R, Dai Z, Zhang Y, *et al*. The incidence and risk factors of low anterior resection syndrome (LARS) after sphincter-preserving surgery of Rectal cancer: a systematic review and meta-analysis. *Support Care Cancer* 2021;29:7249–58.
- 7 Keane C, Fearnhead NS, Bordeianou LG, et al. International consensus definition of low anterior resection syndrome. *Diseases of* the Colon & Rectum 2020;63:274–84.
- 8 Pachler J, Wille-Jørgensen P. Quality of life after Rectal resection for cancer, with or without permanent Colostomy. *Cochrane Database Syst Rev* 2012;12:CD004323.
- 9 Trenti L, Galvez A, Biondo S, *et al.* Quality of life and anterior resection syndrome after surgery for mid to low Rectal cancer: A cross-sectional study. *Eur J Surg Oncol* 2018;44:1031–9.
- 10 Battersby NJ, Bouliotis G, Emmertsen KJ, et al. Development and external validation of a Nomogram and online tool to predict bowel dysfunction following restorative Rectal cancer resection: the POLARS score. Gut 2018;67:688–96.
- 11 Parc Y, Ruppert R, Fuerst A, et al. Better function with a Colonic Jpouch or a side-to-end anastomosis?: A randomized controlled trial to compare the complications, functional outcome, and quality of life in patients with low Rectal cancer after a J-pouch or a side-to-end anastomosis. Ann Surg 2019;269:815–26.
- 12 Allgayer H, Dietrich CF, Rohde W, et al. Prospective comparison of Short- and long-term effects of pelvic floor exercise/Biofeedback training in patients with fecal Incontinence after surgery plus irradiation versus surgery alone for colorectal cancer: clinical, functional and endoscopic/Endosonographic findings. Scand J Gastroenterol 2005;40:1168–75.
- 13 Sun V, Crane TE, Slack SD, et al. Rationale, development, and design of the altering intake, managing symptoms (AIMS) dietary intervention for bowel dysfunction in Rectal cancer survivors. *Contemp Clin Trials* 2018;68:61–6.
- 14 Christensen P, Im Baeten C, Espín-Basany E, et al. Management guidelines for low anterior resection syndrome – the MANUEL project. Colorectal Dis 2021;23:461–75.
- 15 Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Syst Rev 2015;4:1.
- 16 Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ 2015;350:g7647.
- 17 Higgins JPT, Altman DG, Gøtzsche PC, et al. The Cochrane collaboration's tool for assessing risk of bias in randomised trials. BMJ 2011;343:d5928.
- 18 Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of Nonrandomized studies in metaanalyses. *Eur J Epidemiol* 2010;25:603–5.
- 19 Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. BMJ 2008;336:924–6.
- 20 Pucciani F. A review on functional results of sphincter-saving surgery for Rectal cancer: the anterior resection syndrome. *Updates Surg* 2013;65:257–63.
- 21 Burch J, Swatton A, Taylor C, *et al.* Managing bowel symptoms after sphincter-saving Rectal cancer surgery: A Scoping review. *J Pain Symptom Manage* 2021;62:1295–307.
- 22 Pape E, Burch J, van Ramshorst GH, et al. Intervention pathways for low anterior resection syndrome after Sphincter-Saving Rectal cancer surgery: A systematic Scoping review. *Colorectal Dis* 2023;25:538–48.

# **Search Strategies**

# 1. Medline (via PubMed) Search Strategy

No.	Search Term
#1	"Low Anterior Resection Syndrome"[MeSH] OR "Low Anterior Resection Syndrome"[Title/Abstract] OR "Anterior Resection Syndrome"[Title/Abstract]
#2	"Transanal Endoscopic Surgery"[MeSH] OR "Transanal Endoscopic Microsurgery"[MeSH] OR "Sphincter-Preserving Surgery"[Title/Abstract] OR "Sphincter Saving Surgery"[Title/Abstract] OR "Low Anterior Resection"[Title/Abstract] OR "Intersphincteric Resection"[Title/Abstract] OR "Transanal Local Resection"[Title/Abstract] OR "Transanal Excision"[Title/Abstract] OR "Transanal Endoscopic Microsurgery"[Title/Abstract] OR "Transanal Minimally Invasive Surgery"[Title/Abstract] OR "Natural Orifice Transluminal Endoscopic Surgery"[Title/Abstract] OR "Anterior Perineal Plane for Ultra-Low Anterior Resection of the Rectum"[Title/Abstract] OR "Transanal Total Mesorectal Excision"[Title/Abstract] OR "Coloanal Anastomosis"[Title/Abstract]
#3	"Diarrhea"[MeSH] OR "Fecal Incontinence"[MeSH] OR Constipation[MeSH] OR Diarrhea*[Title/Abstract] OR Diarrhoea[Title/Abstract] OR "Bowel Incontinence"[Title/Abstract] OR "Fecal Incontinence"[Title/Abstract] OR "Faecal Incontinence"[Title/Abstract] OR "Faeces Incontinence"[Title/Abstract] OR "Rectal Incontinence"[Title/Abstract] OR Constipation[Title/Abstract] OR Obstipation[Title/Abstract] OR Dyschezia[Title/Abstract] OR "Faecal Urgency"[Title/Abstract]
#4	#2 AND #3
#5	#1 OR #4

# 2. Embase Search Strategy

No.	Search Term
#1	'low anterior resection syndrome':ti,ab,kw OR 'anterior resection syndrome':ti,ab,kw
#2	'transanal endoscopic surgery'/exp OR 'transanal endoscopic microsurgery'/exp OR 'sphincter-preserving surgery':ti,ab,kw OR 'sphincter saving surgery':ti,ab,kw OR 'low anterior resection':ti,ab,kw OR 'intersphincteric resection':ti,ab,kw OR 'transanal local resection':ti,ab,kw OR 'transanal excision':ti,ab,kw OR 'transanal endoscopic microsurgery':ti,ab,kw OR 'transanal minimally invasive surgery':ti,ab,kw OR 'natural orifice transluminal endoscopic surgery':ti,ab,kw OR 'anterior perineal plane for ultra-low anterior resection of the rectum':ti,ab,kw OR 'transanal total mesorectal excision':ti,ab,kw OR 'coloanal anastomosis':ti,ab,kw
#3	'diarrhea'/exp OR 'feces incontinence'/exp OR 'constipation'/exp OR diarrhea:ti,ab,kw OR diarrhea:ti,ab,kw OR 'bowel incontinence':ti,ab,kw OR 'fecal incontinence':ti,ab,kw OR 'faecal incontinence':ti,ab,kw OR 'faeces incontinence':ti,ab,kw OR 'rectal incontinence':ti,ab,kw OR constipation:ti,ab,kw OR obstipation:ti,ab,kw OR dyschezia:ti,ab,kw OR 'faecal urgency':ti,ab,kw
#4	#2 AND #3
#5	#1 OR #4

# **3.** Cochrane Library Search Strategy

No.	Search Term
#1	MeSH descriptor: [Low Anterior Resection Syndrome] explode all trees
#2	(Low Anterior Resection Syndrome):ti,ab,kw OR (Anterior Resection Syndrome):ti,ab,kw
#3	#1 OR #2
#4	MeSH descriptor: [Transanal Endoscopic Surgery] explode all trees
#5	MeSH descriptor: [Transanal Endoscopic Microsurgery] explode all trees
#6	(Sphincter-Preserving Surgery):ti,ab,kw OR (Sphincter Saving Surgery):ti,ab,kw OR (Low Anterior Resection):ti,ab,kw OR (Intersphincteric Resection):ti,ab,kw OR (Transanal Local Resection):ti,ab,kw OR (Transanal Excision):ti,ab,kw OR (Transanal Endoscopic Microsurgery):ti,ab,kw OR (Transanal Minimally Invasive Surgery):ti,ab,kw OR (Natural Orifice Transluminal Endoscopic Surgery):ti,ab,kw OR (Anterior Perineal Plane for Ultra-Low Anterior Resection of the Rectum):ti,ab,kw OR (Transanal Total Mesorectal Excision):ti,ab,kw OR (Coloanal Anastomosis):ti,ab,kw
#7	#4 OR #5 OR #6
<b>#8</b>	MeSH descriptor: [Diarrhea] explode all trees
<b>#9</b>	MeSH descriptor: [Fecal Incontinence] explode all trees
#10	MeSH descriptor: [Constipation] explode all trees
#11	(Diarrhea*):ti,ab,kw OR (Diarrhoea):ti,ab,kw OR (Bowel Incontinence):ti,ab,kw OR (Fecal Incontinence):ti,ab,kw OR (Faecal Incontinence):ti,ab,kw OR (Faeces Incontinence):ti,ab,kw OR (Rectal Incontinence):ti,ab,kw OR (Constipation):ti,ab,kw OR (Obstipation):ti,ab,kw OR (Dyschezia):ti,ab,kw OR (Faecal Urgency):ti,ab,kw
#12	#8 OR #9 OR #10 OR #11
#13	#7 AND #12
#14	#3 OR #13