



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## The optimal concentration of ropivacaine for peripheral nerve blocks in adult patients: A protocol for systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-077876
Article Type:	Protocol
Date Submitted by the Author:	18-Jul-2023
Complete List of Authors:	Li, Jing; Affiliated Hospital of Zunyi Medical University Pan, Jiamei; Affiliated Hospital of Zunyi Medical University Xu, Ying; The Second Affiliated Hospital of Zunyi Medical University Wang, Yi; Affiliated Hospital of Zunyi Medical University Wei, Yiyong; Longgang District Maternity & Child Healthcare Hospital of Shenzhen City (Longgang Maternity and Child Institute of Shantou University Medical College), Department of Anesthesiology Zhang, Donghang; West China Hospital, Sichuan University, Chengdu, Sichuan, China, Department of Anesthesiology
Keywords:	Adult anaesthesia < ANAESTHETICS, PAIN MANAGEMENT, Adult surgery < SURGERY

SCHOLARONE™  
Manuscripts

**The optimal concentration of ropivacaine for peripheral nerve blocks in adult patients: A protocol for systematic review and meta-analysis**

Jing Li <sup>1, #</sup>, Jiamei Pan <sup>1, #</sup>, Ying Xu <sup>2</sup>, Yi Wang <sup>1</sup>, Yiyong Wei <sup>3, \*</sup>, Donghang Zhang <sup>4</sup>,  
\*

<sup>1</sup> Department of Anesthesiology, Affiliated Hospital of Zuni Medical University, Zunyi, 563000, China;

<sup>2</sup> Department of Oncology, The Second Affiliated Hospital of Zunyi Medical University, Zunyi, 563000, China;

<sup>3</sup> Department of Anesthesiology, Longgang District Maternity & Child Healthcare Hospital of Shenzhen City (Longgang Maternity and Child Institute of Shantou University Medical College), Shenzhen, 518100, China;

<sup>4</sup> Department of Anesthesiology, West China Hospital, Sichuan University, Chengdu, 610041, China.

# These authors contributed equally to this work.

**\*Address corresponding to:**

Dr. Donghang Zhang,

E-mail: zhangdhscu@163.com

Dr. Yiyong Wei,

E-mail: 295502476@qq.com

## Abstract

**Introduction:** Ropivacaine is the most widely used local anaesthetic for peripheral nerve blocks (PNB). The effects of various concentrations of ropivacaine in PNB have been investigated and compared by many randomized controlled trials (RCTs). This protocol aims to identify the optimal concentration of ropivacaine for PNB in adult patients.

**Methods and analysis:** PubMed, EMBASE, the Cochrane library, and Web of science will be searched from their inception to July 10, 2023. RCTs that compared the analgesic effects of different concentrations of ropivacaine for PNB will be included. Retrospective studies, meta-analysis, reviews, case reports, letters, conference abstracts, and pediatric studies will be excluded. The duration of analgesia will be named as the primary outcome. Secondary outcomes will include the onset time of motor and sensory blockade, postoperative pain scores, analgesics requirement over 24 hours, and the incidence of adverse effects. The study selection, data extraction, and quality assessment will be performed by two independent reviewers. Data processing and analysis will be performed by RevMan 5.4. The evidence quality will be assessed by the Grading of Recommendations Assessment, Development, and Evaluation approach (GRADE) approach.

**Ethics and dissemination:** Ethical approval is not applicable. The results of this study will be submitted to peer-reviewed journals.

**PROSPERO registration number:** CRD42023406362

**Keywords:** Ropivacaine; Concentration; Peripheral nerve blocks; RCTs; Meta-analysis; Protocol

**Strengths and limitations of this study**

- The results from subgroup analysis will provide evidence to guide ropivacaine use in certain surgical type or specific approach of PNB.
- We will use the GRADE approach to assess the quality of evidence for primary outcomes.
- Significant heterogeneity may exist among the included RCTs due to several factors, such as the type of surgery or nerve block, the usage of general anesthetics, opioids, or adjuvants, and the volumes of ropivacaine.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

## Introduction

Peripheral nerve blocks (PNB) are widely used to provide perioperative analgesia for various types of surgeries<sup>1-3</sup>. In addition to pain relief, PNB can also reduce the consumption of general anesthetics and/or opioids, decrease the incidence of postoperative complications, as well as improve the recovery quality<sup>4-6</sup>. Currently, ropivacaine is the most commonly used local anesthetic for PNB due to its lower toxicity in the central nervous system and hearts<sup>7,8</sup>. The concentration of ropivacaine used for PNB are various, and the efficacy and safety of different concentration of ropivacaine has been compared in several RCTs<sup>9-14</sup>, but the results were inconsistent. Therefore, it is meaningful to perform a systematic review and meta-analysis to determine the optimal concentration of ropivacaine for PNB, which may provide longer analgesia without increasing the incidence of adverse effects. Furthermore, we will perform subgroup analysis to find the recommended ropivacaine concentration for specific type of nerve block.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Methods and analysis**

***Study registration***

This protocol has been registered in the International Prospective Register of Systematic Reviews (PROSPERO; registration number: CRD42023406362). This study is conducted according to the Preferred Reporting Items for Systematic Evaluation and Meta-Analysis (PRISMA-P) Protocols. Ethical approval is not required.

***Search strategy***

We will search PubMed, the Cochrane library, EMBASE, and Web of science from their inception to July 10, 2023 to identify RCTs that compared the analgesic effects of different concentration of ropivacaine for PNB. The keywords for search will include “ropivacaine”, “nerve block”, and “randomized controlled trials”. The search strategy for PubMed is presented in Table 1.

***Inclusion and exclusion***

Inclusion criteria: (1) Study type: RCTs, (2) Participants: Adult patients (> 18 years) underwent surgeries with PNB, (3) Comparisons: Different concentration of ropivacaine for PNB, and (4) Primary outcomes: Duration of analgesia; Secondary outcomes: the onset time of motor and sensory blockade, postoperative pain scores, analgesics requirement over 24 hours, and the incidence of adverse effects. Retrospective studies, meta-analysis, reviews, case reports, letters, and conference abstracts will be excluded.

***Study selection***

Two authors will independently select eligible studies by screening their title, abstract, as well as the full-text. Disagreement will be resolved by discussion with a third author. The flowchart for study selection is shown in Figure 1.

***Data extraction***

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.  
Enseignement Supérieur (ABES).

The following information will be extracted: author, publication year, countries, sample, characteristics of participants, surgical type, type of anesthesia and nerve blocks, adjuvants, comparisons, outcomes, and perioperative analgesia.

### ***Risk of bias assessment***

We will assess the risk of bias for included studies using the Cochrane Collaboration's tool <sup>15</sup>. Six items will be focused: random sequence generation (selection bias); allocation concealment (selection bias); blinding of participants and personnel (performance bias); blinding of outcome assessment (detection bias); incomplete outcome data (attrition bias); and selective reporting (reporting bias). The estimated results for each item will be graded as 'unclear', 'low' or 'high'.

### ***Statistical analysis***

Data analysis will be performed by RevMan 5.4. Continuous data will be summarized using mean differences (MD) with 95% confidence intervals (CI). Dichotomous data will be summarized by risk ratios (RR) with 95% CI. Statistical heterogeneity will be assessed by the  $I^2$  test. Data will be synthesized using fixed-effect model if  $I^2 < 50\%$ . Significant heterogeneity will be considered to be existed when  $I^2 > 50\%$ , then a random-effect model will be applied. Subgroup analysis and meta-regression will be further conducted to explore the heterogeneity source. We will also perform sensitivity analysis to test whether the results are robust and reliable.  $P < 0.05$  means statistically significant. The GRADE approach will assess the quality of evidence for each outcome, and the evidence will be rated as 'very low', 'low', 'moderate', or 'high'.

### ***Patient and Public Involvement***

None.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Ethics and dissemination*

Ethical approval is not applicable. The results of this study will be submitted to peer-reviewed journals.

For peer review only

Enseignement Supérieur (ABES) .  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

## Discussion

This protocol for a systematic review and meta-analysis aims to identify the optimal concentration of ropivacaine for PNB in adult patients. Optimal concentration of ropivacaine will offer longer analgesia but not bring higher incidence of adverse effects. However, significant heterogeneity may exist among the included RCTs due to several factors, such as the type of surgery or nerve block, the usage of general anesthetics, opioids, or adjuvants, and the volumes of ropivacaine. Therefore, we will use subgroup analysis and meta-regression to explore the source of heterogeneity. We will also perform sensitivity analysis to test whether the pooled results are robust and reliable. Furthermore, subgroup analysis will recommend the optimal concentration of ropivacaine for specific nerve block or general anesthesia. Additionally, we will use the GRADE approach to assess the quality of evidence for primary outcomes. Therefore, this study may provide evidence to guide clinical use of ropivacaine for PNB in adult patients.

**Author contributions**

Conceptualization: Yiyong Wei and Donghang Zhang.

Data curation: Jing Li and Jiamei Pan.

Formal analysis: Jing Li, Jiamei Pan, Ying Xu, and Yi Wang.

Methodology: Jing Li and Yiyong Wei.

Validation: Jing Li, Jiamei Pan, and Yiyong Wei..

Writing – original draft: Jing Li, Jiamei Pan, Ying Xu, Yi Wang, and Yiyong Wei.

Writing – review & editing: Yiyong Wei and Donghang Zhang.

**Competing interests**

None.

**Funding**

This study was supported by Grant No. 2022NSFSC1399 (To Donghang Zhang) from the Natural Science Foundation of Sichuan Province.

**Acknowledgments**

None.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

## References

1. Hussain N, Brull R, Vannabouathong C, et al. Network meta-analysis of the analgesic effectiveness of regional anaesthesia techniques for anterior cruciate ligament reconstruction. *Anaesthesia* 2023;78(2):207-24. doi: 10.1111/anae.15873 [published Online First: 2022/11/04]
2. Crutchfield CR, Schweppe EA, Padaki AS, et al. A Practical Guide to Lower Extremity Nerve Blocks for the Sports Medicine Surgeon. *Am J Sports Med* 2023;51(1):279-97. doi: 10.1177/03635465211051757 [published Online First: 2022/04/20]
3. Memtsoudis SG, Cozowicz C, Bekeris J, et al. Peripheral nerve block anesthesia/analgesia for patients undergoing primary hip and knee arthroplasty: recommendations from the International Consensus on Anesthesia-Related Outcomes after Surgery (ICAROS) group based on a systematic review and meta-analysis of current literature. *Reg Anesth Pain Med* 2021;46(11):971-85. doi: 10.1136/rapm-2021-102750 [published Online First: 2021/08/27]
4. Guerra-Londono CE, Privorotskiy A, Cozowicz C, et al. Assessment of Intercostal Nerve Block Analgesia for Thoracic Surgery: A Systematic Review and Meta-analysis. *JAMA Netw Open* 2021;4(11):e2133394. doi: 10.1001/jamanetworkopen.2021.33394 [published Online First: 2021/11/16]
5. Grape S, Kirkham KR, Akiki L, et al. Transversus abdominis plane block versus local anesthetic wound infiltration for optimal analgesia after laparoscopic cholecystectomy: A systematic review and meta-analysis with trial sequential analysis. *J Clin Anesth* 2021;75:110450. doi: 10.1016/j.jclinane.2021.110450 [published Online First: 2021/07/10]
6. A NE, Singleton BN, Moorthy A, et al. Regional and neuraxial anaesthesia techniques for spinal surgery: a scoping review. *Br J Anaesth* 2022;129(4):598-611. doi: 10.1016/j.bja.2022.05.028 [published Online First: 2022/07/12]
7. Vainionpää VA, Haavisto ET, Huha TM, et al. A clinical and pharmacokinetic

- comparison of ropivacaine and bupivacaine in axillary plexus block. *Anesth Analg* 1995;81(3):534-8. doi: 10.1097/00000539-199509000-00019 [published Online First: 1995/09/01]
8. McClellan KJ, Faulds D. Ropivacaine: an update of its use in regional anaesthesia. *Drugs* 2000;60(5):1065-93. doi: 10.2165/00003495-200060050-00007 [published Online First: 2000/12/29]
9. Klein SM, Greengrass RA, Steele SM, et al. A comparison of 0.5% bupivacaine, 0.5% ropivacaine, and 0.75% ropivacaine for interscalene brachial plexus block. *Anesth Analg* 1998;87(6):1316-9. doi: 10.1097/00000539-199812000-00019 [published Online First: 1998/12/08]
10. Bertini L, Tagariello V, Mancini S, et al. 0.75% and 0.5% ropivacaine for axillary brachial plexus block: a clinical comparison with 0.5% bupivacaine. *Reg Anesth Pain Med* 1999;24(6):514-8. doi: 10.1016/s1098-7339(99)90041-x [published Online First: 1999/12/10]
11. Casati A, Fanelli G, Aldegheri G, et al. Interscalene brachial plexus anaesthesia with 0.5%, 0.75% or 1% ropivacaine: a double-blind comparison with 2% mepivacaine. *Br J Anaesth* 1999;83(6):872-5. doi: 10.1093/bja/83.6.872 [published Online First: 2000/03/04]
12. Wang S, Fang H, Qin J, et al. Comparison of the efficacy of costoclavicular space brachial plexus blockade with 0.5% versus 0.375% ropivacaine: a randomized, double-blind, single-centre, noninferiority clinical trial. *Can J Anaesth* 2023;70(1):106-15. doi: 10.1007/s12630-022-02327-9 [published Online First: 2022/09/16]
13. Jian C, Shen Y, Fu H, et al. Effects of ultrasound-guided erector spinae plane block with dexmedetomidine combined with ropivacaine of the same dose and different concentrations on analgesic effect and rehabilitation quality of patients undergoing thoracoscopic wedge resection of the lung: a prospective, randomized, controlled trial. *BMC Anesthesiol* 2022;22(1):225. doi: 10.1186/s12871-022-01768-5 [published Online First: 2022/07/17]
14. Tian Y, Zhan Y, Liu K, et al. Analgesic effects of different concentrations of


ropivacaine in transversalis fascia plane block during laparotomy. *BMC Anesthesiol* 2022;22(1):54. doi: 10.1186/s12871-022-01595-8 [published Online First: 2022/02/28]

15. Higgins J, Green SR. Cochrane Handbook for Systematic Review of Interventions Version 5.1.0. 2011

For peer review only

**Table 1. The search strategy in PubMed.**

Number	Search terms
#1	Ropivacaine [Mesh]
#2	Ropivacaine [Title/Abstract]
#3	#1 OR #2
#4	Nerve block [Title/Abstract]
#5	Nerve blockade [Title/Abstract]
#6	Peripheral nerve block [Title/Abstract]
#7	Peripheral nerve blockade [Title/Abstract]
#8	#4 OR #5 OR #6 OR #7
#9	Randomized controlled trial [Title/Abstract]
#10	Randomized [Title/Abstract]
#11	Clinical study [Title/Abstract]
#12	Clinical trial [Title/Abstract]
#13	Controlled clinical trial [Title/Abstract]
#14	#9 OR #10 OR #11 OR #12 OR #13
#15	#3 AND #8 AND #14



Flowchart of study selection

153x148mm (300 x 300 DPI)



# BMJ Open

## The optimal concentration of ropivacaine for peripheral nerve blocks in adult patients: A protocol for systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-077876.R1
Article Type:	Protocol
Date Submitted by the Author:	01-Nov-2023
Complete List of Authors:	Li, Jing; Affiliated Hospital of Zunyi Medical University Pan, Jiamei; Affiliated Hospital of Zunyi Medical University Xu, Ying; The Second Affiliated Hospital of Zunyi Medical University Wang, Yi; Affiliated Hospital of Zunyi Medical University Zhang, Donghang; West China Hospital, Sichuan University, Chengdu, Sichuan, China, Department of Anesthesiology Wei, Yiyong; Longgang District Maternity & Child Healthcare Hospital of Shenzhen City (Longgang Maternity and Child Institute of Shantou University Medical College), Department of Anesthesiology
<b>Primary Subject Heading</b>:	Anaesthesia
Secondary Subject Heading:	Surgery
Keywords:	Adult anaesthesia < ANAESTHETICS, PAIN MANAGEMENT, Adult surgery < SURGERY

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**The optimal concentration of ropivacaine for peripheral nerve blocks in adult patients: A protocol for systematic review and meta-analysis**

Jing Li <sup>1, #</sup>, Jiamei Pan <sup>1, #</sup>, Ying Xu <sup>2</sup>, Yi Wang <sup>1</sup>, Donghang Zhang <sup>3, \*</sup>, Yiyong Wei <sup>4</sup>,  
\*

<sup>1</sup> Department of Anesthesiology, Affiliated Hospital of Zuni Medical University, Zunyi, 563000, China;

<sup>2</sup> Department of Oncology, The Second Affiliated Hospital of Zunyi Medical University, Zunyi, 563000, China;

<sup>3</sup> Department of Anesthesiology, West China Hospital, Sichuan University, Chengdu, 610041, China;

<sup>4</sup> Department of Anesthesiology, Longgang District Maternity & Child Healthcare Hospital of Shenzhen City (Longgang Maternity and Child Institute of Shantou University Medical College), Shenzhen, 518100, China.

# These authors contributed equally to this work.

**\*Address corresponding to:**

Dr. Donghang Zhang,

E-mail: zhangdhscu@163.com

Dr. Yiyong Wei,

E-mail: 295502476@qq.com

## Abstract

**Introduction:** Ropivacaine is the most widely used local anaesthetic for peripheral nerve blocks (PNB). The effects of various concentrations of ropivacaine in PNB have been investigated and compared by many randomized controlled trials (RCTs). This protocol aims to identify the optimal concentration of ropivacaine for PNB in adult patients.

**Methods and analysis:** PubMed, EMBASE, the Cochrane library, and Web of science will be searched from their inception to July 10, 2023. RCTs that compared the analgesic effects of different concentrations of ropivacaine for PNB will be included. Retrospective studies, meta-analysis, reviews, case reports, letters, conference abstracts, and pediatric studies will be excluded. The duration of analgesia will be named as the primary outcome. Secondary outcomes will include the onset time of motor and sensory blockade, postoperative pain scores, analgesics requirement over 24 hours, and the incidence of adverse effects. The study selection, data extraction, and quality assessment will be performed by two independent reviewers. Data processing and analysis will be performed by RevMan 5.4. The evidence quality will be assessed by the Grading of Recommendations Assessment, Development, and Evaluation approach (GRADE) approach.

**Ethics and dissemination:** Ethical approval is not applicable. The results of this study will be submitted to peer-reviewed journals.

**PROSPERO registration number:** CRD42023406362

**Keywords:** Ropivacaine; Concentration; Peripheral nerve blocks; RCTs; Meta-analysis; Protocol

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Strengths and limitations of this study**

- Subgroup analysis will be used to explore the heterogeneity resource and provide evidence to guide ropivacaine use in certain surgical type or specific approach of PNB.
- We will use the GRADE approach to assess the quality of evidence for primary outcomes.
- Significant heterogeneity may exist among the included RCTs due to several factors, such as the type of surgery or nerve block, the usage of general anesthetics, opioids, or adjuvants, and the volumes of ropivacaine

Enseignement Supérieur (ABES) .  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

## 1 Introduction

Peripheral nerve blocks (PNB) are widely used to provide perioperative analgesia for various types of surgeries (1-3). In addition to pain relief, PNB can also reduce the consumption of general anesthetics and/or opioids, decrease the incidence of postoperative complications, as well as improve the recovery quality (4-6). Currently, ropivacaine is the most commonly used local anesthetic for PNB due to its lower toxicity in the central nervous system and hearts (7-8). The concentration of ropivacaine used for PNB are various, and the efficacy and safety of different concentration of ropivacaine has been compared in several RCTs (9-14), but the results were inconsistent. Therefore, it is meaningful to perform a systematic review and meta-analysis to determine the optimal concentration of ropivacaine for PNB, which may provide longer analgesia without increasing the incidence of adverse effects. Furthermore, we will perform subgroup analysis to find the recommended ropivacaine concentration for specific type of nerve block.



## ***Data extraction***

The following information will be extracted: author, publication year, countries, sample, characteristics of participants, surgical type, type of anesthesia and nerve blocks, adjuvants, comparisons, outcomes, and perioperative analgesia.

## ***Risk of bias assessment***

We will assess the risk of bias for included studies using the Cochrane Collaboration's tool (15). Six items will be focused: random sequence generation (selection bias); allocation concealment (selection bias); blinding of participants and personnel (performance bias); blinding of outcome assessment (detection bias); incomplete outcome data (attrition bias); and selective reporting (reporting bias). The estimated results for each item will be graded as 'unclear', 'low' or 'high'.

## ***Statistical analysis***

Data analysis will be performed by RevMan 5.4. Continuous data will be summarized using mean differences (MD) with 95% confidence intervals (CI). Dichotomous data will be summarized by risk ratios (RR) with 95% CI. Statistical heterogeneity will be assessed by the  $I^2$  test. Data will be synthesized using fixed-effect model if  $I^2 < 50\%$ . Significant heterogeneity will be considered to be existed when  $I^2 > 50\%$ , then a random-effect model will be applied. Subgroup analysis and meta-regression will be further conducted to explore the heterogeneity source. We will also perform sensitivity analysis to test whether the results are robust and reliable.  $P < 0.05$  means statistically significant. The GRADE approach will assess the quality of evidence for each outcome, and the evidence will be rated as 'very low', 'low', 'moderate', or 'high'.

## ***Patient and Public Involvement***

None.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1     ***Ethics and dissemination***

2     Ethical approval is not applicable. The results of this study will be submitted to  
3     peer-reviewed journals.

For peer review only

Enseignement Supérieur (ABES) .  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.



## Discussion

Ropivacaine is the most commonly used local anesthetic for PNB due to its lower central nervous system and cardiac toxicity. Currently, the concentration of ropivacaine used for PNB mainly varies from 0.25% to 1%. Emerging RCTs have compared the effects of different concentration of ropivacaine for PNB, but the optimal concentration remains unclear. This protocol for a systematic review and meta-analysis aims to identify the optimal concentration of ropivacaine for PNB in adult patients. Optimal concentration of ropivacaine will offer longer analgesia but not bring higher incidence of adverse effects. However, several limitations should be noticed. Firstly, significant heterogeneity may exist among the included RCTs due to several factors, such as the type of surgery or nerve block, the usage of general anesthetics, opioids, or adjuvants, and the volumes of ropivacaine. Secondly, it should be considered that the definition of the duration of analgesia among studies may be different, which will influence the results. It is better to minimize the heterogeneity by categorizing the analgesia duration differently according to their definition in individual study. Therefore, we will use subgroup analysis and meta-regression to explore the source of heterogeneity. We will also perform sensitivity analysis to test whether the pooled results are robust and reliable. Another limitation is that the number of studies that comparing the effects of different concentration of ropivacaine for PNB may be relatively small, especially for subgroup analysis. Therefore, well-designed, large sample RCTs may be needed to determine the optimal concentration of ropivacaine for PNB.

According to our protocol, subgroup analysis will recommend the optimal concentration of ropivacaine for specific nerve block or general anesthesia. Additionally, we will use the GRADE approach to assess the quality of evidence for primary outcomes. Therefore, this study may provide evidence to guide clinical use of ropivacaine for PNB in adult patients.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Author contributions**

Conceptualization: Yiyong Wei and Donghang Zhang.  
Methodology: Jing Li and Yiyong Wei.  
Validation: Jing Li, Jiamei Pan, and Yiyong Wei..  
Writing – original draft: Jing Li, Jiamei Pan, Ying Xu, Yi Wang, and Yiyong Wei.  
Writing – review & editing: Yiyong Wei and Donghang Zhang.

**Competing interests**

None.

**Funding**

This study was supported by Grant No. 2022NSFSC1399 (To Donghang Zhang) from the Natural Science Foundation of Sichuan Province.

**Acknowledgments**

None.

1  
2  
3  
4 **Figure 1.** The flowchart for study selection.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

References

1. Hussain N, Brull R, Vannabouathong C, et al. Network meta-analysis of the analgesic effectiveness of regional anaesthesia techniques for anterior cruciate ligament reconstruction. *Anaesthesia* 2023;78(2):207-24. doi: 10.1111/anae.15873 [published Online First: 2022/11/04]

2. Crutchfield CR, Schweppe EA, Padaki AS, et al. A Practical Guide to Lower Extremity Nerve Blocks for the Sports Medicine Surgeon. *Am J Sports Med* 2023;51(1):279-97. doi: 10.1177/03635465211051757 [published Online First: 2022/04/20]

3. Memtsoudis SG, Cozowicz C, Bekeris J, et al. Peripheral nerve block anesthesia/analgesia for patients undergoing primary hip and knee arthroplasty: recommendations from the International Consensus on Anesthesia-Related Outcomes after Surgery (ICAROS) group based on a systematic review and meta-analysis of current literature. *Reg Anesth Pain Med* 2021;46(11):971-85. doi: 10.1136/rapm-2021-102750 [published Online First: 2021/08/27]

4. Guerra-Londono CE, Privorotskiy A, Cozowicz C, et al. Assessment of Intercostal Nerve Block Analgesia for Thoracic Surgery: A Systematic Review and Meta-analysis. *JAMA Netw Open* 2021;4(11):e2133394. doi: 10.1001/jamanetworkopen.2021.33394 [published Online First: 2021/11/16]

5. Grape S, Kirkham KR, Akiki L, et al. Transversus abdominis plane block versus local anesthetic wound infiltration for optimal analgesia after laparoscopic cholecystectomy: A systematic review and meta-analysis with trial sequential analysis. *J Clin Anesth* 2021;75:110450. doi: 10.1016/j.jclinane.2021.110450 [published Online First: 2021/07/10]

6. A NE, Singleton BN, Moorthy A, et al. Regional and neuraxial anaesthesia techniques for spinal surgery: a scoping review. *Br J Anaesth* 2022;129(4):598-611. doi: 10.1016/j.bja.2022.05.028 [published Online First: 2022/07/12]

7. Vainionpää VA, Haavisto ET, Huha TM, et al. A clinical and pharmacokinetic

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

- 1 comparison of ropivacaine and bupivacaine in axillary plexus block. *Anesth*
- 2 *Analg* 1995;81(3):534-8. doi: 10.1097/00000539-199509000-00019
- 3 [published Online First: 1995/09/01]
- 4 8. McClellan KJ, Faulds D. Ropivacaine: an update of its use in regional anaesthesia.
- 5 *Drugs* 2000;60(5):1065-93. doi: 10.2165/00003495-200060050-00007
- 6 [published Online First: 2000/12/29]
- 7 9. Klein SM, Greengrass RA, Steele SM, et al. A comparison of 0.5% bupivacaine,
- 8 0.5% ropivacaine, and 0.75% ropivacaine for interscalene brachial plexus
- 9 block. *Anesth Analg* 1998;87(6):1316-9. doi:
- 10 10.1097/00000539-199812000-00019 [published Online First: 1998/12/08]
- 11 10. Bertini L, Tagariello V, Mancini S, et al. 0.75% and 0.5% ropivacaine for axillary
- 12 brachial plexus block: a clinical comparison with 0.5% bupivacaine. *Reg*
- 13 *Anesth Pain Med* 1999;24(6):514-8. doi: 10.1016/s1098-7339(99)90041-x
- 14 [published Online First: 1999/12/10]
- 15 11. Casati A, Fanelli G, Aldegheri G, et al. Interscalene brachial plexus anaesthesia
- 16 with 0.5%, 0.75% or 1% ropivacaine: a double-blind comparison with 2%
- 17 mepivacaine. *Br J Anaesth* 1999;83(6):872-5. doi: 10.1093/bja/83.6.872
- 18 [published Online First: 2000/03/04]
- 19 12. Wang S, Fang H, Qin J, et al. Comparison of the efficacy of costoclavicular space
- 20 brachial plexus blockade with 0.5% versus 0.375% ropivacaine: a randomized,
- 21 double-blind, single-centre, noninferiority clinical trial. *Can J Anaesth*
- 22 2023;70(1):106-15. doi: 10.1007/s12630-022-02327-9 [published Online First:
- 23 2022/09/16]
- 24 13. Jian C, Shen Y, Fu H, et al. Effects of ultrasound-guided erector spinae plane
- 25 block with dexmedetomidine combined with ropivacaine of the same dose and
- 26 different concentrations on analgesic effect and rehabilitation quality of
- 27 patients undergoing thoracoscopic wedge resection of the lung: a prospective,
- 28 randomized, controlled trial. *BMC Anesthesiol* 2022;22(1):225. doi:
- 29 10.1186/s12871-022-01768-5 [published Online First: 2022/07/17]
- 30 14. Tian Y, Zhan Y, Liu K, et al. Analgesic effects of different concentrations of

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


1           ropivacaine in transversalis fascia plane block during laparotomy. *BMC*  
2           *Anesthesiol* 2022;22(1):54. doi: 10.1186/s12871-022-01595-8 [published  
3           Online First: 2022/02/28]  
4   15. Higgins J, Green SR. Cochrane Handbook for Systematic Review of  
5           InterventionsVersion 5.1.0. 2011

For peer review only

Enseignement Supérieur (ABES) .  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

**Table 1. The search strategy in PubMed.**

Number	Search terms
#1	Ropivacaine [Mesh]
#2	Ropivacaine [Title/Abstract]
#3	#1 OR #2
#4	Nerve block [Title/Abstract]
#5	Nerve blockade [Title/Abstract]
#6	Peripheral nerve block [Title/Abstract]
#7	Peripheral nerve blockade [Title/Abstract]
#8	#4 OR #5 OR #6 OR #7
#9	Randomized controlled trial [Title/Abstract]
#10	Randomized [Title/Abstract]
#11	Clinical study [Title/Abstract]
#12	Clinical trial [Title/Abstract]
#13	Controlled clinical trial [Title/Abstract]
#14	#9 OR #10 OR #11 OR #12 OR #13
#15	#3 AND #8 AND #14



The flowchart for study selection

153x148mm (300 x 300 DPI)



## Search strategies

### EMBASE

#1 'Ropivacaine'/exp OR 'Ropivacaine':ti,ab,kw

#2 'Nerve block'/exp OR 'Nerve block':ti,ab,kw OR 'Nerve blockade':ti,ab,kw OR  
'Peripheral nerve block':ti,ab,kw OR 'Peripheral nerve blockade':ti,ab,kw

#3 'Randomized controlled trial'/exp OR 'Randomized controlled trial':ti,ab,kw OR  
'Controlled clinical trial':ti,ab,kw OR 'Clinical trial':ti,ab,kw OR 'Clinical  
study':ti,ab,kw OR 'Randomized':ti,ab,kw

#4 #1 AND #2 AND #3

### Cochrane library Trials

#1 MeSH descriptor: (Ropivacaine) explode all trees

#2 (Ropivacaine):ti,ab,kw

#3 #1 OR #2

#4 (Nerve block):ti,ab,kw

#5 (Nerve blockade):ti,ab,kw

#6 (Peripheral nerve block):ti,ab,kw

#7 (Peripheral nerve blockade):ti,ab,kw

#8 #4 OR #5 OR #6 OR #7

#9 (Randomized controlled trial):ti,ab,kw

#10 (Controlled clinical trial):ti,ab,kw

#11 (Clinical trial):ti,ab,kw

#12 (Clinical study):ti,ab,kw

#13 (Randomized):ti,ab,kw

#14 #9 OR #10 OR #11 OR #12 OR #13

#15 #3 AND #8 AND #14

### Web of science

#1 Ropivacaine (Topic)

#2 Nerve block (Topic) OR Nerve blockade (Topic) OR Peripheral nerve block (Topic)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

OR Peripheral nerve blockade (Topic)  
#3 Randomized controlled trial (Topic) OR Controlled clinical trial (Topic) OR  
Clinical trial (Topic) OR Clinical study (Topic) OR Randomized (Topic)  
#4 #1 AND #2 AND #3

For peer review only

Enseignement Supérieur (ABES) .  
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

# PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol\*

Section and topic	Item No	Checklist item	Reported on page number
<b>ADMINISTRATIVE INFORMATION</b>			
Title:			
Identification	1a	Identify the report as a protocol of a systematic review	Page 1
Update	1b	If the protocol is for an update of a previous systematic review, identify as such	NA
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number	Page 5
Authors:			
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author	Page 1
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review	Page 9
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments	NA
Support:			
Sources	5a	Indicate sources of financial or other support for the review	Page 9
Sponsor	5b	Provide name for the review funder and/or sponsor	Page 9
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol	Page 9
<b>INTRODUCTION</b>			
Rationale	6	Describe the rationale for the review in the context of what is already known	Page 4
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	Page 4
<b>METHODS</b>			
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review	Page 5
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage	Page 5
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits such that it could be repeated	Page 5

Study records:			
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review	Page 5
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)	Page 5
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently in duplicate), any processes for obtaining and confirming data from investigators	Page 6
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications	Page 6
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale	Page 5
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis	Page 6
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised	Page 6
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I <sup>2</sup> and Kendall's $\tau$ )	Page 6
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression)	Page 6
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned	Page 6
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)	Page 6
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)	Page 6

**\* It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

*From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.*