



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Temporal changes in the burden of leukaemia and lymphoma in the Australasia and Oceania regions, 2010-2019: an analysis of the Global Burden of Disease Study 2019

Journal:	BMJ Open
Manuscript ID	bmjopen-2024-084943
Article Type:	Original research
Date Submitted by the Author:	01-Feb-2024
Complete List of Authors:	Ho, Thi Quynh Anh; Deakin University School of Health and Social Development, Deakin Health Economics Lee, Peter; Deakin University, School of Health and Social Development; Gao, Lan; Deakin University School of Health and Social Development, Deakin Health Economics
Keywords:	Lymphoma < HAEMATOLOGY, Leukaemia < HAEMATOLOGY, EPIDEMIOLOGIC STUDIES

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Title page

Title

Temporal changes in the burden of leukaemia and lymphoma in the Australasia and Oceania regions, 2010-2019: an analysis of the Global Burden of Disease Study 2019

Authors

Thi Quynh Anh Ho*^{#1} - tqho@deakin.edu.au

Peter Lee*¹ - peter.lee@deakin.edu.au

Lan Gao¹ - lan.gao@deakin.edu.au

¹ Deakin University, Institute for Health Transformation, Deakin Health Economics, School of Health and Social Development, Melbourne, Victoria, Australia

*Joint first authors

Corresponding author

T.Q.A Ho

221 Burwood Highway, Burwood, VIC 3125, Australia

E-mail: tqho@deakin.edu.au

Phone: +61 403 844 808

Word count: 3 552 (excluding abstract, keywords, and references)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives

Leukaemias and lymphomas are among the most prevalent and significant cancers in Australasia and Oceania. This study aims to examine the burden of leukaemias/lymphomas and its temporal trend in Australasia and Oceania from 2010 to 2019.

Design

Epidemiological study

Methods

Data from the Global Burden of Disease (GBD) 2019 was used to examine the burden of leukaemia/lymphoma key subtypes (acute lymphocytic leukaemia (ALL), acute myeloid leukaemia (AML), chronic lymphocytic leukaemia (CLL), chronic myeloid leukaemia (CML), Hodgkin-lymphoma (HL), and non-Hodgkin lymphoma (NHL)) by sex and age group, in terms of incidence, prevalence, disability-adjusted-life-years (DALYs), and deaths. Estimated average percentage changes (EAPCs) were calculated to assess the temporal trends of leukaemia/lymphoma burden (incidence/prevalence/DALYs/deaths) from 2010 to 2019 in Australasia and Oceania.

Results

AML and NHL were the leading causes of leukaemia/lymphoma burden in both regions. From 2010 to 2019, Australasia observed an increasing trend in incidence/prevalence/deaths across most leukaemias/lymphomas and increasing/stable trend in DALYs for AML/CLL/NHL, while Oceania observed increasing trends in incidence/prevalence/DALYs for CLL/NHL and stable trends in all outcomes (except for prevalence (stable)) for AML. Contrasting mortality trends for ALL/CML/HL were observed between the two regions (increasing/stable in Australasia and decreasing in Oceania). Statistically significant

differences were observed in disease burden trends between sexes, with males experiencing a greater increase (or smaller decrease) in the burden for AML in both regions.

Conclusions

Different temporal trends in leukaemia/lymphoma burden observed in two closely situated geographic regions with different socio-demographic indices highlights the necessity for region-specific intervention strategies to enhance the access to innovative disease treatments, reducing leukaemia/lymphoma burden.

Keywords: Burden of disease, lymphoma, leukaemia

Strengths and limitations of this study:

- We compared the disease burden and temporal trends in two geographically similar regions with differing socio-demographic indices (SDI) and performed a sensitivity analysis using the world standard population, that facilitated the comparison of trends between Australasia and Oceania, and to other regions.
- Using region-specific data helps reflect disease burden distributions and trends across sexes and age groups more comprehensively and accurately.
- A direct comparison between Australasia and Oceania is limited due to considerable differences in population distributions and risk factors between the two regions.
- Analysing aggregate data made it challenging to discern the impact of various exposures to disease outcomes, thus trends should be interpreted with caution.

INTRODUCTION

Haematological malignancies, including leukaemias and lymphomas, arise from the uncontrolled proliferation of cells in the lymphatic or circulatory systems. Based on the Global Burden of Disease, Injuries and Risk Factors Study 2019 (GBD 2019), which provides the most comprehensive estimates of global disease and injury burden to date, haematological malignancies contribute to a considerable proportion of the global disease burden attributed to cancer ¹⁻³. Globally, leukaemias and lymphomas contributed to 11·7 million and 8·2 million disability-adjusted life years (DALYs) in 2019, respectively ². Studies exploring the temporal trend in haematological malignancies across countries using data from the GBD 2019 study have found that over a 30-year period, age-standardised mortality/DALYs have declined, against a background of increasing incident/prevalent burden. However, the distribution of disease burden and temporal trends in leukaemias/lymphomas varies across geographic regions and varying levels of socioeconomic development ^{1 2 4 5}. Differences in disease burden across regions of high/low socioeconomic development were largely attributed to social and environmental factors including poverty, educational attainment, and access to health care ^{1 2 4}. These large disparities in the health care system highlight the need for population-based epidemiological studies in low-and middle-income countries (LMICs) to inform public health policy and healthcare delivery planning ^{1 2 4 5}. Importantly, no studies were identified that explored trends in disease incidence/prevalence or burden for Australasia and Oceania ^{1 2}. Australasia and Oceania may be classified as regions with high and low socioeconomic development, respectively, despite being in the same geographic region of the Pacific ocean ^{3 3}. Furthermore, there are considerable discrepancies in the ethnic composition between Oceania and Australasia which may contribute to disparities in health outcomes ⁶. Therefore, a comparison of leukaemia/lymphoma trends between the two regions may facilitate the

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

understanding of the impacts attributed to differences in socio-demographic indices (SDIs) on disease occurrence and outcomes.

Hence, this study aims to (1) examine the prevalence, incidence, mortality, and DALYs attributed to leukaemias and lymphomas by sex and age groups and (2) explore the temporal trend in prevalence, incidence, mortality, and DALYs for leukaemias and lymphomas from 2010 to 2019 in Oceania and Australasia regions using GBD 2019 data.

METHODS

Data source

We extracted data from the GBD 2019 and performed a secondary analysis in the current study. All data was collected using Global Health Data Exchange (GHDx) query tool (<http://ghdx.healthdata.org/gbd-results-tool>)³⁷. Details pertaining to the collection, processing, and generation of the GBD 2019 study dataset have been described elsewhere²³.

Case definition

The definition of leukaemias and lymphomas used in the GBD 2019 study has been defined previously²³. Leukaemia is typically classified by the type of white blood affected (lymphocytic or myeloid) and disease progression (acute or chronic)⁸. Key leukaemia subtypes include acute lymphocytic leukaemia (ALL), acute myeloid leukaemia (AML), chronic lymphocytic leukaemia (CLL), and chronic myeloid leukaemia (CML)⁸. Lymphomas are categorised based on the type of lymphocyte affected, with key lymphoma subtypes including non-Hodgkin's lymphomas (NHL) and Hodgkin's lymphoma (HL)⁹.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Population and outcome

In this study, we assessed the burden of disease (including prevalence, incidence, deaths, and DALYs) of leukaemias and lymphomas by subtype in Australasia and Oceania from 2010 to 2019. In line with GBD 2019 definitions, Australasia was defined as Australia and New Zealand, and Oceania (18 countries) was defined as American Samoa, Cook Islands, Fiji, Guam, Kiribati, Marshall Islands, Micronesia, Nauru, Niue, Northern Mariana Islands, Palau, Papua New Guinea, Samoa, Solomon Islands, Tokelau, Tonga, Tuvalu, and Vanuatu ³.

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Patient and public involvement

Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Statistical analysis

A descriptive analysis was performed to characterise the regional burden of leukaemias and lymphomas (by subtype). The number of prevalent and incident cases, deaths, and DALYs were used to derive age-standardised rates (ASRs) (per 100 000 population) in Australasia and Oceania from 2010 to 2019. ASRs were calculated to exclude the impact of age structure on overall population prevalence, incidence, mortality, and DALYs as the number of cases, deaths and DALYs of cancers varies greatly across different age groups ¹⁷. The ASR per 100

000 population is the weighted-average of age-group-specific rates, that is, $ASR = \frac{\sum_{i=1}^A a_i w_i}{\sum_{i=1}^A w_i}$

$\times 100\,000$, where a_i is the age-group specific rate i and w_i is the weight of age group i of the population ⁷. For the purposes of the base-case analysis, the number of outcomes (incidence, prevalence, deaths or DALYs) occurring for each five-year age group (< 5 years to 85+), were divided by the number of individuals estimated in each age group for

Australasia (or Oceania) to estimate age-group-specific rates for each year (a_i). This was multiplied by the proportional distribution (w_i) of individuals in each age group to estimate the ASR.

The estimated average percentage change (EAPC) in ASRs was estimated to assess temporal trends in prevalence, incidence, mortality, and DALYs of leukaemias and lymphomas (by subtype) over a 10-year period (2010 to 2019, inclusive). EAPCs were calculated through generalised linear regression modelling (GLM) with a gaussian family and log-link to estimate trends: $y = \alpha + \beta * x + \varepsilon$, where y is $\ln(\text{ASR})$ and x is the calendar year. The EAPC was expressed as $100 * (e^{\beta} - 1)$ ¹. If the estimated β -coefficient and its 95% CI were both positive, it indicated an upward trend in ASR, whereas a negative β -coefficient and its 95% CI indicated a downward trend in ASR¹⁷. Otherwise, the temporal trend in outcomes were assumed to be stable. P-values < 0.05 were considered statistically significant.

All calculations were performed using Stata 17.0 statistical software (Stata Corp, College Station, TX).

Sensitivity analysis

As described above, ASRs were calculated in our base case analysis based on the estimated population size across 5-year age groups for Australasia and Oceania. To allow for comparability between trends in Australasia and Oceania, a sensitivity analysis was performed by applying the world standard population weights to age-group-specific rates for key outcomes estimated in our base case analysis¹⁰. Results of the sensitivity analysis are presented in Appendix B.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Table 1 summarises the leukaemia and lymphoma burden in terms of incidence, prevalence, DALYs, and deaths across Australasia and Oceania and associated temporal trends. Table 2 summarises trends in leukaemia and lymphoma burden by sex. Figures 1 summarises the leukaemia and lymphoma burden in terms of incidence, prevalence, DALYs, and deaths across Australasia and Oceania, by age group, and sex in 2019, respectively.

[Table 1 around here]

[Table 2 around here]

[Figure 1 around here]

Overview of disease burden

Leukaemia and lymphoma burden in Australasia

Across Australasia, the greatest contributor to incidence of leukaemia and lymphoma in 2019 was CLL (ASR: 5·22 per 100 000) and NHL (ASR: 19·06 per 100 000), respectively. ALL accounted for the largest share in paediatric leukaemia incidence, whereas CLL accounted for the largest share of incident cases for adult leukaemias. NHL (ASRs: 161·68 DALYs and 8·02 deaths per 100 000), AML (ASRs: 89·01 DALYs and 4·15 deaths per 100 000), followed by CLL, contributed to the greatest proportion in terms of DALYs and mortality burden in 2019. NHL was the most prevalent haematological cancer in 2019. Following stratification of disease burden by five-year age band and sex, the incidence, prevalence, DALY, and mortality rates across five-year age groups generally increased with ascending age across leukaemia/lymphoma subtypes for Australasia (see Figure 1 and Appendix A).

Age-specific incidence/prevalence and DALY rates for ALL and HL followed a bimodal pattern.

Leukaemia and lymphoma burden in Oceania

In Oceania, AML and ALL drove leukaemia burden (incident/prevalent/DALYs/deaths) and NHL was the key contributor to the incident lymphoma burden (ASR: 1·08 per 100 000) in 2019. HL (ASR: 0·81 per 100 000) drove prevalent lymphoma burden. Regarding the burden of disease (DALYs/deaths), AML (ASRs: 49·16 DALYs and 0·91 deaths per 100 000) and NHL (ASRs: 38·30 DALYs and 0·98 deaths per 100 000) were the largest contributors in 2019. Disease burden across age-groups and sexes for Oceania were broadly comparable with Australasia. That is, incident/prevalent burden, as well as DALYs/deaths, increased with increasing age and for male sex. (Table 1)

Temporal trends in leukaemia and lymphomas (base-case analysis)

Australasia

The incidence and prevalence of haematological malignancies across Australasia increased significantly over a 10-year period (2010 to 2019, inclusive) ($P<0\cdot001$). Temporal increasing trends in incident/prevalent cases were largely driven by CLL, ALL, and CML (EAPCs ranging from 2·16% to 2·87%) (all $P<0\cdot001$). Notably, the EAPC and associated 95% CI for leukaemia and lymphoma incidence was broadly comparable between sexes with the exception of CML and HL (greater for females), and AML (greater for males). Regarding prevalent burden, EAPCs in males were found to be higher for AML, and lower for HL relative to females.

DALY burden (per 100 000) increased significantly for CLL (EAPC: 1·74%) and AML (EAPC: 1·15%) over time, whereas significant reductions in DALY burden attributed to

CML (EAPC: -1·23%), HL (EAPC: -0·40%) and ALL (EAPC: -0·36%) were observed (all $P<0·05$). DALY burden attributed to NHL remained stable over time (both sexes). Male sex drove temporal reductions in DALYs (ALL, CML, and HL) whereas the only significant reduction in DALY burden for females was for HL. Although AML and CLL DALY burden increased for both sexes, the EAPC estimated for AML was of greater magnitude for males relative to females (EAPCs: 1·51% versus 0·69%). Lastly, mortality from leukaemias/lymphomas significantly increased over time (EAPCs ranging from 0·75% for ALL to 2·11% for CLL), with the exception of CML and HL (stable). Overall, trends in mortality were comparable between sexes, with the exception of greater increases in mortality over time for AML and NHL for males compared with females (EAPCs: 2·55% versus 1·46%). (Table 2)

Oceania

The increased incidence in CLL (EAPC: 1·38%) and NHL (EAPC: 0·92%) was balanced with significantly decreases in ALL (EAPC: -0·45%) and CML (EAPC: -0·58%) incidence. The overall stable trend in HL incidence over time was a result of opposing temporal trends in males (decreasing with EAPC: -0·19%) and females (increasing with EAPC: 0·31%). The overall incidence of AML was stable, driven by females (stable), despite increasing over time for males (EAPC: 0·17%). Although CLL incidence increased over time for both sexes, the EAPC was higher for females. Significant increases in the prevalence of NHL (EAPC: 3·24%), CLL (EAPC: 1·99%) and HL (EAPC: 0·68%) were balanced with reductions in the prevalence of CML (EAPC: -0·76%), ALL (EAPC: -0·56%) and AML (EAPC: -0·21%). Although temporal trends for ALL, CML CLL, HL and NHL prevalence were in the same direction across both sexes (decreasing for ALL/CML and increasing for CLL/HL/NHL), the decline in AML prevalence was driven by females (stable for males). Ultimately, the overall

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

incidence and prevalence of leukaemias/lymphomas across Oceania had remained stable over time.

In terms of disease burden, modest but statistically significant annual reductions in DALYs were observed for ALL, CML and HL (range in EAPCs: -0.55% for HL to -0.88% for CML) across both sexes, whereas the DALY burden attributed to CLL (EAPC: 0.92%) and NHL increased over time (EAPC: 0.26%) (all $P < 0.001$). AML DALY burden over time was stable, and largely driven by males (stable in males while decreasing in females). Lastly, with the exception of AML, CLL and NHL, the mortality burden attributed to leukaemias/lymphomas declined over time (range in EAPCs: -0.48% for CML to -0.35% for HL). Between 2010 and 2019, significant declines in mortality burden attributed to ALL and CML, and significantly increased CLL mortality, were estimated across both sexes. However, AML mortality burden increased significantly for males, and remained stable over time for females. Furthermore, the increase in CLL mortality over time was greater for females relative to males. Regarding lymphoma subtypes and sex, the overall decline in HL mortality was driven by males (stable for females), while NHL mortality increased over time for both sexes. (Table 2)

DISCUSSION

Our study is the first to report trends across a variety of outcomes between Australasia and Oceania over a 10-year period (2010 – 2019, inclusive) in which considerable changes in clinical practice occurred for cancer [1-3](#). Discrepancies in disease burden (ASRs) across regions were observed, which may be attributed to differences in age distribution [11](#) as well as differences in health care and cancer surveillance/registration systems between high-SDI and low-SDI countries [12](#). NHL was the key contributor to overall disease burden (DALYs/deaths) across both Australasia and Oceania. Despite the high relative incidence of AML, prevalence remained low across Australasia and likely reflects poor patient survival

(relative 5-year survival rate of 27%) [12](#). In contrast, the considerable DALY/mortality burden attributed to NHL likely reflects on high disease incidence, as 5-year survival for patients with NHL is considerably higher relative to AML [12 13](#). Notably, recent advances in the treatment of NHL and AML have contributed to improved patient survival, including the development of immune-based cellular and antibody therapies, and small molecule inhibitors [14 15](#). However, despite improved management of patients with leukaemias/lymphomas in Australasia, the considerable disease burden for both NHL and AML warrants further research to address unmet needs.

Upon stratification to explore the distribution of age (5-year age group) and sex (male/female) burden, age-specific leukaemia/lymphoma burden generally increased with increasing age and for males across both regions. Notably, disparate patient outcomes were identified between sexes for both regions with respect to AML; that is, disease burden (DALYs/deaths) was predominately borne by males despite incidence/prevalence rates for AML being similar among females in Australasia and Oceania. This is in line with other studies exploring patient outcomes in AML, and has been attributed to differences in biological risk factors, lifestyle, or environmental factors, and differences in disease treatment and management [16 17](#). Issues with health systems reporting, differential access to health services between females and males, differences in age distribution, and greater gender inequality have been implicated in contributing to discrepancies in disease burden between sexes across low-and high-SDI countries [1 5 18](#). However, further studies are recommended to establish and address sex-based drivers of leukaemia/lymphoma burden in countries with low-or-middle SDIs [1 5 7 18](#).

Based on trend analysis (EAPCs), leukaemia/lymphoma incidence/prevalence/mortality increased significantly over a 10-year period across Australasia (excepting stable mortality inCML and HL). Following stratification by sex, the trend of increasing incident/prevalent

burden was maintained across both sexes, with the exception of CML incidence which was stable for males. The trend of increasing mortality for AML was driven by male sex, whereas mortality trends across leukaemias/lymphomas were broadly comparable between sexes (excepting CML and HL). This is consistent with existing studies exploring the leukaemia/lymphoma burden for high-SDI countries between sexes [17](#). That is, the risks for attributable leukaemia/lymphoma burden were slightly different between sexes, with tobacco use and occupational exposure to carcinogens being more commonly observed in males and increased high body mass index (BMI) being among the top five risk exposure in females [19](#). Notably, significant reductions in the CML/HL/ALL DALY burden were observed and NHL DALY burden remained stable over time. This is reflective of improved outcomes due to changes in patient management and treatment across Australasia [20-23](#). However, the discrepancy between reductions/stable DALY burden (ALL and NHL) against mortality and incidence/prevalence increases over time may highlight disparate outcomes across age groups [20 21 24 25](#). ALL follows a bimodal age distribution, with the number of incident/prevalent cases peaking during early childhood and later adulthood (≥ 50 years), while NHL cases and the risk of mortality (ALL and NHL) increase with increasing age [16 26 27](#). Therefore, it is likely that improved disease management and treatment over time have led to a greater reduction in disease burden (DALYs) for younger patients relative to the change in mortality burden among older patients [20](#). This is supported by recent studies exploring ALL and NHL outcomes across Australasia, which found improvements in the management or treatment of disease coincided with greater gains in survival outcomes for younger relative to older patients [20 21 24 25 28](#).

In comparison with Australasia, the overall incident/prevalent burden across Oceania remained stable over time. Differing trends between regions highlights potential differences in exposure to environmental and lifestyle risk factors such as benzene exposure, tobacco

consumption, BMI and metabolic disease, and the disparity in health infrastructure and resources allocated for the detection for leukaemia/lymphomas between the two regions [29 30](#). Contrasting mortality trends for ALL/CML/HL were observed in the two regions (increasing/stable in Australasia versus decreasing in Oceania) may be attributed to the relatively older population residing in Australasia [11](#). Different trends were observed in AML burden (incidence/prevalence/DALYs/deaths) across males/females. Notably, a *stable* DALY/prevalent AML burden over time against a background of *increasing* mortality/incident burden time for male patients with AML across Oceania was observed. In contrast to AML trends across outcomes in Australia (all increased), this likely reflects a higher risk of mortality relative to patients with AML in Oceania, and suggests potential disparities in access to treatment, particularly novel treatments and clinical trials in AML, between the two regions [24](#). This aligns with studies exploring the association between DALY burden in cancers, and SDI [24](#). That is, DALYs vary with socioeconomic development, with improved patient survival increasing years-lived with disability (and subsequently, DALYs) for regions with a high-SDI [24](#). Although the overall burden (DALY/deaths) of the majority of leukaemias/lymphomas remained stable or slightly decreased over time, the DALY/death burden for CLL and NHL increased across Oceania for both sexes. In comparison with Australasia, the EAPC for CLL trends (incidence/deaths) was greater for females relative to males in Oceania. Furthermore, female NHL burden (DALYs/deaths) had remained stable over time in Australia, while increasing in Oceania. That is, the discrepancy in trend in CLL/NHL burden suggests potential sex disparities in the management or treatment of disease in these regions [1 5 30](#). Lastly, discrepancies in temporal trends were identified in sensitivity analyses using the world standard population. Notably, disease burden attributed to AML and NHL reduced across both regions in comparison with base-case estimates. This is likely attributed to the

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

variation in population structure in each region, which differs from the standard population ³
¹⁰. As sex and age are associated with cancer outcomes, it is possible that EAPCs estimated
using the standard population would inadequately capture the impact of region-specific
differences in age/sex, and ultimately, overestimate patient survival trends.

A key strength of our study lies in using data from the GBD 2019 study, which provides the
latest regional epidemiological distributions and trends of leukaemias and lymphomas ³.

Besides, the comprehensive analysis of trends across a variety of key outcomes facilitates an
understanding of the impacts of changes in patient management/treatment on patient
outcomes and potential areas of unmet need. Furthermore, disease burden and temporal
trends in our study were estimated based on region-specific population, which may better
reflect disease burden distribution and trend across age groups and sexes more
comprehensively and accurately, particularly when regional population structure is
incomparable to the world standard population. Lastly, restricting the analysis to two
geographically similar regions with differing SDI, and performing a sensitivity analysis using
the world standard population facilitates the comparison of trends between the two regions
(base-case), and to other regions or nations at the global level (sensitivity analysis).

However, several limitations to our analyses warrant mention. Limitations associated with
GBD 2019 data have been described previously ¹⁻⁴. In brief, it is likely that key outcomes for
Oceania were underestimated, as the capacity to collect reliable data on haematological
malignancies is often lacking in low-SDI regions. However, this does not considerably
change our findings of disparities in disease burden and temporal trends observed between
Australasia and Oceania, as well as between male/female sex. Second, GBD data includes
disease classifications by sex and age groups, but lacks ethnicity data, which limits the
analysis of genetic susceptibility to disease. Third, a direct comparison between Australasia
and Oceania could be difficult as there are considerable differences in the population

distribution and risk factors between the two regions ¹⁻⁴. As such, to explore the potential impacts attributed to age and sex distribution, our analyses used both region-specific populations (base-case) and standard population weights (sensitivity analyses) to explore changes in age-standardised outcomes. Fourth, it was not possible to discern the impact of various exposures on disease outcomes using aggregate data; as such, trends should be interpreted with caution due to potential confounding ¹⁻⁴. Lastly, it was not possible to capture the impacts of the COVID-19 pandemic on leukaemia/lymphoma burden using GBD 2019 data ^{2 3 31}.

Ultimately, our findings highlight disparities in the management and treatment of haematological malignancies based on SDI and sex. Despite the emergence of novel therapies and improved treatment/management over time, the consistent and considerable burden of AML and NHL in both regions warrants further research to mitigate the gap attributed to socioeconomic disadvantage. Further research is also recommended to explore factors that contribute to disparate outcomes for sex (male/female).

CONCLUSION

This study captures the disease burden of leukaemias/lymphomas and its temporal trends in two closely situated geographic regions with different SDI. The considerable disparity in disease burden observed between the two regions suggests the need for early diagnosis and better management strategies tailored to each region. Further study is required to explore the underlying factors behind the epidemiological trends of different leukaemia and lymphoma subtypes in each region and the male predominance in the majority of leukaemia and lymphoma burden of disease.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Ethics and dissemination

Ethics approval is not required for this study as the study involve information freely available in an open source.

Patient consent

Non patient consent required.

Data availability statement

All data relevant to the study are included in the article and uploaded as supplementary information.

References

1. Zhang N, Wu J, Wang Q, et al. Global burden of hematologic malignancies and evolution patterns over the past 30 years. *Blood Cancer J* 2023;13(1):82. doi: 10.1038/s41408-023-00853-3 [published Online First: 20230517]
2. Global Burden of Disease Cancer Collaboration. Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life Years for 29 Cancer Groups From 2010 to 2019: A Systematic Analysis for the Global Burden of Disease Study 2019. *JAMA Oncology* 2022;8(3):420-44. doi: 10.1001/jamaoncol.2021.6987
3. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020;396(10258):1204-22. doi: 10.1016/s0140-6736(20)30925-9

4. Tran KB, Lang JJ, Compton K, et al. The global burden of cancer attributable to risk factors, 2010–19: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2022;400(10352):563-91. doi: [https://doi.org/10.1016/S0140-6736\(22\)01438-6](https://doi.org/10.1016/S0140-6736(22)01438-6)

5. Yao Y, Lin X, Li F, et al. The global burden and attributable risk factors of chronic lymphocytic leukemia in 204 countries and territories from 1990 to 2019: analysis based on the global burden of disease study 2019. *BioMedical Engineering OnLine* 2022;21(1):4. doi: 10.1186/s12938-021-00973-6

6. Horwood PF, Tarantola A, Goarant C, et al. Health Challenges of the Pacific Region: Insights From History, Geography, Social Determinants, Genetics, and the Microbiome. *Front Immunol* 2019;10:2184. doi: 10.3389/fimmu.2019.02184 [published Online First: 20190913]

7. Lin X, Wang J, Huang X, et al. Global, regional, and national burdens of leukemia from 1990 to 2017: a systematic analysis of the global burden of disease 2017 study. *Aging (Albany NY)* 2021;13(7):10468.

8. Bispo JAB, Pinheiro PS, Kobetz EK. Epidemiology and Etiology of Leukemia and Lymphoma. *Cold Spring Harb Perspect Med* 2020;10(6) doi: 10.1101/cshperspect.a034819 [published Online First: 20200601]

9. Alaggio R, Amador C, Anagnostopoulos I, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Lymphoid Neoplasms.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

- 1
2
3 *Leukemia* 2022;36(7):1720-48. doi: 10.1038/s41375-022-01620-2 [published Online
4
5 First: 20220622]
6
7
8
9
- 10 10. Ahmad OB, Boschi Pinto C, Lopez A, et al. Age Standardization of Rates: A New WHO
11
12 Standard. *GPE Discussion Paper Series, EIP/GPE/EBD, World Health Organization*
13
14 2001;No.31
15
16
- 17 11. Skirbekk V, Dieleman JL, Stonawski M, et al. The health-adjusted dependency ratio as a
18
19 new global measure of the burden of ageing: a population-based study. *Lancet*
20
21 *Healthy Longev* 2022;3(5):e332-e38. doi: 10.1016/s2666-7568(22)00075-7 [published
22
23 Online First: 20220504]
24
25
26
- 27 12. Australian Institute of Health and Welfare. Cancer data in Australia. Canberra: AIHW,
28
29 2022.
30
31
32
33
- 34 13. Howlader N, Noone, A.M., Krapcho, M., Miller, D., Brest, A., Yu, M., Ruhl, J.,
35
36 Tatalovich, Z., Mariotto, A., Lewis, D.R., Chen, H.S., Feuer, E.J. and Cronin, K.A. .
37
38 SEER Cancer Statistics Review, 1975-2017. National Cancer Institute, Bethesda,
39
40 MD., 2020.
41
42
43
44
45
- 46 14. Winer ES, Stone RM. Novel therapy in Acute myeloid leukemia (AML): moving toward
47
48 targeted approaches. *Ther Adv Hematol* 2019;10:2040620719860645. doi:
49
50 10.1177/2040620719860645 [published Online First: 20190710]
51
52
53
54
- 55 15. Sawalha Y, Maddocks K. Novel treatments in B cell non-Hodgkin's lymphomas. *BMJ*
56
57 2022;377:e063439. doi: 10.1136/bmj-2020-063439
58
59
60

16. Dong Y, Shi O, Zeng Q, et al. Leukemia incidence trends at the global, regional, and national level between 1990 and 2017. *Exp Hematol Oncol* 2020;9:14. doi: 10.1186/s40164-020-00170-6 [published Online First: 20200619]

17. Rubin JB, Lagas JS, Broestl L, et al. Sex differences in cancer mechanisms. *Biol Sex Differ* 2020;11(1):17. doi: 10.1186/s13293-020-00291-x [published Online First: 20200415]

18. Lin Q, Mao L, Shao L, et al. Global, Regional, and National Burden of Chronic Myeloid Leukemia, 1990-2017: A Systematic Analysis for the Global Burden of Disease Study 2017. *Front Oncol* 2020;10:580759. doi: 10.3389/fonc.2020.580759 [published Online First: 20201215]

19. Murray CJL, Aravkin AY, Zheng P, et al. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2020;396(10258):1223-49. doi: 10.1016/S0140-6736(20)30752-2

20. Beckmann K, Kearney BA, Yeung D, et al. Changes in five-year survival for people with acute leukaemia in South Australia, 1980-2016. *Med J Aust* 2022;216(6):296-302. doi: 10.5694/mja2.51423 [published Online First: 20220216]

21. Wong SHC, Offner M, Hu R, et al. Treatment outcomes of adult acute lymphoblastic leukaemia in Auckland, New Zealand. *N Z Med J* 2023;136(1572):10-18. [published Online First: 20230324]

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

22. Cortes J, Pavlovsky C, Saußele S. Chronic myeloid leukaemia. *The Lancet* 2021;398(10314):1914-26. doi: 10.1016/S0140-6736(21)01204-6
23. Nguyen J, Wellard C, Chung E, et al. Clinical characteristics of Australian treatment-naïve patients with classical Hodgkin lymphoma from the lymphoma and related diseases registry. *Eur J Haematol* 2023;110(4):386-95. doi: 10.1111/ejh.13915 [published Online First: 20221228]
24. Anderson MA, Berkahn L, Cheah C, et al. Improving outcomes for patients with lymphoma: design and development of the Australian and New Zealand Lymphoma and Related Diseases Registry. *BMC Medical Research Methodology* 2022;22(1):266. doi: 10.1186/s12874-022-01728-0
25. Australian Institute of Health and Welfare. Cancer in Australia 2021. Canberra: AIHW, 2021.
26. Terwilliger T, Abdul-Hay M. Acute lymphoblastic leukemia: a comprehensive review and 2017 update. *Blood Cancer Journal* 2017;7(6):e577-e77. doi: 10.1038/bcj.2017.53
27. Sun H, Xue L, Guo Y, et al. Global, regional and national burden of non-Hodgkin lymphoma from 1990 to 2017: estimates from global burden of disease study in 2017. *Annals of Medicine* 2022;54(1):633-45. doi: 10.1080/07853890.2022.2039957

28. Wright F, Hapgood G, Loganathan A, et al. Relative survival of patients with lymphoma in Queensland according to histological subtype. *Med J Aust* 2018;209(4):166-72. doi: 10.5694/mja17.00937 [published Online First: 20180813]

29. Collaboration GBoDC. Global, Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life-years for 32 Cancer Groups, 1990 to 2015: A Systematic Analysis for the Global Burden of Disease Study. *JAMA Oncology* 2017;3(4):524-48. doi: 10.1001/jamaoncol.2016.5688

30. Cai W, Zeng Q, Zhang X, et al. Trends Analysis of Non-Hodgkin Lymphoma at the National, Regional, and Global Level, 1990-2019: Results From the Global Burden of Disease Study 2019. *Front Med (Lausanne)* 2021;8:738693. doi: 10.3389/fmed.2021.738693 [published Online First: 20210923]

31. Luo Q, O'Connell DL, Yu XQ, et al. Cancer incidence and mortality in Australia from 2020 to 2044 and an exploratory analysis of the potential effect of treatment delays during the COVID-19 pandemic: a statistical modelling study. *Lancet Public Health* 2022;7(6):e537-e48. doi: 10.1016/s2468-2667(22)00090-1

Authors' contributions

Ho TQA. — conceptualisation, data collection, formal analysis, investigation, methodology, visualisation, writing - original draft, and writing-review & editing.

Lee P. — conceptualisation, data collection, formal analysis, investigation, methodology, visualisation, writing-original draft, and writing-review & editing.

Gao L. — conceptualisation, investigation, methodology, supervision, writing-review & editing.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests

Non declared.

For peer review only

Tables

Table 1 – Results of ASRs and EAPCs for key outcomes across leukaemia and lymphoma subtypes in 2010 and 2019

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-C coefficient (95% CI)	P-value
	2010	2019			
AUSTRALASIA					
Incidence	32.59 (28.7, 37.18)	36.52 (28.56, 46.37)	-	-	-
ALL	2.38 (2.02, 3.01)	2.91 (2.24, 3.86)	2.37	0.02 (0.019, 0.027)	<0.001
AML	4.07 (3.34, 4.39)	4.72 (3.67, 5.9)	1.93	0.019 (0.017, 0.021)	<0.001
CLL	4.22 (3.77, 5.21)	5.22 (4.07, 6.87)	2.40	0.02 (0.022, 0.025)	<0.001
CML	1.58 (1.35, 1.95)	1.82 (1.37, 2.45)	2.16	0.02 (0.013, 0.029)	<0.001
HL	2.70 (2.30, 3.26)	2.79 (2.14, 3.66)	0.44	0.00 (0.003, 0.005)	<0.001
NHL	17.64 (15.92, 19.38)	19.06 (15.08, 23.62)	0.95	0.00 (0.005, 0.014)	<0.001
Prevalence	175.46 (150.04, 208.28)	196.64 (150.07, 256.58)	-	-	-
ALL	17.24 (14.33, 21.87)	21.44 (16.25, 28.53)	2.57	0.02 (0.021, 0.030)	<0.001
AML	4.46 (3.64, 5.11)	4.75 (3.74, 5.89)	0.90	0.009 (0.007, 0.011)	<0.001
CLL	27.34 (24.51, 33.8)	33.84 (25.92, 45.03)	2.40	0.024 (0.022, 0.025)	<0.001
CML	6.74 (5.45, 8.49)	8.26 (6.06, 11.42)	2.87	0.028 (0.022, 0.035)	<0.001

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-Coefficient (95% CI)	P-value
	2010	2019			
HL	21.45 (18.19, 25.9)	22.05 (16.83, 29.28)	0.40	0.003 (0.003, 0.005)	<0.001
NHL	98.23 (83.92, 113.11)	106.3 (81.27, 136.44)	1.07	0.015 (0.005, 0.016)	<0.001
DALYs	312.84 (281.93, 341.76)	324.03 (281.64, 366.04)	-	-	-
ALL	21.79 (19.98, 25.93)	21.08 (18.6, 24.78)	-0.36	-0.003 (-0.005, -0.003)	<0.001
AML	81.57 (68.56, 86.23)	89.01 (73.7, 97.38)	1.15	0.015 (0.009, 0.013)	<0.001
CLL	24.99 (21.98, 31)	29.39 (25.11, 36.76)	1.74	0.015 (0.015, 0.020)	<0.001
CML	12.02 (10.94, 14.49)	10.49 (9.15, 12.65)	-1.23	-0.001 (-0.0024, 0.000)	0.042
HL	12.77 (10.9, 15.22)	12.38 (10.44, 15.16)	-0.40	-0.001 (-0.005, -0.003)	<0.001
NHL	159.7 (149.57, 168.89)	161.68 (144.64, 179.31)	0.18	0.000 (-0.004, 0.007)	0.536
Deaths	13.87 (12.2, 15.21)	15.27 (12.97, 17.36)	-	-	-
ALL	0.47 (0.42, 0.57)	0.50 (0.44, 0.60)	0.75	0.006 (0.006, 0.009)	<0.001
AML	3.51 (2.87, 3.77)	4.15 (3.32, 4.65)	2.07	0.021 (0.019, 0.022)	<0.001
CLL	1.4 (1.21, 1.71)	1.71 (1.43, 2.13)	2.11	0.021 (0.019, 0.023)	<0.001
CML	0.55 (0.49, 0.68)	0.52 (0.44, 0.65)	-0.27	-0.003 (-0.014, 0.008)	0.631
HL	0.37 (0.32, 0.44)	0.38 (0.32, 0.45)	0.05	0.000 (-0.001, 0.002)	0.399

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-Coefficient (95% CI)	P-value
	2010	2019			
NHL	7.57 (6.9, 8.04)	8.02 (7.03, 8.87)	1.07	0.01 (-0.005, 0.016)	<0.001
OCEANIA					
Incidence	3.55 (2.65, 4.9)	3.62 (2.68, 4.96)	-	-	-
ALL	0.55 (0.33, 0.93)	0.53 (0.33, 0.84)	-0.45	-0.006 (-0.006, -0.003)	<0.001
AML	1.36 (1.01, 1.88)	1.36 (0.98, 1.89)	-0.02	0.000 (-0.002, 0.001)	0.816
CLL	0.03 (0.02, 0.04)	0.04 (0.03, 0.05)	1.38	0.01 (-0.012, 0.016)	<0.001
CML	0.32 (0.2, 0.51)	0.3 (0.19, 0.46)	-0.58	-0.005 (-0.008, -0.004)	<0.001
HL	0.3 (0.24, 0.39)	0.3 (0.23, 0.41)	-0.05	-0.001 (-0.002, 0.001)	0.405
NHL	0.99 (0.85, 1.15)	1.08 (0.91, 1.31)	0.92	0.000 (0.008, 0.010)	<0.001
Prevalence	5.97 (3.95, 9.17)	6.09 (4.05, 9.14)	-	-	-
ALL	1.46 (0.85, 2.52)	1.39 (0.86, 2.26)	-0.56	-0.005 (-0.007, -0.004)	<0.001
AML	2.68 (1.81, 4.08)	2.63 (1.75, 4)	-0.21	-0.001 (-0.004, 0.000)	0.026
CLL	0.12 (0.09, 0.16)	0.14 (0.1, 0.19)	1.99	0.020 (0.017, 0.022)	<0.001
CML	0.51 (0.3, 0.85)	0.47 (0.28, 0.74)	-0.76	-0.008 (-0.010, -0.005)	<0.001
HL	0.75 (0.59, 0.94)	0.81 (0.6, 1.07)	0.68	0.007 (0.003, 0.011)	0.001

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-Coefficient (95% CI)	P-value
	2010	2019			
NHL	0.45 (0.31, 0.62)	0.64 (0.45, 0.88)	3.24	0.03 (-0.21, 0.043)	<0.001
DALYs	140.07 (100.83, 201.45)	137.73 (98.65, 196.41)	-	-	-
ALL	29.25 (16.74, 51.13)	27.86 (16.96, 45.85)	-0.56	-0.007 (-0.007, -0.004)	<0.001
AML	49.78 (36.68, 70.38)	49.16 (35.01, 71.49)	-0.12	-0.003 (-0.003, 0.000)	0.115
CLL	0.83 (0.6, 1.12)	0.91 (0.65, 1.28)	0.92	0.008 (0.008, 0.010)	<0.001
CML	12.79 (7.25, 21.8)	11.82 (6.88, 18.89)	-0.88	-0.001 (-0.011, -0.007)	<0.001
HL	10.11 (7.85, 13.12)	9.67 (7.29, 13.07)	-0.55	-0.006 (-0.007, -0.005)	<0.001
NHL	37.31 (31.71, 43.9)	38.3 (31.86, 45.84)	0.26	0.001 (0.001, 0.004)	<0.001
Deaths	2.77 (2.11, 3.72)	2.81 (2.1, 3.8)	-	-	-
ALL	0.42 (0.26, 0.7)	0.41 (0.26, 0.64)	-0.39	-0.004 (-0.005, -0.002)	<0.001
AML	0.9 (0.68, 1.21)	0.91 (0.66, 1.27)	0.08	0.000 (-0.001, 0.002)	0.244
CLL	0.02 (0.02, 0.03)	0.03 (0.02, 0.04)	0.99	0.010 (0.009, 0.011)	<0.001
CML	0.29 (0.18, 0.44)	0.28 (0.18, 0.41)	-0.48	-0.005 (-0.007, -0.003)	<0.001
HL	0.22 (0.17, 0.29)	0.21 (0.16, 0.28)	-0.35	-0.003 (-0.004, -0.003)	<0.001
NHL	0.92 (0.8, 1.05)	0.98 (0.82, 1.17)	0.65	0.007 (0.005, 0.008)	<0.001

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia;
CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin
lymphoma; NHL = non-Hodgkin lymphoma

For peer review only

Table 2 - Trends for key outcomes across leukaemia and lymphoma subtypes in 2010 and 2019 by sex

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
AUSTRALASIA						
Incidence						
ALL	2.45	0.024 (0.020, 0.028)	<0.001	2.29	0.023 (0.018, 0.027)	<0.001
AML	2.38	0.024 (0.021, 0.026)	<0.001	1.42	0.011 (0.011, 0.017)	<0.001
CLL	2.48	0.025 (0.02, 0.029)	<0.001	2.28	0.023 (0.016, 0.029)	<0.001
CML	0.64	0.006 (-0.005, 0.018)	0.292	2.84	0.023 (0.022, 0.034)	<0.001
HL	0.30	0.003 (0.002, 0.004)	<0.001	0.61	0.001 (0.005, 0.007)	<0.001
NHL	1.14	0.011 (0.008, 0.015)	<0.001	0.70	0.001 (0.001, 0.013)	0.018
Prevalence						
ALL	2.67	0.026 (0.022, 0.031)	<0.001	2.46	0.023 (0.019, 0.029)	<0.001
AML	1.48	0.015 (0.013, 0.017)	<0.001	0.39	0.004 (0.001, 0.007)	0.006
CLL	2.40	0.024 (0.019, 0.029)	<0.001	2.42	0.024 (0.017, 0.031)	<0.001
CML	1.94	0.019 (0.009, 0.029)	<0.001	3.07	0.03 (0.024, 0.036)	<0.001

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
HL	0.25	0.002 (0.001, 0.004)	<0.001	0.58	0.000 (0.004, 0.007)	<0.001
NHL	1.23	0.012 (0.008, 0.017)	<0.001	0.87	0.009 (0.002, 0.016)	0.016
DALYs						
ALL	-0.56	-0.006 (-0.008, -0.003)	<0.001	-0.08	-0.000 (-0.004, 0.002)	0.589
AML	1.51	0.015 (0.013, 0.017)	<0.001	0.69	0.000 (0.005, 0.009)	<0.001
CLL	1.85	0.018 (0.013, 0.023)	<0.001	1.57	0.019 (0.009, 0.022)	<0.001
CML	-1.86	-0.019 (-0.033, -0.004)	0.011	-0.24	-0.000 (-0.01, 0.006)	0.552
HL	-0.53	-0.005 (-0.006, -0.004)	<0.001	-0.23	-0.002 (-0.004, -0.001)	<0.001
NHL	0.34	0.003 (-0.001, 0.008)	0.165	-0.04	0 (-0.000, 0.007)	0.915
Deaths						
ALL	0.64	0.006 (0.005, 0.008)	<0.001	0.90	0.000 (0.006, 0.012)	<0.001
AML	2.55	0.025 (0.023, 0.027)	<0.001	1.46	0.015 (0.012, 0.017)	<0.001
CLL	2.40	0.024 (0.020, 0.027)	<0.001	1.66	0.016 (0.001, 0.023)	<0.001
CML	-0.72	-0.007 (-0.02, 0.005)	0.263	0.50	0.005 (-0.003, 0.013)	0.221

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
HL	-0.08	-0.001 (-0.002, 0.001)	0.358	0.21	0.000 (-0.001, 0.004)	0.007
NHL	0.86	0.009 (0.005, 0.012)	<0.001	0.28	0.000 (-0.003, 0.009)	0.345
OCEANIA						
Incidence						
ALL	-0.37	-0.004 (-0.005, -0.003)	<0.001	-0.54	-0.000 (-0.008, -0.003)	<0.001
AML	0.17	0.002 (0.001, 0.003)	<0.001	-0.27	-0.000 (-0.006, 0)	0.072
CLL	0.96	0.01 (0.008, 0.011)	<0.001	1.75	0.01 (-0.015, 0.019)	<0.001
CML	-0.61	-0.006 (-0.009, -0.003)	<0.001	-0.58	-0.006 (-0.008, -0.004)	<0.001
HL	-0.19	-0.002 (-0.003, -0.001)	0.005	0.31	0.000 (0.002, 0.004)	<0.001
NHL	0.86	0.009 (0.007, 0.010)	<0.001	1.00	0.01 (-0.009, 0.011)	<0.001
Prevalence						
ALL	-0.46	-0.005 (-0.006, -0.003)	<0.001	-0.66	-0.007 (-0.009, -0.004)	<0.001
AML	-0.04	0 (-0.001, 0.001)	0.509	-0.39	-0.004 (-0.007, -0.001)	0.01
CLL	1.54	0.015 (0.013, 0.018)	<0.001	2.32	0.023 (0.02, 0.026)	<0.001

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
CML	-0.85	-0.009 (-0.011, -0.006)	<0.001	-0.71	-0.009 (-0.009, -0.005)	<0.001
HL	0.47	0.005 (0.001, 0.008)	0.012	1.12	0.010 (0.007, 0.016)	<0.001
NHL	2.60	0.026 (0.021, 0.031)	<0.001	3.73	0.030 (0.021, 0.053)	<0.001
DALYs						
ALL	-0.49	-0.005 (-0.006, -0.004)	<0.001	-0.64	-0.006 (-0.009, -0.004)	<0.001
AML	0.06	0.001 (0, 0.001)	0.136	-0.37	-0.001 (-0.007, -0.001)	0.014
CLL	0.64	0.006 (0.005, 0.008)	<0.001	1.19	0.011 (0.001, 0.013)	<0.001
CML	-0.95	-0.01 (-0.012, -0.007)	<0.001	-0.84	-0.008 (-0.01, -0.007)	<0.001
HL	-0.67	-0.007 (-0.008, -0.006)	<0.001	-0.20	-0.003 (-0.003, -0.001)	<0.001
NHL	0.24	0.002 (0.001, 0.004)	0.001	0.29	0.001 (0.002, 0.004)	<0.001
Deaths						
ALL	-0.32	-0.003 (-0.004, -0.002)	<0.001	-0.46	-0.005 (-0.007, -0.003)	<0.001
AML	0.27	0.003 (0.002, 0.004)	<0.001	-0.19	-0.002 (-0.005, 0.001)	0.175
CLL	0.62	0.006 (0.005, 0.007)	<0.001	1.33	0.013 (0.012, 0.015)	<0.001

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
CML	-0.48	-0.005 (-0.008, -0.002)	0.001	-0.50	-0.006 (-0.007, -0.003)	<0.001
HL	-0.45	-0.004 (-0.006, -0.003)	<0.001	-0.06	-0.001 (-0.001, 0)	0.08
NHL	0.62	0.006 (0.005, 0.008)	<0.001	0.70	0.006 (0.006, 0.008)	<0.001

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures

Figure 1 - Age-specific rates at regional level by sex, in Australasia, and Oceania in 2019

[The figure is attached in a separate file]

For peer review only

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES) .

Appendix A

Table A1 - Age-specific rates in Australasia and Oceania by age groups and by sex (2010)

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
AUSTRALASIA																		
Incidence																		
<5 years	3·68	3·49	3·88	0·64	0·65	0·62	0·00	0·00	0·00	0·10	0·09	0·11	0·22	0·21	0·23	0·00	0·00	0·00
5-9 years	3·19	3·64	2·71	0·36	0·30	0·42	0·00	0·00	0·00	0·05	0·03	0·07	0·36	0·41	0·31	1·26	1·85	0·64
10-14 years	1·50	1·64	1·34	0·43	0·38	0·49	0·00	0·00	0·00	0·06	0·05	0·00	0·72	0·85	0·59	1·23	1·61	0·83
15-19 years	1·93	2·49	1·33	0·94	0·94	0·94	0·00	0·00	0·00	0·12	0·10	0·15	2·66	2·49	2·83	2·47	3·06	1·85
20-24 years	1·25	1·58	0·90	0·89	0·83	0·95	0·11	0·09	0·13	0·25	0·20	0·31	4·60	4·13	5·09	2·02	2·41	1·62
25-29 years	0·92	0·98	0·86	0·93	0·92	0·93	0·10	0·09	0·11	0·38	0·34	0·42	4·79	4·73	4·85	2·09	2·55	1·63
30-34 years	0·63	0·74	0·52	0·80	0·79	0·82	0·10	0·08	0·11	0·35	0·34	0·35	3·71	4·02	3·42	2·66	3·49	1·86
35-39 years	0·58	0·64	0·54	1·01	0·94	1·07	0·17	0·25	0·10	0·53	0·45	0·61	3·01	3·60	2·45	3·78	4·80	2·80
40-44 years	0·63	0·65	0·62	1·26	1·15	1·36	0·38	0·56	0·21	0·58	0·48	0·67	2·68	3·37	2·03	3·57	4·55	2·64
45-49 years	0·90	0·83	0·98	1·77	1·68	1·86	1·04	1·55	0·55	0·67	0·55	0·78	2·19	2·59	1·80	6·07	8·01	4·20
50-54 years	1·55	1·43	1·68	2·69	2·66	2·73	2·65	4·08	1·26	0·97	0·70	1·23	2·14	2·65	1·66	10·58	13·36	7·90
55-59 years	2·46	2·28	2·63	3·96	3·98	3·95	5·46	7·88	3·11	1·41	0·90	1·90	2·34	2·90	1·79	18·17	23·93	12·60
60-64 years	3·89	3·57	4·21	6·47	7·10	5·86	9·24	13·38	5·16	1·91	1·44	2·38	2·64	2·89	2·39	27·29	34·81	19·86
65-69 years	5·54	5·70	5·37	10·74	12·57	8·95	13·75	18·98	8·64	2·78	2·18	3·36	3·03	3·49	2·58	46·37	59·69	33·34

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	7·57	8·71	6·50	17·29	20·26	14·51	22·80	32·32	13·91	5·80	4·16	7·32	3·31	3·76	2·88	74·78	95·43	55·52
75-79 years	7·25	9·05	5·71	21·47	26·59	17·06	27·79	39·45	17·74	7·75	5·91	9·33	4·04	4·31	3·81	113·53	136·31	93·89
80-84 years	6·55	8·05	5·43	26·59	31·89	22·61	22·93	32·71	15·58	12·47	9·85	14·44	4·91	5·63	4·36	125·51	153·47	104·50
85+ years	6·51	7·83	5·82	34·60	41·17	31·19	35·57	49·93	28·11	20·89	17·36	22·72	7·05	8·57	6·25	168·07	211·16	145·67
Prevalence																		
<5 years	30·07	28·42	31·82	1·66	1·37	1·97	0·00	0·00	0·00	0·52	0·40	0·64	1·95	1·85	2·05	0·00	0·00	0·00
5-9 years	24·86	28·16	21·38	0·85	0·58	1·15	0·00	0·00	0·00	0·28	0·14	0·42	3·21	3·64	2·75	5·13	9·99	0·00
10-14 years	10·64	11·41	9·82	0·91	0·64	1·19	0·00	0·00	0·00	0·31	0·20	0·41	6·39	7·48	5·23	4·62	9·00	0·00
15-19 years	12·97	16·52	9·24	2·93	2·48	3·40	0·00	0·00	0·00	0·65	0·43	0·87	3·51	22·07	25·02	17·21	21·49	12·71
20-24 years	8·64	10·57	6·66	2·26	1·75	2·78	0·96	0·80	1·12	1·30	0·83	1·79	10·96	36·68	45·35	12·50	15·87	9·04
25-29 years	6·41	6·59	6·22	1·95	1·55	2·34	0·86	0·75	0·98	1·84	1·30	2·38	12·25	41·62	42·87	13·95	17·06	10·88
30-34 years	4·11	4·67	3·56	1·37	1·04	1·70	0·81	0·70	0·92	1·59	1·26	1·91	12·47	35·09	29·92	18·27	23·69	13·03
35-39 years	3·93	4·06	3·81	1·56	1·17	1·93	1·43	2·09	0·81	2·48	1·59	3·33	16·15	31·21	21·33	26·28	33·18	19·71
40-44 years	4·43	4·27	4·59	1·74	1·27	2·19	3·08	4·46	1·76	2·62	1·60	3·58	13·13	29·05	17·48	21·29	26·71	16·13
45-49 years	6·48	5·61	7·31	2·68	2·08	3·26	8·48	12·55	4·54	3·11	1·95	4·21	8·50	21·84	15·28	35·73	47·17	24·68
50-54 years	11·88	10·19	13·51	4·02	3·16	4·85	21·44	32·72	10·52	4·55	2·45	6·58	7·99	22·05	14·06	66·28	86·34	46·89
55-59 years	19·16	17·02	21·23	4·89	3·87	5·88	44·09	63·13	25·66	6·42	2·50	10·23	8·72	23·11	14·47	112·26	153·67	72·19
60-64 years	29·74	27·06	32·39	6·79	6·51	7·07	72·83	104·29	41·72	8·59	3·85	13·27	10·48	22·25	18·73	176·67	228·63	125·30
65-69 years	42·41	43·57	41·28	10·17	10·88	9·48	104·63	141·82	68·29	12·02	5·29	18·60	11·05	24·11	18·06	303·28	387·86	220·62

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	58·96	65·83	52·55	14·38	15·72	13·12	164·89	229·69	104·43	25·72	9·60	40·76	18·43	20·60	16·40	480·04	609·76	359·01
75-79 years	50·48	60·87	41·53	16·85	19·34	14·70	186·37	256·10	126·25	33·69	14·00	50·67	20·50	20·12	20·83	704·29	825·50	599·80
80-84 years	35·88	44·74	29·22	25·62	30·12	22·24	103·78	139·63	76·85	52·18	22·05	74·81	24·65	26·60	23·19	546·69	653·16	466·71
85+ years	28·29	34·45	25·08	27·22	29·73	25·92	107·51	137·41	91·97	76·75	35·62	98·13	29·74	31·23	28·97	584·51	728·80	509·52
DALYs																		
<5 years	33·57	34·38	32·72	32·33	36·71	27·71	0·00	0·00	0·00	2·28	3·10	1·41	0·79	0·79	0·79	9·81	11·49	8·04
5-9 years	41·77	50·96	32·08	17·54	16·11	19·05	0·00	0·00	0·00	0·97	1·08	0·86	1·31	1·53	1·07	13·04	19·17	6·59
10-14 years	29·70	34·46	24·68	20·93	20·13	21·78	0·00	0·00	0·00	1·35	1·70	0·91	2·75	3·45	2·01	12·63	16·52	8·52
15-19 years	44·58	60·09	28·25	35·59	38·33	32·69	0·00	0·00	0·00	1·94	2·42	1·45	7·98	7·62	8·36	25·11	31·09	18·82
20-24 years	26·17	36·41	15·66	34·85	35·54	34·14	0·80	0·78	0·83	4·06	4·95	3·14	3·92	13·00	14·86	32·22	38·55	25·74
25-29 years	18·70	21·78	15·67	36·99	40·25	33·77	0·71	0·73	0·70	6·53	8·72	4·36	8·66	19·00	18·33	31·77	38·63	24·98
30-34 years	12·69	16·50	9·00	32·40	35·05	29·84	0·69	0·71	0·69	6·29	8·97	3·70	5·58	17·69	13·53	38·10	49·78	26·81
35-39 years	10·59	12·98	8·32	38·59	39·47	37·75	1·32	2·12	0·56	8·50	11·08	6·05	3·96	17·13	10·95	50·22	63·84	37·25
40-44 years	9·82	11·54	8·19	45·96	46·23	45·70	2·81	4·46	1·24	8·92	11·41	6·54	3·31	16·79	9·99	65·38	83·14	48·45
45-49 years	10·29	11·39	9·23	56·19	58·20	54·25	6·83	10·99	2·81	8·87	11·34	6·42	1·62	14·15	9·18	100·43	132·28	69·66
50-54 years	13·20	14·47	11·99	76·51	82·43	70·79	15·87	26·15	5·92	10·92	12·82	9·09	0·92	13·72	8·21	156·47	196·83	117·43
55-59 years	14·77	16·26	13·34	103·10	112·22	94·27	28·96	45·36	13·09	15·02	18·12	12·02	4·32	17·88	10·87	236·72	310·98	164·85
60-64 years	16·92	18·47	15·39	153·16	177·90	128·70	50·95	78·81	23·42	18·94	25·61	12·34	6·51	18·83	14·22	323·60	412·20	236·00
65-69 years	19·95	22·65	17·33	223·90	273·52	175·40	77·41	114·56	41·10	25·37	36·43	14·56	1·87	25·61	18·22	468·61	603·00	337·26

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	24·34	29·50	19·53	311·90	378·56	249·71	128·63	194·23	67·42	41·61	58·00	26·33	36·91	30·38	23·68	632·89	808·40	469·13
75-79 years	24·21	30·78	18·55	323·26	414·37	244·72	150·67	225·50	86·17	46·15	65·79	29·22	38·46	32·00	25·41	781·55	940·58	644·47
80-84 years	23·76	29·38	19·53	305·46	378·02	250·97	172·03	253·91	110·53	56·89	80·97	38·81	46·56	30·73	23·43	866·29	1063·97	717·82
85+ years	21·99	26·97	19·40	247·01	336·78	200·36	225·66	332·13	170·32	69·92	106·41	50·95	62·14	28·86	18·64	835·96	1072·56	712·99
Deaths																		
<5 years	0·36	0·37	0·35	0·37	0·42	0·32	0·00	0·00	0·00	0·03	0·04	0·02	0·01	0·01	0·01	0·11	0·13	0·09
5-9 years	0·49	0·60	0·37	0·21	0·20	0·23	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·02	0·01	0·15	0·22	0·08
10-14 years	0·38	0·44	0·31	0·27	0·26	0·28	0·00	0·00	0·00	0·02	0·02	0·01	0·03	0·04	0·02	0·16	0·20	0·11
15-19 years	0·61	0·82	0·38	0·49	0·53	0·45	0·00	0·00	0·00	0·03	0·03	0·02	0·09	0·09	0·09	0·33	0·41	0·25
20-24 years	0·38	0·53	0·23	0·52	0·53	0·51	0·01	0·01	0·01	0·06	0·07	0·05	0·17	0·16	0·18	0·47	0·56	0·37
25-29 years	0·29	0·34	0·24	0·60	0·65	0·54	0·01	0·01	0·01	0·10	0·14	0·07	0·26	0·26	0·26	0·49	0·60	0·39
30-34 years	0·22	0·28	0·15	0·57	0·62	0·52	0·01	0·01	0·01	0·11	0·16	0·06	0·24	0·27	0·21	0·64	0·84	0·45
35-39 years	0·20	0·24	0·15	0·74	0·76	0·73	0·02	0·04	0·01	0·16	0·21	0·11	0·24	0·29	0·19	0·92	1·17	0·69
40-44 years	0·20	0·24	0·17	0·98	0·98	0·97	0·05	0·09	0·02	0·19	0·24	0·13	0·25	0·32	0·19	1·35	1·71	1·00
45-49 years	0·23	0·26	0·20	1·33	1·38	1·28	0·14	0·23	0·06	0·21	0·27	0·11	0·25	0·30	0·20	2·30	3·02	1·60
50-54 years	0·33	0·36	0·29	2·04	2·20	1·88	0·37	0·61	0·13	0·28	0·34	0·23	0·26	0·32	0·19	4·01	5·05	3·02
55-59 years	0·40	0·45	0·35	3·15	3·43	2·87	0·75	1·19	0·33	0·45	0·55	0·34	0·40	0·49	0·30	6·88	9·03	4·80
60-64 years	0·51	0·57	0·44	5·45	6·33	4·57	1·56	2·43	0·70	0·65	0·90	0·40	0·53	0·61	0·46	10·91	13·88	7·98
65-69 years	0·69	0·80	0·58	9·48	11·59	7·42	2·81	4·18	1·46	1·04	1·53	0·56	0·85	0·99	0·70	18·56	23·81	13·43

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	1·00	1·23	0·78	16·14	19·60	12·92	5·69	8·63	2·94	2·05	2·97	1·20	1·31	1·47	1·16	29·99	38·15	22·38
75-79 years	1·26	1·63	0·95	21·14	27·10	16·00	8·36	12·56	4·73	2·84	4·24	1·64	1·72	1·95	1·52	45·52	54·53	37·75
80-84 years	1·66	2·08	1·35	25·91	32·06	21·29	13·45	19·88	8·62	4·45	6·71	2·74	1·96	2·26	1·73	67·50	82·33	56·36
85+ years	2·24	2·76	1·96	29·33	39·43	24·08	25·75	37·40	19·69	7·63	12·29	5·21	1·98	2·63	1·64	90·34	113·20	78·46
OCEANIA																		
Incidence																		
<5 years	1·18	1·31	1·03	1·42	1·62	1·20	0·00	0·00	0·00	0·19	0·26	0·10	0·02	0·02	0·02	0·00	0·00	0·00
5-9 years	0·74	0·63	0·86	0·74	0·64	0·85	0·00	0·00	0·00	0·09	0·13	0·04	0·14	0·25	0·03	0·34	0·49	0·17
10-14 years	0·65	0·43	0·90	0·79	0·33	1·30	0·00	0·00	0·00	0·14	0·04	0·25	0·09	0·14	0·02	0·55	0·32	0·81
15-19 years	0·52	0·66	0·36	1·65	2·04	1·22	0·00	0·00	0·00	0·03	0·04	0·03	0·20	0·30	0·08	0·65	0·72	0·57
20-24 years	0·34	0·38	0·30	0·98	1·37	0·56	0·02	0·02	0·03	0·11	0·17	0·04	0·35	0·35	0·35	0·49	0·62	0·35
25-29 years	0·37	0·43	0·30	1·39	1·35	1·44	0·02	0·02	0·03	0·36	0·22	0·51	0·35	0·46	0·23	0·70	0·90	0·49
30-34 years	0·11	0·17	0·05	0·84	1·19	0·50	0·02	0·02	0·02	0·24	0·17	0·31	0·46	0·60	0·32	0·64	0·48	0·80
35-39 years	0·26	0·18	0·34	1·33	1·55	1·12	0·03	0·02	0·03	0·43	0·37	0·49	0·35	0·61	0·09	0·93	1·20	0·67
40-44 years	0·14	0·18	0·09	1·17	1·66	0·65	0·03	0·02	0·04	0·12	0·15	0·07	0·36	0·59	0·11	1·23	1·29	1·17
45-49 years	0·34	0·32	0·36	1·93	1·50	2·41	0·03	0·02	0·04	0·68	0·16	1·23	0·53	0·74	0·31	1·49	1·78	1·19
50-54 years	0·49	0·47	0·52	1·79	1·40	2·22	0·08	0·06	0·10	0·88	0·44	1·37	0·72	1·21	0·17	2·04	2·65	1·36
55-59 years	0·69	0·76	0·61	2·83	3·06	2·57	0·24	0·26	0·22	1·04	1·13	0·93	0·61	0·74	0·45	3·19	3·55	2·78
60-64 years	0·38	0·55	0·19	2·84	3·61	1·96	0·24	0·28	0·20	1·63	0·55	2·87	0·93	0·93	0·94	4·10	3·96	4·27

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
65-69 years	0.51	0.71	0.28	2.84	3.56	2.01	0.32	0.21	0.44	1.57	1.10	2.10	0.98	1.45	0.43	6.05	7.42	4.49
70-74 years	0.37	0.54	0.19	2.82	3.20	2.42	0.22	0.18	0.27	2.60	1.11	4.19	1.44	1.54	1.32	8.43	9.13	7.67
75-79 years	0.70	0.31	1.08	6.31	6.90	5.73	0.30	0.34	0.27	0.78	1.35	0.23	1.30	1.86	0.75	9.80	7.49	12.05
80-84 years	0.20	0.24	0.17	4.52	6.56	2.84	0.34	0.54	0.17	1.79	0.52	2.84	0.88	1.03	0.76	11.67	16.54	7.63
85+ years	0.62	0.22	0.87	8.55	16.30	3.56	0.23	0.20	0.25	2.19	0.43	3.33	1.04	1.56	0.70	8.75	10.90	7.36
Prevalence																		
<5 years	3.46	3.86	3.04	3.68	3.74	3.62	0.00	0.00	0.00	0.44	0.62	0.24	0.09	0.09	0.09	0.00	0.00	0.00
5-9 years	2.08	1.76	2.42	1.89	1.42	2.40	0.00	0.00	0.00	0.20	0.29	0.10	0.52	0.88	0.12	0.00	0.00	0.00
10-14 years	1.50	0.99	2.08	1.72	0.58	3.01	0.00	0.00	0.00	0.27	0.09	0.48	0.27	0.42	0.10	1.00	0.00	2.12
15-19 years	1.85	2.36	1.29	5.18	5.90	4.38	0.00	0.00	0.00	0.09	0.10	0.08	0.91	1.33	0.46	1.22	1.13	1.31
20-24 years	1.04	1.13	0.95	2.40	3.15	1.62	0.14	0.13	0.15	0.24	0.38	0.08	1.49	1.31	1.68	0.11	0.21	0.00
25-29 years	0.97	1.14	0.80	3.00	2.50	3.51	0.11	0.09	0.14	0.71	0.42	1.00	1.09	1.32	0.86	0.28	0.49	0.06
30-34 years	0.22	0.35	0.09	1.30	1.65	0.96	0.07	0.07	0.08	0.40	0.29	0.51	1.16	1.37	0.95	0.27	0.00	0.54
35-39 years	0.49	0.34	0.64	1.89	1.91	1.88	0.12	0.09	0.15	0.66	0.57	0.75	0.72	1.23	0.21	0.62	0.81	0.43
40-44 years	0.23	0.30	0.15	1.35	1.75	0.94	0.12	0.07	0.17	0.16	0.21	0.10	0.66	1.09	0.22	0.23	0.33	0.12
45-49 years	0.62	0.56	0.69	2.72	1.72	3.81	0.12	0.07	0.17	1.00	0.24	1.82	1.00	1.42	0.56	0.25	0.49	0.00
50-54 years	0.91	0.81	1.02	2.53	1.60	3.56	0.33	0.22	0.45	1.27	0.64	1.99	1.28	2.17	0.29	0.35	0.67	0.00
55-59 years	1.19	1.29	1.09	3.15	3.00	3.32	0.96	0.99	0.94	1.43	1.55	1.31	0.85	1.08	0.60	0.49	0.75	0.20
60-64 years	0.58	0.83	0.28	2.67	3.18	2.08	0.83	0.92	0.73	2.12	0.69	3.76	1.09	1.13	1.04	0.41	0.22	0.63

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
65-69 years	0.64	0.88	0.36	2.50	3.01	1.92	0.91	0.54	1.34	1.83	1.27	2.48	0.92	1.39	0.37	0.87	1.46	0.20
70-74 years	0.45	0.66	0.22	2.28	2.54	2.01	0.52	0.39	0.65	2.70	1.13	4.38	1.11	1.26	0.95	2.48	2.65	2.30
75-79 years	0.72	0.27	1.17	4.82	5.11	4.53	0.53	0.56	0.50	0.71	1.23	0.20	0.90	1.34	0.47	3.98	1.72	6.19
80-84 years	0.15	0.17	0.14	4.44	6.53	2.70	0.37	0.58	0.19	1.45	0.40	2.31	0.57	0.73	0.44	2.72	4.66	1.11
85+ years	0.34	0.16	0.46	6.31	11.67	2.86	0.19	0.16	0.22	1.64	0.29	2.50	0.75	1.26	0.42	1.37	1.89	1.04
DALYs																		
<5 years	72.64	81.01	63.64	69.80	83.78	54.75	0.00	0.00	0.00	11.58	16.52	6.26	0.98	1.21	0.74	33.59	38.44	28.37
5-9 years	44.00	37.52	51.13	34.63	31.94	37.60	0.00	0.00	0.00	5.41	7.94	2.64	6.71	11.90	1.00	10.73	16.18	4.73
10-14 years	39.90	26.46	55.03	36.75	16.80	59.23	0.00	0.00	0.00	8.54	2.73	15.07	3.98	6.71	0.90	16.90	9.26	25.51
15-19 years	23.94	30.52	16.70	61.11	78.40	42.12	0.00	0.00	0.00	1.55	1.73	1.35	6.83	10.71	2.56	20.14	23.20	16.76
20-24 years	16.30	17.90	14.63	37.62	54.42	20.22	0.94	0.95	0.93	5.02	8.21	1.71	2.03	12.80	11.23	23.69	29.93	17.22
25-29 years	17.45	20.59	14.29	52.57	53.46	51.68	0.82	0.70	0.94	17.33	10.29	24.42	2.89	17.76	8.00	32.83	42.47	23.15
30-34 years	5.22	8.24	2.24	33.17	48.02	18.51	0.59	0.60	0.58	11.82	8.40	15.20	7.50	23.36	11.72	27.14	20.00	34.20
35-39 years	12.24	8.57	15.93	49.78	60.27	39.24	1.00	0.84	1.16	20.15	17.30	23.01	3.28	23.36	3.16	37.11	47.73	26.44
40-44 years	6.06	8.13	3.88	42.70	62.31	22.07	1.04	0.70	1.40	5.23	6.99	3.33	2.88	21.33	3.99	52.71	54.89	50.42
45-49 years	13.10	12.41	13.84	59.13	48.82	70.29	0.86	0.59	1.16	26.46	6.33	48.26	7.06	23.51	10.07	58.35	69.26	46.52
50-54 years	16.80	16.09	17.59	48.79	40.80	57.72	2.11	1.62	2.65	30.63	15.38	47.66	0.62	34.58	5.02	72.76	94.38	48.61
55-59 years	21.01	23.24	18.46	73.33	82.51	62.89	5.71	6.33	5.01	31.95	34.79	28.72	6.52	19.97	12.59	103.92	115.52	90.72
60-64 years	10.41	15.05	5.09	67.72	88.12	44.30	5.37	6.37	4.23	45.07	15.21	79.36	3.14	22.62	23.73	120.13	116.02	124.85

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
65-69 years	12·24	17·14	6·64	59·01	75·55	40·08	6·38	4·30	8·77	38·20	26·84	51·21	11·76	32·16	9·86	153·15	187·54	113·80
70-74 years	7·79	11·41	3·93	50·31	57·99	42·13	4·06	3·39	4·78	55·07	23·54	88·64	8·29	29·79	26·71	179·60	195·13	163·07
75-79 years	12·65	5·54	19·57	93·83	104·77	83·20	4·84	5·56	4·14	14·30	24·69	4·19	11·69	30·54	13·09	173·43	135·34	210·45
80-84 years	3·06	3·61	2·60	51·30	75·67	31·08	4·79	7·69	2·39	26·85	7·78	42·67	2·01	13·45	10·81	185·78	262·67	121·95
85+ years	6·46	2·63	8·93	76·36	147·75	30·39	2·63	2·30	2·84	26·76	5·09	40·71	0·23	14·70	7·36	102·42	125·73	87·41
Deaths																		
<5 years	0·84	0·93	0·73	0·80	0·96	0·63	0·00	0·00	0·00	0·13	0·19	0·07	0·01	0·01	0·01	0·39	0·33	0·45
5-9 years	0·53	0·46	0·62	0·42	0·39	0·46	0·00	0·00	0·00	0·07	0·10	0·01	0·08	0·14	0·01	0·13	0·06	0·20
10-14 years	0·52	0·34	0·72	0·48	0·22	0·77	0·00	0·00	0·00	0·11	0·04	0·20	0·05	0·09	0·01	0·22	0·33	0·12
15-19 years	0·33	0·42	0·23	0·85	1·09	0·58	0·00	0·00	0·00	0·02	0·02	0·02	0·09	0·15	0·04	0·28	0·23	0·32
20-24 years	0·24	0·27	0·22	0·56	0·81	0·30	0·01	0·01	0·01	0·07	0·12	0·03	0·18	0·19	0·17	0·35	0·26	0·45
25-29 years	0·28	0·33	0·23	0·85	0·86	0·83	0·01	0·01	0·02	0·28	0·17	0·39	0·21	0·29	0·13	0·53	0·37	0·68
30-34 years	0·09	0·14	0·04	0·58	0·84	0·32	0·01	0·01	0·01	0·21	0·15	0·27	0·31	0·41	0·21	0·48	0·60	0·35
35-39 years	0·24	0·16	0·31	0·96	1·16	0·75	0·02	0·02	0·02	0·39	0·33	0·44	0·26	0·45	0·06	0·71	0·51	0·92
40-44 years	0·13	0·17	0·08	0·91	1·32	0·47	0·02	0·01	0·03	0·11	0·15	0·01	0·27	0·45	0·08	1·12	1·07	1·17
45-49 years	0·31	0·29	0·33	1·40	1·16	1·66	0·02	0·01	0·03	0·63	0·15	1·14	0·40	0·56	0·24	1·38	1·10	1·64
50-54 years	0·45	0·43	0·47	1·30	1·09	1·54	0·06	0·04	0·07	0·82	0·41	1·27	0·55	0·92	0·13	1·94	1·30	2·52
55-59 years	0·64	0·71	0·56	2·24	2·52	1·92	0·17	0·19	0·15	0·98	1·06	0·88	0·50	0·61	0·38	3·17	2·77	3·52
60-64 years	0·37	0·53	0·18	2·40	3·13	1·57	0·19	0·22	0·15	1·60	0·54	2·82	0·82	0·80	0·84	4·26	4·43	4·12

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
65-69 years	0.52	0.72	0.28	2.49	3.18	1.69	0.27	0.18	0.36	1.61	1.13	2.16	0.92	1.35	0.42	6.46	4.81	7.91
70-74 years	0.40	0.59	0.20	2.59	2.98	2.17	0.21	0.17	0.24	2.84	1.21	4.57	1.46	1.53	1.38	9.25	8.41	10.03
75-79 years	0.82	0.36	1.27	6.07	6.78	5.39	0.31	0.36	0.26	0.93	1.60	0.27	1.41	1.98	0.85	11.25	13.67	8.76
80-84 years	0.26	0.30	0.22	4.29	6.33	2.60	0.40	0.64	0.20	2.26	0.65	3.60	1.01	1.13	0.91	15.68	10.31	22.15
85+ years	1.05	0.31	1.52	8.72	17.07	3.34	0.30	0.27	0.32	2.97	0.60	4.50	1.19	1.66	0.88	12.11	10.23	15.03

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Table A2 - Age-specific rates in Australasia and Oceania in 2019 by age groups and by sex

Age groups	ALL			AML			CLL			CML			LIL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
AUSTRALASIA																		
Incidence																		
<5 years	3·41	3·42	3·40	0·49	0·52	0·46	0·00	0·00	0·00	0·08	0·08	0·08	0·23	0·21	0·24	0·00	0·00	0·00
5-9 years	3·34	3·98	2·66	0·32	0·30	0·35	0·00	0·00	0·00	0·05	0·03	0·07	0·34	0·35	0·32	0·97	1·45	0·47
10-14 years	1·59	1·75	1·43	0·39	0·35	0·42	0·00	0·00	0·00	0·06	0·05	0·08	0·69	0·82	0·56	1·02	1·33	0·70
15-19 years	1·74	2·15	1·31	0·73	0·73	0·74	0·00	0·00	0·00	0·10	0·07	0·13	2·43	2·06	2·82	1·92	2·39	1·41
20-24 years	1·26	1·56	0·95	0·77	0·77	0·78	0·11	0·10	0·12	0·22	0·17	0·27	4·55	3·11	4·99	1·49	1·90	1·08
25-29 years	1·02	1·08	0·96	0·84	0·90	0·79	0·10	0·09	0·11	0·35	0·31	0·39	4·90	3·79	5·00	1·93	2·54	1·34
30-34 years	0·71	0·79	0·63	0·74	0·71	0·77	0·10	0·08	0·11	0·33	0·30	0·37	3·77	3·94	3·62	2·51	3·23	1·81
35-39 years	0·73	0·77	0·70	0·90	0·81	1·00	0·19	0·27	0·11	0·51	0·39	0·62	3·03	3·57	2·52	3·58	4·65	2·53
40-44 years	0·74	0·70	0·79	1·16	1·04	1·28	0·41	0·58	0·24	0·55	0·40	0·69	2·78	3·46	2·12	3·44	4·48	2·44
45-49 years	1·15	1·01	1·29	1·65	1·53	1·77	1·13	1·67	0·62	0·63	0·48	0·77	2·26	3·62	1·92	5·55	7·13	4·04
50-54 years	1·90	1·64	2·15	2·74	2·65	2·82	2·68	3·94	1·48	0·88	0·57	1·18	2·19	3·69	1·71	9·66	12·13	7·31
55-59 years	3·15	2·75	3·53	3·87	3·78	3·96	5·66	7·64	3·78	1·42	0·72	2·08	2·39	3·98	1·82	16·45	21·25	11·89
60-64 years	5·14	4·80	5·45	6·42	7·27	5·62	9·58	13·73	5·68	1·83	1·13	2·49	2·75	3·06	2·45	26·27	34·77	18·29
65-69 years	7·18	7·47	6·91	10·43	12·24	8·74	14·40	19·77	9·36	2·63	1·59	3·60	2·95	3·30	2·63	42·41	54·14	31·43
70-74 years	8·86	10·29	7·52	17·73	21·37	14·32	22·50	31·04	14·47	5·00	2·88	6·98	3·16	3·44	2·89	67·82	86·44	50·32

Age groups	ALL			AML			CLL			CML			NHL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
75-79 years	8·54	10·31	6·96	24·48	30·33	19·21	31·40	42·78	21·16	8·06	5·16	10·67	4·08	2·27	3·91	108·54	131·59	87·78
80-84 years	7·57	9·62	5·89	28·81	36·06	22·89	26·29	36·47	17·98	13·42	9·24	16·83	5·17	2·86	4·62	122·55	148·57	101·32
85+ years	7·24	8·79	6·27	39·29	47·14	34·42	41·49	55·39	32·88	25·02	17·88	29·44	7·92	2·45	6·97	176·74	215·23	152·88
Prevalence																		
<5 years	28·18	28·22	28·13	1·18	1·02	1·36	0·00	0·00	0·00	0·43	0·36	0·50	2·0	0·91	2·12	0·00	0·00	0·00
5-9 years	26·60	31·56	21·38	0·78	0·60	0·96	0·00	0·00	0·00	0·29	0·15	0·44	2·98	0·12	2·84	0·00	0·00	0·00
10-14 years	11·66	12·58	10·68	0·75	0·54	0·97	0·00	0·00	0·00	0·32	0·19	0·46	6·14	0·20	5·02	0·00	0·00	0·00
15-19 years	12·25	15·12	9·24	2·23	1·89	2·60	0·00	0·00	0·00	0·55	0·35	0·76	21·5	0·33	24·98	9·01	17·19	0·43
20-24 years	9·10	10·95	7·20	1·85	1·50	2·21	0·94	0·85	1·04	1·19	0·77	1·63	40·53	3·58	44·59	6·44	12·69	0·00
25-29 years	7·38	7·61	7·17	1·68	1·43	1·91	0·86	0·78	0·95	1·78	1·28	2·28	43·29	4·20	44·36	9·22	16·56	1·99
30-34 years	4·84	5·27	4·44	1·18	0·88	1·47	0·83	0·70	0·96	1·64	1·18	2·08	33·1	3·55	31·84	17·29	22·21	12·58
35-39 years	5·16	5·22	5·11	1·29	0·90	1·67	1·59	2·26	0·94	2·53	1·52	3·52	26·4	3·06	21·99	25·19	32·21	18·36
40-44 years	5·46	4·87	6·02	1·47	1·04	1·87	3·31	4·65	2·02	2·69	1·49	3·85	24·1	2·98	18·44	20·56	26·52	14·82
45-49 years	8·56	7·15	9·90	2·31	1·67	2·91	9·30	13·62	5·20	3·13	1·84	4·34	19·32	2·37	16·42	33·54	42·47	25·07
50-54 years	14·93	12·12	17·60	3·81	2·85	4·71	21·81	31·76	12·35	4·43	2·20	6·55	18·48	2·53	14·63	57·00	70·32	44·32
55-59 years	24·96	20·99	28·72	4·73	3·69	5·72	45·98	61·47	31·28	7·08	2·28	11·63	19·31	2·97	14·90	103·35	139·06	69·49
60-64 years	39·62	36·89	42·19	6·63	6·46	6·80	75·96	107·63	46·21	9·10	3·45	14·41	21·62	2·89	19·49	172·89	231·60	117·75
65-69 years	55·54	57·82	53·41	9·62	10·15	9·12	110·60	149·02	74·65	12·87	4·51	20·69	20·99	2·26	18·86	282·65	357·97	212·18
70-74 years	69·60	78·51	61·24	14·64	16·52	12·88	164·84	223·50	109·76	24·59	7·93	40·24	18·62	1·86	17·45	447·32	566·06	335·79

Age groups	ALL			AML			CLL			CML			HLL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
75-79 years	59·89	69·96	50·81	19·02	22·01	16·33	214·12	282·82	152·26	38·46	14·32	60·20	22·43	22·68	22·24	691·00	821·62	573·39
80-84 years	42·02	54·42	31·90	27·69	33·99	22·55	124·12	163·07	92·34	60·31	23·98	89·96	28·12	33·14	26·48	560·72	665·53	475·20
85+ years	30·93	38·08	26·50	30·38	34·02	28·12	126·15	154·80	108·40	95·23	41·63	128·46	33·93	33·65	32·84	620·56	753·81	537·97
DALYs																		
<5 years	26·87	28·83	24·80	26·12	30·77	21·22	0·00	0·00	0·00	1·63	2·34	0·88	0·76	0·77	0·75	7·22	9·14	5·19
5-9 years	37·00	46·31	27·21	15·79	15·74	15·85	0·00	0·00	0·00	0·83	0·89	0·76	1·16	1·28	1·03	9·94	14·92	4·70
10-14 years	28·97	33·68	24·01	19·49	19·50	19·49	0·00	0·00	0·00	1·08	1·33	0·81	2·33	1·93	1·78	10·21	13·34	6·90
15-19 years	36·05	45·15	26·51	28·09	30·08	26·00	0·00	0·00	0·00	1·31	1·58	1·03	6·77	1·90	7·65	18·97	23·74	13·96
20-24 years	23·51	31·94	14·82	31·04	33·65	28·35	0·76	0·77	0·75	3·06	3·83	2·26	12·74	1·97	13·54	23·07	29·63	16·31
25-29 years	18·38	21·55	15·25	34·34	40·03	28·74	0·68	0·72	0·64	5·14	6·96	3·34	17·48	1·98	16·99	28·49	37·48	19·66
30-34 years	12·86	15·84	10·01	30·41	32·27	28·63	0·66	0·67	0·66	4·96	6·82	3·18	14·22	1·77	12·74	34·80	45·01	25·01
35-39 years	11·82	13·86	9·85	35·36	34·92	35·78	1·37	2·15	0·60	6·86	8·74	5·04	12·68	1·45	9·99	46·07	60·10	32·43
40-44 years	10·03	10·85	9·23	43·59	43·25	43·92	2·81	4·36	1·31	6·95	8·52	5·43	12·56	1·85	9·38	60·79	79·45	42·78
45-49 years	11·33	12·07	10·62	53·23	54·28	52·23	6·92	11·10	2·95	6·88	8·61	5·24	10·79	1·83	8·86	88·79	114·51	64·39
50-54 years	13·79	14·55	13·07	78·87	84·07	73·93	14·91	23·87	6·38	8·18	9·20	7·22	10·36	1·93	7·80	138·78	174·67	104·62
55-59 years	16·47	17·17	15·80	100·63	106·91	94·68	27·89	41·71	14·79	11·89	12·96	10·88	13·74	1·39	10·28	208·42	269·62	150·37
60-64 years	19·48	21·18	17·88	152·58	184·02	123·06	49·79	77·19	24·06	14·28	18·02	10·77	16·02	1·61	13·59	303·09	402·15	210·04
65-69 years	23·22	26·09	20·53	219·54	270·80	171·58	76·22	113·23	41·58	18·23	23·92	12·91	20·21	2·11	17·49	417·20	534·39	307·54
70-74 years	26·37	31·64	21·41	321·35	401·70	245·89	120·42	178·24	66·12	28·22	35·48	21·40	24·19	2·31	22·20	558·49	714·78	411·71

Age groups	ALL			AML			CLL			CML			ALL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
75-79 years	27·39	33·23	22·12	370·48	475·03	276·34	163·00	234·27	98·82	39·59	51·78	28·63	26·66	22·68	24·84	728·74	888·41	584·96
80-84 years	26·80	34·20	20·76	333·08	430·72	253·40	190·92	274·71	122·54	51·62	68·15	38·13	25·87	22·58	22·85	826·89	1009·26	678·08
85+ years	24·19	29·81	20·71	267·36	366·38	205·98	253·23	355·58	189·78	70·86	96·23	55·14	21·93	22·24	18·05	838·42	1046·68	709·32
Deaths																		
<5 years	0·28	0·31	0·26	0·30	0·35	0·24	0·00	0·00	0·00	0·02	0·03	0·01	0·00	0·01	0·01	0·08	0·11	0·06
5-9 years	0·43	0·54	0·31	0·19	0·19	0·19	0·00	0·00	0·00	0·01	0·01	0·01	0·00	0·01	0·01	0·11	0·17	0·05
10-14 years	0·37	0·43	0·30	0·25	0·25	0·25	0·00	0·00	0·00	0·01	0·02	0·01	0·00	0·03	0·02	0·13	0·16	0·08
15-19 years	0·49	0·61	0·36	0·39	0·42	0·36	0·00	0·00	0·00	0·02	0·02	0·01	0·00	0·07	0·09	0·25	0·31	0·18
20-24 years	0·34	0·46	0·21	0·46	0·50	0·42	0·01	0·01	0·01	0·04	0·06	0·03	0·15	0·15	0·16	0·33	0·43	0·24
25-29 years	0·29	0·34	0·24	0·55	0·65	0·46	0·01	0·01	0·01	0·08	0·11	0·05	0·24	0·25	0·23	0·44	0·58	0·31
30-34 years	0·22	0·27	0·17	0·53	0·57	0·50	0·01	0·01	0·01	0·09	0·12	0·05	0·22	0·24	0·19	0·59	0·76	0·42
35-39 years	0·22	0·26	0·18	0·68	0·67	0·69	0·02	0·04	0·01	0·13	0·17	0·09	0·22	0·26	0·17	0·84	1·10	0·60
40-44 years	0·20	0·22	0·19	0·93	0·92	0·93	0·05	0·08	0·02	0·14	0·18	0·11	0·23	0·30	0·18	1·25	1·64	0·88
45-49 years	0·25	0·27	0·23	1·26	1·29	1·24	0·15	0·23	0·06	0·16	0·20	0·12	0·23	0·27	0·19	2·03	2·61	1·47
50-54 years	0·34	0·36	0·31	2·10	2·25	1·97	0·34	0·56	0·14	0·21	0·24	0·18	0·24	0·30	0·18	3·55	4·46	2·68
55-59 years	0·44	0·47	0·41	3·07	3·27	2·88	0·72	1·09	0·37	0·35	0·39	0·31	0·38	0·48	0·28	6·05	7·81	4·37
60-64 years	0·57	0·64	0·50	5·43	6·55	4·37	1·50	2·35	0·71	0·49	0·63	0·35	0·51	0·59	0·43	10·19	13·50	7·08
65-69 years	0·78	0·90	0·67	9·30	11·47	7·26	2·74	4·11	1·45	0·73	1·00	0·48	0·78	0·89	0·67	16·47	21·03	12·20
70-74 years	1·06	1·30	0·84	16·64	20·80	12·73	5·27	7·84	2·86	1·36	1·81	0·94	1·17	1·27	1·07	26·36	33·61	19·55

Age groups	ALL			AML			CLL			CML			HLL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
75-79 years	1·43	1·77	1·12	24·27	31·13	18·08	8·93	12·88	5·38	2·39	3·31	1·55	1·59	1·72	1·47	42·20	51·22	34·09
80-84 years	1·86	2·41	1·42	28·28	36·59	21·51	14·82	21·39	9·46	3·94	5·61	2·58	1·86	1·13	1·64	64·02	77·68	52·88
85+ years	2·52	3·13	2·14	32·43	43·74	25·42	29·48	40·86	22·42	7·80	11·30	5·64	1·9	1·49	1·54	92·20	112·39	79·68
OCEANIA																		
Incidence																		
<5 years	1·06	1·19	0·92	1·36	1·55	1·16	0·00	0·00	0·00	0·15	0·20	0·08	0·02	0·02	0·02	0·00	0·00	0·00
5-9 years	0·73	0·64	0·83	0·74	0·67	0·81	0·00	0·00	0·00	0·08	0·12	0·04	0·13	0·23	0·03	0·34	0·49	0·18
10-14 years	0·68	0·45	0·93	0·79	0·34	1·28	0·00	0·00	0·00	0·13	0·04	0·22	0·08	0·14	0·03	0·62	0·35	0·92
15-19 years	0·54	0·68	0·38	1·71	2·12	1·26	0·00	0·00	0·00	0·03	0·04	0·03	0·2	0·30	0·10	0·75	0·81	0·68
20-24 years	0·36	0·41	0·30	1·02	1·46	0·56	0·03	0·03	0·03	0·10	0·17	0·03	0·3	0·36	0·37	0·55	0·69	0·40
25-29 years	0·34	0·40	0·29	1·34	1·34	1·34	0·02	0·02	0·03	0·31	0·19	0·43	0·33	0·44	0·23	0·73	0·95	0·51
30-34 years	0·11	0·17	0·04	0·82	1·18	0·47	0·02	0·02	0·02	0·21	0·16	0·27	0·4	0·57	0·32	0·64	0·49	0·79
35-39 years	0·26	0·18	0·33	1·33	1·58	1·08	0·03	0·02	0·04	0·39	0·34	0·43	0·33	0·58	0·09	0·96	1·24	0·69
40-44 years	0·13	0·18	0·08	1·10	1·62	0·58	0·03	0·02	0·04	0·10	0·14	0·06	0·34	0·56	0·11	1·20	1·26	1·14
45-49 years	0·32	0·29	0·35	1·78	1·40	2·19	0·03	0·02	0·04	0·60	0·15	1·08	0·5	0·70	0·29	1·47	1·71	1·22
50-54 years	0·46	0·45	0·48	1·67	1·36	2·01	0·08	0·06	0·11	0·79	0·40	1·21	0·66	0·12	0·16	2·01	2·58	1·38
55-59 years	0·66	0·72	0·58	2·75	3·02	2·46	0·24	0·24	0·24	0·94	1·00	0·87	0·57	0·71	0·42	3·25	3·61	2·84
60-64 years	0·38	0·54	0·19	2·79	3·56	1·91	0·25	0·27	0·22	1·49	0·52	2·60	0·90	0·89	0·90	4·28	4·09	4·50
65-69 years	0·51	0·70	0·29	2·77	3·48	1·96	0·34	0·21	0·49	1·46	1·00	1·99	0·93	0·39	0·41	6·33	7·77	4·67

Age groups	ALL			AML			CLL			CML			HLL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	0.37	0.54	0.19	2.72	3.14	2.28	0.23	0.18	0.28	2.36	1.00	3.80	1.35	1.46	1.22	8.37	9.13	7.57
75-79 years	0.68	0.31	1.04	6.12	6.85	5.41	0.31	0.34	0.27	0.73	1.24	0.22	1.27	0.74	0.70	10.14	7.64	12.60
80-84 years	0.22	0.26	0.18	4.97	7.20	3.01	0.38	0.61	0.18	1.67	0.52	2.69	0.83	1.00	0.72	13.29	18.01	9.12
85+ years	0.70	0.24	1.02	9.71	18.32	3.70	0.26	0.23	0.28	1.98	0.43	3.07	1.06	1.60	0.69	10.92	13.30	9.26
Prevalence																		
<5 years	3.12	3.51	2.71	3.52	3.56	3.48	0.00	0.00	0.00	0.34	0.48	0.19	0.00	0.08	0.10	0.00	0.00	0.00
5-9 years	2.06	1.79	2.35	1.87	1.47	2.30	0.00	0.00	0.00	0.18	0.26	0.09	0.53	0.89	0.14	0.00	0.00	0.00
10-14 years	1.57	1.04	2.16	1.72	0.60	2.95	0.00	0.00	0.00	0.25	0.08	0.44	0.30	0.46	0.13	1.60	0.00	3.37
15-19 years	1.91	2.41	1.35	5.36	6.13	4.52	0.00	0.00	0.00	0.09	0.09	0.08	1.02	1.44	0.56	1.65	1.41	1.91
20-24 years	1.10	1.24	0.96	2.50	3.35	1.59	0.16	0.15	0.17	0.23	0.38	0.07	1.70	1.47	1.94	0.11	0.21	0.00
25-29 years	0.92	1.07	0.77	2.87	2.48	3.27	0.12	0.10	0.15	0.62	0.37	0.86	1.15	1.37	0.92	0.34	0.57	0.12
30-34 years	0.21	0.34	0.09	1.26	1.62	0.89	0.08	0.07	0.09	0.35	0.26	0.45	1.23	1.44	1.03	0.60	0.19	1.01
35-39 years	0.47	0.34	0.61	1.88	1.93	1.83	0.13	0.10	0.17	0.60	0.53	0.67	0.75	0.27	0.22	1.17	1.51	0.83
40-44 years	0.21	0.29	0.14	1.27	1.70	0.84	0.12	0.07	0.18	0.14	0.20	0.09	0.66	1.10	0.22	0.13	0.21	0.06
45-49 years	0.57	0.51	0.63	2.51	1.61	3.46	0.13	0.07	0.19	0.89	0.23	1.59	0.90	1.40	0.56	0.19	0.36	0.00
50-54 years	0.84	0.77	0.92	2.35	1.55	3.22	0.36	0.22	0.51	1.14	0.58	1.75	1.27	1.15	0.29	0.33	0.64	0.00
55-59 years	1.15	1.25	1.03	3.06	2.97	3.17	1.03	0.99	1.07	1.31	1.39	1.22	0.88	1.12	0.60	0.63	0.97	0.23
60-64 years	0.58	0.83	0.31	2.63	3.14	2.04	0.91	0.94	0.88	1.95	0.65	3.43	1.14	1.19	1.08	0.95	0.63	1.31
65-69 years	0.66	0.90	0.39	2.45	2.95	1.89	1.06	0.58	1.62	1.72	1.16	2.37	0.95	1.45	0.37	1.84	2.30	1.30

Age groups	ALL			AML			CLL			CML			MPL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	0·46	0·67	0·23	2·20	2·49	1·90	0·55	0·40	0·71	2·47	1·02	4·00	1·07	2·23	0·90	2·61	2·70	2·50
75-79 years	0·70	0·28	1·13	4·70	5·09	4·31	0·56	0·57	0·54	0·66	1·14	0·19	0·87	1·30	0·45	4·47	1·88	7·02
80-84 years	0·16	0·18	0·15	4·93	7·23	2·89	0·44	0·69	0·21	1·36	0·40	2·21	0·58	0·75	0·44	3·28	5·43	1·38
85+ years	0·39	0·17	0·53	7·14	13·07	3·00	0·23	0·19	0·26	1·49	0·29	2·33	0·86	1·45	0·46	2·50	3·02	2·14
DALYs																		
<5 years	65·28	73·52	56·44	66·76	80·06	52·47	0·00	0·00	0·00	9·03	12·71	5·07	0·88	1·02	0·73	28·87	33·79	23·60
5-9 years	43·79	38·18	49·90	34·72	33·57	35·97	0·00	0·00	0·00	4·77	6·99	2·34	5·88	1·34	1·02	10·18	15·28	4·63
10-14 years	41·60	27·66	57·05	36·87	17·57	58·25	0·00	0·00	0·00	7·83	2·60	13·63	3·77	1·24	0·97	17·67	9·37	26·87
15-19 years	24·75	31·22	17·59	63·59	81·75	43·48	0·00	0·00	0·00	1·60	1·73	1·45	6·78	1·43	2·74	21·83	24·51	18·86
20-24 years	17·23	19·67	14·61	39·69	58·14	19·98	1·04	1·09	0·99	4·90	8·02	1·56	11·94	1·83	10·99	25·15	31·77	18·06
25-29 years	16·52	19·25	13·76	51·00	53·51	48·47	0·85	0·72	0·97	15·11	9·22	21·06	12·07	1·41	7·58	32·18	42·06	22·19
30-34 years	5·10	8·07	2·18	32·35	47·75	17·19	0·60	0·61	0·60	10·46	7·64	13·23	16·19	2·34	11·12	25·93	19·41	32·36
35-39 years	12·07	8·55	15·58	49·73	61·66	37·83	1·03	0·87	1·19	18·23	16·04	20·42	12·27	2·47	2·98	36·30	46·92	25·70
40-44 years	5·84	8·00	3·65	40·47	60·93	19·74	1·02	0·70	1·34	4·68	6·50	2·84	11·87	1·81	3·71	51·69	54·13	49·21
45-49 years	12·36	11·30	13·49	54·55	45·69	63·95	0·87	0·58	1·18	23·67	6·01	42·41	15·56	2·49	9·28	57·90	67·17	48·07
50-54 years	15·77	15·29	16·29	45·61	39·61	52·19	2·08	1·51	2·70	27·37	14·08	41·95	18·55	3·26	4·59	71·52	91·71	49·36
55-59 years	19·96	22·03	17·61	71·28	81·06	60·22	5·45	5·66	5·22	29·00	31·00	26·75	15·13	1·33	11·50	105·11	116·15	92·61
60-64 years	10·25	14·68	5·22	66·33	86·80	43·11	5·29	5·90	4·59	41·20	14·37	71·64	21·63	2·18	22·15	124·00	118·70	130·01
65-69 years	12·15	16·79	6·81	57·68	73·91	38·99	6·61	4·24	9·34	35·49	24·41	48·26	20·32	3·04	9·12	156·91	192·39	116·06

Age groups	ALL			AML			CLL			CML			HLL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	7·79	11·37	4·02	48·49	56·93	39·62	4·05	3·28	4·86	50·08	21·15	80·47	26·22	22·82	24·52	176·00	192·72	158·45
75-79 years	12·22	5·55	18·81	91·08	103·88	78·46	4·79	5·41	4·18	13·25	22·69	3·95	19·95	22·00	12·00	174·11	135·20	212·49
80-84 years	3·21	3·86	2·63	56·40	82·90	33·03	5·42	8·67	2·55	25·08	7·78	40·34	11·28	11·73	10·00	209·09	284·14	142·90
85+ years	7·28	2·83	10·39	86·54	165·84	31·16	2·90	2·58	3·12	24·07	5·04	37·36	9·75	10·00	6·78	122·12	146·93	104·80
Deaths																		
<5 years	0·75	0·85	0·65	0·77	0·92	0·60	0·00	0·00	0·00	0·10	0·15	0·06	0·00	0·01	0·01	0·34	0·39	0·27
5-9 years	0·53	0·46	0·61	0·42	0·41	0·44	0·00	0·00	0·00	0·06	0·09	0·03	0·00	0·13	0·01	0·12	0·18	0·06
10-14 years	0·54	0·36	0·74	0·48	0·23	0·76	0·00	0·00	0·00	0·10	0·03	0·18	0·00	0·08	0·01	0·23	0·12	0·35
15-19 years	0·34	0·43	0·24	0·88	1·14	0·60	0·00	0·00	0·00	0·02	0·02	0·02	0·09	0·14	0·04	0·30	0·34	0·26
20-24 years	0·26	0·29	0·22	0·59	0·87	0·30	0·02	0·02	0·01	0·07	0·12	0·02	0·18	0·19	0·16	0·38	0·47	0·27
25-29 years	0·27	0·31	0·22	0·82	0·86	0·78	0·01	0·01	0·02	0·24	0·15	0·34	0·19	0·26	0·12	0·52	0·68	0·36
30-34 years	0·09	0·14	0·04	0·57	0·84	0·30	0·01	0·01	0·01	0·18	0·13	0·23	0·28	0·37	0·19	0·45	0·34	0·57
35-39 years	0·23	0·16	0·30	0·96	1·19	0·73	0·02	0·02	0·02	0·35	0·31	0·39	0·23	0·41	0·06	0·70	0·90	0·49
40-44 years	0·12	0·17	0·08	0·86	1·29	0·42	0·02	0·01	0·03	0·10	0·14	0·06	0·23	0·42	0·08	1·10	1·15	1·05
45-49 years	0·29	0·27	0·32	1·29	1·08	1·51	0·02	0·01	0·03	0·56	0·14	1·00	0·36	0·51	0·22	1·37	1·59	1·14
50-54 years	0·42	0·41	0·43	1·22	1·06	1·39	0·05	0·04	0·07	0·73	0·38	1·12	0·49	0·83	0·12	1·91	2·45	1·32
55-59 years	0·61	0·67	0·54	2·17	2·47	1·83	0·16	0·17	0·16	0·89	0·95	0·82	0·46	0·56	0·35	3·21	3·54	2·83
60-64 years	0·36	0·52	0·18	2·35	3·08	1·53	0·18	0·21	0·16	1·46	0·51	2·54	0·77	0·75	0·79	4·40	4·21	4·62
65-69 years	0·51	0·71	0·29	2·43	3·12	1·64	0·27	0·18	0·39	1·50	1·03	2·04	0·86	0·26	0·39	6·62	8·11	4·90

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	0·40	0·58	0·21	2·49	2·93	2·04	0·21	0·17	0·25	2·58	1·09	4·15	1·33	1·43	1·27	9·06	9·91	8·17
75-79 years	0·79	0·36	1·22	5·90	6·73	5·09	0·31	0·35	0·27	0·86	1·47	0·26	1·29	1·81	0·78	11·29	8·75	13·79
80-84 years	0·27	0·32	0·22	4·72	6·93	2·77	0·45	0·72	0·21	2·12	0·65	3·41	0·93	1·07	0·84	17·66	23·98	12·09
85+ years	1·19	0·33	1·79	9·93	19·25	3·43	0·33	0·30	0·35	2·68	0·60	4·13	1·14	1·59	0·82	14·59	17·78	12·36

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Appendix B

Table B1: Results of sensitivity analysis of temporal trends using World standard population


Measure	Base case			Sensitivity analysis			Change in trend
	EAPC (%)	B-Coefficient (95% CI)	P-value	EAPC (%)	B-Coefficient (95% CI)	P-value	
Australasia							
Incidence	-	-	-	-	-	-	
ALL	2.37	0.023 (0.019, 0.027)	<0.001	1.34	0.013 (0.010, 0.017)	<0.001	N/A
AML	1.93	0.019 (0.017, 0.021)	<0.001	0.27	0.003 (0.001, 0.005)	0.015	N/A
CLL	2.40	0.024 (0.022, 0.025)	<0.001	0.68	0.007 (0.005, 0.009)	<0.001	N/A
CML	2.16	0.021 (0.013, 0.029)	<0.001	0.45	0.004 (-0.004, 0.013)	0.300	Stable
HL	0.44	0.004 (0.003, 0.005)	<0.001	0.11	0.001 (0.000, 0.002)	0.027	N/A
NHL	0.95	0.009 (0.005, 0.014)	<0.001	-0.63	-0.006 (-0.012, -0.001)	0.018	Decreasing
Prevalence	-	-	-	-	-	-	
ALL	2.57	0.025 (0.021, 0.030)	<0.001	1.57	0.016 (0.012, 0.020)	<0.001	N/A
AML	0.90	0.009 (0.007,0.011)	<0.001	-0.60	-0.006 (-0.008, -0.004)	<0.001	Decreasing
CLL	2.40	0.024 (0.022, 0.025)	<0.001	0.70	0.007 (0.005, 0.009)	<0.001	N/A
CML	2.87	0.028 (0.022, 0.035)	<0.001	1.27	0.013 (0.006, 0.020)	<0.001	N/A
HL	0.40	0.004 (0.003,0.005)	<0.001	0.19	0.002 (0.001, 0.003)	<0.001	N/A
NHL	1.07	0.011 (0.005, 0.016)	<0.001	-0.68	-0.007 (-0.013, 0.000)	0.046	Decreasing
DALYs	-	-	-	-	-	-	
ALL	-0.36	-0.004 (-0.005, -0.003)	<0.001	-0.13	-0.001(-0.003, 0.001)	0.207	Stable
AML	1.15	0.011 (0.009, 0.013)	<0.001	-0.71	-0.007 (-0.008, -0.006)	<0.001	Decreasing
CLL	1.74	0.017 (0.015, 0.020)	<0.001	0.04	0.000 (-0.002, 0.003)	0.784	Stable
CML	-1.23	-0.012 (-0.024, 0.000)	0.042	-2.43	-0.025 (-0.037, -0.013)	<0.001	N/A
HL	-0.40	-0.004 (-0.005, -0.003)	<0.001	-0.93	-0.009 (-0.011, -0.008)	<0.001	N/A
NHL	0.18	0.002 (-0.004, 0.007)	0.536	-1.12	-0.011 (-0.017, -0.005)	<0.001	Decreasing
Deaths	-	-	-	-	-	-	
ALL	0.75	0.007 (0.006, 0.009)	<0.001	-0.26	-0.003 (-0.004, -0.002)	<0.001	Decreasing
AML	2.07	0.020 (0.019, 0.022)	<0.001	0.42	0.004 (0.003, 0.006)	<0.001	N/A
CLL	2.11	0.021 (0.019, 0.023)	<0.001	0.30	0.003 (0.001, 0.005)	0.002	N/A
CML	-0.27	-0.003 (-0.014, 0.008)	0.631	-2.13	-0.022 (-0.033, -0.010)	<0.001	N/A
HL	0.05	0.000 (-0.001, 0.002)	0.399	-1.02	-0.010 (-0.011, -0.009)	<0.001	Decreasing
NHL	1.07	0.011 (0.005, 0.016)	<0.001	-0.98	-0.010 (-0.015, -0.005)	<0.001	Decreasing
OCEANIA							
Incidence	-	-	-	-	-	-	

ALL	-0.45	-0.005 (-0.006, -0.003)	<0.001	-0.32	-0.003 (-0.005, -0.002)	<0.001	N/A
AML	-0.02	0.000 (-0.002, 0.001)	0.816	-0.18	-0.002 (-0.003, -0.000)	0.012	Decreasing
CLL	1.38	0.014 (0.012, 0.016)	<0.001	0.37	0.004 (0.002, 0.005)	<0.001	N/A
CML	-0.58	-0.006 (-0.008, -0.004)	<0.001	-1.13	-0.011 (-0.013, -0.009)	<0.001	N/A
HL	-0.05	-0.001 (-0.002, 0.001)	0.405	-0.55	-0.005 (-0.007, -0.004)	<0.001	Decreasing
NHL	0.92	0.009 (0.008, 0.010)	<0.001	0.37	0.004 (0.002, 0.005)	<0.001	N/A
Prevalence	-	-	-	-	-	-	-
ALL	-0.56	-0.006 (-0.007, -0.004)	<0.001	-0.33	-0.003 (-0.005, -0.001)	0.001	N/A
AML	-0.21	-0.002 (-0.004, 0.000)	0.026	-0.14	-0.001 (-0.003, 0.001)	0.150	Stable
CLL	1.99	0.020 (0.017, 0.022)	<0.001	0.99	0.010 (0.007, 0.012)	<0.001	N/A
CML	-0.76	-0.008 (-0.010, -0.005)	<0.001	-1.15	-0.012 (-0.014, -0.009)	<0.001	N/A
HL	0.68	0.007 (0.003, 0.011)	0.001	0.42	0.004 (0.000, 0.008)	0.039	N/A
NHL	3.24	0.032 (0.021, 0.043)	<0.001	3.19	0.031 (0.023, 0.040)	<0.001	N/A
DALYs	-	-	-	-	-	-	-
ALL	-0.56	-0.006 (-0.007, -0.004)	<0.001	-0.34	-0.003 (-0.005, -0.002)	<0.001	N/A
AML	-0.12	-0.001 (-0.003, 0.000)	0.115	-0.18	-0.002 (-0.003, -0.000)	0.014	Decreasing
CLL	0.92	0.009 (0.008, 0.010)	<0.001	0.00	0.000 (-0.001, 0.001)	0.993	Stable
CML	-0.88	-0.009 (-0.011, -0.007)	<0.001	-1.21	-0.012 (-0.014, -0.010)	<0.001	N/A
HL	-0.55	-0.006 (-0.007, -0.005)	<0.001	-0.89	-0.009 (-0.010, -0.008)	<0.001	N/A
NHL	0.26	0.003 (0.001, 0.004)	<0.001	-0.05	0.000 (-0.002, 0.001)	0.587	Stable
Deaths	-	-	-	-	-	-	-
ALL	-0.39	-0.004 (-0.005, -0.002)	<0.001	-0.29	-0.003 (-0.004, -0.002)	<0.001	N/A
AML	0.08	0.001 (-0.001, 0.002)	0.244	-0.16	-0.002 (-0.003, -0.000)	0.005	Decreasing
CLL	0.99	0.010 (0.009, 0.011)	<0.001	0.04	0.000 (-0.001, 0.001)	0.499	Stable
CML	-0.48	-0.005 (-0.007, -0.003)	<0.001	-1.11	-0.011 (-0.013, -0.009)	<0.001	N/A
HL	-0.35	-0.003 (-0.004, -0.003)	<0.001	-0.90	-0.009 (-0.010, -0.008)	<0.001	N/A
NHL	0.65	0.007 (0.005, 0.008)	<0.001	0.15	0.001 (0.000, 0.003)	0.140	Stable

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; N/A = not assessed; NHL = non-Hodgkin lymphoma

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Figure 1 - Age-specific rates at regional level by sex, in Australasia and Oceania in 2019



This figure represents age-specific rates (per 100 000 population) of incidence, prevalence, DALYs, and deaths in Australasia (a,b,c,d), and in Oceania (e,f,g,h) by sex in 2019.

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

BMJ Open

Temporal changes in the burden of leukaemia and lymphoma in the Australasia and Oceania regions, 2010-2019: an analysis of the Global Burden of Disease Study 2019

Journal:	BMJ Open
Manuscript ID	bmjopen-2024-084943.R1
Article Type:	Original research
Date Submitted by the Author:	22-Oct-2024
Complete List of Authors:	Ho, Thi Quynh Anh; Deakin University School of Health and Social Development, Deakin Health Economics Lee, Peter; Deakin University, School of Health and Social Development; Gao, Lan; Deakin University School of Health and Social Development, Deakin Health Economics
Primary Subject Heading:	Haematology (incl blood transfusion)
Secondary Subject Heading:	Epidemiology
Keywords:	Lymphoma < HAEMATOLOGY, Leukaemia < HAEMATOLOGY, EPIDEMIOLOGIC STUDIES

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Title page**Title**

Temporal changes in the burden of leukaemia and lymphoma in the Australasia and Oceania regions, 2010-2019: an analysis of the Global Burden of Disease Study 2019

Authors

Thi Quynh Anh Ho*^{#1} - tqho@deakin.edu.au

Peter Lee*¹ - peter.lee@deakin.edu.au

Lan Gao¹ - lan.gao@deakin.edu.au

¹ Deakin University, Institute for Health Transformation, Deakin Health Economics, School of Health and Social Development, Melbourne, Victoria, Australia

Joint first authors*Corresponding author**

T.Q.A Ho

221 Burwood Highway, Burwood, VIC 3125, Australia

E-mail: tqho@deakin.edu.au

Phone: +61 403 844 808

Word count: 3 552 (excluding abstract, keywords, and references)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives

Leukaemias and lymphomas are among the most prevalent and significant cancers in Australasia and Oceania. This study aims to examine the burden of leukaemias/lymphomas and its temporal trend in Australasia and Oceania from 2010 to 2019.

Design

Epidemiological study

Methods

Data from the Global Burden of Disease (GBD) 2019 was used to examine the burden of leukaemia/lymphoma key subtypes (acute lymphocytic leukaemia (ALL), acute myeloid leukaemia (AML), chronic lymphocytic leukaemia (CLL), chronic myeloid leukaemia (CML), Hodgkin-lymphoma (HL), and non-Hodgkin lymphoma (NHL)) by sex and 5-year age groups (from <5yrs to 85yrs+), in terms of incidence, prevalence, disability-adjusted-life-years (DALYs), and deaths. Estimated average percentage changes (EAPCs) were calculated to assess the temporal trends of leukaemia/lymphoma burden (incidence/prevalence/DALYs/deaths) from 2010 to 2019 in Australasia and Oceania.

Results

AML and NHL were the leading causes of leukaemia/lymphoma burden in both regions. Age-standardised rates (ASRs) for AML vs NHL in Australasia were: incidence 4.72 vs. 19.06, DALYs 89.01 vs. 161.68, and deaths 4.15 vs. 8.02 per 100 000 population. ASRs for AML vs. NHL in Oceania were: incidence 1.36 vs 1.08, DALYs 49.16 vs 38.3, and deaths 0.94 vs. 0.98 per 100 000 population. From 2010 to 2019, Australasia observed an increasing trend in incidence/prevalence/deaths across most leukaemias/lymphomas and increasing/stable trend in DALYs for AML/CLL/NHL, while Oceania observed increasing trends in incidence/prevalence/DALYs for CLL/NHL and stable trends in all outcomes (except for prevalence (stable)) for AML. Contrasting mortality trends for ALL/CML/HL were observed between the two regions (increasing/stable in Australasia and decreasing in Oceania). Statistically significant differences were observed in disease burden trends between sexes, with males experiencing a greater increase (or smaller decrease) in the burden for AML in both regions.

Conclusions

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

1 Different temporal trends in leukaemia/lymphoma burden observed in two closely situated
2 geographic regions with different socio-demographic indices highlights the necessity for
3 region-specific intervention strategies to enhance the access to innovative disease treatments,
4 reducing leukaemia/lymphoma burden.

5
6 **Keywords:** Burden of disease, lymphoma, leukaemia
7

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations of this study:

- We compared the disease burden and temporal trends in two geographically similar regions with differing socio-demographic indices (SDI) and performed a sensitivity analysis using the world standard population, that facilitated the comparison of trends between Australasia and Oceania, and to other regions.
- Using region-specific data helps reflect disease burden distributions and trends across sexes and age groups more comprehensively and accurately.
- A direct comparison between Australasia and Oceania is limited due to considerable differences in population distributions and risk factors between the two regions.
- Analysing aggregate data made it challenging to discern the impact of various exposures to disease outcomes, thus trends should be interpreted with caution.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES).

INTRODUCTION

Haematological malignancies, including leukaemias and lymphomas, arise from the uncontrolled proliferation of cells in the lymphatic or circulatory systems. Based on the Global Burden of Disease, Injuries and Risk Factors Study (GBD) 2019, which provides the most comprehensive estimates of global disease and injury burden to date, haematological malignancies contribute to a considerable proportion of the global disease burden attributed to cancer¹⁻³. Globally, leukaemias and lymphomas contributed to 11·7 million and 8·2 million disability-adjusted life years (DALYs) in 2019, respectively². Studies exploring the temporal trend in haematological malignancies across countries using data from the GBD 2019 study have found that over a 30-year period, age-standardised mortality/DALYs have declined, against a background of increasing incident/prevalent burden. However, the distribution of disease burden and temporal trends in leukaemias/lymphomas varies across geographic regions and varying levels of socioeconomic development^{1 2 4 5}. Differences in disease burden across regions of high/low socioeconomic development were largely attributed to social and environmental factors including poverty, educational attainment, and access to health care^{1 2 4}. These large disparities in the health care system highlight the need for population-based epidemiological studies in both high and low-and middle-income countries (LMICs) to inform public health policy and healthcare delivery planning^{1 2 4 5}. Importantly, no studies have systematically explored trends in disease incidence/prevalence or burden of leukaemias/lymphomas for Australasia and Oceania^{1 2}. Epidemiological studies comparing these two Pacific regions are particularly beneficial given the considerable socioeconomic, cultural and ethnic differences between these regions³. As such, a comparison of contemporaneous leukaemia/lymphoma trends between Australasia and Oceania may facilitate the understanding of healthcare disparities, the impacts of sociodemographic factors on disease occurrence and outcomes, and the role of healthcare infrastructure in managing these cancers. Moreover, although data on leukaemia and lymphoma burden are often reported in regional cancer registry reports, and also publicly available in the GBD data set, research specifically focusing on trends in haematological malignancies in Australasia and Oceania are scarce⁶⁻⁹. This gap in the literature underscores the importance of region-specific research to better understand these trends and inform policies tailored to these regions. Ultimately, such a study would inform future research, public healthcare planning strategies, and policies aimed at reducing the burden related to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

leukaemia/lymphoma in Australasia and Oceania – the two regions populated with Indigenous people, closely geographically located but varied sociodemographic factors.⁶⁻⁹Hence, this study aims to (1) examine the prevalence, incidence, mortality, and DALYs attributed to leukaemias and lymphomas by sex and age groups and (2) explore the temporal trend in these metrics for leukaemias and lymphomas from 2010 to 2019 in Oceania and Australasia regions using GBD 2019 data.

METHODS

Data source

We extracted data from the GBD 2019 and performed a secondary analysis in the current study¹⁰. All data was collected using Global Health Data Exchange (GHDx) query tool (<http://ghdx.healthdata.org/gbd-results-tool>)^{3 11}. Details pertaining to the collection, processing, and generation of the GBD 2019 study dataset have been described elsewhere^{2 3}.

Case definition

The definition of leukaemias and lymphomas used in the GBD 2019 study has been defined previously using International Classification of Diseases (ICD) codes (Appendix A)^{2 3}. Leukaemia is typically classified by the type of white blood affected (lymphocytic or myeloid) and disease progression (acute or chronic)¹². Key leukaemia subtypes include acute lymphocytic leukaemia (ALL), acute myeloid leukaemia (AML), chronic lymphocytic leukaemia (CLL), and chronic myeloid leukaemia (CML)¹². Lymphomas are categorised based on the type of lymphocyte affected, with key lymphoma subtypes including non-Hodgkin’s lymphomas (NHL) and Hodgkin’s lymphoma (HL)¹³.

Population and outcome

In this study, we assessed the burden of disease (including prevalence, incidence, deaths, and DALYs) of leukaemias and lymphomas by subtype in Australasia and Oceania from 2010 to 2019. In line with GBD 2019 definitions, Australasia was defined as Australia and New Zealand, and Oceania (18 countries) was defined as American Samoa, Cook Islands, Fiji, Guam, Kiribati, Marshall Islands, Micronesia, Nauru, Niue, Northern Mariana Islands, Palau, Papua New Guinea, Samoa, Solomon Islands, Tokelau, Tonga, Tuvalu, and Vanuatu³.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Patient and public involvement

Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Statistical analysis

A descriptive analysis was performed to characterise the regional burden of leukaemias and lymphomas (by subtype). The number of prevalent and incident cases, deaths, and DALYs were used to derive age-standardised rates (ASRs) (per 100 000 population) in Australasia and Oceania from 2010 to 2019. ASRs were calculated to exclude the impact of age structure on overall population prevalence, incidence, mortality, and DALYs as the number of cases, deaths and DALYs of cancers varies greatly across different age groups¹¹. The ASR per 100 000 population is the weighted-average of age-group-specific rates, that is, $ASR = \frac{\sum_{i=1}^A a_i w_i}{\sum_{i=1}^A w_i} \times 100\,000$, where a_i is the age-group specific rate i and w_i is the weight of age group i of the population¹¹. For the purposes of the base-case analysis, the number of outcomes (incidence, prevalence, deaths or DALYs) occurring for each five-year age group (< 5 years to 85+), were divided by the number of individuals estimated in each age group for Australasia (or Oceania) to estimate age-group-specific rates for each year (a_i). This was multiplied by the proportional distribution (w_i) of individuals in each age group to estimate the ASR.

The estimated average percentage change (EAPC) in ASRs was estimated to assess temporal trends in prevalence, incidence, mortality, and DALYs of leukaemias and lymphomas (by subtype) over a 10-year period (2010 to 2019, inclusive). EAPCs were calculated through generalised linear regression modelling (GLM) with a gaussian family and log-link to estimate trends: $y = \alpha + \beta * x + \varepsilon$, where y is $\ln(ASR)$ and x is the calendar year. The EAPC was expressed as $100 * (e^{\beta} - 1)$ ¹. If the estimated β -coefficient and its 95% CI were both positive, it indicated an upward trend in ASR, whereas a negative β -coefficient and its 95% CI indicated a downward trend in ASR¹¹. Otherwise, the temporal trend in outcomes were assumed to be stable. P-values <0.05 were considered statistically significant.

All calculations were performed using Stata 17.0 statistical software (Stata Corp, College Station, TX).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Sensitivity analysis**

2 As described above, ASRs were calculated in our base case analysis based on the estimated
3 population size across 5-year age groups for Australasia and Oceania. To allow for
4 comparability between trends in Australasia and Oceania, a sensitivity analysis was
5 performed by applying the world standard population weights to age-group-specific rates for
6 key outcomes estimated in our base case analysis ¹⁴.

8 **RESULTS**

9 The leukaemia and lymphoma burden in terms of incidence, prevalence, DALYs, and deaths
10 across Australasia and Oceania and associated temporal trends are summarised in Table 1 and
11 2, respectively. Table 3 and 4 summarises trends in leukaemia and lymphoma burden by sex.
12 Figures 1 summarises the leukaemia and lymphoma burden in terms of incidence, prevalence,
13 DALYs, and deaths across Australasia and Oceania, by age group, and sex in 2019,
14 respectively. Figure 2 illustrates the trends in leukaemia and lymphoma burden by subtypes
15 during the period 2010-2019.

17 [Table 1 around here]

19 [Table 2 around here]

21 [Table 3 around here]

23 [Table 4 around here]

26 [Figure 1 around here]

28 [Figure 2 around here]

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES) .

Overview of disease burden

Leukaemia and lymphoma burden in Australasia

Across Australasia, the greatest contributor to incidence of leukaemia and lymphoma in 2019 was CLL (ASR: 5·22 per 100 000) and NHL (ASR: 19·06 per 100 000), respectively. ALL accounted for the largest share in paediatric leukaemia incidence, whereas CLL accounted for the largest share of incident cases for adult leukaemias. NHL (ASRs: 161·68 DALYs and 8·02 deaths per 100 000), AML (ASRs: 89·01 DALYs and 4·15 deaths per 100 000), followed by CLL, contributed to the greatest proportion in terms of DALYs and mortality burden in 2019. NHL was the most prevalent haematological cancer in 2019. Following stratification of disease burden by five-year age band and sex, the incidence, prevalence, DALY, and mortality rates across five-year age groups generally increased with ascending age across leukaemia/lymphoma subtypes for Australasia (see Figure 1 and Appendix B). Age-specific incidence/prevalence and DALY rates for ALL and HL followed a bimodal pattern.

Leukaemia and lymphoma burden in Oceania

In Oceania, AML and ALL drove leukaemia burden (incident/prevalent/DALYs/deaths) and NHL was the key contributor to the incident lymphoma burden (ASR: 1·08 per 100 000) in 2019. HL (ASR: 0·81 per 100 000) drove prevalent lymphoma burden. Regarding the burden of disease (DALYs/deaths), AML (ASRs: 49·16 DALYs and 0·91 deaths per 100 000) and NHL (ASRs: 38·30 DALYs and 0·98 deaths per 100 000) were the largest contributors in 2019. Disease burden across age-groups and sexes for Oceania were broadly comparable with Australasia. That is, incident/prevalent burden, as well as DALYs/deaths, increased with increasing age and for male sex. (Table 1)

Temporal trends in leukaemia and lymphomas (base-case analysis)

Australasia

The incidence and prevalence of haematological malignancies across Australasia increased significantly over a 10-year period (2010 to 2019, inclusive) ($P<0\cdot001$). Temporal increasing trends in incident/prevalent cases were largely driven by CLL, ALL, and CML (EAPCs ranging from 2·16% to 2·87%) (all $P<0\cdot001$). Notably, the EAPC and associated 95% CI for leukaemia and lymphoma incidence was broadly comparable between sexes with the exception of CML and HL (greater for females), and AML (greater for males). Regarding

1 prevalent burden, EAPCs in males were found to be higher for AML, and lower for HL
2 relative to females.
3 DALY burden (per 100 000) increased significantly for CLL (EAPC: 1·74%) and AML
4 (EAPC: 1·15%) over time, whereas significant reductions in DALY burden attributed to
5 CML (EAPC: -1·23%), HL (EAPC: -0·40%) and ALL (EAPC: -0·36%) were observed (all
6 $P<0\cdot05$). DALY burden attributed to NHL remained stable over time (both sexes). Male sex
7 drove temporal reductions in DALYs (ALL, CML, and HL) whereas the only significant
8 reduction in DALY burden for females was for HL. Although AML and CLL DALY burden
9 increased for both sexes, the EAPC estimated for AML was of greater magnitude for males
10 relative to females (EAPCs: 1·51% versus 0·69%). Lastly, mortality from
11 leukaemias/lymphomas significantly increased over time (EAPCs ranging from 0·75% for
12 ALL to 2·11% for CLL), with the exception of CML and HL (stable). Overall, trends in
13 mortality were comparable between sexes, with the exception of greater increases in
14 mortality over time for AML and NHL for males compared with females (EAPCs: 2·55%
15 versus 1·46%). (Table 2)

16
17 *Oceania*

18 The increased incidence in CLL (EAPC: 1·38%) and NHL (EAPC: 0·92%) was balanced
19 with significantly decreases in ALL (EAPC: -0·45%) and CML (EAPC: -0·58%) incidence.
20 The overall stable trend in HL incidence over time was a result of opposing temporal trends
21 in males (decreasing with EAPC: -0·19%) and females (increasing with EAPC: 0·31%). The
22 overall incidence of AML was stable, driven by females (stable), despite increasing over time
23 for males (EAPC: 0·17%). Although CLL incidence increased over time for both sexes, the
24 EAPC was higher for females. Significant increases in the prevalence of NHL (EAPC:
25 3·24%), CLL (EAPC: 1·99%) and HL (EAPC: 0·68%) were balanced with reductions in the
26 prevalence of CML (EAPC: -0·76%), ALL (EAPC: -0·56%) and AML (EAPC: -0·21%).
27 Although temporal trends for ALL, CML CLL, HL and NHL prevalence were in the same
28 direction across both sexes (decreasing for ALL/CML and increasing for CLL/HL/NHL), the
29 decline in AML prevalence was driven by females (stable for males). Ultimately, the overall
30 incidence and prevalence of leukaemias/lymphomas across Oceania had remained stable over
31 time.
32 In terms of disease burden, modest but statistically significant annual reductions in DALYs
33 were observed for ALL, CML and HL (range in EAPCs: -0·55% for HL to -0·88% for CML)
34 across both sexes, whereas the DALY burden attributed to CLL (EAPC: 0·92%) and NHL

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES)

increased over time (EAPC: 0·26%) (all $P < 0\cdot001$). AML DALY burden over time was stable, and largely driven by males (stable in males while decreasing in females). Lastly, with the exception of AML, CLL and NHL, the mortality burden attributed to leukaemias/lymphomas declined over time (range in EAPCs: -0·48% for CML to -0·35% for HL). Between 2010 and 2019, significant declines in mortality burden attributed to ALL and CML, and significantly increased CLL mortality, were estimated across both sexes. However, AML mortality burden increased significantly for males, and remained stable over time for females. Furthermore, the increase in CLL mortality over time was greater for females relative to males. Regarding lymphoma subtypes and sex, the overall decline in HL mortality was driven by males (stable for females), while NHL mortality increased over time for both sexes. (Table 2)

Sensitivity analysis

Results of the sensitivity analysis are presented in Appendix C. While broadly comparable with base-case estimates, discrepancies in the direction of the trend (decreasing, stable or increasing) over time were identified for 20 (42%) out of 48 estimable trends following application of the World Standard population.

DISCUSSION

Our study is the first to report trends across a variety of outcomes between Australasia and Oceania over a 10-year period (2010 – 2019, inclusive) in which considerable changes in clinical practice occurred for cancer¹⁻³. Discrepancies in disease burden (as indicated by ASRs) across regions, and across age groups and sexes were observed. This is likely attributed to differences in age distribution^{1 15}, differences in health care and cancer surveillance/registration systems, as well as differences in accessing the social determinants of health between high socio-demographic indices (SDI) and low-SDI countries^{1 2}. For example, recent advances in the treatment of NHL and AML have contributed to improved patient survival, including the development of immune-based cellular and antibody therapies, and small molecule inhibitors^{15 16}. However, despite improved management of patients with leukaemias/lymphomas in Australasia, the considerable disease burden for both NHL and AML warrants further research to address unmet need. Moreover, NHL contributed the most to mortality burden in Oceania, which may highlight issues in the effective management of NHL in the region. Upon stratification to explore the distribution of age (5-year age group) and sex (male/female) burden, the observed leukaemia/lymphoma burden was greater for

1
2
3 1 older (versus younger) persons and males (versus females) across both regions. This is in line
4
5 2 with other studies exploring patient outcomes in AML, and has been attributed to differences
6
7 3 in biological risk factors, lifestyle, or environmental factors, and differences in disease
8
9 4 treatment and management [17 18](#). Factors including issues with health systems reporting,
10
11 5 differential access to health services between females and males, the interaction between sex
12
13 6 and age distribution, and gender inequality have also been implicated in contributing to
14
15 7 discrepancies in disease burden between sexes across low-and high-SDI countries [1 5 19](#).
16
17 8 However, further studies are recommended to establish and address sex-based drivers of
18
19 9 leukaemia/lymphoma burden in countries with low-or-middle SDIs [1 5 10 19](#). The greater
20
21 10 burden observed in adults and older age groups also highlights the need for age-period-cohort
22
23 11 effect analysis to explore age-specific risk factors and examine the effects of age, time period,
24
25 12 and birth cohort on leukaemia/lymphoma incidence and mortality in these regions [20](#).
26
27 13 The interaction between factors for differences in leukaemia/lymphoma burden trends
28
29 14 between Australasia and Oceanian is similarly complex. First, there are considerable
30
31 15 differences in ethnic and genetic distribution across regions. For example, CLL is relatively
32
33 16 rare among Pacific Islanders (who comprise the majority of Oceanian populations) compared
34
35 17 with Caucasian or European peoples (Australasia) [21](#). Second, the majority of countries in
36
37 18 Oceania have limited healthcare infrastructure and access to healthcare specialists, as well as
38
39 19 limited access to early detection and advanced medical treatments relative to Australasia [6 22](#).
40
41 20 Moreover, although there are several cancer registries in Oceania, the quality and
42
43 21 completeness of registration varies [23](#). For example, cancer registries in Fiji have reported
44
45 22 cases under-registration, gender miscoding, and variations in coding causes of deaths. Issues
46
47 23 regarding the completeness of data and validity of diagnoses have also been reported in
48
49 24 Tonga, Cook Islands, and Niue. In the US-affiliated islands such as American Samoa and
50
51 25 Guam, there is the potential for missing cases due to historical barriers and the lack of
52
53 26 resources for diagnosis and staging [6](#). In contrast, Australasia has well-developed healthcare
54
55 27 systems with advanced medical technologies, leading to earlier detection, more effective
56
57 28 treatment options and improved survival rates [22](#). Notably, significant reductions in the
58
59 29 CML/HL/ALL DALY burden were observed, and NHL DALY burden remained stable over
60
30 time which reflect improved outcomes due to changes in patient management and treatment
31
32 across Australasia [24-27](#). However, overall, our findings highlight the need to improve
33
34 prevention strategies to mitigate exposure to lifestyle factors associated with
leukaemia/lymphoma, and the efficient allocation of resources among vulnerable population
groups for the treatment for leukaemias/lymphomas [28](#).

First, while trends were comparable across both sexes; a greater increase in AML mortality for males was estimated relative to females. As with disparate disease burden between sexes, disparate trends between sexes is consistent with existing studies exploring the leukaemia/lymphoma burden for high-SDI countries [1 10](#). That is, differences be attributed to differences in tobacco use, occupational exposure to carcinogens and high body mass index (BMI) [29](#). Second, the discrepancy between reductions/stable DALY burden (ALL and NHL) against increasing mortality/incidence/prevalence over time may highlight disparate outcomes across age groups [8 24 25 30](#). ALL follows a bimodal age distribution, with incident/prevalent cases peaking during early childhood and later adulthood (≥ 50 years), while NHL cases and the risk of mortality (ALL and NHL) increase with age [17 31 32](#). Therefore, it is likely that while improved disease management and treatment over time have reduced disease burden (DALYs) for younger patients, these changes may not have had an equal impact on reducing mortality burden among older patients [24](#). This is supported by recent studies exploring ALL and NHL outcomes across Australasia, which found improvements in the management or treatment of disease coincided with greater gains in survival outcomes for younger relative to older patients [8 24 25 30 33](#). To effectively address the ongoing burden of leukaemias/lymphomas, further research should prioritise exploring trends in disease burden and treatment outcomes across different age groups, with a focus on identifying and addressing the factors contributing to the observed disparities among these groups. Although causes of most leukaemias/lymphomas are unknown, general cancer prevention strategies can target lifestyle factors, such as avoiding tobacco, having a healthy diet, and reducing exposure to hazards such as radiation and toxic chemicals to reduce the risk of leukaemias/lymphomas [1](#). Lastly, discrepancies in temporal trends were identified in sensitivity analyses using the world standard population. Notably, disease burden attributed to AML and NHL reduced across both regions in comparison with base-case estimates. This is likely attributed to the variation in population structure in each region, which differs from the standard population [3 13](#). As sex and age are associated with cancer outcomes, it is possible that EAPCs estimated using the standard population would inadequately capture the impact of region-specific differences in age/sex, and ultimately, overestimate patient survival trends.

A key strength of our study lies in using data from the GBD 2019 study, which to date, provides the latest regional epidemiological distributions and trends of leukaemias and lymphomas [3](#). Besides, the comprehensive analysis of trends across a variety of key outcomes facilitates an understanding of the impacts of changes in patient management/treatment on

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 patient outcomes and potential areas of unmet need. Furthermore, disease burden and
2 temporal trends in our study were estimated based on region-specific population, which may
3 better reflect disease burden distribution and trend across age groups and sexes more
4 comprehensively and accurately, particularly when regional population structure is
5 incomparable to the world standard population. Lastly, restricting the analysis to two
6 geographically similar regions with differing SDI, and performing a sensitivity analysis using
7 the world standard population facilitates the comparison of trends between the two regions
8 (base-case), and to other regions or nations at the global level (sensitivity analysis).
9 However, several limitations to our analyses warrant mention. First, this study used data from
10 GBD 2019, which does not include data in 2020 and 2021 (which will be available in the
11 latest GBD update). As a result, it may not capture the most recent trends in disease burden
12 up to 2021. Other limitations associated with GBD 2019 data have been described previously
13 [1-4](#). In brief, it is likely that key outcomes for Oceania were underestimated, as the capacity to
14 collect reliable data on haematological malignancies is often lacking in low-SDI regions.
15 However, this does not considerably change our findings of disparities in disease burden and
16 temporal trends observed between Australasia and Oceania, as well as between male/female
17 sex. Moreover, data variation among the included countries in both regions, such as data
18 quality, accuracy and the degree of missing data, might contribute to the deviation in the
19 estimates, leading to discrepancies between regions [1](#). Second, GBD data includes disease
20 classifications by sex and age groups, but lacks ethnicity data, which limits the analysis of
21 genetic susceptibility to disease. Third, a direct comparison between Australasia and Oceania
22 could be difficult as there are considerable differences in the population distribution and risk
23 factors between the two regions [1-4](#). As such, to explore the potential impacts attributed to age
24 and sex distribution, our analyses used both region-specific populations (base-case) and
25 standard population weights (sensitivity analyses) to explore changes in age-standardised
26 outcomes. Fourth, it was not possible to discern the impact of various exposures on disease
27 outcomes using aggregate data; as such, trends should be interpreted with caution due to
28 potential confounding [1-4](#). Additionally, our EAPC model were limited to detect constant
29 linear trends during the 10-year period. Future research could consider capturing nonlinear
30 trends during this explored period. Moreover, trends of leukaemia/lymphoma burden were
31 predicted based on a single measure, age-standardised rate, for all age groups. This is likely
32 to overlook the differences in trends between children/adolescents and adult population, who
33 might have distinct characteristics and exposures to leukaemias/lymphomas. Lastly, it was

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

not possible to capture the impacts of the COVID-19 pandemic on leukaemia/lymphoma burden using GBD 2019 data [2334](#). Ultimately, our findings highlight disparities in the management and treatment of haematological malignancies based on SDI and sex. Despite the emergence of novel therapies and improved treatment/management over time, the consistent and considerable burden of AML and NHL in both regions warrants further research to mitigate the gap attributed to socioeconomic disadvantage. Further research is also recommended to explore factors that contribute to disparate outcomes for sex (male/female).

CONCLUSION

This study captures the disease burden of leukaemias/lymphomas and its temporal trends in two closely situated geographic regions with different SDI. The considerable disparity in disease burden observed between the two regions suggests the need for early diagnosis and better management strategies tailored to each region. Further study is required to explore the underlying factors behind the epidemiological trends of different leukaemia and lymphoma subtypes in each region and the male predominance in the majority of leukaemia and lymphoma burden of disease.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Ethics and dissemination

Ethics approval is not required for this study as the study involve information freely available in an open source.

Patient consent

Non patient consent required.

Data availability statement

All data relevant to the study are included in the article and uploaded as supplementary information.

Patient and public involvement

Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Authors' contributions

Ho TQA. — conceptualisation, data collection, formal analysis, investigation, methodology, visualisation, writing - original draft, and writing-review & editing.
Lee P. — conceptualisation, data collection, formal analysis, investigation, methodology, visualisation, writing-original draft, and writing-review & editing.
Gao L. — conceptualisation, investigation, methodology, supervision, writing-review & editing. Gao L. is the guarantor.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests

Non declared.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES).

References

References

1. Zhang N, Wu J, Wang Q, et al. Global burden of hematologic malignancies and evolution patterns over the past 30 years. *Blood Cancer J* 2023;13(1):82. doi: 10.1038/s41408-023-00853-3 [published Online First: 20230517]
2. Global Burden of Disease Cancer Collaboration. Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life Years for 29 Cancer Groups From 2010 to 2019: A Systematic Analysis for the Global Burden of Disease Study 2019. *JAMA Oncology* 2022;8(3):420-44. doi: 10.1001/jamaoncol.2021.6987
3. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020;396(10258):1204-22. doi: 10.1016/s0140-6736(20)30925-9
4. Tran KB, Lang JJ, Compton K, et al. The global burden of cancer attributable to risk factors, 2010–19: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2022;400(10352):563-91. doi: [https://doi.org/10.1016/S0140-6736\(22\)01438-6](https://doi.org/10.1016/S0140-6736(22)01438-6)
5. Yao Y, Lin X, Li F, et al. The global burden and attributable risk factors of chronic lymphocytic leukemia in 204 countries and territories from 1990 to 2019: analysis based on the global burden of disease study 2019. *BioMedical Engineering OnLine* 2022;21(1):4. doi: 10.1186/s12938-021-00973-6

1
2
3 1 6. Van Dyne EA, Saraiya M, White A, et al. Cancer Mortality in the US-Affiliated Pacific
4
5 2 Islands, 2008-2013. *Hawaii J Health Soc Welf* 2020;79(6 Suppl 2):99-107.
6
7
8
9
10 3 7. Moore MA, Baumann F, Foliaki S, et al. Cancer epidemiology in the pacific islands - past,
11
12 4 present and future. *Asian Pac J Cancer Prev* 2010;11 Suppl 2(0 2):99-106.
13
14
15
16 5 8. Australian Institute of Health and Welfare. Cancer in Australia 2021. Canberra: AIHW,
17
18 6 2021.
19
20
21
22
23 7 9. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204
24
25 8 countries and territories, 1990–2019: a systematic analysis for the Global Burden of
26
27 9 Disease Study 2019. *The Lancet* 2020;396(10258):1204-22. doi:
28
29 10 [https://doi.org/10.1016/S0140-6736\(20\)30925-9](https://doi.org/10.1016/S0140-6736(20)30925-9)
30
31
32
33
34 11 10. The Global Burden of Disease Collaborator Network. Global Burden of Disease
35
36 12 Collaborative Network. Global Burden of Disease Study 2019 (GBD 2019) Disease
37
38 13 and Injury Burden 1990-2019. 2019 ed. Seattle. USA., 2020.
39
40
41
42
43 14 11. Lin X, Wang J, Huang X, et al. Global, regional, and national burdens of leukemia from
44
45 15 1990 to 2017: a systematic analysis of the global burden of disease 2017 study. *Aging*
46
47 16 *(Albany NY)* 2021;13(7):10468.
48
49
50
51
52 17 12. Bispo JAB, Pinheiro PS, Kobetz EK. Epidemiology and Etiology of Leukemia and
53
54 18 Lymphoma. *Cold Spring Harb Perspect Med* 2020;10(6) doi:
55
56 19 10.1101/cshperspect.a034819 [published Online First: 20200601]
57
58
59
60

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

13. Alaggio R, Amador C, Anagnostopoulos I, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Lymphoid Neoplasms. *Leukemia* 2022;36(7):1720-48. doi: 10.1038/s41375-022-01620-2 [published Online First: 20220622]
14. Ahmad OB, Boschi Pinto C, Lopez A, et al. Age Standardization of Rates: A New WHO Standard. *GPE Discussion Paper Series, EIP/GPE/EBD, World Health Organization* 2001;No.31
15. Skirbekk V, Dieleman JL, Stonawski M, et al. The health-adjusted dependency ratio as a new global measure of the burden of ageing: a population-based study. *Lancet Healthy Longev* 2022;3(5):e332-e38. doi: 10.1016/s2666-7568(22)00075-7 [published Online First: 20220504]
16. Winer ES, Stone RM. Novel therapy in Acute myeloid leukemia (AML): moving toward targeted approaches. *Ther Adv Hematol* 2019;10:2040620719860645. doi: 10.1177/2040620719860645 [published Online First: 20190710]
17. Sawalha Y, Maddocks K. Novel treatments in B cell non-Hodgkin's lymphomas. *BMJ* 2022;377:e063439. doi: 10.1136/bmj-2020-063439
18. Dong Y, Shi O, Zeng Q, et al. Leukemia incidence trends at the global, regional, and national level between 1990 and 2017. *Exp Hematol Oncol* 2020;9:14. doi: 10.1186/s40164-020-00170-6 [published Online First: 20200619]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

19. Rubin JB, Lagas JS, Broestl L, et al. Sex differences in cancer mechanisms. *Biol Sex Differ* 2020;11(1):17. doi: 10.1186/s13293-020-00291-x [published Online First: 20200415]

20. Lin Q, Mao L, Shao L, et al. Global, Regional, and National Burden of Chronic Myeloid Leukemia, 1990-2017: A Systematic Analysis for the Global Burden of Disease Study 2017. *Front Oncol* 2020;10:580759. doi: 10.3389/fonc.2020.580759 [published Online First: 20201215]

21. Rosenberg PS, Check DP, Anderson WF. A web tool for age-period-cohort analysis of cancer incidence and mortality rates. *Cancer Epidemiol Biomarkers Prev* 2014;23(11):2296-302. doi: 10.1158/1055-9965.Epi-14-0300 [published Online First: 20140821]

22. Mukkamalla S, Taneja A, Malipeddi D, et al. Chronic Lymphocytic Leukemia. [Updated 2023 Mar 7]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470433/> StatPearls: Treasure Island (FL): StatPearls Publishing; 2023 [updated 7 Mar 2023. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470433/>.

23. Eric C. Schneider, Arnav Shah, Michelle M. Doty, et al. Mirror, Mirror 2021 - Reflecting Poorly: Health Care in the U.S. Compared to Other High-Income Countries 2021.

24. Tervonen H, Foliaki S, Bray F, et al. Cancer epidemiology in the small nations of Pacific Islands. *Cancer Epidemiology* 2017;50:184-92. doi: <https://doi.org/10.1016/j.canep.2017.09.002>

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

- 1 25. Beckmann K, Kearney BA, Yeung D, et al. Changes in five-year survival for people with
2 acute leukaemia in South Australia, 1980-2016. *Med J Aust* 2022;216(6):296-302.
3 doi: 10.5694/mja2.51423 [published Online First: 20220216]
- 4 26. Wong SHC, Offner M, Hu R, et al. Treatment outcomes of adult acute lymphoblastic
5 leukaemia in Auckland, New Zealand. *N Z Med J* 2023;136(1572):10-18. [published
6 Online First: 20230324]
- 7 27. Cortes J, Pavlovsky C, Sauße S. Chronic myeloid leukaemia. *The Lancet*
8 2021;398(10314):1914-26. doi: 10.1016/S0140-6736(21)01204-6
- 9 28. Nguyen J, Wellard C, Chung E, et al. Clinical characteristics of Australian treatment-
10 naïve patients with classical Hodgkin lymphoma from the lymphoma and related
11 diseases registry. *Eur J Haematol* 2023;110(4):386-95. doi: 10.1111/ejh.13915
12 [published Online First: 20221228]
- 13 29. Coughlin SS, Datta B, Majeed B. Preventive Behaviors Among Leukemia and
14 Lymphoma Cancer Survivors: Results From the 2020 Behavioral Risk Factor
15 Surveillance System Survey. *AJPM Focus* 2023;2(1) doi:
16 10.1016/j.focus.2022.100041 [published Online First: 20221024]
- 17 30. Murray CJL, Aravkin AY, Zheng P, et al. Global burden of 87 risk factors in 204
18 countries and territories, 1990–2019: a systematic analysis for the Global
19 Burden of Disease Study 2019. *The Lancet* 2020;396(10258):1223-49. doi:
20 10.1016/S0140-6736(20)30752-2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

31. Anderson MA, Berkahn L, Cheah C, et al. Improving outcomes for patients with lymphoma: design and development of the Australian and New Zealand Lymphoma and Related Diseases Registry. *BMC Medical Research Methodology* 2022;22(1):266. doi: 10.1186/s12874-022-01728-0

32. Terwilliger T, Abdul-Hay M. Acute lymphoblastic leukemia: a comprehensive review and 2017 update. *Blood Cancer Journal* 2017;7(6):e577-e77. doi: 10.1038/bcj.2017.53

33. Sun H, Xue L, Guo Y, et al. Global, regional and national burden of non-Hodgkin lymphoma from 1990 to 2017: estimates from global burden of disease study in 2017. *Annals of Medicine* 2022;54(1):633-45. doi: 10.1080/07853890.2022.2039957

34. Wright F, Hapgood G, Loganathan A, et al. Relative survival of patients with lymphoma in Queensland according to histological subtype. *Med J Aust* 2018;209(4):166-72. doi: 10.5694/mja17.00937 [published Online First: 20180813]

35. Luo Q, O'Connell DL, Yu XQ, et al. Cancer incidence and mortality in Australia from 2020 to 2044 and an exploratory analysis of the potential effect of treatment delays during the COVID-19 pandemic: a statistical modelling study. *Lancet Public Health* 2022;7(6):e537-e48. doi: 10.1016/s2468-2667(22)00090-1

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Tables

Table 1 – Results of ASRs and EAPCs for key outcomes across leukaemia and lymphoma subtypes in Australasia in 2010 and 2019

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-C Coefficient (95% CI)	P-value
	2010	2019			
Incidence	32.59 (28.7, 37.18)	36.52 (28.56, 46.37)	-	-	-
ALL	2.38 (2.02, 3.01)	2.91 (2.24, 3.86)	2.37	0.029 (-0.19, 0.027)	<0.001
AML	4.07 (3.34, 4.39)	4.72 (3.67, 5.9)	1.93	0.019 (-0.17, 0.021)	<0.001
CLL	4.22 (3.77, 5.21)	5.22 (4.07, 6.87)	2.40	0.029 (-0.22, 0.025)	<0.001
CML	1.58 (1.35, 1.95)	1.82 (1.37, 2.45)	2.16	0.029 (-0.13, 0.029)	<0.001
HL	2.70 (2.30, 3.26)	2.79 (2.14, 3.66)	0.44	0.004 (-0.03, 0.005)	<0.001
NHL	17.64 (15.92, 19.38)	19.06 (15.08, 23.62)	0.95	0.004 (-0.05, 0.014)	<0.001
Prevalence	175.46 (150.04, 208.28)	196.64 (150.07, 256.58)	-	-	-
ALL	17.24 (14.33, 21.87)	21.44 (16.25, 28.53)	2.57	0.029 (-0.21, 0.030)	<0.001
AML	4.46 (3.64, 5.11)	4.75 (3.74, 5.89)	0.90	0.009 (-0.07, 0.011)	<0.001
CLL	27.34 (24.51, 33.8)	33.84 (25.92, 45.03)	2.40	0.029 (-0.22, 0.025)	<0.001
CML	6.74 (5.45, 8.49)	8.26 (6.06, 11.42)	2.87	0.029 (-0.22, 0.035)	<0.001
HL	21.45 (18.19, 25.9)	22.05 (16.83, 29.28)	0.40	0.004 (-0.03, 0.005)	<0.001
NHL	98.23 (83.92, 113.11)	106.3 (81.27, 136.44)	1.07	0.019 (-0.05, 0.016)	<0.001
DALYs	312.84 (281.93, 341.76)	324.03 (281.64, 366.04)	-	-	-
ALL	21.79 (19.98, 25.93)	21.08 (18.6, 24.78)	-0.36	-0.004 (-0.005, -0.003)	<0.001
AML	81.57 (68.56, 86.23)	89.01 (73.7, 97.38)	1.15	0.011 (0.009, 0.013)	<0.001
CLL	24.99 (21.98, 31)	29.39 (25.11, 36.76)	1.74	0.017 (0.015, 0.020)	<0.001

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-Coefficient (95% CI)	P-value
	2010	2019			
CML	12·02 (10·94, 14·49)	10·49 (9·15, 12·65)	-1·23	-0·00 (-0·024, 0·000)	0·042
HL	12·77 (10·9, 15·22)	12·38 (10·44, 15·16)	-0·40	-0·00 (-0·005, -0·003)	<0·001
NHL	159·7 (149·57, 168·89)	161·68 (144·64, 179·31)	0·18	0·00 (0·004, 0·007)	0·536
Deaths	13·87 (12·2, 15·21)	15·27 (12·97, 17·36)	-	-	-
ALL	0·47 (0·42, 0·57)	0·50 (0·44, 0·60)	0·75	0·00 (0·006, 0·009)	<0·001
AML	3·51 (2·87, 3·77)	4·15 (3·32, 4·65)	2·07	0·02 (0·019, 0·022)	<0·001
CLL	1·4 (1·21, 1·71)	1·71 (1·43, 2·13)	2·11	0·02 (0·019, 0·023)	<0·001
CML	0·55 (0·49, 0·68)	0·52 (0·44, 0·65)	-0·27	-0·00 (-0·014, 0·008)	0·631
HL	0·37 (0·32, 0·44)	0·38 (0·32, 0·45)	0·05	0·00 (-0·001, 0·002)	0·399
NHL	7·57 (6·9, 8·04)	8·02 (7·03, 8·87)	1·07	0·01 (0·005, 0·016)	<0·001

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Table 2 – Results of ASRs and EAPCs for key outcomes across leukaemia and lymphoma subtypes in Oceania in 2010 and 2019

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-Coefficient (95% CI)	P-value
	2010	2019			
Incidence	3.55 (2.65, 4.9)	3.62 (2.68, 4.96)	-	-	-
ALL	0.55 (0.33, 0.93)	0.53 (0.33, 0.84)	-0.45	-0.006 (-0.006, -0.003)	<0.001
AML	1.36 (1.01, 1.88)	1.36 (0.98, 1.89)	-0.02	0.000 (-0.002, 0.001)	0.816
CLL	0.03 (0.02, 0.04)	0.04 (0.03, 0.05)	1.38	0.012 (0.012, 0.016)	<0.001
CML	0.32 (0.2, 0.51)	0.3 (0.19, 0.46)	-0.58	-0.008 (-0.008, -0.004)	<0.001
HL	0.3 (0.24, 0.39)	0.3 (0.23, 0.41)	-0.05	-0.002 (-0.002, 0.001)	0.405
NHL	0.99 (0.85, 1.15)	1.08 (0.91, 1.31)	0.92	0.008 (0.008, 0.010)	<0.001
Prevalence	5.97 (3.95, 9.17)	6.09 (4.05, 9.14)	-	-	-
ALL	1.46 (0.85, 2.52)	1.39 (0.86, 2.26)	-0.56	-0.007 (-0.007, -0.004)	<0.001
AML	2.68 (1.81, 4.08)	2.63 (1.75, 4)	-0.21	-0.001 (-0.004, 0.000)	0.026
CLL	0.12 (0.09, 0.16)	0.14 (0.1, 0.19)	1.99	0.022 (0.017, 0.022)	<0.001
CML	0.51 (0.3, 0.85)	0.47 (0.28, 0.74)	-0.76	-0.008 (-0.010, -0.005)	<0.001
HL	0.75 (0.59, 0.94)	0.81 (0.6, 1.07)	0.68	0.003 (0.003, 0.011)	0.001
NHL	0.45 (0.31, 0.62)	0.64 (0.45, 0.88)	3.24	0.032 (0.021, 0.043)	<0.001
DALYs	140.07 (100.83, 201.45)	137.73 (98.65, 196.41)	-	-	-
ALL	29.25 (16.74, 51.13)	27.86 (16.96, 45.85)	-0.56	-0.006 (-0.007, -0.004)	<0.001
AML	49.78 (36.68, 70.38)	49.16 (35.01, 71.49)	-0.12	-0.001 (-0.003, 0.000)	0.115
CLL	0.83 (0.6, 1.12)	0.91 (0.65, 1.28)	0.92	0.009 (0.008, 0.010)	<0.001
CML	12.79 (7.25, 21.8)	11.82 (6.88, 18.89)	-0.88	-0.009 (-0.011, -0.007)	<0.001
HL	10.11 (7.85, 13.12)	9.67 (7.29, 13.07)	-0.55	-0.006 (-0.007, -0.005)	<0.001

Measure	ASRs per 100 000 (95% CI)		EAPC (%)	B-Coefficient (95% CI)	P-value
	2010	2019			
NHL	37.31 (31.71, 43.9)	38.3 (31.86, 45.84)	0.26	0.00 (0.001, 0.004)	<0.001
Deaths	2.77 (2.11, 3.72)	2.81 (2.1, 3.8)	-	-	-
ALL	0.42 (0.26, 0.7)	0.41 (0.26, 0.64)	-0.39	-0.00 (-0.005, -0.002)	<0.001
AML	0.9 (0.68, 1.21)	0.91 (0.66, 1.27)	0.08	0.00 (-0.001, 0.002)	0.244
CLL	0.02 (0.02, 0.03)	0.03 (0.02, 0.04)	0.99	0.01 (-0.009, 0.011)	<0.001
CML	0.29 (0.18, 0.44)	0.28 (0.18, 0.41)	-0.48	-0.00 (-0.007, -0.003)	<0.001
HL	0.22 (0.17, 0.29)	0.21 (0.16, 0.28)	-0.35	-0.00 (-0.004, -0.003)	<0.001
NHL	0.92 (0.8, 1.05)	0.98 (0.82, 1.17)	0.65	0.00 (-0.005, 0.008)	<0.001

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Table 3 - Trends for key outcomes across leukaemia and lymphoma subtypes in Australasia in 2010 and 2019 by sex

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
Incidence						
ALL	2.45	0.024 (0.020, 0.028)	<0.001	2.29	0.023 (0.018, 0.027)	<0.001
AML	2.38	0.024 (0.021, 0.026)	<0.001	1.42	0.011 (0.011, 0.017)	<0.001
CLL	2.48	0.025 (0.02, 0.029)	<0.001	2.28	0.023 (0.016, 0.029)	<0.001
CML	0.64	0.006 (-0.005, 0.018)	0.292	2.84	0.022 (0.022, 0.034)	<0.001
HL	0.30	0.003 (0.002, 0.004)	<0.001	0.61	0.005 (0.005, 0.007)	<0.001
NHL	1.14	0.011 (0.008, 0.015)	<0.001	0.70	0.001 (0.001, 0.013)	0.018
Prevalence						
ALL	2.67	0.026 (0.022, 0.031)	<0.001	2.46	0.023 (0.019, 0.029)	<0.001
AML	1.48	0.015 (0.013, 0.017)	<0.001	0.39	0.001 (0.001, 0.007)	0.006
CLL	2.40	0.024 (0.019, 0.029)	<0.001	2.42	0.023 (0.017, 0.031)	<0.001
CML	1.94	0.019 (0.009, 0.029)	<0.001	3.07	0.031 (0.024, 0.036)	<0.001
HL	0.25	0.002 (0.001, 0.004)	<0.001	0.58	0.001 (0.004, 0.007)	<0.001
NHL	1.23	0.012 (0.008, 0.017)	<0.001	0.87	0.001 (0.002, 0.016)	0.016
DALYs						
ALL	-0.56	-0.006 (-0.008, -0.003)	<0.001	-0.08	-0.001 (-0.004, 0.002)	0.589
AML	1.51	0.015 (0.013, 0.017)	<0.001	0.69	0.007 (0.005, 0.009)	<0.001
CLL	1.85	0.018 (0.013, 0.023)	<0.001	1.57	0.016 (0.009, 0.022)	<0.001

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
CML	-1·86	-0·019 (-0·033, -0·004)	0·011	-0·24	-0·002 (-0·01, 0·006)	0·552
HL	-0·53	-0·005 (-0·006, -0·004)	<0·001	-0·23	-0·003 (-0·004, -0·001)	<0·001
NHL	0·34	0·003 (-0·001, 0·008)	0·165	-0·04	0 (-0·001, 0·007)	0·915
Deaths						
ALL	0·64	0·006 (0·005, 0·008)	<0·001	0·90	0·009 (0·006, 0·012)	<0·001
AML	2·55	0·025 (0·023, 0·027)	<0·001	1·46	0·013 (0·012, 0·017)	<0·001
CLL	2·40	0·024 (0·020, 0·027)	<0·001	1·66	0·013 (0·011, 0·023)	<0·001
CML	-0·72	-0·007 (-0·02, 0·005)	0·263	0·50	0·003 (-0·003, 0·013)	0·221
HL	-0·08	-0·001 (-0·002, 0·001)	0·358	0·21	0·002 (0·001, 0·004)	0·007
NHL	0·86	0·009 (0·005, 0·012)	<0·001	0·28	0·003 (-0·003, 0·009)	0·345

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Table 4 - Trends for key outcomes across leukaemia and lymphoma subtypes in Oceania in 2010 and 2019 by sex

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
Incidence						
ALL	-0.37	-0.004 (-0.005, -0.003)	<0.001	-0.54	-0.006 (-0.008, -0.003)	<0.001
AML	0.17	0.002 (0.001, 0.003)	<0.001	-0.27	-0.003 (-0.006, 0)	0.072
CLL	0.96	0.01 (0.008, 0.011)	<0.001	1.75	0.015 (0.015, 0.019)	<0.001
CML	-0.61	-0.006 (-0.009, -0.003)	<0.001	-0.58	-0.003 (-0.008, -0.004)	<0.001
HL	-0.19	-0.002 (-0.003, -0.001)	0.005	0.31	0.002 (0.002, 0.004)	<0.001
NHL	0.86	0.009 (0.007, 0.010)	<0.001	1.00	0.01 (0.009, 0.011)	<0.001
Prevalence						
ALL	-0.46	-0.005 (-0.006, -0.003)	<0.001	-0.66	-0.007 (-0.009, -0.004)	<0.001
AML	-0.04	0 (-0.001, 0.001)	0.509	-0.39	-0.004 (-0.007, -0.001)	0.01
CLL	1.54	0.015 (0.013, 0.018)	<0.001	2.32	0.025 (0.02, 0.026)	<0.001
CML	-0.85	-0.009 (-0.011, -0.006)	<0.001	-0.71	-0.007 (-0.009, -0.005)	<0.001
HL	0.47	0.005 (0.001, 0.008)	0.012	1.12	0.01 (0.007, 0.016)	<0.001
NHL	2.60	0.026 (0.021, 0.031)	<0.001	3.73	0.03 (0.021, 0.053)	<0.001
DALYs						
ALL	-0.49	-0.005 (-0.006, -0.004)	<0.001	-0.64	-0.006 (-0.009, -0.004)	<0.001
AML	0.06	0.001 (0, 0.001)	0.136	-0.37	-0.004 (-0.007, -0.001)	0.014
CLL	0.64	0.006 (0.005, 0.008)	<0.001	1.19	0.012 (0.01, 0.013)	<0.001

Measure	Males			Females		
	EAPC (%)	B-coefficient (95%CI)	P-value	EAPC (%)	B-coefficient (95%CI)	P-value
CML	-0.95	-0.01 (-0.012, -0.007)	<0.001	-0.84	-0.001 (-0.001, -0.007)	<0.001
HL	-0.67	-0.007 (-0.008, -0.006)	<0.001	-0.20	-0.003 (-0.003, -0.001)	<0.001
NHL	0.24	0.002 (0.001, 0.004)	0.001	0.29	0.002 (0.002, 0.004)	<0.001
Deaths						
ALL	-0.32	-0.003 (-0.004, -0.002)	<0.001	-0.46	-0.003 (-0.007, -0.003)	<0.001
AML	0.27	0.003 (0.002, 0.004)	<0.001	-0.19	-0.001 (-0.005, 0.001)	0.175
CLL	0.62	0.006 (0.005, 0.007)	<0.001	1.33	0.012 (0.012, 0.015)	<0.001
CML	-0.48	-0.005 (-0.008, -0.002)	0.001	-0.50	-0.003 (-0.007, -0.003)	<0.001
HL	-0.45	-0.004 (-0.006, -0.003)	<0.001	-0.06	-0.001 (-0.001, 0)	0.08
NHL	0.62	0.006 (0.005, 0.008)	<0.001	0.70	0.006 (0.006, 0.008)	<0.001

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma


Figures (attached in separate files)

Figure 1 – Age-specific rates at regional level by sex in Australasia and Oceania in 2010 and 2019

Figure 2 - Trends of disease burden based on age-standardised rates in Australasia and Oceania from 2010 to 2019


For peer review only

Figure 1. Age-specific rates at regional level by sex in Australasia and Oceania in 2010 and 2019



This figure represents age-specific rates (per 100,000 population) of incidence, prevalence, disability-adjusted life years (DALYs), and deaths in Australasia (a, b, c, d, e, f, g, h), and in Oceania (i, j, k, l, m, n, o, p) by sex in 2010 and 2019, respectively. ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Figure 2 – Trends of disease burden based on age-standardised rates in Australasia and Oceania from 2010 to 2019



*: statistically significant at 5%; ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Appendices

Appendix A

“The International Classification of Diseases (ICD) definition of leukaemias and lymphomas used in the GBD 2019 study has been defined previously ¹². ICD codes mapped to the GBD cause list for leukaemia or lymphoma incidence data are as follows: AML (C92.0–C92.02, C92.3–C92.62, C93.0–C93.02, C94.0–C94.02, C94.2–C94.22), ALL (C91.0–C91.02), CML (C92.1–C92.12), CLL (C91.1–C91.12), HL (C81–C81.49, C81.7–C81.79, C81.9–C81.99, Z85.71–Z85.72), and NHL (C82–C85.29, C85.7–C86.6, C96–C96.9) ¹². ICD codes mapped to the GBD cause list for leukaemia or lymphoma mortality data are as follows: AML (C92.0, C92.3–C92.6, C93.0, C94.0, C94.2, C94.4–C94.5), ALL (C91.0), CML (C92.1), CLL (C91.1), HL (C81–C81.9), and NHL (C82–C86.6, C96–C96.9) ¹².

References

1. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020;396(10258):1204-22. doi: 10.1016/s0140-6736(20)30925-9

2. Global Burden of Disease Cancer Collaboration. Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life Years for 29 Cancer Groups From 2010 to 2019: A Systematic Analysis for the Global Burden of Disease Study 2019. *JAMA Oncology* 2022;8(3):420-44. doi: 10.1001/jamaoncol.2021.6987

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

Appendix B

Table B1 - Age-specific rates in Australasia and Oceania by age groups and by sex (2010)

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
AUSTRALASIA																		
Incidence																		
<5 years	3·68	3·49	3·88	0·64	0·65	0·62	0·00	0·00	0·00	0·10	0·09	0·11	0·22	0·21	0·23	0·00	0·00	0·00
5-9 years	3·19	3·64	2·71	0·36	0·30	0·42	0·00	0·00	0·00	0·05	0·03	0·07	0·36	0·41	0·31	1·26	1·85	0·64
10-14 years	1·50	1·64	1·34	0·43	0·38	0·49	0·00	0·00	0·00	0·06	0·05	0·07	0·72	0·85	0·59	1·23	1·61	0·83
15-19 years	1·93	2·49	1·33	0·94	0·94	0·94	0·00	0·00	0·00	0·12	0·10	0·15	2·66	2·49	2·83	2·47	3·06	1·85
20-24 years	1·25	1·58	0·90	0·89	0·83	0·95	0·11	0·09	0·13	0·25	0·20	0·31	4·60	4·13	5·09	2·02	2·41	1·62
25-29 years	0·92	0·98	0·86	0·93	0·92	0·93	0·10	0·09	0·11	0·38	0·34	0·42	4·79	4·73	4·85	2·09	2·55	1·63
30-34 years	0·63	0·74	0·52	0·80	0·79	0·82	0·10	0·08	0·11	0·35	0·34	0·35	3·71	4·02	3·42	2·66	3·49	1·86
35-39 years	0·58	0·64	0·54	1·01	0·94	1·07	0·17	0·25	0·10	0·53	0·45	0·61	3·01	3·60	2·45	3·78	4·80	2·80
40-44 years	0·63	0·65	0·62	1·26	1·15	1·36	0·38	0·56	0·21	0·58	0·48	0·67	2·68	3·37	2·03	3·57	4·55	2·64
45-49 years	0·90	0·83	0·98	1·77	1·68	1·86	1·04	1·55	0·55	0·67	0·55	0·78	2·19	2·59	1·80	6·07	8·01	4·20
50-54 years	1·55	1·43	1·68	2·69	2·66	2·73	2·65	4·08	1·26	0·97	0·70	1·23	2·14	2·65	1·66	10·58	13·36	7·90
55-59 years	2·46	2·28	2·63	3·96	3·98	3·95	5·46	7·88	3·11	1·41	0·90	1·90	2·34	2·90	1·79	18·17	23·93	12·60
60-64 years	3·89	3·57	4·21	6·47	7·10	5·86	9·24	13·38	5·16	1·91	1·44	2·38	2·64	2·89	2·39	27·29	34·81	19·86
65-69 years	5·54	5·70	5·37	10·74	12·57	8·95	13·75	18·98	8·64	2·78	2·18	3·36	3·03	3·49	2·58	46·37	59·69	33·34
70-74 years	7·57	8·71	6·50	17·29	20·26	14·51	22·80	32·32	13·91	5·80	4·16	7·32	3·31	3·76	2·88	74·78	95·43	55·52
75-79 years	7·25	9·05	5·71	21·47	26·59	17·06	27·79	39·45	17·74	7·75	5·91	9·33	4·04	4·31	3·81	113·53	136·31	93·89
80-84 years	6·55	8·05	5·43	26·59	31·89	22·61	22·93	32·71	15·58	12·47	9·85	14·44	4·91	5·63	4·36	125·51	153·47	104·50
85+ years	6·51	7·83	5·82	34·60	41·17	31·19	35·57	49·93	28·11	20·89	17·36	22·72	7·05	8·57	6·25	168·07	211·16	145·67
Prevalence																		
<5 years	30·07	28·42	31·82	1·66	1·37	1·97	0·00	0·00	0·00	0·52	0·40	0·64	1·95	1·85	2·05	0·00	0·00	0·00

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
5-9 years	24·86	28·16	21·38	0·85	0·58	1·15	0·00	0·00	0·00	0·28	0·14	0·42	3·21	3·64	2·75	5·13	9·99	0·00
10-14 years	10·64	11·41	9·82	0·91	0·64	1·19	0·00	0·00	0·00	0·31	0·20	0·42	6·39	7·48	5·23	4·62	9·00	0·00
15-19 years	12·97	16·52	9·24	2·93	2·48	3·40	0·00	0·00	0·00	0·65	0·43	0·87	3·51	22·07	25·02	17·21	21·49	12·71
20-24 years	8·64	10·57	6·66	2·26	1·75	2·78	0·96	0·80	1·12	1·30	0·83	1·79	0·96	36·68	45·35	12·50	15·87	9·04
25-29 years	6·41	6·59	6·22	1·95	1·55	2·34	0·86	0·75	0·98	1·84	1·30	2·38	2·25	41·62	42·87	13·95	17·06	10·88
30-34 years	4·11	4·67	3·56	1·37	1·04	1·70	0·81	0·70	0·92	1·59	1·26	1·91	2·47	35·09	29·92	18·27	23·69	13·03
35-39 years	3·93	4·06	3·81	1·56	1·17	1·93	1·43	2·09	0·81	2·48	1·59	3·33	6·15	31·21	21·33	26·28	33·18	19·71
40-44 years	4·43	4·27	4·59	1·74	1·27	2·19	3·08	4·46	1·76	2·62	1·60	3·58	3·13	29·05	17·48	21·29	26·71	16·13
45-49 years	6·48	5·61	7·31	2·68	2·08	3·26	8·48	12·55	4·54	3·11	1·95	4·23	8·50	21·84	15·28	35·73	47·17	24·68
50-54 years	11·88	10·19	13·51	4·02	3·16	4·85	21·44	32·72	10·52	4·55	2·45	6·58	7·99	22·05	14·06	66·28	86·34	46·89
55-59 years	19·16	17·02	21·23	4·89	3·87	5·88	44·09	63·13	25·66	6·42	2·50	10·23	8·72	23·11	14·47	112·26	153·67	72·19
60-64 years	29·74	27·06	32·39	6·79	6·51	7·07	72·83	104·29	41·72	8·59	3·85	13·27	0·48	22·25	18·73	176·67	228·63	125·30
65-69 years	42·41	43·57	41·28	10·17	10·88	9·48	104·63	141·82	68·29	12·02	5·29	18·60	1·05	24·11	18·06	303·28	387·86	220·62
70-74 years	58·96	65·83	52·55	14·38	15·72	13·12	164·89	229·69	104·43	25·72	9·60	40·76	8·43	20·60	16·40	480·04	609·76	359·01
75-79 years	50·48	60·87	41·53	16·85	19·34	14·70	186·37	256·10	126·25	33·69	14·00	50·67	0·50	20·12	20·83	704·29	825·50	599·80
80-84 years	35·88	44·74	29·22	25·62	30·12	22·24	103·78	139·63	76·85	52·18	22·05	74·81	4·65	26·60	23·19	546·69	653·16	466·71
85+ years	28·29	34·45	25·08	27·22	29·73	25·92	107·51	137·41	91·97	76·75	35·62	98·13	9·74	31·23	28·97	584·51	728·80	509·52
DALYs																		
<5 years	33·57	34·38	32·72	32·33	36·71	27·71	0·00	0·00	0·00	2·28	3·10	1·41	0·79	0·79	0·79	9·81	11·49	8·04
5-9 years	41·77	50·96	32·08	17·54	16·11	19·05	0·00	0·00	0·00	0·97	1·08	0·86	1·31	1·53	1·07	13·04	19·17	6·59
10-14 years	29·70	34·46	24·68	20·93	20·13	21·78	0·00	0·00	0·00	1·35	1·70	0·97	2·75	3·45	2·01	12·63	16·52	8·52
15-19 years	44·58	60·09	28·25	35·59	38·33	32·69	0·00	0·00	0·00	1·94	2·42	1·45	7·98	7·62	8·36	25·11	31·09	18·82
20-24 years	26·17	36·41	15·66	34·85	35·54	34·14	0·80	0·78	0·83	4·06	4·95	3·14	3·92	13·00	14·86	32·22	38·55	25·74
25-29 years	18·70	21·78	15·67	36·99	40·25	33·77	0·71	0·73	0·70	6·53	8·72	4·36	8·66	19·00	18·33	31·77	38·63	24·98
30-34 years	12·69	16·50	9·00	32·40	35·05	29·84	0·69	0·71	0·69	6·29	8·97	3·70	5·58	17·69	13·53	38·10	49·78	26·81

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
35-39 years	10·59	12·98	8·32	38·59	39·47	37·75	1·32	2·12	0·56	8·50	11·08	6·05	13·96	17·13	10·95	50·22	63·84	37·25
40-44 years	9·82	11·54	8·19	45·96	46·23	45·70	2·81	4·46	1·24	8·92	11·41	6·54	13·31	16·79	9·99	65·38	83·14	48·45
45-49 years	10·29	11·39	9·23	56·19	58·20	54·25	6·83	10·99	2·81	8·87	11·34	6·49	11·62	14·15	9·18	100·43	132·28	69·66
50-54 years	13·20	14·47	11·99	76·51	82·43	70·79	15·87	26·15	5·92	10·92	12·82	9·09	10·92	13·72	8·21	156·47	196·83	117·43
55-59 years	14·77	16·26	13·34	103·10	112·22	94·27	28·96	45·36	13·09	15·02	18·12	12·02	14·32	17·88	10·87	236·72	310·98	164·85
60-64 years	16·92	18·47	15·39	153·16	177·90	128·70	50·95	78·81	23·42	18·94	25·61	12·34	16·51	18·83	14·22	323·60	412·20	236·00
65-69 years	19·95	22·65	17·33	223·90	273·52	175·40	77·41	114·56	41·10	25·37	36·43	14·56	11·87	25·61	18·22	468·61	603·00	337·26
70-74 years	24·34	29·50	19·53	311·90	378·56	249·71	128·63	194·23	67·42	41·61	58·00	26·33	16·91	30·38	23·68	632·89	808·40	469·13
75-79 years	24·21	30·78	18·55	323·26	414·37	244·72	150·67	225·50	86·17	46·15	65·79	29·22	18·46	32·00	25·41	781·55	940·58	644·47
80-84 years	23·76	29·38	19·53	305·46	378·02	250·97	172·03	253·91	110·53	56·89	80·97	38·81	16·56	30·73	23·43	866·29	1063·97	717·82
85+ years	21·99	26·97	19·40	247·01	336·78	200·36	225·66	332·13	170·32	69·92	106·41	50·95	12·14	28·86	18·64	835·96	1072·56	712·99
Deaths																		
<5 years	0·36	0·37	0·35	0·37	0·42	0·32	0·00	0·00	0·00	0·03	0·04	0·02	0·01	0·01	0·01	0·11	0·13	0·09
5-9 years	0·49	0·60	0·37	0·21	0·20	0·23	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·02	0·01	0·15	0·22	0·08
10-14 years	0·38	0·44	0·31	0·27	0·26	0·28	0·00	0·00	0·00	0·02	0·02	0·01	0·03	0·04	0·02	0·16	0·20	0·11
15-19 years	0·61	0·82	0·38	0·49	0·53	0·45	0·00	0·00	0·00	0·03	0·03	0·02	0·09	0·09	0·09	0·33	0·41	0·25
20-24 years	0·38	0·53	0·23	0·52	0·53	0·51	0·01	0·01	0·01	0·06	0·07	0·05	0·17	0·16	0·18	0·47	0·56	0·37
25-29 years	0·29	0·34	0·24	0·60	0·65	0·54	0·01	0·01	0·01	0·10	0·14	0·07	0·26	0·26	0·26	0·49	0·60	0·39
30-34 years	0·22	0·28	0·15	0·57	0·62	0·52	0·01	0·01	0·01	0·11	0·16	0·06	0·24	0·27	0·21	0·64	0·84	0·45
35-39 years	0·20	0·24	0·15	0·74	0·76	0·73	0·02	0·04	0·01	0·16	0·21	0·11	0·24	0·29	0·19	0·92	1·17	0·69
40-44 years	0·20	0·24	0·17	0·98	0·98	0·97	0·05	0·09	0·02	0·19	0·24	0·13	0·25	0·32	0·19	1·35	1·71	1·00
45-49 years	0·23	0·26	0·20	1·33	1·38	1·28	0·14	0·23	0·06	0·21	0·27	0·15	0·25	0·30	0·20	2·30	3·02	1·60
50-54 years	0·33	0·36	0·29	2·04	2·20	1·88	0·37	0·61	0·13	0·28	0·34	0·23	0·26	0·32	0·19	4·01	5·05	3·02
55-59 years	0·40	0·45	0·35	3·15	3·43	2·87	0·75	1·19	0·33	0·45	0·55	0·34	0·40	0·49	0·30	6·88	9·03	4·80
60-64 years	0·51	0·57	0·44	5·45	6·33	4·57	1·56	2·43	0·70	0·65	0·90	0·40	0·53	0·61	0·46	10·91	13·88	7·98

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
65-69 years	0·69	0·80	0·58	9·48	11·59	7·42	2·81	4·18	1·46	1·04	1·53	0·56	0·85	0·99	0·70	18·56	23·81	13·43
70-74 years	1·00	1·23	0·78	16·14	19·60	12·92	5·69	8·63	2·94	2·05	2·97	1·20	1·31	1·47	1·16	29·99	38·15	22·38
75-79 years	1·26	1·63	0·95	21·14	27·10	16·00	8·36	12·56	4·73	2·84	4·24	1·64	1·72	1·95	1·52	45·52	54·53	37·75
80-84 years	1·66	2·08	1·35	25·91	32·06	21·29	13·45	19·88	8·62	4·45	6·71	2·74	1·96	2·26	1·73	67·50	82·33	56·36
85+ years	2·24	2·76	1·96	29·33	39·43	24·08	25·75	37·40	19·69	7·63	12·29	5·21	1·98	2·63	1·64	90·34	113·20	78·46
OCEANIA																		
Incidence																		
<5 years	1·18	1·31	1·03	1·42	1·62	1·20	0·00	0·00	0·00	0·19	0·26	0·10	0·02	0·02	0·02	0·00	0·00	0·00
5-9 years	0·74	0·63	0·86	0·74	0·64	0·85	0·00	0·00	0·00	0·09	0·13	0·04	0·14	0·25	0·03	0·34	0·49	0·17
10-14 years	0·65	0·43	0·90	0·79	0·33	1·30	0·00	0·00	0·00	0·14	0·04	0·25	0·09	0·14	0·02	0·55	0·32	0·81
15-19 years	0·52	0·66	0·36	1·65	2·04	1·22	0·00	0·00	0·00	0·03	0·04	0·03	0·20	0·30	0·08	0·65	0·72	0·57
20-24 years	0·34	0·38	0·30	0·98	1·37	0·56	0·02	0·02	0·03	0·11	0·17	0·04	0·35	0·35	0·35	0·49	0·62	0·35
25-29 years	0·37	0·43	0·30	1·39	1·35	1·44	0·02	0·02	0·03	0·36	0·22	0·51	0·35	0·46	0·23	0·70	0·90	0·49
30-34 years	0·11	0·17	0·05	0·84	1·19	0·50	0·02	0·02	0·02	0·24	0·17	0·31	0·46	0·60	0·32	0·64	0·48	0·80
35-39 years	0·26	0·18	0·34	1·33	1·55	1·12	0·03	0·02	0·03	0·43	0·37	0·49	0·35	0·61	0·09	0·93	1·20	0·67
40-44 years	0·14	0·18	0·09	1·17	1·66	0·65	0·03	0·02	0·04	0·12	0·15	0·07	0·36	0·59	0·11	1·23	1·29	1·17
45-49 years	0·34	0·32	0·36	1·93	1·50	2·41	0·03	0·02	0·04	0·68	0·16	1·23	0·53	0·74	0·31	1·49	1·78	1·19
50-54 years	0·49	0·47	0·52	1·79	1·40	2·22	0·08	0·06	0·10	0·88	0·44	1·37	0·72	1·21	0·17	2·04	2·65	1·36
55-59 years	0·69	0·76	0·61	2·83	3·06	2·57	0·24	0·26	0·22	1·04	1·13	0·93	0·61	0·74	0·45	3·19	3·55	2·78
60-64 years	0·38	0·55	0·19	2·84	3·61	1·96	0·24	0·28	0·20	1·63	0·55	2·87	0·93	0·93	0·94	4·10	3·96	4·27
65-69 years	0·51	0·71	0·28	2·84	3·56	2·01	0·32	0·21	0·44	1·57	1·10	2·10	0·98	1·45	0·43	6·05	7·42	4·49
70-74 years	0·37	0·54	0·19	2·82	3·20	2·42	0·22	0·18	0·27	2·60	1·11	4·19	1·44	1·54	1·32	8·43	9·13	7·67
75-79 years	0·70	0·31	1·08	6·31	6·90	5·73	0·30	0·34	0·27	0·78	1·35	0·23	1·30	1·86	0·75	9·80	7·49	12·05
80-84 years	0·20	0·24	0·17	4·52	6·56	2·84	0·34	0·54	0·17	1·79	0·52	2·84	0·88	1·03	0·76	11·67	16·54	7·63
85+ years	0·62	0·22	0·87	8·55	16·30	3·56	0·23	0·20	0·25	2·19	0·43	3·33	1·04	1·56	0·70	8·75	10·90	7·36

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
Prevalence																		
<5 years	3·46	3·86	3·04	3·68	3·74	3·62	0·00	0·00	0·00	0·44	0·62	0·24	0·09	0·09	0·09	0·00	0·00	0·00
5-9 years	2·08	1·76	2·42	1·89	1·42	2·40	0·00	0·00	0·00	0·20	0·29	0·10	0·52	0·88	0·12	0·00	0·00	0·00
10-14 years	1·50	0·99	2·08	1·72	0·58	3·01	0·00	0·00	0·00	0·27	0·09	0·48	0·27	0·42	0·10	1·00	0·00	2·12
15-19 years	1·85	2·36	1·29	5·18	5·90	4·38	0·00	0·00	0·00	0·09	0·10	0·08	0·91	1·33	0·46	1·22	1·13	1·31
20-24 years	1·04	1·13	0·95	2·40	3·15	1·62	0·14	0·13	0·15	0·24	0·38	0·08	1·49	1·31	1·68	0·11	0·21	0·00
25-29 years	0·97	1·14	0·80	3·00	2·50	3·51	0·11	0·09	0·14	0·71	0·42	1·00	1·09	1·32	0·86	0·28	0·49	0·06
30-34 years	0·22	0·35	0·09	1·30	1·65	0·96	0·07	0·07	0·08	0·40	0·29	0·51	1·16	1·37	0·95	0·27	0·00	0·54
35-39 years	0·49	0·34	0·64	1·89	1·91	1·88	0·12	0·09	0·15	0·66	0·57	0·75	0·72	1·23	0·21	0·62	0·81	0·43
40-44 years	0·23	0·30	0·15	1·35	1·75	0·94	0·12	0·07	0·17	0·16	0·21	0·10	0·66	1·09	0·22	0·23	0·33	0·12
45-49 years	0·62	0·56	0·69	2·72	1·72	3·81	0·12	0·07	0·17	1·00	0·24	1·82	1·00	1·42	0·56	0·25	0·49	0·00
50-54 years	0·91	0·81	1·02	2·53	1·60	3·56	0·33	0·22	0·45	1·27	0·64	1·99	1·28	2·17	0·29	0·35	0·67	0·00
55-59 years	1·19	1·29	1·09	3·15	3·00	3·32	0·96	0·99	0·94	1·43	1·55	1·31	0·85	1·08	0·60	0·49	0·75	0·20
60-64 years	0·58	0·83	0·28	2·67	3·18	2·08	0·83	0·92	0·73	2·12	0·69	3·76	1·09	1·13	1·04	0·41	0·22	0·63
65-69 years	0·64	0·88	0·36	2·50	3·01	1·92	0·91	0·54	1·34	1·83	1·27	2·48	0·92	1·39	0·37	0·87	1·46	0·20
70-74 years	0·45	0·66	0·22	2·28	2·54	2·01	0·52	0·39	0·65	2·70	1·13	4·38	1·11	1·26	0·95	2·48	2·65	2·30
75-79 years	0·72	0·27	1·17	4·82	5·11	4·53	0·53	0·56	0·50	0·71	1·23	0·20	0·90	1·34	0·47	3·98	1·72	6·19
80-84 years	0·15	0·17	0·14	4·44	6·53	2·70	0·37	0·58	0·19	1·45	0·40	2·31	0·57	0·73	0·44	2·72	4·66	1·11
85+ years	0·34	0·16	0·46	6·31	11·67	2·86	0·19	0·16	0·22	1·64	0·29	2·50	0·75	1·26	0·42	1·37	1·89	1·04
DALYs																		
<5 years	72·64	81·01	63·64	69·80	83·78	54·75	0·00	0·00	0·00	11·58	16·52	6·26	0·98	1·21	0·74	33·59	38·44	28·37
5-9 years	44·00	37·52	51·13	34·63	31·94	37·60	0·00	0·00	0·00	5·41	7·94	2·62	6·71	11·90	1·00	10·73	16·18	4·73
10-14 years	39·90	26·46	55·03	36·75	16·80	59·23	0·00	0·00	0·00	8·54	2·73	15·07	3·98	6·71	0·90	16·90	9·26	25·51
15-19 years	23·94	30·52	16·70	61·11	78·40	42·12	0·00	0·00	0·00	1·55	1·73	1·35	6·83	10·71	2·56	20·14	23·20	16·76
20-24 years	16·30	17·90	14·63	37·62	54·42	20·22	0·94	0·95	0·93	5·02	8·21	1·71	2·03	12·80	11·23	23·69	29·93	17·22

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
25-29 years	17·45	20·59	14·29	52·57	53·46	51·68	0·82	0·70	0·94	17·33	10·29	24·42	12·89	17·76	8·00	32·83	42·47	23·15
30-34 years	5·22	8·24	2·24	33·17	48·02	18·51	0·59	0·60	0·58	11·82	8·40	15·20	7·50	23·36	11·72	27·14	20·00	34·20
35-39 years	12·24	8·57	15·93	49·78	60·27	39·24	1·00	0·84	1·16	20·15	17·30	23·01	13·28	23·36	3·16	37·11	47·73	26·44
40-44 years	6·06	8·13	3·88	42·70	62·31	22·07	1·04	0·70	1·40	5·23	6·99	3·37	2·88	21·33	3·99	52·71	54·89	50·42
45-49 years	13·10	12·41	13·84	59·13	48·82	70·29	0·86	0·59	1·16	26·46	6·33	48·26	7·06	23·51	10·07	58·35	69·26	46·52
50-54 years	16·80	16·09	17·59	48·79	40·80	57·72	2·11	1·62	2·65	30·63	15·38	47·66	0·62	34·58	5·02	72·76	94·38	48·61
55-59 years	21·01	23·24	18·46	73·33	82·51	62·89	5·71	6·33	5·01	31·95	34·79	28·72	6·52	19·97	12·59	103·92	115·52	90·72
60-64 years	10·41	15·05	5·09	67·72	88·12	44·30	5·37	6·37	4·23	45·07	15·21	79·36	13·14	22·62	23·73	120·13	116·02	124·85
65-69 years	12·24	17·14	6·64	59·01	75·55	40·08	6·38	4·30	8·77	38·20	26·84	51·21	1·76	32·16	9·86	153·15	187·54	113·80
70-74 years	7·79	11·41	3·93	50·31	57·99	42·13	4·06	3·39	4·78	55·07	23·54	88·64	8·29	29·79	26·71	179·60	195·13	163·07
75-79 years	12·65	5·54	19·57	93·83	104·77	83·20	4·84	5·56	4·14	14·30	24·69	4·19	1·69	30·54	13·09	173·43	135·34	210·45
80-84 years	3·06	3·61	2·60	51·30	75·67	31·08	4·79	7·69	2·39	26·85	7·78	42·67	2·01	13·45	10·81	185·78	262·67	121·95
85+ years	6·46	2·63	8·93	76·36	147·75	30·39	2·63	2·30	2·84	26·76	5·09	40·71	0·23	14·70	7·36	102·42	125·73	87·41
Deaths																		
<5 years	0·84	0·93	0·73	0·80	0·96	0·63	0·00	0·00	0·00	0·13	0·19	0·07	0·01	0·01	0·01	0·39	0·33	0·45
5-9 years	0·53	0·46	0·62	0·42	0·39	0·46	0·00	0·00	0·00	0·07	0·10	0·03	0·08	0·14	0·01	0·13	0·06	0·20
10-14 years	0·52	0·34	0·72	0·48	0·22	0·77	0·00	0·00	0·00	0·11	0·04	0·20	0·05	0·09	0·01	0·22	0·33	0·12
15-19 years	0·33	0·42	0·23	0·85	1·09	0·58	0·00	0·00	0·00	0·02	0·02	0·02	0·09	0·15	0·04	0·28	0·23	0·32
20-24 years	0·24	0·27	0·22	0·56	0·81	0·30	0·01	0·01	0·01	0·07	0·12	0·03	0·18	0·19	0·17	0·35	0·26	0·45
25-29 years	0·28	0·33	0·23	0·85	0·86	0·83	0·01	0·01	0·02	0·28	0·17	0·39	0·21	0·29	0·13	0·53	0·37	0·68
30-34 years	0·09	0·14	0·04	0·58	0·84	0·32	0·01	0·01	0·01	0·21	0·15	0·27	0·31	0·41	0·21	0·48	0·60	0·35
35-39 years	0·24	0·16	0·31	0·96	1·16	0·75	0·02	0·02	0·02	0·39	0·33	0·44	0·26	0·45	0·06	0·71	0·51	0·92
40-44 years	0·13	0·17	0·08	0·91	1·32	0·47	0·02	0·01	0·03	0·11	0·15	0·07	0·27	0·45	0·08	1·12	1·07	1·17
45-49 years	0·31	0·29	0·33	1·40	1·16	1·66	0·02	0·01	0·03	0·63	0·15	1·14	0·40	0·56	0·24	1·38	1·10	1·64
50-54 years	0·45	0·43	0·47	1·30	1·09	1·54	0·06	0·04	0·07	0·82	0·41	1·27	0·55	0·92	0·13	1·94	1·30	2·52

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
55-59 years	0.64	0.71	0.56	2.24	2.52	1.92	0.17	0.19	0.15	0.98	1.06	0.88	0.50	0.61	0.38	3.17	2.77	3.52
60-64 years	0.37	0.53	0.18	2.40	3.13	1.57	0.19	0.22	0.15	1.60	0.54	2.82	0.82	0.80	0.84	4.26	4.43	4.12
65-69 years	0.52	0.72	0.28	2.49	3.18	1.69	0.27	0.18	0.36	1.61	1.13	2.16	0.92	1.35	0.42	6.46	4.81	7.91
70-74 years	0.40	0.59	0.20	2.59	2.98	2.17	0.21	0.17	0.24	2.84	1.21	4.57	1.46	1.53	1.38	9.25	8.41	10.03
75-79 years	0.82	0.36	1.27	6.07	6.78	5.39	0.31	0.36	0.26	0.93	1.60	0.27	1.41	1.98	0.85	11.25	13.67	8.76
80-84 years	0.26	0.30	0.22	4.29	6.33	2.60	0.40	0.64	0.20	2.26	0.65	3.60	1.01	1.13	0.91	15.68	10.31	22.15
85+ years	1.05	0.31	1.52	8.72	17.07	3.34	0.30	0.27	0.32	2.97	0.60	4.50	1.19	1.66	0.88	12.11	10.23	15.03

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Table B2 - Age-specific rates in Australasia and Oceania in 2019 by age groups and by sex

Age groups	ALL			AML			CLL			CML			LIL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
AUSTRALASIA																		
Incidence																		
<5 years	3·41	3·42	3·40	0·49	0·52	0·46	0·00	0·00	0·00	0·08	0·08	0·08	0·23	0·21	0·24	0·00	0·00	0·00
5-9 years	3·34	3·98	2·66	0·32	0·30	0·35	0·00	0·00	0·00	0·05	0·03	0·07	0·34	0·35	0·32	0·97	1·45	0·47
10-14 years	1·59	1·75	1·43	0·39	0·35	0·42	0·00	0·00	0·00	0·06	0·05	0·08	0·69	0·82	0·56	1·02	1·33	0·70
15-19 years	1·74	2·15	1·31	0·73	0·73	0·74	0·00	0·00	0·00	0·10	0·07	0·13	2·43	3·06	2·82	1·92	2·39	1·41
20-24 years	1·26	1·56	0·95	0·77	0·77	0·78	0·11	0·10	0·12	0·22	0·17	0·27	4·55	5·11	4·99	1·49	1·90	1·08
25-29 years	1·02	1·08	0·96	0·84	0·90	0·79	0·10	0·09	0·11	0·35	0·31	0·39	4·99	5·79	5·00	1·93	2·54	1·34
30-34 years	0·71	0·79	0·63	0·74	0·71	0·77	0·10	0·08	0·11	0·33	0·30	0·37	3·77	4·94	3·62	2·51	3·23	1·81
35-39 years	0·73	0·77	0·70	0·90	0·81	1·00	0·19	0·27	0·11	0·51	0·39	0·62	3·03	4·57	2·52	3·58	4·65	2·53
40-44 years	0·74	0·70	0·79	1·16	1·04	1·28	0·41	0·58	0·24	0·55	0·40	0·69	2·78	4·46	2·12	3·44	4·48	2·44
45-49 years	1·15	1·01	1·29	1·65	1·53	1·77	1·13	1·67	0·62	0·63	0·48	0·77	2·26	4·62	1·92	5·55	7·13	4·04
50-54 years	1·90	1·64	2·15	2·74	2·65	2·82	2·68	3·94	1·48	0·88	0·57	1·18	2·19	4·69	1·71	9·66	12·13	7·31
55-59 years	3·15	2·75	3·53	3·87	3·78	3·96	5·66	7·64	3·78	1·42	0·72	2·08	2·39	4·98	1·82	16·45	21·25	11·89
60-64 years	5·14	4·80	5·45	6·42	7·27	5·62	9·58	13·73	5·68	1·83	1·13	2·49	2·75	4·06	2·45	26·27	34·77	18·29
65-69 years	7·18	7·47	6·91	10·43	12·24	8·74	14·40	19·77	9·36	2·63	1·59	3·60	2·93	4·30	2·63	42·41	54·14	31·43
70-74 years	8·86	10·29	7·52	17·73	21·37	14·32	22·50	31·04	14·47	5·00	2·88	6·98	3·16	4·44	2·89	67·82	86·44	50·32
75-79 years	8·54	10·31	6·96	24·48	30·33	19·21	31·40	42·78	21·16	8·06	5·16	10·67	4·08	4·27	3·91	108·54	131·59	87·78
80-84 years	7·57	9·62	5·89	28·81	36·06	22·89	26·29	36·47	17·98	13·42	9·24	16·83	5·11	4·86	4·62	122·55	148·57	101·32
85+ years	7·24	8·79	6·27	39·29	47·14	34·42	41·49	55·39	32·88	25·02	17·88	29·44	7·92	4·45	6·97	176·74	215·23	152·88
Prevalence																		
<5 years	28·18	28·22	28·13	1·18	1·02	1·36	0·00	0·00	0·00	0·43	0·36	0·50	2·01	1·91	2·12	0·00	0·00	0·00
5-9 years	26·60	31·56	21·38	0·78	0·60	0·96	0·00	0·00	0·00	0·29	0·15	0·44	2·98	2·12	2·84	0·00	0·00	0·00

Age groups	ALL			AML			CLL			CML			ALL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
10-14 years	11·66	12·58	10·68	0·75	0·54	0·97	0·00	0·00	0·00	0·32	0·19	0·46	6·14	6·20	5·02	0·00	0·00	0·00
15-19 years	12·25	15·12	9·24	2·23	1·89	2·60	0·00	0·00	0·00	0·55	0·35	0·76	21·58	11·33	24·98	9·01	17·19	0·43
20-24 years	9·10	10·95	7·20	1·85	1·50	2·21	0·94	0·85	1·04	1·19	0·77	1·63	40·53	33·58	44·59	6·44	12·69	0·00
25-29 years	7·38	7·61	7·17	1·68	1·43	1·91	0·86	0·78	0·95	1·78	1·28	2·28	43·29	44·20	44·36	9·22	16·56	1·99
30-34 years	4·84	5·27	4·44	1·18	0·88	1·47	0·83	0·70	0·96	1·64	1·18	2·08	33·17	33·55	31·84	17·29	22·21	12·58
35-39 years	5·16	5·22	5·11	1·29	0·90	1·67	1·59	2·26	0·94	2·53	1·52	3·52	26·46	33·06	21·99	25·19	32·21	18·36
40-44 years	5·46	4·87	6·02	1·47	1·04	1·87	3·31	4·65	2·02	2·69	1·49	3·85	24·11	22·98	18·44	20·56	26·52	14·82
45-49 years	8·56	7·15	9·90	2·31	1·67	2·91	9·30	13·62	5·20	3·13	1·84	4·34	19·32	22·37	16·42	33·54	42·47	25·07
50-54 years	14·93	12·12	17·60	3·81	2·85	4·71	21·81	31·76	12·35	4·43	2·20	6·55	18·48	22·53	14·63	57·00	70·32	44·32
55-59 years	24·96	20·99	28·72	4·73	3·69	5·72	45·98	61·47	31·28	7·08	2·28	11·63	19·33	22·97	14·90	103·35	139·06	69·49
60-64 years	39·62	36·89	42·19	6·63	6·46	6·80	75·96	107·63	46·21	9·10	3·45	14·41	21·62	22·89	19·49	172·89	231·60	117·75
65-69 years	55·54	57·82	53·41	9·62	10·15	9·12	110·60	149·02	74·65	12·87	4·51	20·69	20·99	22·26	18·86	282·65	357·97	212·18
70-74 years	69·60	78·51	61·24	14·64	16·52	12·88	164·84	223·50	109·76	24·59	7·93	40·24	18·62	17·86	17·45	447·32	566·06	335·79
75-79 years	59·89	69·96	50·81	19·02	22·01	16·33	214·12	282·82	152·26	38·46	14·32	60·20	22·45	21·68	22·24	691·00	821·62	573·39
80-84 years	42·02	54·42	31·90	27·69	33·99	22·55	124·12	163·07	92·34	60·31	23·98	89·96	28·12	31·14	26·48	560·72	665·53	475·20
85+ years	30·93	38·08	26·50	30·38	34·02	28·12	126·15	154·80	108·40	95·23	41·63	128·46	33·91	33·65	32·84	620·56	753·81	537·97
DALYs																		
<5 years	26·87	28·83	24·80	26·12	30·77	21·22	0·00	0·00	0·00	1·63	2·34	0·88	0·76	0·77	0·75	7·22	9·14	5·19
5-9 years	37·00	46·31	27·21	15·79	15·74	15·85	0·00	0·00	0·00	0·83	0·89	0·76	1·11	1·28	1·03	9·94	14·92	4·70
10-14 years	28·97	33·68	24·01	19·49	19·50	19·49	0·00	0·00	0·00	1·08	1·33	0·81	2·33	2·93	1·78	10·21	13·34	6·90
15-19 years	36·05	45·15	26·51	28·09	30·08	26·00	0·00	0·00	0·00	1·31	1·58	1·03	6·75	6·90	7·65	18·97	23·74	13·96
20-24 years	23·51	31·94	14·82	31·04	33·65	28·35	0·76	0·77	0·75	3·06	3·83	2·26	12·74	12·97	13·54	23·07	29·63	16·31
25-29 years	18·38	21·55	15·25	34·34	40·03	28·74	0·68	0·72	0·64	5·14	6·96	3·34	17·48	17·98	16·99	28·49	37·48	19·66
30-34 years	12·86	15·84	10·01	30·41	32·27	28·63	0·66	0·67	0·66	4·96	6·82	3·18	14·22	14·77	12·74	34·80	45·01	25·01
35-39 years	11·82	13·86	9·85	35·36	34·92	35·78	1·37	2·15	0·60	6·86	8·74	5·04	12·68	12·45	9·99	46·07	60·10	32·43

Age groups	ALL			AML			CLL			CML			HLL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
40-44 years	10·03	10·85	9·23	43·59	43·25	43·92	2·81	4·36	1·31	6·95	8·52	5·43	12·56	11·85	9·38	60·79	79·45	42·78
45-49 years	11·33	12·07	10·62	53·23	54·28	52·23	6·92	11·10	2·95	6·88	8·61	5·24	10·79	11·83	8·86	88·79	114·51	64·39
50-54 years	13·79	14·55	13·07	78·87	84·07	73·93	14·91	23·87	6·38	8·18	9·20	7·22	10·30	11·93	7·80	138·78	174·67	104·62
55-59 years	16·47	17·17	15·80	100·63	106·91	94·68	27·89	41·71	14·79	11·89	12·96	10·88	13·74	15·39	10·28	208·42	269·62	150·37
60-64 years	19·48	21·18	17·88	152·58	184·02	123·06	49·79	77·19	24·06	14·28	18·02	10·77	16·02	18·61	13·59	303·09	402·15	210·04
65-69 years	23·22	26·09	20·53	219·54	270·80	171·58	76·22	113·23	41·58	18·23	23·92	12·91	20·21	22·11	17·49	417·20	534·39	307·54
70-74 years	26·37	31·64	21·41	321·35	401·70	245·89	120·42	178·24	66·12	28·22	35·48	21·40	24·19	27·31	22·20	558·49	714·78	411·71
75-79 years	27·39	33·23	22·12	370·48	475·03	276·34	163·00	234·27	98·82	39·59	51·78	28·63	26·66	27·68	24·84	728·74	888·41	584·96
80-84 years	26·80	34·20	20·76	333·08	430·72	253·40	190·92	274·71	122·54	51·62	68·15	38·13	25·87	27·58	22·85	826·89	1009·26	678·08
85+ years	24·19	29·81	20·71	267·36	366·38	205·98	253·23	355·58	189·78	70·86	96·23	55·14	21·95	22·24	18·05	838·42	1046·68	709·32
Deaths																		
<5 years	0·28	0·31	0·26	0·30	0·35	0·24	0·00	0·00	0·00	0·02	0·03	0·01	0·00	0·01	0·01	0·08	0·11	0·06
5-9 years	0·43	0·54	0·31	0·19	0·19	0·19	0·00	0·00	0·00	0·01	0·01	0·01	0·00	0·01	0·01	0·11	0·17	0·05
10-14 years	0·37	0·43	0·30	0·25	0·25	0·25	0·00	0·00	0·00	0·01	0·02	0·01	0·03	0·03	0·02	0·13	0·16	0·08
15-19 years	0·49	0·61	0·36	0·39	0·42	0·36	0·00	0·00	0·00	0·02	0·02	0·01	0·08	0·07	0·09	0·25	0·31	0·18
20-24 years	0·34	0·46	0·21	0·46	0·50	0·42	0·01	0·01	0·01	0·04	0·06	0·03	0·15	0·15	0·16	0·33	0·43	0·24
25-29 years	0·29	0·34	0·24	0·55	0·65	0·46	0·01	0·01	0·01	0·08	0·11	0·05	0·24	0·25	0·23	0·44	0·58	0·31
30-34 years	0·22	0·27	0·17	0·53	0·57	0·50	0·01	0·01	0·01	0·09	0·12	0·05	0·22	0·24	0·19	0·59	0·76	0·42
35-39 years	0·22	0·26	0·18	0·68	0·67	0·69	0·02	0·04	0·01	0·13	0·17	0·09	0·22	0·26	0·17	0·84	1·10	0·60
40-44 years	0·20	0·22	0·19	0·93	0·92	0·93	0·05	0·08	0·02	0·14	0·18	0·11	0·23	0·30	0·18	1·25	1·64	0·88
45-49 years	0·25	0·27	0·23	1·26	1·29	1·24	0·15	0·23	0·06	0·16	0·20	0·12	0·23	0·27	0·19	2·03	2·61	1·47
50-54 years	0·34	0·36	0·31	2·10	2·25	1·97	0·34	0·56	0·14	0·21	0·24	0·18	0·24	0·30	0·18	3·55	4·46	2·68
55-59 years	0·44	0·47	0·41	3·07	3·27	2·88	0·72	1·09	0·37	0·35	0·39	0·31	0·38	0·48	0·28	6·05	7·81	4·37
60-64 years	0·57	0·64	0·50	5·43	6·55	4·37	1·50	2·35	0·71	0·49	0·63	0·35	0·51	0·59	0·43	10·19	13·50	7·08
65-69 years	0·78	0·90	0·67	9·30	11·47	7·26	2·74	4·11	1·45	0·73	1·00	0·48	0·78	0·89	0·67	16·47	21·03	12·20

Age groups	ALL			AML			CLL			CML			HLL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
70-74 years	1·06	1·30	0·84	16·64	20·80	12·73	5·27	7·84	2·86	1·36	1·81	0·94	1·11	1·27	1·07	26·36	33·61	19·55
75-79 years	1·43	1·77	1·12	24·27	31·13	18·08	8·93	12·88	5·38	2·39	3·31	1·55	1·59	1·72	1·47	42·20	51·22	34·09
80-84 years	1·86	2·41	1·42	28·28	36·59	21·51	14·82	21·39	9·46	3·94	5·61	2·58	1·88	1·13	1·64	64·02	77·68	52·88
85+ years	2·52	3·13	2·14	32·43	43·74	25·42	29·48	40·86	22·42	7·80	11·30	5·64	1·91	1·49	1·54	92·20	112·39	79·68
OCEANIA																		
Incidence																		
<5 years	1·06	1·19	0·92	1·36	1·55	1·16	0·00	0·00	0·00	0·15	0·20	0·08	0·02	0·02	0·02	0·00	0·00	0·00
5-9 years	0·73	0·64	0·83	0·74	0·67	0·81	0·00	0·00	0·00	0·08	0·12	0·04	0·13	0·23	0·03	0·34	0·49	0·18
10-14 years	0·68	0·45	0·93	0·79	0·34	1·28	0·00	0·00	0·00	0·13	0·04	0·22	0·08	0·14	0·03	0·62	0·35	0·92
15-19 years	0·54	0·68	0·38	1·71	2·12	1·26	0·00	0·00	0·00	0·03	0·04	0·03	0·22	0·30	0·10	0·75	0·81	0·68
20-24 years	0·36	0·41	0·30	1·02	1·46	0·56	0·03	0·03	0·03	0·10	0·17	0·03	0·33	0·36	0·37	0·55	0·69	0·40
25-29 years	0·34	0·40	0·29	1·34	1·34	1·34	0·02	0·02	0·03	0·31	0·19	0·43	0·33	0·44	0·23	0·73	0·95	0·51
30-34 years	0·11	0·17	0·04	0·82	1·18	0·47	0·02	0·02	0·02	0·21	0·16	0·27	0·44	0·57	0·32	0·64	0·49	0·79
35-39 years	0·26	0·18	0·33	1·33	1·58	1·08	0·03	0·02	0·04	0·39	0·34	0·43	0·33	0·58	0·09	0·96	1·24	0·69
40-44 years	0·13	0·18	0·08	1·10	1·62	0·58	0·03	0·02	0·04	0·10	0·14	0·06	0·34	0·56	0·11	1·20	1·26	1·14
45-49 years	0·32	0·29	0·35	1·78	1·40	2·19	0·03	0·02	0·04	0·60	0·15	1·08	0·50	0·70	0·29	1·47	1·71	1·22
50-54 years	0·46	0·45	0·48	1·67	1·36	2·01	0·08	0·06	0·11	0·79	0·40	1·21	0·66	0·12	0·16	2·01	2·58	1·38
55-59 years	0·66	0·72	0·58	2·75	3·02	2·46	0·24	0·24	0·24	0·94	1·00	0·87	0·57	0·71	0·42	3·25	3·61	2·84
60-64 years	0·38	0·54	0·19	2·79	3·56	1·91	0·25	0·27	0·22	1·49	0·52	2·60	0·90	0·89	0·90	4·28	4·09	4·50
65-69 years	0·51	0·70	0·29	2·77	3·48	1·96	0·34	0·21	0·49	1·46	1·00	1·99	0·93	0·39	0·41	6·33	7·77	4·67
70-74 years	0·37	0·54	0·19	2·72	3·14	2·28	0·23	0·18	0·28	2·36	1·00	3·80	1·35	0·46	1·22	8·37	9·13	7·57
75-79 years	0·68	0·31	1·04	6·12	6·85	5·41	0·31	0·34	0·27	0·73	1·24	0·22	1·21	0·74	0·70	10·14	7·64	12·60
80-84 years	0·22	0·26	0·18	4·97	7·20	3·01	0·38	0·61	0·18	1·67	0·52	2·69	0·85	0·00	0·72	13·29	18·01	9·12
85+ years	0·70	0·24	1·02	9·71	18·32	3·70	0·26	0·23	0·28	1·98	0·43	3·07	1·06	0·60	0·69	10·92	13·30	9·26

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Age groups	ALL			AML			CLL			CML			HLL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
Prevalence																		
<5 years	3·12	3·51	2·71	3·52	3·56	3·48	0·00	0·00	0·00	0·34	0·48	0·19	0·09	0·08	0·10	0·00	0·00	0·00
5-9 years	2·06	1·79	2·35	1·87	1·47	2·30	0·00	0·00	0·00	0·18	0·26	0·09	0·53	0·89	0·14	0·00	0·00	0·00
10-14 years	1·57	1·04	2·16	1·72	0·60	2·95	0·00	0·00	0·00	0·25	0·08	0·44	0·30	0·46	0·13	1·60	0·00	3·37
15-19 years	1·91	2·41	1·35	5·36	6·13	4·52	0·00	0·00	0·00	0·09	0·09	0·08	1·02	0·44	0·56	1·65	1·41	1·91
20-24 years	1·10	1·24	0·96	2·50	3·35	1·59	0·16	0·15	0·17	0·23	0·38	0·07	1·70	0·47	1·94	0·11	0·21	0·00
25-29 years	0·92	1·07	0·77	2·87	2·48	3·27	0·12	0·10	0·15	0·62	0·37	0·86	1·11	0·37	0·92	0·34	0·57	0·12
30-34 years	0·21	0·34	0·09	1·26	1·62	0·89	0·08	0·07	0·09	0·35	0·26	0·45	1·23	0·44	1·03	0·60	0·19	1·01
35-39 years	0·47	0·34	0·61	1·88	1·93	1·83	0·13	0·10	0·17	0·60	0·53	0·67	0·75	0·27	0·22	1·17	1·51	0·83
40-44 years	0·21	0·29	0·14	1·27	1·70	0·84	0·12	0·07	0·18	0·14	0·20	0·09	0·66	0·10	0·22	0·13	0·21	0·06
45-49 years	0·57	0·51	0·63	2·51	1·61	3·46	0·13	0·07	0·19	0·89	0·23	1·59	0·99	0·40	0·56	0·19	0·36	0·00
50-54 years	0·84	0·77	0·92	2·35	1·55	3·22	0·36	0·22	0·51	1·14	0·58	1·75	1·27	0·15	0·29	0·33	0·64	0·00
55-59 years	1·15	1·25	1·03	3·06	2·97	3·17	1·03	0·99	1·07	1·31	1·39	1·22	0·88	0·12	0·60	0·63	0·97	0·23
60-64 years	0·58	0·83	0·31	2·63	3·14	2·04	0·91	0·94	0·88	1·95	0·65	3·43	1·14	0·19	1·08	0·95	0·63	1·31
65-69 years	0·66	0·90	0·39	2·45	2·95	1·89	1·06	0·58	1·62	1·72	1·16	2·37	0·95	0·45	0·37	1·84	2·30	1·30
70-74 years	0·46	0·67	0·23	2·20	2·49	1·90	0·55	0·40	0·71	2·47	1·02	4·00	1·07	0·23	0·90	2·61	2·70	2·50
75-79 years	0·70	0·28	1·13	4·70	5·09	4·31	0·56	0·57	0·54	0·66	1·14	0·19	0·87	0·30	0·45	4·47	1·88	7·02
80-84 years	0·16	0·18	0·15	4·93	7·23	2·89	0·44	0·69	0·21	1·36	0·40	2·21	0·58	0·75	0·44	3·28	5·43	1·38
85+ years	0·39	0·17	0·53	7·14	13·07	3·00	0·23	0·19	0·26	1·49	0·29	2·33	0·86	0·45	0·46	2·50	3·02	2·14
DALYs																		
<5 years	65·28	73·52	56·44	66·76	80·06	52·47	0·00	0·00	0·00	9·03	12·71	5·07	0·88	0·02	0·73	28·87	33·79	23·60
5-9 years	43·79	38·18	49·90	34·72	33·57	35·97	0·00	0·00	0·00	4·77	6·99	2·34	5·88	0·34	1·02	10·18	15·28	4·63
10-14 years	41·60	27·66	57·05	36·87	17·57	58·25	0·00	0·00	0·00	7·83	2·60	13·63	3·74	0·24	0·97	17·67	9·37	26·87
15-19 years	24·75	31·22	17·59	63·59	81·75	43·48	0·00	0·00	0·00	1·60	1·73	1·45	6·78	1·43	2·74	21·83	24·51	18·86
20-24 years	17·23	19·67	14·61	39·69	58·14	19·98	1·04	1·09	0·99	4·90	8·02	1·56	11·94	1·83	10·99	25·15	31·77	18·06

Age groups	ALL			AML			CLL			CML			ALL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
25-29 years	16·52	19·25	13·76	51·00	53·51	48·47	0·85	0·72	0·97	15·11	9·22	21·06	12·01	11·41	7·58	32·18	42·06	22·19
30-34 years	5·10	8·07	2·18	32·35	47·75	17·19	0·60	0·61	0·60	10·46	7·64	13·23	16·19	22·34	11·12	25·93	19·41	32·36
35-39 years	12·07	8·55	15·58	49·73	61·66	37·83	1·03	0·87	1·19	18·23	16·04	20·42	12·22	22·47	2·98	36·30	46·92	25·70
40-44 years	5·84	8·00	3·65	40·47	60·93	19·74	1·02	0·70	1·34	4·68	6·50	2·84	11·83	11·81	3·71	51·69	54·13	49·21
45-49 years	12·36	11·30	13·49	54·55	45·69	63·95	0·87	0·58	1·18	23·67	6·01	42·41	15·57	22·49	9·28	57·90	67·17	48·07
50-54 years	15·77	15·29	16·29	45·61	39·61	52·19	2·08	1·51	2·70	27·37	14·08	41·95	18·55	33·26	4·59	71·52	91·71	49·36
55-59 years	19·96	22·03	17·61	71·28	81·06	60·22	5·45	5·66	5·22	29·00	31·00	26·75	15·13	11·33	11·50	105·11	116·15	92·61
60-64 years	10·25	14·68	5·22	66·33	86·80	43·11	5·29	5·90	4·59	41·20	14·37	71·64	21·63	22·18	22·15	124·00	118·70	130·01
65-69 years	12·15	16·79	6·81	57·68	73·91	38·99	6·61	4·24	9·34	35·49	24·41	48·26	20·32	23·04	9·12	156·91	192·39	116·06
70-74 years	7·79	11·37	4·02	48·49	56·93	39·62	4·05	3·28	4·86	50·08	21·15	80·47	26·22	23·82	24·52	176·00	192·72	158·45
75-79 years	12·22	5·55	18·81	91·08	103·88	78·46	4·79	5·41	4·18	13·25	22·69	3·95	19·95	23·00	12·00	174·11	135·20	212·49
80-84 years	3·21	3·86	2·63	56·40	82·90	33·03	5·42	8·67	2·55	25·08	7·78	40·34	11·28	11·73	10·00	209·09	284·14	142·90
85+ years	7·28	2·83	10·39	86·54	165·84	31·16	2·90	2·58	3·12	24·07	5·04	37·36	9·75	11·00	6·78	122·12	146·93	104·80
Deaths																		
<5 years	0·75	0·85	0·65	0·77	0·92	0·60	0·00	0·00	0·00	0·10	0·15	0·06	0·01	0·01	0·01	0·34	0·39	0·27
5-9 years	0·53	0·46	0·61	0·42	0·41	0·44	0·00	0·00	0·00	0·06	0·09	0·03	0·01	0·13	0·01	0·12	0·18	0·06
10-14 years	0·54	0·36	0·74	0·48	0·23	0·76	0·00	0·00	0·00	0·10	0·03	0·18	0·05	0·08	0·01	0·23	0·12	0·35
15-19 years	0·34	0·43	0·24	0·88	1·14	0·60	0·00	0·00	0·00	0·02	0·02	0·02	0·09	0·14	0·04	0·30	0·34	0·26
20-24 years	0·26	0·29	0·22	0·59	0·87	0·30	0·02	0·02	0·01	0·07	0·12	0·02	0·18	0·19	0·16	0·38	0·47	0·27
25-29 years	0·27	0·31	0·22	0·82	0·86	0·78	0·01	0·01	0·02	0·24	0·15	0·34	0·19	0·26	0·12	0·52	0·68	0·36
30-34 years	0·09	0·14	0·04	0·57	0·84	0·30	0·01	0·01	0·01	0·18	0·13	0·23	0·28	0·37	0·19	0·45	0·34	0·57
35-39 years	0·23	0·16	0·30	0·96	1·19	0·73	0·02	0·02	0·02	0·35	0·31	0·39	0·23	0·41	0·06	0·70	0·90	0·49
40-44 years	0·12	0·17	0·08	0·86	1·29	0·42	0·02	0·01	0·03	0·10	0·14	0·06	0·25	0·42	0·08	1·10	1·15	1·05
45-49 years	0·29	0·27	0·32	1·29	1·08	1·51	0·02	0·01	0·03	0·56	0·14	1·00	0·37	0·51	0·22	1·37	1·59	1·14
50-54 years	0·42	0·41	0·43	1·22	1·06	1·39	0·05	0·04	0·07	0·73	0·38	1·12	0·49	0·83	0·12	1·91	2·45	1·32

Age groups	ALL			AML			CLL			CML			HL			NHL		
	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female
55-59 years	0·61	0·67	0·54	2·17	2·47	1·83	0·16	0·17	0·16	0·89	0·95	0·82	0·46	0·56	0·35	3·21	3·54	2·83
60-64 years	0·36	0·52	0·18	2·35	3·08	1·53	0·18	0·21	0·16	1·46	0·51	2·54	0·77	0·75	0·79	4·40	4·21	4·62
65-69 years	0·51	0·71	0·29	2·43	3·12	1·64	0·27	0·18	0·39	1·50	1·03	2·04	0·86	0·26	0·39	6·62	8·11	4·90
70-74 years	0·40	0·58	0·21	2·49	2·93	2·04	0·21	0·17	0·25	2·58	1·09	4·15	1·33	0·43	1·27	9·06	9·91	8·17
75-79 years	0·79	0·36	1·22	5·90	6·73	5·09	0·31	0·35	0·27	0·86	1·47	0·26	1·29	0·81	0·78	11·29	8·75	13·79
80-84 years	0·27	0·32	0·22	4·72	6·93	2·77	0·45	0·72	0·21	2·12	0·65	3·41	0·93	0·07	0·84	17·66	23·98	12·09
85+ years	1·19	0·33	1·79	9·93	19·25	3·43	0·33	0·30	0·35	2·68	0·60	4·13	1·14	0·59	0·82	14·59	17·78	12·36

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; HL = Hodgkin lymphoma; NHL = non-Hodgkin lymphoma

Appendix C

Table C1: Results of sensitivity analysis of temporal trends using World standard population

Measure	Base case			Sensitivity analysis			Change in trend
	EAPC (%)	B-Coefficient (95% CI)	P-value	EAPC (%)	B-Coefficient (95% CI)	P-value	
Australasia							
Incidence	-	-	-	-	-	-	
ALL	2.37	0.023 (0.019, 0.027)	<0.001	1.34	0.013 (0.010, 0.017)	<0.001	N/A
AML	1.93	0.019 (0.017, 0.021)	<0.001	0.27	0.003 (0.001, 0.005)	0.015	N/A
CLL	2.40	0.024 (0.022, 0.025)	<0.001	0.68	0.007 (0.005, 0.009)	<0.001	N/A
CML	2.16	0.021 (0.013, 0.029)	<0.001	0.45	0.004 (-0.004, 0.013)	0.300	Stable
HL	0.44	0.004 (0.003, 0.005)	<0.001	0.11	0.001 (0.000, 0.002)	0.027	N/A
NHL	0.95	0.009 (0.005, 0.014)	<0.001	-0.63	-0.006 (-0.012, -0.001)	0.018	Decreasing
Prevalence	-	-	-	-	-	-	
ALL	2.57	0.025 (0.021, 0.030)	<0.001	1.57	0.016 (0.012, 0.020)	<0.001	N/A
AML	0.90	0.009 (0.007,0.011)	<0.001	-0.60	-0.006 (-0.008, -0.004)	<0.001	Decreasing
CLL	2.40	0.024 (0.022, 0.025)	<0.001	0.70	0.007 (0.005, 0.009)	<0.001	N/A
CML	2.87	0.028 (0.022, 0.035)	<0.001	1.27	0.013 (0.006, 0.020)	<0.001	N/A
HL	0.40	0.004 (0.003,0.005)	<0.001	0.19	0.002 (0.001, 0.003)	<0.001	N/A
NHL	1.07	0.011 (0.005, 0.016)	<0.001	-0.68	-0.007 (-0.013, 0.000)	0.046	Decreasing
DALYs	-	-	-	-	-	-	
ALL	-0.36	-0.004 (-0.005, -0.003)	<0.001	-0.13	-0.001(-0.003, 0.001)	0.207	Stable
AML	1.15	0.011 (0.009, 0.013)	<0.001	-0.71	-0.007 (-0.008, -0.006)	<0.001	Decreasing
CLL	1.74	0.017 (0.015, 0.020)	<0.001	0.04	0.000 (-0.002, 0.003)	0.784	Stable
CML	-1.23	-0.012 (-0.024, 0.000)	0.042	-2.43	-0.025 (-0.037, -0.013)	<0.001	N/A
HL	-0.40	-0.004 (-0.005, -0.003)	<0.001	-0.93	-0.009 (-0.011, -0.008)	<0.001	N/A
NHL	0.18	0.002 (-0.004, 0.007)	0.536	-1.12	-0.011 (-0.017, -0.005)	<0.001	Decreasing
Deaths	-	-	-	-	-	-	
ALL	0.75	0.007 (0.006, 0.009)	<0.001	-0.26	-0.003 (-0.004, -0.002)	<0.001	Decreasing
AML	2.07	0.020 (0.019, 0.022)	<0.001	0.42	0.004 (0.003, 0.006)	<0.001	N/A
CLL	2.11	0.021 (0.019, 0.023)	<0.001	0.30	0.003 (0.001, 0.005)	0.002	N/A
CML	-0.27	-0.003 (-0.014, 0.008)	0.631	-2.13	-0.022 (-0.033, -0.010)	<0.001	N/A
HL	0.05	0.000 (-0.001, 0.002)	0.399	-1.02	-0.010 (-0.011, -0.009)	<0.001	Decreasing
NHL	1.07	0.011 (0.005, 0.016)	<0.001	-0.98	-0.010 (-0.015, -0.005)	<0.001	Decreasing
OCEANIA							
Incidence	-	-	-	-	-	-	
ALL	-0.45	-0.005 (-0.006, -0.003)	<0.001	-0.32	-0.003 (-0.005, -0.002)	<0.001	N/A
AML	-0.02	0.000 (-0.002, 0.001)	0.816	-0.18	-0.002 (-0.003, -0.000)	0.012	Decreasing
CLL	1.38	0.014 (0.012, 0.016)	<0.001	0.37	0.004 (0.002, 0.005)	<0.001	N/A
CML	-0.58	-0.006 (-0.008, -0.004)	<0.001	-1.13	-0.011 (-0.013, -0.009)	<0.001	N/A
HL	-0.05	-0.001 (-0.002, 0.001)	0.405	-0.55	-0.005 (-0.007, -0.004)	<0.001	Decreasing
NHL	0.92	0.009 (0.008, 0.010)	<0.001	0.37	0.004 (0.002, 0.005)	<0.001	N/A
Prevalence	-	-	-	-	-	-	
ALL	-0.56	-0.006 (-0.007, -0.004)	<0.001	-0.33	-0.003 (-0.005, -0.001)	0.001	N/A
AML	-0.21	-0.002 (-0.004, 0.000)	0.026	-0.14	-0.001 (-0.003, 0.001)	0.150	Stable
CLL	1.99	0.020 (0.017, 0.022)	<0.001	0.99	0.010 (0.007, 0.012)	<0.001	N/A
CML	-0.76	-0.008 (-0.010, -0.005)	<0.001	-1.15	-0.012 (-0.014, -0.009)	<0.001	N/A

Measure	Base case			Sensitivity analysis			Change in trend
	EAPC (%)	B-Coefficient (95% CI)	P-value	EAPC (%)	B-Coefficient (95% CI)	P-value	
HL	0.68	0.007 (0.003, 0.011)	0.001	0.42	0.004 (0.000, 0.008)	0.039	N/A
NHL	3.24	0.032 (0.021, 0.043)	<0.001	3.19	0.031 (0.023, 0.040)	<0.001	N/A
DALYs	-	-	-	-	-	-	-
ALL	-0.56	-0.006 (-0.007, -0.004)	<0.001	-0.34	-0.003 (-0.005, -0.002)	<0.001	N/A
AML	-0.12	-0.001 (-0.003, 0.000)	0.115	-0.18	-0.002 (-0.003, -0.000)	0.014	Decreasing
CLL	0.92	0.009 (0.008, 0.010)	<0.001	0.00	0.000 (-0.001, 0.001)	0.993	Stable
CML	-0.88	-0.009 (-0.011, -0.007)	<0.001	-1.21	-0.012 (-0.014, -0.010)	<0.001	N/A
HL	-0.55	-0.006 (-0.007, -0.005)	<0.001	-0.89	-0.009 (-0.010, -0.008)	<0.001	N/A
NHL	0.26	0.003 (0.001, 0.004)	<0.001	-0.05	0.000 (-0.002, 0.001)	0.587	Stable
Deaths	-	-	-	-	-	-	-
ALL	-0.39	-0.004 (-0.005, -0.002)	<0.001	-0.29	-0.003 (-0.004, -0.002)	<0.001	N/A
AML	0.08	0.001 (-0.001, 0.002)	0.244	-0.16	-0.002 (-0.003, -0.000)	0.005	Decreasing
CLL	0.99	0.010 (0.009, 0.011)	<0.001	0.04	0.000 (-0.001, 0.001)	0.499	Stable
CML	-0.48	-0.005 (-0.007, -0.003)	<0.001	-1.11	-0.011 (-0.013, -0.009)	<0.001	N/A
HL	-0.35	-0.003 (-0.004, -0.003)	<0.001	-0.90	-0.009 (-0.010, -0.008)	<0.001	N/A
NHL	0.65	0.007 (0.005, 0.008)	<0.001	0.15	0.001 (0.000, 0.003)	0.140	Stable

ALL = acute lymphocytic leukaemia; AML = acute myeloid leukaemia; ASR = age-standardised rate; CLL = chronic lymphocytic leukaemia; CML = chronic myeloid leukaemia; DALY = disability-adjusted life year; EAPC = estimated annual percentage change; HL = Hodgkin lymphoma; N/A = not assessed; NHL = non-Hodgkin lymphoma

1. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204

countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020;396(10258):1204-22. doi: 10.1016/s0140-6736(20)30925-9

2. Global Burden of Disease Cancer Collaboration. Cancer Incidence, Mortality, Years of

Life Lost, Years Lived With Disability, and Disability-Adjusted Life Years for 29 Cancer Groups From 2010 to 2019: A Systematic Analysis for the Global Burden of Disease Study 2019. *JAMA Oncology* 2022;8(3):420-44. doi: 10.1001/jamaoncol.2021.6987