


BMJ Open Comparing the effectiveness of pulsed radiofrequency treatment to lumbar dorsal root ganglion according to application times in patients with lumbar radicular pain: protocol for a randomised controlled trial

Soyoon Park,¹ Jae Ni Jang,¹ Ji-Hoon Park,² Yumin Song,¹ Choi Sooil,¹ Young Uk Kim,¹ Sukhee Park 

To cite: Park S, Jang JN, Park J-H, *et al.* Comparing the effectiveness of pulsed radiofrequency treatment to lumbar dorsal root ganglion according to application times in patients with lumbar radicular pain: protocol for a randomised controlled trial. *BMJ Open* 2024;**14**:e077847. doi:10.1136/bmjopen-2023-077847

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2023-077847>).

Received 17 July 2023

Accepted 02 February 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Catholic Kwandong University College of Medicine, Gangneung, Gangwon-do, Korea

²Keimyung University College of Medicine, Daegu, Daegu, Korea

Correspondence to

Professor Sukhee Park; appealex@gmail.com

ABSTRACT

Introduction Lumbar radicular pain (LRP) is a common symptom characterised by a sharp, shooting or lancinating sensation localised to one or more dermatomes of the lumbar spine. Despite its high prevalence and significant impact on quality of life, the most effective conservative treatment for patients with LRP remains uncertain. When conventional treatment methods do not provide satisfactory results, the option of using epidural steroids and/or pulsed radiofrequency (PRF) treatment may be considered as a secondary approach for managing the condition. Ongoing advances in the field have led to a wide range of PRF parameters being investigated and extensively documented. Therefore, this study will aim to evaluate the treatment efficacy, sustainability and adverse effects of PRF application for different durations in patients with LRP.

Methods and analysis This study will be a double-blind, randomised, controlled trial. Eligible patients with LRP who visit the International St. Mary's Hospital pain clinic in Korea will be assigned to three groups (1:1:1 ratio) based on the duration of PRF application: 240, 360 and 480 s. Outcome measures will include an assessment of radicular pain intensity, physical function, global improvement, treatment satisfaction and adverse events. The primary outcome will be a Numeric Rating Scale (NRS) score 3 months after the procedure. The secondary outcomes will be the number of subjects in each group reporting successful treatment defined as a significant decrease of NRS or improved physical function score or high satisfaction at the 3 and 6 months follow-up. X² or Fisher's exact test and one-way analysis of variance will be used to compare the outcomes.

Ethics and dissemination This trial was approved by the Ethics Committee of Catholic Kwandong University International St. Mary's Hospital (IS23EISE0018). The findings will be disseminated in peer-reviewed journals and at scientific conferences.

Trial registration number KCT00008612.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This study will be the first prospective study to evaluate the effectiveness of different application times of pulsed radiofrequency treatment in patients with lumbar radicular pain.
- ⇒ This will be a double-blind study where the researcher collecting the data and the operator are blinded.
- ⇒ The study has a relatively small sample size per arm and a short follow-up period.

INTRODUCTION

Lumbar radicular pain (LRP) is a common manifestation of lower back pain that presents as a sharp, shooting or lancinating sensation localised to one or more dermatomes in the lumbar spine.^{1 2} Chronic LRP is often associated with lumbar disc herniation, spinal stenosis or degenerative spondylolisthesis and can persist even after lumbar spine surgery.² This condition is often associated with lesions that either directly affect the dorsal root ganglion (DRG) or indirectly affect the spinal nerve and its roots by inducing axonal ischaemia or an inflammatory response.³

LRP is highly prevalent and significantly affects quality of life; however, the most effective conservative treatment for patients with LRP remains uncertain.^{4 5} If conservative treatments such as medication and physical therapy have not yielded satisfactory results, an epidural steroid injection may be tried. If repeated epidural steroid injections do not work or do not last long, pulsed radiofrequency (PRF) treatment may be considered a secondary approach to managing the condition.⁶

PRF is a relatively new neuromodulation technique that has been effective for many types of pain.^{4,5} However, there are conflicting results regarding its effectiveness.⁷⁻⁹ Although 45 V is a commonly used output voltage for PRF to lumbar DRG, but this varies from study to study, and there is no standard for the most effective output voltage and application time. Therefore, it is necessary to identify the variables that need to be adjusted during PRF to maximise its effectiveness.

Previous studies have shown that longer-duration PRF improves pain relief in trigeminal neuralgia and postherpetic neuralgia.^{10,11} The authors of this study aim to evaluate whether the efficacy of PRF in the lumbar DRG of patients with LRP depends on the duration of PRF application. To the best of our knowledge, no study has evaluated the therapeutic effects of PRF according to application time in patients with LRP.

METHODS AND ANALYSIS

Hypothesis

We hypothesised that the longer the duration of standard voltage (45 V) PRF applied to the lumbar DRG, the greater the improvement in pain and the longer the effect would last.

Trial design

We began enrolling subjects on 1 July 2023 with an expected end-of-study date of 30 June 2024. Prior to enrolment, participants will be provided with a written informed consent form as an integral part of the ethical approval process. The study will be a single-centre, double-blind (to both a researcher collecting data and an operator performing RF) randomised controlled trial with three groups in a 1:1:1 allocation ratio. After obtaining informed consent (online supplemental material), the patients will be randomly assigned to one of the three groups using a computer-generated number. The

clinical trial and the items to be examined in this study are presented in [table 1](#).

Inclusion criteria

The inclusion criteria for the patients will be as follows:

1. Age ≥ 20 years old.
2. Chronic LRP lasting ≥ 12 weeks.
3. Mild or moderate lumbar spinal stenosis or disc herniation confirmed by MRI using the Lee grading system^{12,13}
4. Failure of conservative management such as physiotherapy, exercise therapy or analgesic medications.
5. Patients who received conventional fluoroscopy-guided diagnostic/therapeutic transforaminal epidural injections of local anaesthetics and steroids.
6. Patients who reported persistent pain (Numeric Rating Scale (NRS) score ≥ 5) after receiving transforaminal epidural steroid injection (TFESI).

Exclusion criteria

The exclusion criteria for patients are as follows:

1. Patient refusal.
2. Signs of progressive motor weakness or neurological deficits.
3. Severe spinal stenosis in the lumbar spine confirmed by MRI.
4. Allergies to steroids or contrast dyes.
5. Coagulopathy.
6. Epidural steroid injection within the previous 4 weeks.
7. Systemic infection, injection site infection.
8. Malignancy.

Eligible patients for this study will be those who present to our institution and receive two fluoroscopic lumbar epidural steroid injections, 2 weeks apart. Patients who have received two epidural injections will be given a telephone interview 1 week after the second epidural injection.

Table 1 Trial schedule and observation items

Period	Screening (telephone)	Baseline (visit)	During procedure	30 mins after procedure	1 week (telephone)	2 weeks (telephone)	1 month (visit)	3 months (visit)	6 months (visit)
Oral/written consent form	✓	✓							
Demographic survey		✓							
Selection/exclusion criteria	✓								
Randomisation		✓							
NRS	✓	✓	✓	✓	✓	✓	✓	✓	✓
ODI		✓					✓	✓	✓
GPE							✓	✓	✓
Adverse reaction monitoring		✓	✓	✓	✓	✓	✓	✓	✓

NRS, numeric rating score; ODI, Oswestry Disability Index; GPE, Global Perceived Effect.
GPE, Global Perceived Effect; NRS, Numeric Rating Score; ODI, Oswestry Disability Index.

Patients with no pain relief or only transient pain relief from the two epidural injections (NRS pain intensity of ≥ 5 points) will be enrolled. Patient recruitment and enrolment will be conducted by a blinded researcher.

Assignment of interventions and blinding

After collecting baseline data, patients will be randomly assigned to one of the three groups based on the duration of radiofrequency application: 240s (120s \times two cycles), 360s (120s \times three cycles) and 480s (120s \times four cycles). At the beginning of the procedure, a nurse will uncover a sequentially numbered opaque envelope containing the group assignments. The generator is operated by the nurse, and the display is hidden from the patient and attending physician. Data on patient demographics, procedural information, and procedural outcomes were collected by a blinded researcher.

Procedure

The patient will be placed in the prone position with a pillow placed under the lower abdomen. After the aseptic preparation of the needle insertion site, the skin will be anaesthetised with 1% lidocaine. A 22-gauge, 4-inch RF cannula with a 10mm curved active tip (RFK-C101020B; Cosman Medical, Burlington, USA) will then be carefully inserted under fluoroscopic guidance (OEC 9800; General Electric Healthcare, Little Chalfont, Buckinghamshire, UK). The cannula will be directed into the lateral opening of the targeted intervertebral foramen. In the anteroposterior view, the cannula will advance no further medially than the lateral aspect of the pedicle. The final position of the tip of the cannula will be at the 2 o'clock position of the intervertebral foramen in the lateral image. In the anteroposterior view, the tip of the cannula will be at the 6 o'clock position of the pedicle column (figure 1). Further confirmation was obtained by injecting a contrast solution (Bonorex 300; Dai Han Pharm). Once the RF cannula is properly positioned, the stylet will be replaced with the RF probe (CB112-TC;




Figure 1 Fluoroscopic images of pulsed radiofrequency to left L5 dorsal root ganglion. Anteroposterior image (A) confirms needle in the mid pedicle line. Lateral image (B) is used to confirm needle depth and to visualise the needle entering the L4-L5 foramen. Contrasts outline the exiting nerve root and epidural space. Green=pedicle, light blue=intervertebral foramen, red X=needle tip position.

Cosman Medical). Subsequently, the probe will be connected to the PRF generator (Radiofrequency Ablation for Pain Management, G4 RF Generator; Cosman Medical). The final position of the PRF cannula will be determined by whether or not radicular pain occurs in the patient's corresponding nerve segment on sensory stimulation (50Hz) threshold of 0.3 to 0.5 V. The motor stimulus (2Hz) will aim to form at a voltage at least 1.5 times higher than the threshold of the sensory stimulus while ensuring that the impedance remains below 400 Ω . After each treatment cycle, the position of the RF cannula was fine-tuned and motor and sensory stimulations were performed to ensure accuracy and confirmation. The three groups will have different PRF application times: 120s \times two times (240s total), 120s \times three times (360s total) and 120s \times four times (480s total). The PRF output voltage will be 45 V for all three groups.

The PRF generator will keep the temperature below 42°C and will be set to a maximum output voltage of 45 V. The pulse width will be set to 20 ms at a frequency of 2Hz. PRF will be applied for 120s per cycle. Group 1 will be administered two cycles of PRF and two cycles of sham to blind the patient. In Group 2, three cycles of PRF and one cycle of sham will be applied. Group 3 will be administered four cycles of PRF.

Outcome assessment and follow-up

Comprehensive data, including demographic factors, such as age, sex, height, weight and body mass index, will be collected at baseline. Medical history will be assessed for conditions such as diabetes and hypertension. Other baseline information will include the diagnosis, total duration of pain, target level of the affected nerve root and number of previous epidural injections. For intra-operative parameters, we will collect the stimulation voltages used during 50 and 2Hz electrical stimulation positioning, procedure duration, output voltage, output current, impedance values (before and after the procedure) and electric field intensity ((output voltage)²/resistance).

Outcome measures will include the following assessments:

1. Assessment of pain intensity at the treatment site, both during and 30 min after the procedure.
2. Assessment of radicular pain intensity at specific time points: 1 and 2 weeks (telephone visit) and 1, 3 and 6 months after the procedure.
3. Assessment of physical function scores before the procedure and at 1, 3, and 6 months after the procedure.
4. Measurement of global improvement and satisfaction with treatment at 1, 3, and 6 months after the procedure.
5. Monitoring and documentation of any adverse events.

Pain intensity at the treatment site and radicular pain intensity will be assessed using an 11-point NRS, ranging from 0 (no pain) to 10 (unbearable pain). The physical function score will be assessed using the 10-item Korean version of the Oswestry Disability Index (ODI)




Figure 2 Flowchart of study design. PRF, pulsed radiofrequency treatment.

questionnaire, which ranges from 0 to 100, with 0 indicating no disability. Patient satisfaction and improvement will be measured using the 7-point Likert scale: Global Perceived Effect (GPE). Any adverse events occurring during treatment and follow-up will be documented individually. The flowchart of the study is presented in [figure 2](#).

The primary outcome will be the NRS score at 3 months after the procedure. The secondary outcomes will be the number of responders in each group reporting successful treatment after the 3 and 6 months follow-up period. A successful response will be defined as one or more of the following three outcomes:

1. A reduction in pain intensity was measured using the NRS of at least 50% or a reduction of at least 4 points.
2. Decrease in ODI of at least 10 points.
3. A score of at least 6 on the GPE scale on a 7-point Likert scale.

As the secondary outcomes, NRS scores will be assessed at 1 and 2 weeks postprocedure and at 1 and 6 months postprocedure. ODI scores will be assessed at 1, 3 and 6 months after the procedure. GPE scores will be assessed at 1, 3 and 6 months after the procedure. We will also analyse the degree of reduction in NRS pain and ODI scores at 3 and 6 months postprocedure in relation to preprocedure baseline scores. Any complications that occur during the procedure will be reported accordingly. During the 1, 3 and 6 follow-up visits, close attention will be paid to a comprehensive assessment of adverse events.

Sample size calculation and statistical analysis

Our preliminary study showed that when PRF was applied at 45 V for a total of 240s, the mean NRS score after 3 months was 3.2 with a SD of 1.3.¹⁴ We considered a difference in NRS scores of 1 or more points to be a clinically significant difference between groups and applied a 5% significance level and 90% power, requiring 18 subjects in each group. Considering a dropout rate of 10%, we will conduct a study with 20 subjects in each group, for a total of 60 subjects.

The intention-to-treat (ITT) analysis will be applied, and the data of every randomised subject will be analysed at each follow-up moment, regardless of lost to follow-up or withdrawal from the study. We will further perform a per-protocol analysis to show if there is a significant

difference from the ITT analysis. The categorical variables will be presented as numbers and percentages. The continuous variables will be presented as the mean with SD or median and IQR. To compare data from the three groups, the χ^2 or Fisher's exact test will be used for categorical variables, and one-way analysis of variance (ANOVA) or repeated measures ANOVA will be used for continuous variables. When there is missing data or drop-outs, a linear mixed model will be used to analyse the secondary continuous variables. Statistical significance will be set at p value < 0.05. All analyses will be performed using SPSS V.26.0 (IBM Corporation, Chicago, Illinois, USA).

ETHICS AND DISSEMINATION

This trial was approved by the Ethics Committee of Catholic Kwandong University International St. Mary's Hospital (IS23EISE0018). This trial is registered with the Clinical Trial Registry of Korea (<https://cris.nih.go.kr/cris/index/index.do>). The findings will be disseminated in peer-reviewed journals and at scientific conferences.

Contributors CS, J-HP, YS and Sukheep conceived and designed the study. YUK, YS and Soyeon P revised the manuscript. Sukheep will lead to the statistical analysis. JNJ and SoyeonP oversaw the data acquisition. All authors have reviewed and approved the final manuscript.

Funding This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean Government Ministry of Science and ICT (grant number 2022R1F1A1068925).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Sukhee Park <http://orcid.org/0000-0002-8798-7578>

REFERENCES

- 1 Stafford MA, Peng P, Hill DA. Sciatica: a review of history, epidemiology, pathogenesis, and the role of epidural steroid injection in management. *Br J Anaesth* 2007;99:461–73.
- 2 Van Boxem K, Cheng J, Patijn J, et al. 11. Lumbosacral radicular pain. *Pain Pract* 2010;10:339–58.
- 3 Howe JF, Loeser JD, Calvin WH. Mechanosensitivity of dorsal root ganglia and chronically injured axons: a physiological basis for the radicular pain of nerve root compression. *Pain* 1977;3:25–41.

- 4 Sluijter ME. Pulsed radiofrequency. *Anesthesiology* 2005;103:1313;
- 5 Van Zundert J, Patijn J, Kessels A, *et al.* Pulsed radiofrequency adjacent to the cervical dorsal root ganglion in chronic cervical radicular pain: a double blind sham controlled randomized clinical trial. *Pain* 2007;127:173–82.
- 6 Sluijter ME, Cosman ER, Rittman lii WB, *et al.* The effects of pulsed radiofrequency fields applied to the dorsal root ganglion: a preliminary report. *Pain Clinic* 1998;11:109.
- 7 De M, Mohan VK, Bhoi D, *et al.* Transforaminal epidural injection of local anesthetic and dorsal root ganglion pulsed radiofrequency treatment in lumbar radicular pain: a randomized, triple-blind, active-control trial. *Pain Pract* 2020;20:154–67.
- 8 Koh W, Choi S-S, Karm MH, *et al.* Treatment of chronic lumbosacral radicular pain using adjuvant pulsed radiofrequency: a randomized controlled study. *Pain Med* 2015;16:432–41.
- 9 Lee DG, Ahn SH, Lee J. Comparative effectiveness of pulsed radiofrequency and transforaminal steroid injection for radicular pain due to disc Herniation: a prospective randomized trial. *J Korean Med Sci* 2016;31:1324–30.
- 10 Wan C-F, Liu Y, Dong D-S, *et al.* Bipolar high-voltage, long-duration pulsed radiofrequency improves pain relief in postherpetic neuralgia. *Pain Physician* 2016;19:E721–8.
- 11 Wan C, Dong D-S, Song T. Long-duration pulsed radiofrequency on Gasserian ganglion improves acute/subacute Zoster-related trigeminal neuralgia: a randomized, double-blinded. *Pain Physician* 2019;22:361–8.
- 12 Lee GY, Lee JW, Choi HS, *et al.* A new grading system of lumbar central canal stenosis on MRI: an easy and reliable method. *Skeletal Radiol* 2011;40:1033–9.
- 13 Lee S, Lee JW, Yeom JS, *et al.* A practical MRI grading system for lumbar foraminal stenosis. *AJR Am J Roentgenol* 2010;194:1095–8.
- 14 Jang JN, Park S, Park J-H, *et al.* Output current and efficacy of pulsed radiofrequency of the lumbar dorsal root ganglion in patients with lumbar radiculopathy: a prospective, double-blind, randomized pilot study. *Pain Physician* 2023;26:E797–804.

Comparing the effectiveness of pulsed radiofrequency treatment to lumbar dorsal root ganglion according to application times in patients with lumbar radicular pain: Protocol for a randomised controlled trial

Soyoon Park,¹ Jae Ni Jang,¹ Ji-Hoon Park,² Yu min Song,¹ Sooil Choi,¹ Young Uk Kim,¹ and Sukhee Park^{1*}

¹Department of Anesthesiology and Pain Medicine, International St. Mary's Hospital, Catholic Kwandong University School of Medicine, Incheon, Korea

²Department of Anesthesiology and Pain Medicine, Dongsan Medical Center, Keimyung University School of Medicine, Daegu, Korea

*Correspondence: Sukhee Park, MD

Department of Anesthesiology and Pain Medicine, International St. Mary's Hospital, Catholic Kwandong University School of Medicine, Incheon, Korea

Tel.: +82-32-290-3932; Fax: +82-32-290-3879; Email: appealex@gmail.com

1. Title of the research
Comparing the effectiveness of pulsed radiofrequency treatment to lumbar dorsal root ganglion according to application times in patients with lumbar radicular pain: double blind, randomized controlled study
2. Principal Investigator
Sukhee Park, Department of anesthesia and pain medicine
3. What you need to know before deciding to participate in the clinical research
The researchers of this study will follow related regulations when obtaining and documenting your consent to participate in this study, and will follow lawful procedures based on ethical principles from the Declaration of Helsinki. This study has experimental and unproven aspects. One of the researchers will explain the purpose of the research. You can contact us at any time with any questions about this consent form. Your participation in this research is entirely voluntary and you may decide to participate or not to participate. You are encouraged to take time to think about your decision to participate in this research and to discuss your decision with family or others if necessary. No matter what you decide, there is no penalty on you. If you decide to participate, you will be asked to sign this consent form, and you will receive one signed copy of this document.
4. Research objectives
In this research, we aim to investigate the effectiveness of pulsed radiofrequency treatment by varying the time of radiofrequency application to the lumbar posterior root ganglion. Pulsed radiofrequency treatment procedures typically use an output voltage of 45V, which is known as the standard voltage. The purpose of this research is to learn if there is a difference in the effectiveness depending on the application time at standard voltage. No studies have compared the effectiveness of lumbar dorsal root ganglion(DRG) pulsed radiofrequency(PRF) over application time in patients with lumbar radicular pain. However, animal studies have shown that pain relief increases with application time, but a difference in effectiveness is not significant and may cause nerve damage after 12 minutes. Therefore, based on these existing data, this clinical trial aims to evaluate the efficacy of pulsed radiofrequency treatment on the posterior root ganglion for lumbar radiculopathy with different durations of action while not causing nerve damage, and to compare the pain and discomfort during the procedure.
5. Duration of research
A total study duration is 12 months, and your actual participation in the study is 6 months.
6. Number of patients participating in the research
We are expecting a total 60 patients from this hospital to participate.
7. Procedure of the research

<p>You are eligible for this clinical research because you have undergone two lumbar epidural injections at our institution with no pain reduction, and decided to undergo pulsed radiofrequency treatment on the lumbar posterior root ganglion. You will receive same medications and medical devices that are routinely used in pulsed radiofrequency treatment. But pulsed radiofrequency procedure will be applicated for different durations with same voltage. After consent to participate in this study, you will be randomized to one of three groups with different durations of pulsed radiofrequency treatment on the lumbar posterior root ganglion. You are assigned to one of the groups by chance (like flipping a coin) with a 33.3% probability. The research will be double blinded to investigator and participant, which means that neither the investigator nor the participant know which group they are in until the end of the study. All research will be conducted outpatient procedure room in Pain Clinic. Typically, patients undergoing pulsed radiofrequency treatment in our clinic will have outpatient follow-up at 1,3,6 months after the procedure and participation in this research will not require additional outpatient follow-up. However, you will be contacted by phone 1 and 2 weeks after the procedure to ask about lower extremity radiation pain intensity and side effects, which will take less than 5 minutes. At your clinic visit, you will be asked about your radiating pain intensity and satisfaction with treatment, and afterward, you will complete the Korean version of the Oswestry Disability Index(ODI), which is a series of 10 questions and usually takes about 5 minutes to complete. This is a routine procedure for all patients undergoing pulsed radiofrequency treatment and your participation in this study will not take any additional time. To summarize, you will have two phone calls and three clinic visits after your pulsed radiofrequency treatment procedure.</p>
<p>8. What you need to know while participating in a study</p> <p>If you decide to participate in this study, it is important you agree to the following to ensure a successful, safe, and scientific research.</p> <ul style="list-style-type: none">- Please follow the instructions of your principal investigator or research coordinator.- It is important that you visit the clinic on your scheduled appointment. If you are unable to keep your appointment, please contact your research coordinator to schedule a new appointment.- If you suspect any unwanted effects, damage, or adverse reactions, you should notify your investigator immediately and follow instructions.- You must inform the researcher of any other studies in which you are currently participating, and you must not participate in any other clinical research while participating in this research.- If you change your mind about participating in the research, you must tell the investigator
<p>9. Expected risks or inconveniences of participating in this research</p> <p>It is common to experience pain with pulsed radiofrequency treatment, and it is possible that the pain may increase over time. However, in our previous practice, almost all patients found the pain to be tolerable and can be easily detected by patient self-report. If a patient reports pain scoring above 5 during the procedure, we will stop the study. Other than pain and discomfort during the procedure, there are no expected side effects with more application duration.</p>
<p>10. Expected benefits</p>

<p>There is no expected benefit to you for participating in this research, however the information gained from this research will help other patients in the future.</p>
<p>11. Treatments performed when you don't participate in this research</p>
<p>If you do not participate in this study, you will receive a conventional duration of pulsed radiofrequency treatment with a standard voltage.</p>
<p>12. Compensation/Indemnification and treatment in the event of injury related to clinical research</p>
<p>No additional risks other than pain during the procedure are expected for this research. The investigator will make every effort to ensure your safety throughout the research. If you experience any adverse events or other harm as a result of this research, the investigator will treat you as best as he or she can and compensate you according to the protocol for compensation of victims.</p>
<p>13. Discontinuing participation in the research</p>
<p>You may discontinue the study at any time without penalty or loss of benefit. You will be withdrawn from the study if you complain of moderate or severe pain during pulsatile radiofrequency therapy that is greater than or equal to 5 on a numerical rating scale, if the procedure plan is changed during the procedure, or if you are considered to be hemodynamically unstable during the procedure by the investigator. If you would like to discontinue participating in the research, please contact the investigator. You will not get any penalties or loss of benefits if you decide to withdraw. The investigator may also withdraw you from the research when he or she believes it is in your best interest. There is a possibility that the investigator of this research may discontinue the entire study.</p>
<p>14. Honorarium policy</p>
<p>- If you sign to participate in this research, you will receive ₩50,000 to cover your transportation costs, and there are no additional expenses needed with participating in the study.</p> <p>- Even if you participate in this study, you will still be responsible for the cost of any medications you take and any hospitalizations, laboratory tests, and medical care that occur unrelated to this study.</p>
<p>15. Regarding privacy and providing personal information</p>
<p>Your information collected in this research is protected by applicable laws, including the Personal Data Protection Act. Therefore, the investigator would like to explain to you about the items of personal information (including sensitive information) that will be collected for this research study, its purpose, retention period, use, and disclosure, and to obtain your consent. In addition, all data collected from you will be anonymized and any records that could identify you will be kept confidential. Your data may be used for national/international publication of the research result if needed. Even in case of publication, your personal information will still remain confidential.</p>

The purpose of collecting and using your personal and sensitive information is to organize, analyze, and report the results of the reasearch.

The information we collect about you is as follows

Information type	Contents
Personal information	Full name (including initials), age (date of birth), gender, address, contact number, account number, etc.
Sensitive information	Medical records (medical history, concomitant medications, birth control, pregnancy/nursing, test results, etc.)

※ Your personal information is collected for identifying participants, scheduling appointments for research, and paying for transportation costs etc.

Your personal information may be disclosed to limited personnel for research-related purposes as described in section 15 (see "underline" below).

(Your personal information will only be disclosed within the hospital and will not be disclosed or transferred outside of the hospital).

Your personal and sensitive information may be retained and used within the purpose of the research for a period of 3 years from the end date of the research.

After that, it will be immediately destroyed unless there is a special reason.

Consent form

I have been informed about this consent, I have read and understood the contents of this consent, and I have received answers to all my questions. I agree to participate in this research on a voluntary basis, so I will sign the consent form. And I understand that I will be provided with a copy of the consent form.

I also agree that if I stop participating in the study or withdraw from the study, the data collected prior to my withdrawal can be used.

☐ Yes

☐ No

Title of the research: Comparing the effectiveness of pulsed radiofrequency treatment to lumbar dorsal root ganglion according to application times in patients with lumbar radicular pain: double blind, randomized controlled study

■ Research subjects

(Name) (Signature) (Date)

■ Proxy (for minors or when voluntary consent is not possible) (if necessary)

(Name) (Signature) (Date)

(Relationship to research subjects)

- ※ Proxy means: If the subject is incapable of consent, a proxy can consent on behalf of the subject without violating the subject's wishes.
1. legally authorized representative
- : A person whose representation becomes effective by law directly, without the need for a delegation.
2. If there is no legal representative, the spouse, immediate family members, and immediate relatives shall be appointed as a proxy in that order. If there are more than one eligible proxy, they shall be determined by mutual agreement, and if no agreement is reached, the elderly person shall be the proxy.

■ Observer (if the subject or legal representative is unable to read the consent form) (if needed)

(Name) (Signature) (Date)

This observer was involved throughout the consent process, and the information in the consent form and all information about the study was provided accurately and understood. The subject (the subject's proxy) has given voluntary consent.

■ Principal Investigator (or Co-Investigator)

Name) (Signature) (Date)