## Supplementary 13. Research Schedule

Research Phase	Screening Radiotherapy period					Follow up period										
Visits	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Time	-3M-0	1W	2W	3W	4-7W	6M	12M	18M	24M	30M	36M	42M	48M	54M	60M	In the event of a relapse
Informed consent	×															
Inclusion/exclusion	×															
criteria																
Frozen/paraffin tissue	×															
Pathology	×*															
Tumor markers	×					×	×	×	×	×	×	×	×	×	×	×
Breast + LN	×					×	×	×	×	×	×	×	×	×	×	
ultrasound/MR																
Mammography	×*						×		×		×		×		×	
Chest X-ray/CT	×					×	×	×	×	×	×	×	×	×	×	×
Abdominal	×					×	×	×	×	×	×	×	×	×	×	×
ultrasound/CT																
Thyroid + LN	×					×	×	×	×	×	×	×	×	×	×	
ultrasound																
Bone Scan	×*						×		×		×		×		×	×
Cranial CT/MRI																×***
Demographic features	×															
Medical history	×															
Physical examination	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	

Upper and lower arm	×				×		×		×		×		×		×	
circumference																
measurement◆																
Body weight	×				×		×		×		×		×		×	
ECOG score	×				×		×		×		×		×		×	
Blood count	×**	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×
Liver and kidney	×**				×	×	×	×	×	×	×	×	×	×	×	×
function																
Infectious disease	×*															
Serum pregnancy test	×**															
EORTC QLQ-C30	×**				×		×		×		×		×		×	
EORTC QLQ-BR23	×**				×		×		×		×		×		×	
Harris	×**						×		×		×		×		×	
Baker	×**						×		×		×		×		×	
CTCAE V4.03	×**	×	×	×	×		×	×	×		×		×		×	
Acute radiation		×	×	×	×											
reaction																
Late radiation injury							×	×	×		×		×		×	

<sup>◆</sup>Upper and lower arm circumference measurements: measured at the bilateral metacarpophalangeal joints (mid-metacarpal), at the wrist, and at the distal 10 cm and proximal 15 cm of the elbow epicondyle.

<sup>\*</sup> Within 12 months prior to randomization

<sup>\*\*</sup> Within 14 days before randomization

<sup>\*\*\*</sup> When cranial symptoms were present