

Supplementary table 1 Baseline characteristics in 54,568 patients with out of hospital cardiac arrest stratified by pre-existence of type 1 diabetes.			
	Type 1 diabetes	No type 1 diabetes	p
Socioeconomic status - n (%)			
Region of birth			0.370
<i>Sweden</i>	401 (89.7)	46123 (85.6)	
<i>Denmark Finland Norway Iceland</i>	16 (3.6)	2976 (5.5)	
<i>EU</i>	12 (2.7)	1598 (3.0)	
<i>Europe not EU</i>	8 (1.8)	1138 (2.1)	
<i>North America</i>	0 (0.0)	130 (0.2)	
<i>Asia</i>	6 (1.3)	1301 (2.4)	
<i>Africa</i>	2 (0.4)	362 (0.7)	
<i>South America</i>	2 (0.4)	166 (0.3)	
<i>Other</i>	0 (0.0)	75 (0.1)	
Disposable family income - median IQR ¹	2574.50 [1502.75, 3968.25]	2469.00 [1515.00, 3748.00]	0.353
Work or profession- n (%)			
<i>Unemployed</i>	274 (69.0)	37564 (77.2)	
Educational level- n (%)			0.382
<i>Pre gymnasium 9 years</i>	95 (23.8)	13755 (27.9)	
<i>Pre gymnasium 9 years</i>	53 (13.2)	6220 (12.6)	
<i>Gymnasium 3 years</i>	124 (31.0)	13770 (27.9)	
<i>Gymnasium 3 years</i>	58 (14.5)	6110 (12.4)	
<i>Post gymnasium 3 years</i>	28 (7.0)	3811 (7.7)	
<i>Post gymnasium 3 years or longer</i>	28 (7.0)	4191 (8.5)	
<i>Research education</i>	2 (0.5)	322 (0.7)	
<i>Unknown</i>	12 (3.0)	1169 (2.4)	
Marital status- n(%)			0.005
<i>Not married</i>	121 (30.0)	11225 (22.3)	
<i>Married</i>	162 (40.2)	22946 (45.7)	
<i>Surviving partner</i>	0 (0.0)	3 (0.0)	
<i>Registered partner</i>	0 (0.0)	16 (0.0)	
<i>Divorced</i>	75 (18.6)	8407 (16.7)	
<i>Divorced partner</i>	0 (0.0)	11 (0.0)	
<i>Widow widower</i>	45 (11.2)	7632 (15.2)	
Previous conditions - n (%)			
<i>Hypertension</i>	274 (61.2)	24183 (44.7)	<0.001
<i>Heart failure</i>	101 (22.5)	12316 (22.8)	0.960
<i>Chronic ischemic heart disease</i>	104 (23.2)	11142 (20.6)	0.190
<i>Atrial fibrillation</i>	74 (16.5)	11124 (20.6)	0.041

Type 2 diabetes	0 (0.0)	10423 (19.3)	<0.001
Dyslipidemia	128 (28.6)	8432 (15.6)	<0.001
Angina, including unstable angina	81 (18.1)	8263 (15.3)	0.114
Alcohol dependency	76 (17.0)	7683 (14.2)	0.109
Acute myocardial infarction	64 (14.3)	7280 (13.5)	0.656
Affective disorders	62 (13.8)	5649 (10.4)	0.024
Renal failure	68 (15.2)	5366 (9.9)	<0.001
Thrombotic stroke	33 (7.4)	4739 (8.8)	0.340
Alzheimers dementia	27 (6.0)	4059 (7.5)	0.276
Aortic stenosis	25 (5.6)	3303 (6.1)	0.718
Medications prescribed - n (%)			
Anticoagulant or antiplatelet agent ATC ² B01	166 (37.1)	19636 (36.3)	0.773
Beta blockers	165 (36.8)	18153 (33.5)	0.156
ACE inhibitor or ARB	201 (44.9)	17690 (32.7)	<0.001
Diuretics	144 (32.1)	14607 (27.0)	0.017
Lipid lowering drugs	165 (36.8)	12801 (23.7)	<0.001
Drugs for acid related disorders	111 (24.8)	10822 (20.0)	0.014
Calcium channel blockers	110 (24.6)	8604 (15.9)	<0.001
Other cardiovascular drugs ATC C01	55 (12.3)	6421 (11.9)	0.845
Antihypertensive drugs ATC C02	11 (2.5)	610 (1.1)	0.016
Abbreviations: ¹ Interquartile range, ² Anatomical Therapeutic Chemical Classification System			