BMJ Open Retinal and choroidal microvascular features during pregnancy: a systematic review and meta-analysis

Peng Zhang 💿 , Caixia Wang, Yuchen Liang, Qingli Shang

To cite: Zhang P, Wang C, Liang Y, et al. Retinal and choroidal microvascular features during pregnancy: a systematic review and meta-analysis. BMJ Open 2024;14:e087319. doi:10.1136/ bmiopen-2024-087319

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (https://doi.org/10.1136/ bmjopen-2024-087319).

Received 08 April 2024 Accepted 23 July 2024

Check for updates

C Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Department of Ophthalmology, The Second Hospital of Hebei Medical University, Shijiazhuang, Hebei, People's Republic of China

Correspondence to Qingli Shang: qinglishang@hebmu.edu.cn

ABSTRACT

Purpose To analyse retinal and choroidal microvasculature features in healthy pregnant women using optical coherence tomography angiography (OCTA). Methods PubMed, Embase, Ovid, Web of Science and Cochrane Library were searched for published studies that compare retinal and choroidal microvasculature characteristics between pregnant females and nonpregnant females from inception to November 2023. The statistics were continuous variables. They were presented as the mean difference (MD) with a corresponding 95% Cl. We used Review Manager software 5.4.1 for statistical

analysis Results A total of 5 eligible studies involving 219 eves in the pregnancy group and 186 eyes in the control group were included in the meta-analysis. The pooled results showed that the retinal deep capillary plexus vascular density (DCP VD) measured by OCTA was significantly higher in healthy pregnant women than in controls (MD 1.53: 95% CI 0.62 to 2.44: $l^2=0\%$: p=0.0009). However. differences between the two groups in the retinal superficial capillary plexus VD (SCP VD) (MD 0.41;95% Cl -1.26 to 2.09; $l^2=85\%$; p=0.63), the foveal avascular zone (FAZ) (MD 0.01; 95% CI -0.01 to 0.03; I²=14%; p=0.18), the choriocapillaris VD (CC VD) (MD 0.76; 95% CI -1.11 to 2.64; I²=79%; p=0.43) were uncertain.

Conclusions Our meta-analysis found that the DCP VD of healthy pregnant women was higher than that of nonpregnant controls. However, differences in SCP VD, FAZ and CC VD between the healthy pregnant women and the non-pregnancy controls were uncertain. Our findings can help to get a deeper understanding of retinal and choroidal microvascular characteristics during pregnancy.

INTRODUCTION

Pregnancy is a physiological state which causes a variety of anatomic and functional body changes in several organ systems, such as metabolic, cardiovascular, hormonal, immunological and haematological systems. These changes mainly aim to provide favourable conditions for fetal development. The multiorgan changes induced by pregnancy also can affect the eye. Previous studies revealed that meibomian gland, cornea, conjunctiva, crystalline lens, refractive state, choroidal thickness (CT) can be affected in pregnant women.^{2–7} In most cases, the above

STRENGTHS AND LIMITATIONS OF THIS STUDY

- \Rightarrow This study systematically reviewed the retinal and choroidal microvascular features during pregnancy.
- \Rightarrow We assessed the evidence quality using the Grading of Recommendations Assessment. Development and Evaluation approach, which provided essential information for primary outcomes.
- \Rightarrow The total sample size in the included studies was relatively small.
- There exists heterogeneity across the included \Rightarrow studies.

changes are reversible and temporary. They are fully reversed in the months after delivery. However, sometimes they can be pathological and permanent, such as central serous chorioretinopathy, serous retinal detachment during pregnancy and vascular changes due to pre-eclampsia. Some studies speculated that these changes may be attributed to a altered hormonal levels and altered ocular \blacksquare blood flow.⁸⁹

The most significant systemic changes ≥ during pregnancy are haemodynamic state is mainly to meet the demand for increased blood circulation. Plasma volume increases by **9** 50% and reaches its peak at 39 works. tion.¹⁰ Also, there is an increase in cardiac <u>0</u> output, peripheral vasodilation and a fall in systemic vascular resistance.¹¹ In addition, another significant change during pregnancy is hormone levels. Oestrogen, progesterone and renin–angiotensin levels increase markedly in pregnancy, which play an important \underline{G} role in regulating vascular endothelial cells, as well as the systemic and ocular vasculature.¹²

Previous research evaluated ocular vascular changes during pregnancy, which mainly focused on retinal and choroidal microcirculation.^{13 14} Microcirculation is a system consisting of arterioles, venules, capillaries and arteriovenous anastomosis. Normal microcirculation is the basis of retinal and choroidal function. In addition, studying

e

Protected by copyright, including for uses related

the microcirculation of the eve helps to understand the vascular effects of systemic diseases. Optical coherence tomography angiography (OCTA) is a non-invasive technology that allows us to visualise the retinal and choroidal vasculature without the need for intravenous contrast agent and eye-drops. Some parameters can be safely obtained using OCTA, such as retinal superficial capillary plexus (SCP) vascular density (VD), retinal deep capillary plexus (DCP) VD, foveal avascular zone (FAZ) and choriocapillaris (CC) VD.¹⁵ Some studies have observed retinal and choroidal microvasculature features in pregnant women using OCTA. However, because of the limited sample size of the clinical trials, the outcomes were inconsistent. Given the discrepancy, we performed a detailed meta-analysis based on PICO (Population, Intervention, Comparator, Outcome) questions to analyse retinal and choroidal microvasculature features during pregnancy and provide deep understanding of microvasculature changes in pregnant women. The detailed PICO questions are shown in online supplemental material 1.

METHODS

Search strategy

This meta-analysis was performed in compliance with the Cochrane Collaboration recommendations and the results were reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. PubMed, Embase, Ovid, Web of Science and Cochrane Library were searched for published eligible studies from inception to November 2023. The search strategy was based on a combination of terms: (1) "pregnancy" or "pregnant" or "gravid" or "gravids" or "gestation" or "conception"; (2)"retinal microvasculature" or "choroidal microvasculature" or "superficial capillary plexus" or "SCP" or "deep capillary plexus" or "DCP" or "foveal avascular zone" or "FAZ" or "choriocapillaris" or "CC" and (3) "optical coherence tomography angiography" or "OCTA" (online supplemental file 2).

Study selection

In this meta-analysis, the inclusion criteria were as follows: (1) Studies comparing retinal microvasculature features or choroidal microvasculature features between pregnant women and non-pregnant women. (2) The parameters in the studies included SCP-VD, DCP-VD, FAZ and CC-VD. (3) The parameters were measured by OCTA. (4) Casecontrol studies involving human subjects. (5) Articles published in English were considered qualified.

In this meta-analysis, the exclusion criteria were as follows: (1) They were reviews, case reports, animal studies, conference abstracts, comments, posters or duplicate studies. (2) There are no sufficient data or data could not be extracted. (3) The included pregnant women and non-pregnant women were diagnosed with certain systemic diseases, such as hypertension, diabetes, pre-eclampsia, anaemia, etc.

Brought of the set acted by consulting a third author and discussion. According to the Cochrane Handbook, the following data were extracted from the included studies: the first author of each study, year of publication, country, the mean age, the gestation weeks, OCTA device and parameter results. **Dully assessment** of the included studies was performed to the following the Newcastle-Ottawa Scale (NOS). The evaluation or criteria include subject selection (4 points), subject comparability (2 points) and exposure assessment (3 points). The result of the quality assessment is a score range of 0–9 points. Studies with a total score <4 points of outcomes were evaluated using the Grading of Recomparability (2 points) and exposure assessment (3 points). The result of the quality. The evidence quality of uctoomes were evaluated using the Grading of Recomparability (2 points) and publication bias, evidence (GRADE) method. In terms of risk of bias, inconsistency, indirectness, imprecision and publication bias, evidence quality of uctoomes were evaluated using the Grading of Recomparability (2 montes) frisk of bias, inconsistency, indirectness, imprecision and publication bias, evidence quality of uctoomes were evaluated using the Grading of Recomparability (2 montes) frisk of bias, inconsistency, indirectness, imprecision and publication bias, evidence quality was divided into 'high quality', 'moderate quality' is quality was divided into 'high quality', 'moderate quality' and 'very low quality' and 'very low quality' is possible of the generation of the statistic test and 1[°] statistic test. 1[°]>50% was considered to be of the mean ±SD, and mean difference (MDs) with its 95% CI were evaluated for poloting the statistic test and 1[°] statistic test. 1[°]>50% was considered to be statistically significant heterogeneity. A fixed-effect

 χ^2 statistic test and I² statistic test. I²>50% was considered **B** to be statistically significant heterogeneity. A fixed-effect $\overline{\mathbf{Z}}$ model was adopted if there was no significant heterogeneity among studies ($I^2 < 50\%$), otherwise a random-effect \ge model was used. The publication bias was evaluated by funnel plots. A p<0.05 was considered statistically significant. According to the distribution-based methods, we considered the minimal clinically important difference (MCID) as a difference of 0.85% in SCP VD, 0.46% in DCP VD, 0.01 mm^2 in FAZ, 0.96% in CC VD.¹⁶¹⁷ When the differences between healthy pregnant women and nonpregnant controls were smaller than MCIDs, although technologies. p<0.05, we considered the difference as small and not clinically important.

Patient and public involvement

None.

RESULTS **Search results**

The detailed flow chart of retrieval and screening process is shown in figure 1. A total of 173 articles, including 91 studies potentially useful, were obtained from all databases after duplicates were removed. Two reviewers



Figure 1 Flow diagram of study selection and search results.

independently screened the titles and abstracts and excluded 79 articles. Then, 12 articles were assessed for qualified ones. These 12 records were screened for the

text

ar

d data mining

abstract and full text. Among them, five studies focused on pregnant women with systemic complications, one study had incomplete parameters and one study in which statistics were not presented as mean±SD was also excluded. Ultimately, five studies¹⁸⁻²² were selected for inclusion. The five included articles were published from 2019 to 2023 and involved 219 eyes of pregnant women and 186 control eyes of non-pregnant women. All these studies were non-randomised comparative observational studies. The extracted general characteristics of included studies are shown in table 1.

Quality assessment

Protected by copy NOS quality assessment scores of the included studies were 6-8. They were considered to be of fair to good quality. The detailed quality assessment results of included studies are displayed in table 1.

We used the GRADE method to evaluate the quality of evidence for the primary outcomes: SCP VD, DCP VD, FAZ and CC VD. According to five degradation factors, namely risk of bias, inconsistency, indirectness, imprecision and publication bias, the results suggested that the g evidence quality of SCP VD and CC VD was low and the uses related to evidence quality of DCP VD and FAZ was moderate. The detailed assessment results of evidence quality are shown in table 2.

Meta-analysis

The retinal SCP VD analysis in pregnant women and controls

Four studies with a total of 360 eyes (197 eyes in the pregnancy group and 163 eyes in the control group) reported the retinal SCP VD, but with significant heterogeneity across studies (χ^2 =20.54, p=0.0001, I²=85%). The difference between the pregnancy group and the control group was uncertain (MD 0.41; 95% CI -1.26 to 2.09; $I^2=85\%$; p=0.63). The results are shown in figure 2.

The retinal DCP VD analysis in pregnant women and controls

Al train Four studies with a total of 360 eyes (197 eyes in the pregnancy group and 163 eyes in the control group) reported the retinal DCP VD. There was minimal heterogeneity , and across studies (χ^2 =0.14, p=0.99, I²=0%). The retinal DCP VD was significantly higher in pregnant women than in controls (MD 1.53; 95% CI 0.62 to 2.44; $I^2=0\%$; p=0.0009). The results are shown in figure 2.

FAZ analysis in pregnant women and controls

technologi Four studies with a total of 360 eyes (197 eyes in the pregnancy group and 163 eyes in the control group) reported the FAZ, with minimal heterogeneity across studies $(\chi^2=3.50, p=0.32, I^2=14\%)$. The difference between the pregnancy group and the control group was uncertain (MD 0.01; 95% CI -0.01 to 0.03; $I^2=14\%$; p=0.18). The results are shown in figure 2.

CC VD analysis in pregnant women and controls

Three studies with a total of 194 eyes (82 eyes in the pregnancy group and 112 eyes in the control group) reported the CCVD. However, significant heterogeneity still existed

Table T General characteristics of the included studies in the meta-analysis								
Study	Country	Study design	Number of eyes	Age(years)	OCTA device	Scan size	Pregnancy period (weeks)	NOS score
Yildirim <i>et al</i> ¹⁸	Turkey	Cross- sectional study	Pregnancy: 40 Controls: 40	Pregnancy: 28.80±5.27 Controls: 29.23±5.39	Optovue	SCPVD, DCPVD: 6×6mm FAZ: 3×3mm	34 (28–41)	7
Hepokur <i>et al</i> ¹⁹	Turkey	Prospective, cross- sectional study	Pregnancy: 97 controls: 34	Pregnancy: 30.32±6.34 Controls: 29.38±7.03	Optovue	6×6mm	34.47±3.34	6
Pota and Apaydın ²⁰	Turkey	Cross- sectional study	Pregnancy: 41 Controls: 45	Pregnancy: 30.8±5.9 Controls: 29.7±5.9	Topcon	6×6mm	31±2	7
Chanwimol et al ²¹	USA	Prospective, case–control study	Pregnancy: 19 Controls: 44	Pregnancy: 36±7 Controls: 35±8	Optovue	3×3mm	33±3	7
Su et al ²²	USA	Prospective, case–control study	Pregnancy: 22 Controls: 23	Pregnancy: 32.27±6.96 Controls: 30.08±5.49	Heidelberg	10°×10°	33±3.59	7

...

DCP VD, deep capillary plexus vascular density; FAZ, foveal avascular zone; NOS, Newcastle-Ottawa Scale; OCTA, optical coherence tomography angiography; SCP VD, superficial capillary plexus VD.

(χ^2 =9.69, p=0.008, I²=79%). The difference between the pregnancy group and the control group was uncertain (MD 0.76; 95% CI -1.11 to 2.64; I²=79%; p=0.43). The results are shown in figure 2.

DISCUSSION

In the present study, we compared the retinal and choroidal microvascular features in pregnant women and non-pregnant controls using meta-analysis. Some previous studies have observed and elucidated the difference, but the results were controversial.^{18–22} This study proved and expanded previous research to improve the quality of the evidence. As far as we know, this is the first meta-analysis study that reports the retinal and choroidal microvascular features detected by OCTA during pregnancy. This meta-analysis indicates that the DCP VD of the healthy pregnant women was significantly higher than that of the non-pregnant controls. However, uncertain difference

was observed in retinal SCP VD, FAZ and CC VD between healthy pregnant women and non-pregnant controls.

In humans, pregnancy is accompanied by dramatic le X changes in hemodynamics, levels of various hormones and vasoactive metabolites. It has been suggested that haemodynamic changes during pregnancy occur through autonomic control mechanisms.¹¹ Furthermore, the sympathetic nervous system plays a role in haemodynamic homeostasis during pregnancy. Previous studies have proposed that a fall in systemic vascular tone primarily triggered maternal haemodynamic adaptation to pregnancy. Cardiac output rises 40% during pregnancy responding to this decrease in systemic vascular resistance and the increase of blood volume.²³ The peripheral arterial vasculature is maintained in a active vasodilation condition by continuous synthesis of endothelium-derived nitric oxide formed from L-arginine.^{12 24 25} As we know, pregnancy per se can lead to sympathetic activation despite a normal

Table 2 Evaluation of GRADE evidence quality								
Outcome indicators	MD, 95%Cl	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Upgrade quality	Quality
SCP VD	0.41 (-1.26 to 2.09)	Serious*	Serious†	No	No	Undetected	None	Low
DCP VD	1.53 (0.62 to 2.44)	Serious*	No	No	No	Undetected	None	Moderate
FAZ	0.01 (-0.01 to 0.03)	Serious*	No	No	No	Undetected	None	Moderate
CC VD	0.76 (-1.11 to 2.64)	Serious*	Serious†	No	No	Undetected	None	Low

*Absence of allocation concealment and blinding.

†The p value of the heterogeneity test was <0.1 and l^2 >50%.

CC VD, choriocapillaris vascular density; DCP VD, deep capillary plexus VD; FAZ, foveal avascular zone; GRADE, Grading of Recommendations Assessment, Development and Evaluation; MD, mean difference; SCP VD, superficial capillary plexus VD.

	pregnancy		Control		Mean Difference		Mean Difference		
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% Cl	IV, Random, 95% Cl
1.1.1 SCP VD									
Abdulmutalip2020	52.8	2.6	40	52.03	3.31	40	25.1%	0.77 [-0.53, 2.07]	
Cisil2023	42.5	3	41	40	2	45	26.3%	2.50 [1.41, 3.59]	_
Karntida2019	47.9	3.3	19	49.7	3.1	44	22.4%	-1.80 [-3.54, -0.06]	
Mustafa2021	52.78	2.5	97	52.91	2.96	34	26.2%	-0.13 [-1.24, 0.98]	
Subtotal (95% CI)			197			163	100.0%	0.41 [-1.26, 2.09]	
Heterogeneity: Tau ² =	2.45; Chi	² = 20.54	, df = 3	(P = 0.00)	001); I ² = 3	85%			
Test for overall effect: 2	Z = 0.49 (P = 0.63)							
1.1.2 DCP VD									
Abdulmutalip2020	57.73	6.12	40	56.54	5.44	40	12.8%	1.19 [-1.35, 3.73]	
Cisil2023	45.4	3.5	41	43.8	2.1	45	53.9%	1.60 [0.37, 2.83]	
Karntida2019	55.5	3.9	19	53.8	3.5	44	19.8%	1.70 [-0.34, 3.74]	
Mustafa2021	58.82	4.95	97	57.5	6.72	34	13.5%	1.32 [-1.14, 3.78]	
Subtotal (95% CI)			197			163	100.0%	1.53 [0.62, 2.44]	-
Heterogeneity: Tau ² =	0.00; Chi	² = 0.14,	df = 3 (P = 0.99)	; I ² = 0%				
Test for overall effect: 2	Z = 3.31 (P = 0.00	09)						
1.1.3 FAZ									
Abdulmutalip2020	0.33	0.1	40	0.28	0.1	40	19.0%	0.05 [0.01, 0.09]	•
Cisil2023	0.2996	0.0686	41	0.2884	0.0626	45	40.3%	0.01 [-0.02, 0.04]	•
Karntida2019	0.3	0.1	19	0.3	0.1	44	13.1%	0.00 [-0.05, 0.05]	+
Mustafa2021	0.29	0.09	97	0.29	0.09	34	27.7%	0.00 [-0.04, 0.04]	•
Subtotal (95% CI)			197			163	100.0%	0.01 [-0.01, 0.03]	
Heterogeneity: Tau ² =	0.00; Chi	²= 3.50,	df = 3 (P = 0.32)	; I ² = 14%)			
Test for overall effect: 2	Z = 1.35 (P = 0.18)							
1.1.4 CC VD									
Cisil2023	52.1	2.9	41	49.7	2.3	45	37.6%	2.40 [1.29, 3.51]	
Karntida2019	41.14	2.23	19	40.81	2.8	44	35.8%	0.33 [-0.97, 1.63]	
Su2020	54.06	3.95	22	55.02	3.78	23	26.6%	-0.96 [-3.22, 1.30]	
Subtotal (95% CI)			82			112	100.0%	0.76 [-1.11, 2.64]	
Heterogeneity: Tau ² = 2.12; Chi ² = 9.69, df = 2 (P = 0.008); l ² = 79%									
Test for overall effect: Z = 0.80 (P = 0.43)									
									-4 -2 0 2 4
									pregnancy control

Test for subaroup differences: Chi² = 11.57, df = 3 (P = 0.009), l² = 74.1%

Figure 2 Forest plot for superficial capillary plexus vascular density (SCP VD), deep capillary plexus VD (DCP), foveal avascular zone (FAZ) and choriocapillaris (CC) VD between healthy pregnant women and non-pregnant controls.

blood pressure. However, In contrast with the sympathetic activation, peripheral vascular resistance was found to be lower in normal pregnant women than that in nonpregnant women. Oestrogen may weaken the impact of sympathetic activation on vascular resistance through local mechanisms in humans.²⁴

The regulation of the ocular circulation is complex. There are substantial differences between retinal circulation regulation and choroidal circulation regulation. Retinal blood flow is mainly determined by autoregulatory mechanisms and local factors while choroidal blood flow is mainly controlled by autonomic innervation. The flow through a vascular bed is determined by both perfusion pressure and vascular resistance. However, several studies have suggested that the influence of a moderate change in perfusion pressure on retinal blood flow is negligible. So it can be inferred that mechanisms in the retinal circulation that adapt vascular resistance to changes in perfusion pressure make retinal blood flow constant. Unfortunately, the detailed mechanisms are not yet fully understood. In contrast with retina, the choroidal circulation is mainly under neurogenic control

Protected by copyright, including for uses related to text and data mining, Al train and is not autoregulated. Some researches revealed that electrical stimulation of sympathetic and parasympathetic nerves had an effect on choroidal perfusion.²⁴²⁶

, ĝ This meta-analysis indicates that the DCP VD of the healthy pregnant women was significantly higher than that of the non-pregnant controls, but uncertain difference was observed in retinal SCP VD and FAZ between healthy pregnant women and non-pregnant controls. Retinal circulation is characterised by lower blood flow than that of choroid. Retinal circulation lacks autonomic nol innervation, which shows efficient autoregulation and is mainly affected by local factors. In addition, the influence of hormones and neurotransmitters on retinal vascular 8 resistance is generally supposed to be negligible because of the blood-retinal barrier. Although pregnancy is associated with a decrease in vascular resistance and dramatic increase in blood volume, blood flow in retina is always in a constant state by autoregulation. In our meta-analysis, the difference in SCP VD and FAZ between healthy pregnant women and non-pregnant controls was uncertain. This result was consistent with the characteristics of retinal blood flow regulation. On the contrary, DCP VD of the

pregnant women was significantly higher than in the nonpregnant controls, which meant higher DCP blood flow during pregnancy. Histological data from humans and other mammalian species have illustrated great variations in the density and laminar structure of the superficial and deep retinal capillary networks.²⁷⁻²⁹ Chandrasekera et al reported that capillaries composing the SCP were derived directly from precapillary arterioles and drained directly into postcapillary venules while no evidence was found where precapillary arterioles connected directly to capillaries composing the DCP. Furthermore, in the same plane as the SCP, there were major branches of the retinal artery and vein, while at the level of the DCP, segments of retinal arterioles and venules were not seen, and the DCP was characterised by numerous closed vascular loops. Not only that, available data supported a model in which major arterial inflow occurs at the level of the SCP while venous outflow preponderantly tracks through the DCP.³⁰ In pregnancy, there is an increase in DCP perfusion density. The possible reason is that venous dilation secondary to increased blood volume and higher nitrous oxide levels predominantly occurs in DCP. The series and parallel organisation of peripapillary capillary plexuses may also make SCP and DCP different haemodynamics characteristics.³¹

The choroid is a complex structure consisting of vascular and stromal components, which are involved in physiological function and provides oxygen and nutrition for the outer layer of the retina.¹³ In the choroid, the flow per perfused volume is higher than any other human tissue, and choroidal blood supply provides 85% of total ocular blood flow.³² In addition, the ability of the choroidal vasculature to regulate its blood flow is far more restricted than that of retina because of the absence of glial cells, the considerably reduced pericyte ensheathment of the choroidal vasculature and the lack of intermediate filaments in choroidal pericytes.²⁶ Therefore, the anatomical and functional characteristics of choroid may make it susceptible to the changes in haemodynamics during pregnancy. However, uncertain difference was found in CC VD between healthy pregnant women and non-pregnant controls. Jiang *et al*¹³ reported that the CT of the healthy pregnant women was significantly higher than that of the non-pregnant women. However, Su *et al*²² found that there was no significant difference between the pregnant and control groups in subfoveal CTs, Choroidal Vascularity Index or choriocapillaris flow deficits (CC FD). When calculating CC VD, both the vascular and interstitial components of the choroid are taken into consideration. Therefore, CC VD was believed to be more trustworthy and less affected by confounding factors such as intraocular pressure, axial length and systolic blood pressure than CT.²² What's more, pregnancy-induced hormone changes may get involved. Oestrogens exert an influence on vasodilation while progesterone and reninangiotensin increase the resistance of the ophthalmic artery and its branches. Hence, the mixed effects of hormones may lead to uncertain changes in the choroid,

such as hyperperfusion, hypoperfusion of the choroid or no change at all.³¹

Our study has some limitations. First, the total sample size in the included studies was relatively small, which may make the quality of the evidence relatively low. Second, the pregnancy period of the included data was difficult to guarantee the consistency. At different pregnancy periods, changes in the retinal and choroidal microvasculature may be different. If there are enough data that focus on different pregnancy periods, further meta-analysis can be conducted. Third, except for pregnancy period, there still exists heterogeneity across the included studies. The potential causes of the heterogeneity may be study designs, eye condition (such as the dioptres of the eyes and the intraocular pressures), the symplectic of OCTA devices used to obtain the metrics, algorithms used by the different OCTA device and different generation. scan size. The heterogeneity of the included studies may threaten the validity and reliability of results. However, because of possible incomplete reporting of data, we cannot perform meta-regression to illuminate the source of heterogeneity.

In conclusion, our meta-analysis found that the DCP VD of healthy pregnant women was higher than that of nonpregnant controls. However, the SCP VD, FAZ and CC VD came to the opposite conclusion; uncertain difference was found between the healthy pregnant women and the non-pregnancy controls. Our meta-analysis revealed more convincing results by a larger sample size compared with the individual studies. In addition, our findings can help to get a deeper understanding of retinal and choroidal microvascular characteristics during pregnancy.

Contributors PZ and QS designed the study. PZ, CW and YL searched the articles and contributed to data acquisition. PZ was responsible for processing data and writing original manuscript. QS and PZ revised the manuscript. All authors read and approved the final manuscript. QS is the guarantor.

Funding This study was supported by the Natural Science Foundation of Hebei Province (No. H2022206287) and S&T Program of Hebei (22377754D).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants but the ethical approval of this meta-analysis was exempted according to the Institutional Review Board of the second hospital of Hebei Medical University. This meta-analysis used statistically processed data and did not involve data of individual patient. In addition, all included articles were approved by the institutional review board of hospital or university where the clinical trials were conducted. This meta-analysis used statistically processed data which was extracted from the published articles and did not involve data of individual patient.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content

<u>ð</u>

includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iD

Peng Zhang http://orcid.org/0000-0002-6537-8261

REFERENCES

- 1 Ouzounian JG, Elkayam U. Physiologic changes during normal pregnancy and delivery. *Cardiol Clin* 2012;30:317–29.
- 2 Mehdizadehkashi K, Chaichian S, Mehdizadehkashi A, et al. Visual acuity changes during pregnancy and postpartum: a cross-sectional study in Iran. J Pregnancy 2014;2014:675792.
- 3 Anantharaman D, Radhakrishnan A, Anantharaman V. Subjective dry eye symptoms in pregnant women-A SPEED survey. J Pregnancy 2023;2023:3421269.
- 4 Kyei S, Asiedu K, Ephraim RKD, *et al.* Meibomian gland dysfunction and lipid profile: a study among pregnant women. *Eye Contact Lens* 2021;47:598–603.
- 5 Mirzajani A, Narooie-Noori F, Amini Vishteh R, *et al.* Changes in refractive and optometric findings during pregnancy. *Med J Islam Repub Iran* 2022;36:102.
- 6 Bujor IA, lancu RC, Istrate SL, *et al.* Corneal biomechanical changes in third trimester of pregnancy. *Medicina (Kaunas)* 2021;57:600.
- 7 Özkaya D, Usta G, Karaca U, et al. Evaluation of anterior segment parameters during pregnancy. Semin Ophthalmol 2022;37:131–5.
- 8 Schocket LS, Grunwald JE, Tsang AF, et al. The effect of pregnancy on retinal hemodynamics in diabetic versus nondiabetic mothers. Am J Ophthalmol 1999;128:477–84.
- 9 Roos NM, Wiegman MJ, Jansonius NM, *et al*. Visual disturbances in (pre)eclampsia. *Obstet Gynecol Surv* 2012;67:242–50.
- 10 Conrad KP. Emerging role of relaxin in the maternal adaptations to normal pregnancy: implications for preeclampsia. *Semin Nephrol* 2011;31:15–32.
- 11 Fu Q, Levine BD. Autonomic circulatory control during pregnancy in humans. Semin Reprod Med 2009;27:330–7.
- 12 Carbillon L, Uzan M, Uzan S. Pregnancy, vascular tone, and maternal hemodynamics: a crucial adaptation. *Obstet Gynecol Surv* 2000;55:574–81.
- 13 Jiang M-S, Xu X-L, Yang T, et al. Comparison of choroidal thickness in preeclamptic, healthy pregnant, and nonpregnant women: a systematic review and meta-analysis. Ophthalmic Res 2019;62:1–10.
- 14 Kızıltunç PB, Varlı B, Büyüktepe TÇ, et al. Ocular vascular changes during pregnancy: an optical coherence tomography angiography study. Graefe's Arch Clin Exp Ophthalmol 2019.

- 15 Min W. Re-understanding optical coherence tomography angiography. *Chin J Ocul Fundus Dis* 2019;35:1–2.
- 16 Muñoz-Leyva F, El-Boghdadly K, Chan V. Is the minimal clinically important difference (MCID) in acute pain a good measure of analgesic efficacy in regional anesthesia? *Reg Anesth Pain Med* 2020;45:1000–5.
- 17 Levene T, Livingston G, Banerjee S, *et al.* Minimum clinically important difference of the social functioning in dementia scale (SF-DEM): cross-sectional study and Delphi survey. *BMJ Open* 2022;12:e058252.
- 18 Yildirim A, Kurt E, Altinisik M, et al. Evaluation of retinochoroidal tissues in third trimester pregnants: an optical coherence tomography angiography study. Eur J Ophthalmol 2022;32:501–7.
- 19 Hepokur M, Gönen B, Hamzaoğlu K, et al. Investigation of retinal vascular changes during pregnancy using optical coherence tomography angiography. Semin Ophthalmol 2021;36:19–27.
- 20 Pota ČE, Apaydin KC. Retinal and choroidal microvascular changes during pregnancy detected with OCTA. Can J Ophthalmol 2023;1–8.
- 21 Chanwimol K, Balasubramanian S, Nassisi M, *et al.* Retinal vascular changes during pregnancy detected with optical coherence tomography angiography. *Invest Ophthalmol Vis Sci* 2019;60:2726–32.
- 22 Su L, Taweebanjongsin W, Gaw SL, et al. Evaluation of the choroid in women with uncomplicated pregnancy. *Trans Vis Sci Tech* 2020;9:24.
- 23 Gyselaers W, Spaanderman M, International Working Group on Maternal Hemodynamics. Assessment of venous hemodynamics and volume homeostasis during pregnancy: recommendations of the International working group on maternal hemodynamics. *Ultrasound Obstet Gynecol* 2018;52:174–85.
- 24 Delaey C, Van De Voorde J. Regulatory mechanisms in the retinal and choroidal circulation. *Ophthalmic Res* 2000;32:249–56.
- 25 Kuate Defo A, Daskalopoulou SS. Alterations in vessel hemodynamics across uncomplicated pregnancy. *Am J Hypertens* 2023;36:183–91.
- 26 Kur J, Newman EA, Chan-Ling T. Cellular and physiological mechanisms underlying blood flow regulation in the retina and choroid in health and disease. *Prog Retin Eye Res* 2012;31:377–406.
- 27 Chan G, Balaratnasingam C, Yu PK, et al. Quantitative morphometry of perifoveal capillary networks in the human retina. *Invest* Ophthalmol Vis Sci 2012;53:5502–14.
- 28 Tan PEZ, Yu PK, Balaratnasingam C, *et al*. Quantitative confocal imaging of the retinal microvasculature in the human retina. *Invest Ophthalmol Vis Sci* 2012;53:5728–36.
- 29 Yu PK, Mammo Z, Balaratnasingam C, et al. Quantitative study of the macular microvasculature in human donor eyes. *Invest Ophthalmol Vis Sci* 2018;59:108–16.
- 30 Scharf J, Freund KB, Sadda S, et al. Paracentral acute middle maculopathy and the organization of the retinal capillary plexuses. *Prog Retin Eye Res* 2021;81:100884.
- 31 Chandrasekera E, An D, McAllister IL, et al. Three-dimensional microscopy demonstrates series and parallel organization of human peripapillary capillary plexuses. *Invest Ophthalmol Vis Sci* 2018;59:4327–44.
- 32 Sochurek JAM, Gembicki M, Grisanti S, et al. Vascular choroidal alterations in uncomplicated third-trimester pregnancy. *Tomography* 2022;8:2609–17.