



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Factors Influencing Virtual Reality Use in Paediatric Acquired Brain Injury Upper Limb Rehabilitation – A qualitative study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-083120
Article Type:	Original research
Date Submitted by the Author:	12-Dec-2023
Complete List of Authors:	Rathinam, Chandrasekar; University of Birmingham College of Medical and Dental Sciences, Institute of clinical sciences; Birmingham Women's and Children's NHS Foundation Trust, Therapy Service Farr, William; University of Cambridge Ray, Daniel; Birmingham Women's and Children's NHS Foundation Trust Gupta, Rajat; University of Birmingham; Birmingham Women's and Children's NHS Foundation Trust
Keywords:	Virtual Reality, REHABILITATION MEDICINE, QUALITATIVE RESEARCH, Developmental neurology & neurodisability < PAEDIATRICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Title

**Factors Influencing Virtual Reality Use in Paediatric Acquired Brain Injury
Upper Limb Rehabilitation – A qualitative study**

Authors

Chandrasekar Rathinam^{1,2*}

William Farr³

Daniel Ray²

Rajat Gupta^{1,2}

Affiliations

¹ College of Medical and Dental Sciences, University of Birmingham, UK

² Birmingham Women's and Children's NHS Foundation Trust, Birmingham, UK

³ Faculty of education, St Edmunds college, University of Cambridge, UK

*Corresponding author

College of Medical and Dental Sciences

University of Birmingham

Edgbaston

Birmingham

B15 2TJ

United Kingdom

Birmingham Women's and Children's NHS Foundation Trust

Steelhouse Lane

Birmingham

B4 6NH

United Kingdom

Contact (Email)

Chandrasekar Rathinam - c.rathnam@bham.ac.uk

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Chandrasekar.rathinam@nhs.net

William Farr - wjf33@cam.ac.uk

Daniel Ray - Daniel.ray2@nhs.net

Rajat Gupta - rajatgupta@nhs.net

ORCID

Chandrasekar Rathinam: 0000-0002-0049-8430

Rajat Gupta: 0000-0001-5111-7821

Authors contribution

All authors will read, provide feedback and approve the final manuscript.

Funding

This article was written as part of Topol digital fellowship programme, UK (2022 – 2023). The study sponsors were not involved in the study design, data collection, analysis and interpretation of data, and the decision to submit the manuscript for publication. Funding/grant number not applicable.

Disclosure of interest

The authors have stated that they had no interests which might be perceived as posing a conflict or bias.

Word count: 5842

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Abstract

Background

Upper limb movement difficulties in children with acquired brain injury (ABI) result in longer recovery. Intensive neurorehabilitation promotes good long-term functional outcomes. Virtual reality (VR) and video game technology are invaluable adjuncts to traditional neurological rehabilitation. VR benefits children with improved compliance and active engagement in goal-directed therapy. However, VR and video game technology are not routinely used in the NHS; it requires embedding to benefit children and their families.

Objective

VR Intervention requires development, and associated influencing factors need exploring before routine use in rehabilitation. This project aims to understand factors influencing VR use in upper limb rehabilitation.

Methods

An interpretative qualitative study utilised focus groups and 1:1 semi-structured interviews conducted in person and online with three physiotherapists, five occupational therapists, a play worker, and four participants from the young persons' advisory group. Thematic analysis was used to create model participants described as the factors influencing VR use for neurorehabilitation.

Results and discussion

Five closely related major themes and thirty associated subthemes were developed: training, knowledge, promotion, consideration of barriers, and family factors. There is a lack of knowledge and understanding about the use of VR, its limitations, and the clinicians' motivation to use it. Training packages with available VR equipment, clinical indicators, and scientific evidence are required. Staff need frequent training, logistical knowledge (uninterrupted Wi-Fi, software, hardware), and simple instruction manuals. VR needs to be embedded in routine practice to facilitate behavioural change.

Key words

Acquired brain injury; Children; Virtual reality; Rehabilitation; Thematic analysis

Strengths and limitations of this study

- This study gathered therapists' and young people's views on using virtual reality for rehabilitation.
- Technology access, time pressure, and knowledge factors limit therapists' use of technology for rehabilitation.
- Regular training and encouragement are needed to facilitate behaviour change to use virtual reality in clinical settings.
- A 4i model (improving knowledge, investing in training, infrastructure, and implementing changes) is suggested.
- Members checking was not done to improve triangulation.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

1. Introduction

Worldwide, acquired brain injury (ABI) is one of the most common causes of neuro-disability in children and young people (CYP)(1). In the UK, ABI annually accounts for 35,000 childhood presentations to emergency departments(2). Of these, 5% have moderate to severe brain injury, but with complex and unclear outcomes for the other 95%(3). A subsection of children with ABI has stroke-like presentation, and functional loss of their upper limb results in an increased dependency(4).

Following injury, upper limb function usually takes longer to recover than lower limb functions, and children often require early and intensive neurorehabilitation to achieve their maximum physical potential(5). High-intensity and repetitive practice require a high level of labour-intensive therapy, often difficult to provide due to high work demands and capacity issues(6). These children can be passive, and parents struggle to motivate them to do the prescribed exercise(7). Most children, however, do enjoy playing computer games(8). Virtual reality (VR) based computer gaming technology is emerging as a valuable adjunct to traditional neurological rehabilitation(9). It has been recently used to treat CYP with cerebral palsy (CP)(10) and ABI(11).

VR interventions need less professional input and can be programmed to suit an individual child's needs. This type of intervention helps address boredom and compliance problems, facilitates increased therapy time, and does not rely on therapeutic contact time(9). Parents are actively empowered to facilitate the active participation of children. However, VR has not been used routinely in the NHS.

A cross-sectional survey in Canada that examined the use of VR by therapists reported that 46% of therapists had VR experience, and 15% of them had used it to treat people who had suffered a brain injury. This study found that lack of funds, space, time, support staff, and appropriate clients was a barrier to using VR; they found client motivation, therapists' knowledge, and management support as the facilitators to adopt VR in regular practice(12). Farr et al., (2020) examined the paediatric physiotherapists' and occupational therapists' learning needs of VR in the UK and noted that only 11% of therapists reported current use of VR with their patients; managers have little or no influence on VR adaptation, and the lack of IT support is another barrier(9). Farr *et al.*, (2020) recommended that tailored support is required to facilitate VR adaptation.

In rehabilitation, treatment goals have shifted from managing the body structure to managing activity limitations and participation restrictions(13). Our previous co-production work with children who have neurological motor disorders showed that the children found formal exercise boring and lacking in variety; they found that it was hard to fit their exercise programme into their daily life; it competed with academic needs; it isolated them from their peers; and it was harder to do them as they grew older. They recommended play-based activities, video games, and VR-based activities that fit in with their life and would be more acceptable because they could play with their friends/peers and siblings. In line with these outcomes, therapists must evolve from conventional hands-on therapy to next-generation technology-based therapy programmes that facilitate personalised care(6). VR can be seamlessly embedded in integrated services to improve patient care and efficiency and promote functional

Enseignement Supérieur (ABES) .
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

outcomes[14]. For this, VR Intervention requires development, and the associated influencing factors need exploring before it can be used routinely in rehabilitation.

This study aims to understand the factors influencing VR use in upper limb rehabilitation for children with ABI and determine the infrastructure support and training required for therapists to plan and deliver personalised VR programmes.

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2. Methods

2.1 Design

A qualitative approach was used to explore the participants' experience, focusing on their views on VR for upper limb rehabilitation. The authors followed the Standards for Reporting Qualitative Research (SRQR) Checklist (15) (appendix 1).

2.1.1 Ethics

This study was registered with the Birmingham Women's and Children's Hospital Research & Development office and in accordance with the UK National Research Ethics Service guidance (REC: 16/BWC/LA/Rathinam).

2.2 Patient and Public Involvement

Patients and public were not involved as this study aimed to understand the therapists need to use VR in clinical settings.

2.3 Participants

Purposive sampling strategy was used by inviting physiotherapists (PT) and occupational therapists (OT) experienced in treating children with ABI at Birmingham Children's Hospital (BCH) and Oxford Children's Hospital, UK. We also invited play workers (PW) and BCH's Young Persons' Advisory Group (YPAG) members through their coordinator. All the participants had given consent. The participants' (therapists and play worker) professional experience and using virtual reality for treatment in years were given in table-1. Appointments were scheduled to meet the participants

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES)

in person or through an online platform (Microsoft Teams); the coordinators from CBIT and YPAG were present for some of the sessions.

2.4 Procedure

Four focus groups or 1:1 semi-structured in-depth interviews were completed, each session lasting between 60 and 90 minutes. A semi-structured interview guide was used to facilitate discussion. Participants were asked about their VR experience, barriers, adherence factors, and reflections. CR undertook all interviews, and the Interviews were recorded with permission. Written notes were taken simultaneously. Reflexivity was maintained by CR by diary keeping during the interview and analysing the transcript.

2.5 Analysis

Interviews were transcribed in full. Thematic analysis was conducted following Braun and Clark's (2006) guidelines. A multistage approach includes familiarisation, indexing, sorting, developing an initial framework, applying the framework, and charting the data. Interpretation followed to analyse the collected data(16). An initial line-by-line inductive coding was performed verbatim. This enabled themes to evolve from the data related to the project aim. These themes were categorised, and five major themes and associated but interrelated sub-themes were developed. Trustworthiness was ensured by triangulating the data gathered with the previous study data (WF). Analytical rigour was maintained with an audit trail and aligning the coding between the authors (CR and WF).

3. Results

Twelve participants (two PTs, five OTs, a PW, and four participants from YPAG along with their coordinators) consented and participated. All the professionals were aware of the use of VR for rehabilitation, but very few had some experience, and most reported no experience using it for treatment purposes. A model participants described as the factors influencing VR use in neurorehabilitation was created (Figure 1), and verbatim quotes demonstrating pertinent quotes for individual themes are presented below.

3.1 Knowledge

VR is considered a relatively new domain, and lack of knowledge and awareness are two factors that limit therapists' use in the clinical setting.

3.1.1 VR Tools

Participants were aware of the use of VR for treatment purposes. Many assumed that VR was an immersive technology using goggles or a headset. They needed to be made aware of the non-immersive style. They were keen to understand the different VR types currently available for rehabilitation, how they work, and how they can be used for rehabilitation.

"I would need to understand how virtual reality works. Would it be a headpiece? Would it be a kind of feedback thing on a screen?" (EW; PT)

3.1.2 Evidence base

Participants were unaware of the existing evidence supporting the use of VR. Having the evidence will assist them in arriving at an informed decision, mainly if the evidence shows validity, transferability, and generalisability. They asked for specific information about the indications, contraindications, limitations, patient position, and the minimal ability required to engage children in a VR session. They asked for information about different VR techniques, timings, progression mechanisms, expected changes or improvements, desired outcomes, treatment effectiveness measurements, and any other factors that support the clinical utilization of VR.

"....knowing the evidence base, and what the validity of it is, and if they have been a particular success in particular patient groups, any limitations on kind of age or cognitive ability or physical ability as to what they can, who can use it, like knowing how to make a good patient selection." (SH; PT)

3.1.3 Decision making

Choosing any modality for treatment underpins the relationship between clinical reasoning, evidence base, and equipment availability. This is also related to knowing a child's condition and the parents' capability. Participants were keen to know how and when VR should be identified as an intervention. This will be a clinical decision by the treating therapist, following a good-quality assessment of the child and identifying the goals of all concerned. Only then can it be determined whether VR equipment will assist in achieving the goals.

"....a good quality assessment, that means that we've gotten to know that child and their context and their family...." (PW; OT)

3.1.4 Desired games

The participants expressed their views on the type of games to choose and which to avoid. The games should give a positive experience, some of which can be achieved through good graphics, up-to-date software, and equipment, giving them the feeling that they are playing proper computer games. There need to be various games to avoid boredom, which naturally progresses to different levels as the child progresses. They should have the option to change the features so children can easily play without tiring, and a bigger switch control mechanism.

“....variety of things (games) so that they don't just play one game and then they are bored of playing that one, they have a good variety, and they can play.”

(CJ; OT)

Participants had suggested avoiding games that are gender stereotypes, horror games, and car crash games, particularly for car accident victims. Therapists also need to be mindful of not setting up games that can show what they cannot do to avoid giving the child negative feelings, a sense of failure, and a risk of disengagement when they find the games too hard.

“....acquired brain injury (be)cause, let's say it happened in a car crash, then you bring this piece of kit, and you're playing a car game. It's gonna be a bit triggering....” (BD; YPAG)

3. 2 Training

Lack of training has been identified as an essential factor limiting VR use in clinics. Participants highlighted their training needs related to the child’s functional loss following ABI.

Enseignement Supérieur (ABES) .
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

3.2.1 Loss of function

Some children lose their motor function following ABI. This is especially true for those who have a stroke-like presentation or tone abnormalities. These children struggle to use their hands for everyday functional activities, impacting their independence and disrupting their development. At times, they struggle to entertain themselves. Participants were keen to understand the barriers to using VR during the acute phase. These include having a cannula in their hand, frequent seizure episodes, and not having part of their skull following brain surgery.

“....if they have lost the use of their hands, they really struggle with....” (LP; PW)

3.2.2 Emotion

Emotional regulation is often challenging, and these children often get frustrated when they have lost independence or skills following ABI. Although they know what happened to them, they often do not reflect on why. The psychological impact includes feeling low and scared, and they wish to go home instead of being in the hospital. They become aware of what they can play despite cognitive and learning problems but are depressed as they cannot manipulate small controls in their VR consoles.

“....They've just got a brain injury. They feel grotty. They're scared. They don't want to be in hospital, but they don't really quite know what they want....” (AM; OT)

3.2.3 Fatigue

Fatigue is a common problem experienced by children with ABI from the early stage, and most of them tend to have this problem on a long-term basis. The participants want to learn the early signs of fatigue or tiredness to stop the session or make parents recognise fatigue and differentiate between tiredness and boredom. This will also help negate fatigue by pacing, changing the game's complexity, and modifying the VR's intensity.

"most of the time, they have got tired....and I've had to stop." (LP; PW)

3.2.4 Resource person

Participants have suggested a resource person within their team to keep their skills, be responsible for ongoing training, and demonstrate that VR works. The resource person can also take responsibility for reviewing the VR programme regularly and ensure that the family is competent in delivering it while in the ward. The participants commented that they need time to practice VR and a higher training volume to maintain competency so that it can become second nature. They wanted yearly and frequent refresher training programmes to ensure learning reinforcement.

"We would want to always ensure that we've demonstrated and that the person is using it in the correct way...." (PW; OT)

3.2.5 Guidance Workshops & Video

Participants suggested frequent workshops for therapists, parents, and children, along with pre-course materials about VR systems and how they work to educate people. The workshop method was considered helpful as it offers practical opportunities to work with the VR sets, reinforces learning by repetition, and provides

ongoing support. Journal club and keeping abreast of the research and constant refresher will minimise difficulties in using the equipment.

“....having a field training module where people could see this sort of troubleshooting guide. So the basics of how to connect to the Internet, Bluetooth, troubleshooting....” (IW; YPAG)

YPAG participants suggested a YouTube channel containing training videos for parents, children, and clinicians for VR training. Video content includes tutorials such as instructions on setting up the VR, how to play, troubleshooting, and frequently asked questions and answers.

“....could have like a YouTube channel and it's basically like to the people (who) have already used it, find out their most common problems and then you can specifically have a video that (shows) how to fix this.” (BD; YPAG)

3.2.6 Learning content

As part of the training, the participants suggested that the learning content should contain VR's benefits, clinical utility, indications, and suitability for different age ranges, settings, and game variety and complexity levels. Additional information, such as how VR can be used in a therapeutic way to treat other problems, collecting, analyzing, and interpreting data to see what the patient has done, and how VR aligns with other treatments, were also required. The learning content is expected to include further details about ease of use, difficulties, pros and cons, and treatment dosage for optimal outcomes. If specific VR equipment has been used, the participants were interested in learning the manufacturer's data, equipment information, and repetition altering mechanism to avoid fatigue, charging, and readjusting the units. From the patient's

perspective, the participants needed further training in goal setting, how VR fits within their rehabilitation programme, and the functional activities and muscle groups they were exercising. This information was considered beneficial to train families to use VR under the guidance of therapists.

3.2.7 Learning style & walk-throughs

The participants expressed different learning styles and strategies. Although a training manual is available, it does not allow the learner to ask questions, and they prefer learning on the job and training in person. Some expressed the need to learn personally and to use VR frequently, which would help to reinforce their learning and increase their confidence. Demonstrations and sitting with experienced users were the preferred methods because it would allow them to ask questions and practice under supervision. Participants wanted much hand-holding, slowly explaining the functions of it and walking through the game, especially when they started using it.

“I would like just some guidance on what to do with them and how far to push them, and that would be great for my training needs....” (LP; PW)

3.3 Barriers

3.3.1 Habitual practice

Therapists are accustomed to working with minimal resources. Therefore, despite availability, technology does not usually come to mind as a treatment option. For example, some of the therapists still draw their exercise programmes instead of using computer-generated programmes. Any new treatment technique requires time for evaluation and reflection. If the technology is not working, they prefer to return to old treatment techniques as they need more time to investigate and troubleshoot. This

means that the new technology is not used frequently, which leads to a lack of familiarity and a decreased inclination to use it without further training.

"....I am still drawing stickman for the exercise programme because trying to get to the computer to log on to (write an exercise programme is) hard to navigate and crashes all the time. When you know your patient is improving so quickly and you will need to redo this exercise programme into two days actually just stops me using it all together...." (SH; PT)

3.3.2 Generational & Self-perceptions

Some commented that they were old-generation therapists and too old to try VR. They felt that the generational difference between therapists and children is vast, and there is a gap in knowing what the children like and how to play with them using VR. They don't understand VR and don't know how to use it. They lacked the experience and confidence to use it for therapy purposes. Some therapists and YPAG participants who had tried a VR headset reported feeling sick and disliked using it. Those who had tried VR reported that they were not technology literate, felt rubbish, and needed to remember about using them.

"....I'm not a gamer; gaming (has) been around for a long time, but I'm not one. But actually, the immersion in technology and the development of the internet and everything has really only been in my adult life." (EW; PT)

3.3.3 Logistics & Complex manual

The participants reported many logistic issues. For example, communication between part-time staff, such as sharing information and hand-over, is problematic from a staffing perspective. Having a designated safe space is a challenge, as therapists

often need guidance on where to treat these children or where to store the VR equipment. Connecting the VR units to the internet is a challenge, and the participants felt that having easy access to IT services may overcome this issue. Although manuals are available, participants reported that these are complex, and hard to navigate and find the solution. Therapists wanted a simple laminated crib sheet or a quick starter guide explaining troubleshooting, screenshots of what has been shown in the manual, set-up, games, and calibration techniques.

“....the problem with the manuals and things are so long often that you can't find the thing that you're looking for..... having some crib sheet that quickly tells you this would help with this aspect....” (AM; OT)

From the equipment perspective, some need to be set up whenever the children play a different game. Some find it hard to grip the VR sets and need to readjust their handhold for each new game. For the complex VR sets, the children need someone around them to set up the programme, motivate them, and engage in the games for therapy. This is a challenge after their allocated time with the therapists, particularly over the weekend. This usually results in children not doing anything over the weekend. Besides, most VR gadgets were made for typically developing children, and very little equipment is available for children with disabilities. When children with ABI tried to use the usual consoles, it became more challenging for them to manipulate, and they tired quickly. Most children had prior experience playing VR games with rich visual content, but rehabilitation-focused games are generally primitive; hence, children quickly lose interest.

Enseignement Supérieur (ABES) .
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

"...kids these days will have high expectations from the point of view of graphics and action quality because of what they see in films and games." (SH; PT)

3.3.4 Time factor

Lack of time is a common problem expressed by most of the therapists. They acknowledged that they need dedicated time for training to use VR properly. Despite the availability of the systems, therapists need more training time and clinical time to use them appropriately.

"....but often we have to reset it 3, 4, 5 times to make sure that (all) the Wi-Fi connectors are actually connected. So, for one successful session, it's about 2 hours. So that means 100% more time than you (usually spend)...." (AM; OT)

3.3.5 Technology

Many participants reported that poor Wi-Fi connectivity in their hospital limits their chance to use VR. They commented that the connection would not work or the equipment would not pick up the signal. Having reliable network connectivity is vital if VR is used in the hospital.

"I think the practical difficulties in the hospital is lack of time, space, Wi-Fi, general resources (which) make it very difficult to be consistent in anything that needs a set-up" (CJ; OT)

Participants believed that the technology constraints listed above often put them off. It requires a lot more support if it is to be used effectively, which defeats the purpose of why VR was introduced in the first instance.

1
2
3 *"....if the IT set up for this sort of thing was not easy to use and not reliable, that*
4 *would make me use other adjuncts instead.... I'll spend an hour setting it up*
5 *and then it won't work anyway" (SH; PT)*
6
7
8
9

10
11
12 **3.3.6 Change management**
13

14 Participants acknowledged that people take the path of least resistance and that they
15 generally do not like change. They require convincing and demonstrating the utility,
16 as people are not tech-savvy. They suggested example videos of VR in practice,
17 showing how children used it and the outcome achieved.
18
19

20 *"....it's a change in practice, so people don't like change. Particularly and will*
21 *only engage if the change is managed, almost like a change management*
22 *way....it's about saying, 'this could help; why don't you give a (try)'." (EW; PT)*
23
24
25
26
27
28
29

30
31
32
33 **3.3.7 Accessibility**
34

35 Lack of space is an essential factor that limits the use of VR. Leaving the system close
36 to the bedside in some wards takes a lot of work. In some places, expensive units are
37 available, but therapists have yet to be trained to use them. Sometimes, more minor
38 issues, such as missing cables or sets needing to be charged more, caused
39 embarrassment when the session was disrupted. Many therapists felt excited initially
40 to use VR but became reluctant because of accessibility issues; therefore, sustaining
41 enthusiasm is difficult.
42
43
44
45
46
47
48
49

50 *"....it's also about how we've gotta make it accessible to everybody. In our*
51 *department, we find all the expensive kits that once were the pride and joy and*
52 *are now stuck at the back of a cupboard because nobody can remember how*
53
54
55
56
57
58
59
60

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

to set up the Wii or nobody's got the key to get the search insert. So we've got all these pieces of kit, but I've never seen them used." (MP; OT)

3.4 Promotion & Consideration

3.4.1 Motivation for staff & children

The YPAG suggested different strategies to promote VR for rehabilitation purposes. These included age-appropriate games, offering taster sessions, encouraging games with parents or siblings, challenging others, or beating their previous scores. Watching other children enjoy VR may enthuse disinterested children. Promoting VR success stories online, getting people to talk about their experience, having competitions by playing against each other with a leader board, and playing interactively can give reassurance and motivation.

"....you should get other kids to try it out in front of kids that don't wanna do it yet...." (FH; YPAG)

3.4.2 Reward/feedback

From the therapists' perspective, VR is not a universal treatment provision but an additional tool that can be used with their usual treatment. Positive feedback is always a good motivational factor. This can be the visual displays on the game, a certificate for reaching certain milestones, or purely verbal feedback.

"I definitely like that when they get the feedback as well, like party, pop sound effects or something just to kind of go well, you know, it's that big 'Well done! You've just done that level'." (AM; OT)

3.4.2 Perceived benefits

All the participants acknowledged that children love computer games and VR. They can spend hours playing as it is enjoyable, fun, engaging, and interactive. Therapists felt that VR helps to improve the child's engagement, which can facilitate movements, increase movement control, and facilitate stability and function. The end products are improved performance, grip strength, muscle length, satisfaction, spasticity management, and function improvement underpinned by component skill development. VR helps to distract the children, get them moving and enable them to do their therapy without realizing that they are doing it.

"Children love gaming, so I think it's a great way to distract them and actually get them a bit more moving." (FH, YPAG)

VR can also be used for motivation instead of therapy, especially for fun if they do not want to do anything in the ward. VR also has the potential to make hospitals more interesting, turning a 'no' person into a 'yes' person by trusting their therapists. It empowers parents to improve their independence.

"He was smiling and laughing, and that was the first time I'd seen that.... you're not going to do anything horrible and that just playing computer games and that did have a knock-on effect as well to his positivity in other sessions." (AM; OT)

3.4.3 Confidence

Participants who had prior experience using VR were confident in turning on Bluetooth, adjusting tables to use VR, deciding suitable VR games for different age groups, selecting games' complexity, progressing to the next level if the child achieved the target, using different movements and techniques to control VR and helped them to use both hands.

"I'd like you to guide on how to do it. So I've been shown now, but I wouldn't ever use equipment or do something with a child unless I've been shown." (LP; PW)

3.4.4 Prior experience

Some therapists had experience in using VR and noted that it was beneficial. It was used to improve attention and concentration, engagement, movements, and strength. They learned that a child does not need much movement to operate VR to use it at a low level. However, the lack of Bluetooth connectivity limited the VR unit's use for treatment and led to frustration. Children lost interest quickly when the system did not work, and it compromised other aspects of therapy. Apart from connectivity, they have noted other challenges, including children being competitive and making it too difficult for themselves. Therapists had assumed that parents understood the treatment goals and could follow what they were seeing, but this was only sometimes the case.

"I've had that a few times, and then they lose interest really quickly, then they lose their faith in you knowing what you're doing because the whole system is just not working. So, if you're not careful, that can rub off on other aspects of your therapy, and you don't want that." (CJ; OT)

3.4.5 Deliberation

Participants from the YPAG group suggested that the commercially available VR games they are playing are fast-moving, high-motion simulation, and have loud sound effects. These games will be challenging to play for children with ABI due to their slow reactions; the effects of flash and sound effects on eyes and ears, especially when

most already suffer from headaches. Therapists deliberated about whether they could advocate screen time for the acute ABI children, when it would be appropriate to engage them in VR, and what games and equipment would kindle a child's interest and maintain their motivation to exercise and facilitate them taking responsibility. They would want to ensure that the child and their family perceived it as therapeutic and that barriers to its use for rehabilitation were minimised.

“....it's not good for an injured and healing brain to have a lot of screen time because of the way it fires up all your neurones....” (EW; PT)

3.4.6 Grading & Adaptation

Children with ABI experience various complex challenges, including slow reaction speed; therefore, adapting VR games is vital to make it inclusive for everyone. Adaptation of VR requires consideration of a child's age range, physical and cognitive ability, concentration and engagement, communication issues, sensory impairment, and hearing and visual impairment. In the acute phase, the children struggle to be in a busy ward environment where the noise and other distractions limit their focus in therapy sessions. These children will require both the games and their environment to be adapted. Instructions should be clear and concise for all parents to understand.

“In terms of understanding families and making sure that they can understand the instructions.... you need to adapt how you gave the instructions or how the interface worked....” (EW; PT)

3.5 Family factors

3.5.1 Family Needs

Digital illiteracy and digital poverty are essential factors, as some families and parents may need access to technology. During the acute rehabilitation phase, families struggle to cope with sudden changes, and it was considered hard to expect them to engage their children in VR programmes.

"...thinking about high incidences of and younger parents, high incidences of parents who might not have access to technology, high incidences of parents who might have needs, learning disability, physical disabilities themselves. How are we expecting them to carry over...." (BH; OT)

3.5.2 Parenting culture and negotiation

Parenting covers a massive spectrum that is influenced by cultural and historical context; any technology intervention should fit meaningfully around the family culture and belief system. Having cultural sensitivity around parenting and boundaries is required. One aspect of this that therapists observed was the varying approaches to technology access and screen time. Parents must see the value and feel confident that VR will support their child's rehabilitation. Therapists suggested negotiating a set-up programme with the parents.

"....different parents might have different parenting approaches relating to access to technology and screen time, and one of the jobs we'd have to do to set up a treatment program is negotiate that with them." (MP; OT)

3.5.3 Goals

Engaging family and children around personalised goal setting is considered paramount. The therapist's role is to decide whether VR helps achieve the agreed goals.

“....we don't want to set people up with a device that measures their failure because that's really disheartening.” (PW; OT)

3.5.4 Structure to family

Families need to use VR for the well-being of their children, and they require a structure around the use of VR. This includes clarity about how VR will be used, a timetable for VR games, and limiting the play time and frequency when playing complex games. Participants initially recommended playing for 10 minutes and building up if the children were eager to play further. They have suggested playing between 5 and 45 minutes, morning and evening, and having a break between sessions.

4. Discussion

This scoping work is based on the perceptions of therapists, clinicians, YPAG members and people in an NHS setting when using VR for rehabilitation. Five major themes relating to issues around knowledge, training, and implementation emerged. Four key principles ('4i') were developed, which need consideration by professionals when using VR in upper limb rehabilitation (Figure 2). Those four principles structure the discussion.

4.1 Improving knowledge

Evidence-based intervention is the current principle that enables clinicians to use interventions with confidence(17). To facilitate VR intervention, the NHS workforce requires extensive investment to improve awareness, knowledge, and convincing evidence base in VR use. Although VR technology has been used for rehabilitation for more than two decades, there are not enough high-quality studies encouraging clinicians to utilise VR in routine practice(9). Commercially available VR equipment has not been used by the ABI population and requires adaptation of VR equipment. It may not be possible to conduct research to the level of RCTs, especially under individual conditions; therefore, translating the knowledge gained from other conditions, such as adult stroke and children with CP should be considered within the ABI population.

4.2 Invest in training

Therapists have trained to be 'hands-on' and require considerable and extensive training to adapt technology for treatment. Therapists often feel uncomfortable trying

new technology as part of their practice due to their age perceptions(12). Introducing new technology requires considerable training to change behaviour[12]. Various learning and training models and descriptors of available VR equipment, clinical indicators, and participants' experience need consideration when designing training packages. VR needs to be embedded in routine practice to facilitate behaviour change. Tact *et al.*, (2021) have suggested a digital adaptation framework for the NHS workforce, and it can be considered for technology-related training and education(18). Technology-based intervention must be a core skill gained through graduate training so future therapists can become early adopters(19,20).

4.3 Infrastructure

Basic infrastructure provision will eliminate many barriers to using VR for rehabilitation(21). Early adopters or technophobes within the workforce are at risk of abandoning VR if the required resources, such as uninterrupted Wi-Fi, updated software, and user-friendly hardware, are not readily available. Participants perceive that much of the available rehabilitation-focused VR equipment was heavy, came in universal/uniform sizes only, was hard to operate, unstable, and time-consuming. Finally, attrition of technology limits use due to poor longevity, and a lot of equipment needs upgrading in healthcare settings due to ongoing cost factors associated with degradation over time(22). The introduction of digital technology, including VR, should take place alongside investment in the stability of the approach to rehabilitation, including the required infrastructure should be in place.

4.4 Implementing change

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Our findings suggest that human factors play an essential role in digital adaptation. Various factors related to affordability, digital poverty, and the family's readiness to try technology as a tool significantly limit the uptake of VR for rehabilitation. The NHS has a digital technology framework for allied health professionals(23) and a long-term plan for digitally enabled care(24). However, the embodied reality is far behind and in relative infancy. If NHS organizations commit to embedding digital interventions as a high priority, and investing heavily in their workforce and resources, technology-based interventions such as VR will likely reach end users. Therapists are also responsible for changing their rehabilitation strategies by moving away from traditional models and adopting positive approaches of early adopters, which will lead to the evolution of future leaders who will crystallise further embedding of change. For example, a group of practitioners have used VR for ABI rehabilitation and noted the improvement and engagement from children. A recent survey indicates therapists' higher confidence and motivation when using digital technology(18). This indicates either a step change in therapists' age or a sea change in the UK's commitment to health services. Either way, it is a positive sign.

Our findings enable us to understand factors that need addressing in the implementation of VR for ABI rehabilitation. VR intervention can be developed by addressing the issues identified in this work, and a future feasibility study is needed to see if VR can be used for upper limb rehabilitation in children with ABI.

Limitations

This study did not report the parents' views of using VR, and this could add value to the themes developed. Members checking was not tried due to limited time availability

for the clinicians, which compromised the triangulation. Only two therapists and a play worker who took part in this project had some experience, and the remaining therapists' views appear to be their perception rather than their experience of using VR. Generalisability is not possible

Conclusion

The use of VR technology for upper limb rehabilitation of children with ABI depends on the investment in training for health professionals to improve their knowledge and thereby implement change in rehabilitation techniques. We must invest in improving knowledge, repeated training, and positive behaviour change among health professionals to use VR technology for upper limb rehabilitation of children with Acquired Brain Injury.

Authors Contribution

- CR, WF, DR and RG: involved in study conceptualisation and preparing final manuscript
- CR and WF: responsible for study design
- CR: responsible for data collection
- CR, WF: responsible for data interpreting and coding
- CR: guarantor of the study
- All authors read, provided feedback and approved the final manuscript

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Acknowledgement

The authors would like to thank Sharon Machin (Coordinator, Child Brain Injury Trust); Ian Wood (Coordinator, YPAG, Birmingham Children's Hospital); parents; Bethany Dennis, Fathih Hassan, Kirsten Duffie and Taffi Matts (YPAG participants, Birmingham Children's Hospital); Elizabeth Wright, Shirley Hallett, Ben Harris, Poppy Walters, Madeline Paul and Lisa Barrett (Birmingham Children's Hospital); Anne Myhill and Charlotte Jones (Oxford Children's Hospital) for their help and participation in interviews/focus groups. The authors also thank Ms. Janet Peirson, Retired Physiotherapist, Cambridge and Dr Vikram Mohan, Senior Lecturer, Department of Rehabilitation and Sports Sciences, Bournemouth University for their constructive comments and preparation of this script.

References:

1. Dunford C, Cobbold C, Ray I, Wales L. The Information Gap for Children and Young People with Acquired Brain Injury. *Dev Neurorehabil*. 2020 Jan;23(1):1–8.
2. Rathinam C, Mohan V, Bill P, Yates D, Gupta R, Peirson J. Impact of transcranial magnetic stimulation on motor function in children with acquired brain injury: a scoping review protocol. *BMJ Paediatr open*. 2023 May;7(1).
3. Trefan L, Houston R, Pearson G, Edwards R, Hyde P, Maconochie I, et al. Epidemiology of children with head injury: a national overview. *Arch Dis Child*. 2016 Jun;101(6):527–32.
4. Abgottspon S, Steiner L, Slavova N, Steinlin M, Grunt S, Everts R. Relationship between motor abilities and executive functions in patients after pediatric stroke. *Appl Neuropsychol Child*. 2022;11(4):618–28.
5. J. S, S. J, C. C, H. X. Virtual Reality for Pediatric Traumatic Brain Injury Rehabilitation: A Systematic Review. *Am J Lifestyle Med* [Internet]. 2018; Available from: <http://europepmc.org/articles/pmc6933564?pdf=render>
6. Rathinam C. Virtual reality for neurorehabilitation: hopes and challenges for therapists. *Dev Med Child Neurol*. 2021 Apr;63(4):370–1.
7. Rathinam C, Mohan V, Peirson J, Skinner J, Nethaji KS, Kuhn I. Effectiveness of virtual reality in the treatment of hand function in children with cerebral palsy: A systematic review. *J Hand Ther*. 2018 Jul;
8. Choi JY, Yi S-H, Ao L, Tang X, Xu X, Shim D, et al. Virtual reality rehabilitation in children with brain injury: a randomized controlled trial. *Dev Med Child Neurol*. 2021 Apr;63(4):480–7.
9. Farr W, Green D. Lab to Rehab: Is virtual reality being adopted clinically for

- children? In: 13th ICDVRAT with ITAG. 2020.
10. Choi JY, Yi S, Ao L, Tang X, Xu X, Dain S, et al. Upper extremity rehabilitation using a virtual reality system in children with brain injury including cerebral palsy: randomized controlled trial. *Dev Med Child Neurol*. 2020;
 11. Shen J, Johnson S, Chen C, Xiang H. Virtual Reality for Pediatric Traumatic Brain Injury Rehabilitation: A Systematic Review. *Am J Lifestyle Med*. 2020;14(1):6–15.
 12. Levac D, Glegg S, Colquhoun H, Miller P, Noubary F. Virtual Reality and Active Videogame-Based Practice, Learning Needs, and Preferences: A Cross-Canada Survey of Physical Therapists and Occupational Therapists. *Games Health J*. 2017 Aug;6(4):217–28.
 13. Ryan JM, Cassidy EE, Noorduyt SG, O'Connell NE. Exercise interventions for cerebral palsy. *Cochrane Database Syst Rev* [Internet]. 2017;(6). Available from: <http://dx.doi.org/10.1002/14651858.CD011660.pub2>
 14. Haun JN, Cotner BA, Melillo C, Panaite V, Messina W, Patel-Teague S, et al. Proactive integrated virtual healthcare resource use in primary care. *BMC Health Serv Res* [Internet]. 2021;21(1):802. Available from: <https://doi.org/10.1186/s12913-021-06783-9>
 15. O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for reporting qualitative research: a synthesis of recommendations. *Acad Med*. 2014 Sep;89(9):1245–51.
 16. Byrne D. A worked example of Braun and Clarke's approach to reflexive thematic analysis. *Qual Quant* [Internet]. 2022;56(3):1391–412. Available from: <https://doi.org/10.1007/s11135-021-01182-y>
 17. Kumah EA, McSherry R, Bettany-Saltikov J, van Schaik P. Evidence-informed

- practice: simplifying and applying the concept for nursing students and academics. *Br J Nurs*. 2022 Mar;31(6):322–30.
18. Tack C, Holdsworth L, Wilson A, McComiskie E, McCabe P, Wilkinson W, et al. Digital competency: a survey of UK allied health professionals. *Br J Healthc Manag* [Internet]. 2022 Jul 28;28(8):1–15. Available from: <https://doi.org/10.12968/bjhc.2021.0123>
19. Aungst TD, Patel R. Integrating Digital Health into the Curriculum- Considerations on the Current Landscape and Future Developments. *J Med Educ Curric Dev*. 2020;7:2382120519901275.
20. Parsons D, MacCallum K, Schofield L, Johnstone A, Coulter S-K. Next-Generation Digital Curricula for Future Teaching and Learning BT - Emerging Technologies and Pedagogies in the Curriculum. In: Yu S, Ally M, Tsinakos A, editors. Singapore: Springer Singapore; 2020. p. 3–19. Available from: https://doi.org/10.1007/978-981-15-0618-5_1
21. Glegg SMN, Levac DE. Barriers, Facilitators and Interventions to Support Virtual Reality Implementation in Rehabilitation: A Scoping Review. *PM R*. 2018 Nov;10(11):1237-1251.e1.
22. Williams MG, Stott R, Bromwich N, Oblak SK, Espie CA, Rose JB. Determinants of and barriers to adoption of digital therapeutics for mental health at scale in the NHS. *BMJ Innov* [Internet]. 2020 Jul 1;6(3):92 LP – 98. Available from: <http://innovations.bmj.com/content/6/3/92.abstract>
23. England NHS. A digital framework for allied health professionals. London: NHS England; 2019.
24. Glasper A. A long-term plan for embracing digital healthcare technology. *Br J Nurs*. 2019;28(3):204–5.

Initial	Profession	Experience (years)	
		Professional	VR use
BH	OT	10	6
PW	OT	6	0
MP	OT	20	0
AM	OT	20	0
CJ	OT	4	0
EW	PT	32	4
SH	PT	17	0
LP	PW	4	1

Table-1: Number of professional and virtual reality (VR) use experience among the participants in years (OT – Occupational Therapist; PT – Physiotherapist; PW - Play Worker




Figure 1: Model for factors influencing virtual reality use for upper limb rehabilitation for children with acquired brain injury




Figure 2: 4i model to use implement VR for upper limb rehabilitation

Appendix – 1: Factors Influencing Virtual Reality Use in Paediatric Acquired Brain Injury Upper Limb Rehabilitation - A qualitative study

Standards for Reporting Qualitative Research (SRQR) Check list

O'Brien B.C., Harris, I.B., Beckman, T.J., Reed, D.A., & Cook, D.A. (2014). Standards for reporting qualitative research: a synthesis of recommendations. *Academic Medicine*, 89(9), 1245-1251.

No.	Topic	Item	Page
	Title and abstract		
S1	Title	Concise description of the nature and topic of the study identifying the study as qualitative or indicating the approach (e.g., ethnography, grounded theory) or data collection methods (e.g., interview, focus group) is recommended	Title page-1
S2	Abstract	Summary of key elements of the study using the abstract format of the intended publication; typically includes objective, methods, results, and conclusions	Title page - 3
	Introduction		
S3	Problem formulation	Description and significance of the problem/phenomenon studied; review of relevant theory and empirical work; problem statement	1 - 2
S4	Purpose or research question	Purpose of the study and specific objectives or questions	3
	Methods		
S5	Qualitative approach and research paradigm	Qualitative approach (e.g., ethnography, grounded theory, case study, phenomenology, narrative research) and guiding theory if appropriate; identifying the research paradigm (e.g., positivist, constructivist/interpretivist) is also recommended	4
S6	Researcher characteristics and reflexivity	Researchers' characteristics that may influence the research, including personal attributes, qualifications/experience, relationship with participants, assumptions, or presuppositions; potential or actual interaction between researchers' characteristics and the research questions, approach, methods, results, or transferability	Not given
S7	Context	Setting/site and salient contextual factors; rationale ^a	
S8	Sampling strategy	How and why research participants, documents, or events were selected; criteria for deciding when no further sampling was necessary (e.g., sampling saturation); rationale ^a	4

S9 Ethical issues pertaining to human subjects	Documentation of approval by an appropriate ethics review board and participant consent, or explanation for lack thereof; other confidentiality and data security issues	4
S10 Data collection methods	Types of data collected; details of data collection procedures including (as appropriate) start and stop dates of data collection and analysis, iterative process, triangulation of sources/methods, and modification of procedures in response to evolving study findings; rationale ^a	4 - 5
S11 Data collection instruments and technologies	Description of instruments (e.g., interview guides, questionnaires) and devices (e.g., audio recorders) used for data collection; if/how the instrument(s) changed over the course of the study	5
S12 Units of study	Number and relevant characteristics of participants, documents, or events included in the study; level of participation (could be reported in results)	6
S13 Data processing	Methods for processing data prior to and during analysis, including transcription, data entry, data management and security, verification of data integrity, data coding, and anonymization/deidentification of excerpts	5
S14 Data analysis	Process by which inferences, themes, etc., were identified and developed, including researchers involved in data analysis; usually references a specific paradigm or approach; rationale ^a	5
S15 Techniques to enhance trustworthiness	Techniques to enhance trustworthiness and credibility of data analysis (e.g., member checking, audit trail, triangulation); rationale ^a	5
Results/Findings		
S16 Synthesis and interpretation	Main findings (e.g., interpretations, inferences, and themes); might include development of a theory or model, or integration with prior research or theory	6 - 22
S17 Links to empirical data	Evidence (e.g., quotes, field notes, text excerpts, photographs) to substantiate analytic findings	6 - 22
Discussion		
S18 Integration with prior work, implications, transferability, and contribution(s) to the field	Short summary of main findings; explanation of how findings and conclusions connect to, support, elaborate on, or challenge conclusions of earlier scholarship; discussion of scope of application/generalizability; identification of unique contribution(s) to scholarship in a discipline or field	23 - 25
S19 Limitations	Trustworthiness and limitations of findings	25 -26
Other		

S20	Conflicts of interest	Potential sources of influence or perceived influence on study conduct and conclusions; how these were managed	n/a
S21	Funding	Sources of funding and other support; role of funders in data collection, interpretation, and reporting	Title page - 2

^aThe rationale should briefly discuss the justification for choosing that theory, approach, method, or technique rather than other options available, the assumptions and limitations implicit in those choices, and how those choices influence study conclusions and transferability. As appropriate, the rationale for several items might be discussed together.

For peer review only

BMJ Open

Factors Influencing Virtual Reality Use in Paediatric Acquired Brain Injury Upper Limb Rehabilitation – A qualitative study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-083120.R1
Article Type:	Original research
Date Submitted by the Author:	04-Dec-2024
Complete List of Authors:	Rathinam, Chandrasekar; University of Birmingham College of Medical and Dental Sciences, Institute of clinical sciences; Birmingham Women's and Children's NHS Foundation Trust, Therapy Service Farr, William; University of Cambridge Ray, Daniel; Birmingham Women's and Children's NHS Foundation Trust Gupta, Rajat; University of Birmingham; Birmingham Women's and Children's NHS Foundation Trust
Primary Subject Heading:	Qualitative research
Secondary Subject Heading:	Paediatrics, Rehabilitation medicine, Qualitative research
Keywords:	Virtual Reality, REHABILITATION MEDICINE, QUALITATIVE RESEARCH, Developmental neurology & neurodisability < PAEDIATRICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Title

**Factors Influencing Virtual Reality Use in Paediatric Acquired Brain Injury
Upper Limb Rehabilitation – A Qualitative Study**

Authors

Chandrasekar Rathinam^{1,2*}

William Farr³

Daniel Ray²

Rajat Gupta^{1,2}

Affiliations

¹ College of Medical and Dental Sciences, University of Birmingham, UK

² Birmingham Women's and Children's NHS Foundation Trust, Birmingham, UK

³ Faculty of Education, St Edmunds College, University of Cambridge, UK

***Corresponding author**

College of Medical and Dental Sciences

University of Birmingham

Edgbaston

Birmingham

B15 2TJ

United Kingdom

Birmingham Women's and Children's NHS Foundation Trust

Steelhouse Lane

Birmingham

B4 6NH

United Kingdom

Contact (Email)

Chandrasekar Rathinam - c.rathinam@bham.ac.uk

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

32Chandrasekar.rathinam@nhs.net

33William Farr - wjf33@cam.ac.uk

34Daniel Ray - Daniel.ray2@nhs.net

35Rajat Gupta - rajatgupta@nhs.net

36

37

38**ORCID**

39Chandrasekar Rathinam: 0000-0002-0049-8430

40Rajat Gupta: 0000-0001-5111-7821

41Willam Farr 0000-0003-3644-5311

42

43

44**Authors contribution**

45All authors read, provided feedback and approved the final manuscript.

46

47

48**Data availability statement**

49The participants of this work did not give consent for their data to be shared publicly.

50Therefore, the research supporting data is not available.

51

52**Funding**

53This article was written as part of Topol digital fellowship programme, UK (2022 –

542023). The study sponsors were not involved in the study design, data collection,

55analysis and interpretation of data, and the decision to submit the manuscript for

56publication. Funding/grant number not applicable.

57

58**Disclosure of interest**

59The authors have stated that they had no interests which might be perceived as

60posing a conflict or bias.

61

62**Word count: 6183**

Abstract

Objective

Upper limb movement difficulties in children with acquired brain injury (ABI) result in longer recovery times compared to lower limb. Intensive neurorehabilitation promotes a good long-term functional outcome. Virtual reality (VR) and video game technologies are invaluable adjuncts to traditional neurological rehabilitation as they help to motivate, engage and gain children's compliance in goal-directed therapy. However, this technology is not routinely used in the National Health Services, UK; it requires embedding to benefit children and their families.

VR implementation in rehabilitation practice requires development. The associated influencing factors require further exploration before routine use can be established. This project aims to understand the factors influencing the use of VR in upper limb rehabilitation in children.

Design

An interpretative qualitative study utilised focus groups and 1:1 semi-structured interviews conducted in person and online to explore participants' experiences. These were analysed for inductive overarching themes, particularly focusing on the views of professionals and young people regarding the use of VR in upper limb rehabilitation.

Setting

Two neuro-rehabilitation services located in two children's hospitals in England, UK.

Participants

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

88 Three physiotherapists, five occupational therapists, a play worker, and four members
89 from the young persons’ advisory group took part. Four focus groups with 2 – 4
90 participants in each group and two 1:1 semi-structured interviews were conducted.
91 Thematic analysis was used to create the model participants described as the factors
92 that influenced the use of VR in neurorehabilitation.

94 **Results**

95 Five closely related major themes and thirty associated subthemes were developed:
96 training, knowledge, promotion, consideration of barriers, and family factors. There is
97 a lack of knowledge and understanding about the use of VR, its limitations, and the
98 clinicians’ motivation to use it. Training packages with available VR equipment, clinical
99 indicators, and scientific evidence are required. Staff need frequent training, logistics
100 (uninterrupted Wi-Fi, software, hardware), and simple instruction manuals.

102 **Conclusion**

103 To introduce VR into the routine rehabilitation of children with ABI, investment in
104 improving knowledge, frequent training, and positive behaviour change among
105 health professionals is needed.

107 **Keywords**

108 Acquired brain injury; Children; Virtual reality; Rehabilitation; Thematic analysis

Strengths and limitations of this study

- This study gathered therapists' and young people's views on using virtual reality for rehabilitation.
- Participation was broad and inclusive of therapists and young people's groups.
- A 4i model (improving knowledge, investing in training, infrastructure, and implementing changes) is suggested.
- Members checking was not done to improve triangulation.

136 **1. Introduction**

137
138 Worldwide, acquired brain injury (ABI) is one of the most common causes of neuro-
139 disability in children and young people (CYP)[1]. In the UK, ABI accounts for 35,000
140 childhood presentations to emergency departments annually[2]. Of these, 5% have a
141 moderate to severe brain injury[3]. A subsection of children with ABI have a stroke-
142 like presentation and have a functional loss of an upper limb that results in an
143 increased dependency[4].

144
145 Evidence indicates that upper limb function takes longer to recover than lower limb
146 function in the adult stroke population[5]. Neuroplasticity is vital in regaining motor
147 skills following brain injury[6]. Dendritic growth and synaptic changes in the brain are
148 linked to neural plasticity[7]. Repetitive, purposeful, and goal-directed movements are
149 required to induce neural plasticity. Following ABI, children often require early and
150 intensive neurorehabilitation to achieve their maximum potential[8]. High-intensity and
151 repetitive practice require a high level of labour-intensive therapy, often difficult to
152 provide due to high work demands and capacity issues[9]. These children can be
153 passive, and parents struggle to motivate them to do the prescribed exercise[10].
154 Most children, however, do enjoy playing computer games[11]. Virtual reality (VR)
155 based computer gaming technology is emerging as a valuable adjunct to traditional
156 neurological rehabilitation[12]. It has been recently used to treat CYP with cerebral
157 palsy (CP)[11] and ABI[13]. VR has been shown to improve upper limb functions in
158 children with CP when used with conventional therapy[14]. Motivation is a crucial
159 element for active participation in neuro-rehabilitation. VR has been shown to improve

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

the upper limb functions of children with CP Manual Ability Classification System levels III and IV[11].

VR interventions need less professional input and can be programmed to suit an individual child's needs. This type of intervention helps address boredom and adherence to interventions whilst facilitating increased therapy time and may not rely on therapeutic contact time[12]. Parents are empowered to facilitate the active participation of children. However, VR has not been used routinely in the UK National Health Service (NHS).

A cross-sectional survey in Canada that examined the use of VR by therapists reported that 46% of therapists had VR experience, and 15% of them had used it to treat people who had suffered a brain injury. This study found that lack of funds, space, time, support staff, and appropriate clients were barriers to using VR; they found that client motivation, therapists' knowledge, and management support were the facilitators to enable VR to be used in regular practice[15]. Farr *et al.*, (2020) examined the paediatric physiotherapists' and occupational therapists' learning needs of VR in the UK and noted that only 11% of therapists reported current use of VR as a rehabilitation intervention with their patients; managers have little or no influence on VR adaptation, and the lack of IT support is another barrier[12]. Farr *et al.*, (2020) recommended that tailored support is required to facilitate VR adaptation[12].

In rehabilitation, treatment goals have shifted from managing the body structure to managing activity limitations and participation restrictions[16]. Our previous co-production work with children who have neurological motor disorders showed that the

children found formal exercise lacking in salience and variety; they found that it was hard to fit their exercise programme into their daily life; it competed with academic needs; it isolated them from their peers; and it was harder to do the programmes as they grew older. They recommended play-based activities, video games, and VR-based activities that fit in with their life and would be more acceptable because they could play with their friends/peers and siblings. In line with these outcomes, therapists must evolve from conventional hands-on therapy to next-generation technology-based therapy programmes that facilitate personalised care[9]. VR can be seamlessly embedded in integrated services to improve patient care and efficiency and promote functional outcomes[15]. For this, VR Intervention requires development, and the associated influencing factors need exploring before it can be used routinely in rehabilitation.

This study aims to understand the factors influencing VR use in upper limb rehabilitation for children with ABI and to determine the infrastructure support and training required for therapists to plan and deliver personalised VR programmes.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

2. Methods

2.1 Design

A series of qualitative semi-structured interviews was used to explore participants' experiences and analysed for inductive overarching themes using inductive thematic analysis. The focus was on professional and young people's views of VR for upper limb rehabilitation. The authors followed the Standards for Reporting Qualitative Research (SRQR) Checklist [17] (Appendix 1).

2.1.1 Ethics

This study was registered with the Birmingham Women's and Children's Hospital Research & Development office and in accordance with the UK National Research Ethics Service guidance (REC: 16/BWC/LA/Rathinam).

2.2 Patient and Public Involvement

Patients and the public were not involved as this study aimed to understand the therapists' need to use VR in clinical settings.

2.3 Participants

All the physiotherapy and occupational therapy team members of the Birmingham Children's Hospital (BCH) and Oxford Children's Hospital, treating children with ABI were invited. Purposive sampling strategy was used. Three physiotherapists (PT), five occupational therapists (OT) from both centres and a play worker (PW) from BCH agreed to take part and gave verbal consent. We also invited BCH's Young Persons' Advisory Group (YPAG); four members (13 - 16 years) consented through their

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

coordinator. The therapists' and PW's professional experience and use of VR for treatment over the years are given in Table 1. Appointments were scheduled to meet the participants in person or through an online platform (Microsoft Teams); the YPAG coordinator was present when meeting the YPAG group.

2.4 Procedure

Between 2 and 4 participants per group attended four focus groups, and two 1:1 semi-structured in-depth interviews were completed, each session lasting between 60 and 90 minutes according to staff availability. Participants were offered both options of a 1:1 interview or a focus group, and take-up was according to personal preference. There was no remuneration for participants. A semi-structured interview guide was used to facilitate discussion (Appendix 2). Participants were asked about their VR experience, barriers, adherence factors, and reflections. CR undertook all interviews, and the Interviews were recorded with permission. Written notes were taken simultaneously. CR is an experienced clinician working with children with neuro disability, and positionality has been developed through systematic review published [10] in this area. CR maintained reflexivity by diary keeping during the interview and analysing the transcript.

2.5 Analysis

Interviews were transcribed in full. All transcripts were deidentified. Pilot coding and training took place between WF and CR. Thematic analysis was conducted following Braun and Clark's (2006) guidelines, and was conducted by hand. A multistage approach included familiarisation, indexing, sorting, developing an initial framework, applying the framework, and charting the data, and interviews were read and coded

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

260 until the saturation point was achieved. Interpretation followed to analyse the collected
261 data[18]. An initial line-by-line inductive coding was performed verbatim. This
262 enabled themes to evolve from the data related to the project aim. These themes were
263 categorised, and five major themes and associated but interrelated sub-themes were
264 developed. Trustworthiness was ensured by triangulating the data gathered with the
265 previous study data (WF). Analytical rigour was maintained with an audit trail, and the
266 coding between the authors was aligned across independent coding of transcripts and
267 multiple reviews to align emerging themes (CR and WF).

3. Results

Thirteen participants (three PTs, five OTs, a PW, and four participants from YPAG, along with their coordinators) consented and participated. All the professionals were aware of using VR for rehabilitation, but very few had any experience, and most reported no experience using it for treatment purposes. A model describing the factors influencing VR use in neurorehabilitation was created (Figure 1), and verbatim quotes demonstrating pertinent quotes for individual themes are presented below.

3.1 Knowledge

VR is considered a relatively new domain, and lack of knowledge and awareness are two factors that limit any therapists' use in the clinical setting.

3.1.1 VR Tools

Participants were all aware of the use of VR for treatment purposes. Many assumed that VR was an immersive technology using goggles or a headset. They needed to be made aware of the non-immersive style. They were keen to understand the different types currently available for rehabilitation, how they work, and how they can be used for rehabilitation.

"I would need to understand how virtual reality works. Would it be a headpiece? Would it be a kind of feedback thing on a screen?" (Participant-1)

3.1.2 Evidence base

Participants were unaware of the existing evidence supporting the use of VR. If the evidence shows validity, transferability, and generalisability, it will enable them to

make informed decisions. They asked for specific information about the indications, contraindications, limitations, patient position, and the minimal ability required to engage children in a VR session. They asked for details about different VR techniques, timings, progression mechanisms, expected changes or improvements, desired outcomes, treatment effectiveness measurements, and other factors supporting VR's clinical utilisation.

"....knowing the evidence base, and what the validity of it is, and if they have been a particular success in particular patient groups, any limitations on kind of age or cognitive ability or physical ability as to what they can, who can use it, like knowing how to make a good patient selection." (Participant-2)

3.1.3 Decision making

Choosing any modality for treatment underpins the relationship between clinical reasoning, evidence base, and equipment availability. This is also related to knowing a child's condition and the parents' capability. Participants were keen to know how and when VR should be identified as an intervention. This will be a clinical decision by the treating therapist, following a good-quality assessment of the child and identifying the goals of all concerned. Only then can it be determined whether VR equipment will assist in achieving the goals.

"....a good quality assessment, that means that we've gotten to know that child and their context and their family...." (Participant-3)

3.1.4 Desired games

The participants expressed their views on the type of games to choose and which to avoid. The games should give a positive experience, some of which can be achieved

through good graphics, up-to-date software, and equipment, giving them the feeling that they are playing proper computer games. There need to be various games to avoid boredom, which naturally moves to different levels as the child progresses. They should have the option to change the features so children can easily play without tiring, and a bigger switch control mechanism.

“....variety of things (games) so that they don't just play one game and then they are bored of playing that one, they have a good variety, and they can play.”

(Participant-4)

Participants had suggested avoiding games that are gender stereotypical, horror games, and car crash games, particularly for car accident victims. Therapists also need to be mindful of not setting up games that can show what the child cannot do to avoid giving them negative feelings, a sense of failure, and a risk of disengagement when they find the games too hard.

“....acquired brain injury (be)cause, let's say it happened in a car crash, then you bring this piece of kit, and you're playing a car game. It's gonna be a bit triggering....” (YPAG-1)

3. 2 Training

Lack of training has been identified as an essential factor limiting VR use in clinics. Participants highlighted their training needs related to the child's functional loss.

3.2.1 Loss of function

Some children lose their motor function following ABI. This is especially true for those who have a stroke-like presentation or tone abnormalities. These children struggle to use their hands for everyday functional activities, impacting their independence and disrupting their development. At times, they struggle to entertain themselves. Participants were keen to understand the barriers to using VR during the acute phase. These include having a cannula in their hand, frequent seizure episodes, and not having part of their skull following brain surgery.

“....if they have lost the use of their hands, they really struggle with....”

(Participant-8)

3.2.2 Emotion

Emotional regulation is often challenging, and these children often get frustrated when they have lost independence or skills following ABI. Although they know what happened to them, they often do not reflect on why. The psychological impact includes feeling low and scared, and they wish to go home instead of being in the hospital. They become aware of what they can play despite cognitive and learning problems but are depressed as they cannot manipulate small controls in their VR consoles.

“....They've just got a brain injury. They feel grotty. They're scared. They don't want to be in hospital, but they don't really quite know what they want....”

(Participant-5)

3.2.3 Fatigue

Fatigue is a common problem experienced by children with ABI from the early stage, and most of them tend to have this problem on a long-term basis. The participants

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

want to learn the early signs of fatigue or tiredness, so stop the session at an appropriate time or help parents recognise fatigue and differentiate between tiredness and boredom. This will also help negate fatigue by pacing, changing the game's complexity, and modifying the VR's intensity.

"most of the time, they have got tired....and I've had to stop." (Participant-8)

3.2.4 Resource person

Participants have suggested a resource person within their team to keep their skills, be responsible for ongoing training, and demonstrate that VR works. The resource person can also take responsibility for reviewing the VR programmes regularly and ensure that the family is competent in delivering it while in the ward. The participants commented that they need time to practice VR and a higher training volume to maintain competency so that it can become second nature. They wanted yearly and frequent refresher training programmes to ensure learning reinforcement.

"We would want to always ensure that we've demonstrated and that the person is using it in the correct way...." (Participant-3)

3.2.5 Guidance Workshops & Video

Participants suggested frequent workshops for therapists, parents, and children, along with pre-course materials about VR systems and how they work to educate people. The workshop method was considered helpful as it offers practical opportunities to work with the VR sets, reinforces learning by repetition, and provides ongoing support. Journal club, keeping abreast of the research, and constant refresher will minimise difficulties in using VR.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

“....having a field training module where people could see this sort of troubleshooting guide. So the basics of how to connect to the Internet, Bluetooth, troubleshooting....” (YPAG-2)

YPAG participants suggested a YouTube channel containing training videos for parents, children, and clinicians for VR training. Video content includes tutorials such as instructions on setting up the VR, how to play, troubleshooting, and frequently asked questions and answers.

“....could have like a YouTube channel and it's basically like to the people (who) have already used it, find out their most common problems and then you can specifically have a video that (shows) how to fix this.” (YPAG-1)

3.2.6 Learning content

As part of the training, the participants suggested that the learning content should contain VR's benefits, clinical utility, indications, and suitability for different age ranges, settings, and game variety and complexity levels. Additional information, such as how VR can be used in a therapeutic way to treat other problems, collecting, analysing, and interpreting data to see what the patient has done, and how VR aligns with other treatments, were also required. The learning content is expected to include further details about ease of use, difficulties, pros and cons, and treatment dosage for optimal outcomes. If specific VR equipment has been used, the participants were interested in learning the manufacturer's data, equipment information, and repetition altering mechanism to avoid fatigue, charging, and readjusting the units. From the patient's perspective, the participants needed further training in goal setting, how VR fits within their rehabilitation programme, and the functional activities and muscle groups they

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

431 were exercising. This information was considered beneficial to training families to use
432 VR under the guidance of therapists.

434 *3.2.7 Learning style & walk-throughs*

435 The participants expressed different learning styles and strategies. Although a training
436 manual is available, it does not allow the learner to ask questions, and they prefer
437 learning on the job and training in person. Some expressed the need to learn
438 personally and to use VR frequently, which would help to reinforce their learning and
439 increase their confidence. Demonstrations and sitting with experienced users were
440 the preferred methods because it would allow them to ask questions and practice
441 under supervision. Participants wanted much handholding, slow explanations of the
442 functions, and to be walked through the game, especially when they began using it.

443 *“I would like just some guidance on what to do with them and how far to push*
444 *them, and that would be great for my training needs....”* (Participant-8)

446 **3.3 Barriers**

447 *3.3.1 Habitual practice*

448 Therapists are accustomed to working with minimal resources. Therefore, technology
449 is not usually considered a treatment option despite its availability. For example, some
450 of the therapists still draw their exercise programmes instead of using computer-
451 generated programmes. Any new treatment technique requires time for evaluation
452 and reflection. If the technology is not working, they prefer to return to old treatment
453 techniques as they need more time to investigate and troubleshoot. This means that
454 the new technology is not used frequently, which leads to a lack of familiarity and a
455 decreased inclination to use it without further training.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

1
2
3 456 *"....I am still drawing stickman for the exercise programme because trying to*
4
5 457 *get to the computer to log on to (write an exercise programme is) hard to*
6
7
8 458 *navigate and crashes all the time. When you know your patient is improving so*
9
10 459 *quickly and you will need to redo this exercise programme in two days, it*
11
12 460 *actually just stops me using it all together...."* (Participant-2)
13
14
15 461

16 17 462 3.3.2 Generational & Self-perceptions

18
19 463 Some commented that they were old-generation therapists and too old to try VR. They
20
21 464 felt that the generational difference between therapists and children is vast, and there
22
23 465 is a gap in knowing what the children like and how to play with them using VR. They
24
25 466 lacked the experience and confidence to use it for therapy purposes. Some therapists
26
27 467 and YPAG participants who had tried a VR headset reported feeling sick and disliked
28
29 468 using it. Those who had tried VR reported that they were not technology literate, felt
30
31 469 rubbish, and needed to remember about using them.
32
33

34
35 470 *"....I'm not a gamer; gaming (has) been around for a long time, but I'm not*
36
37 471 *one. But actually, the immersion in technology and the development of the*
38
39 472 *internet and everything has really only been in my adult life.* (Participant-1)
40
41
42 473

43 44 474 3.3.3 Logistics & Complex manual

45
46 475 The participants reported many logistical issues. For example, communication
47
48 476 between part-time staff, such as sharing information and hand-over, is problematic
49
50 477 from a staffing perspective. Having a designated safe space is a challenge, as
51
52 478 therapists often need guidance on where to treat these children or where to store the
53
54 479 VR equipment. Connecting the VR units to the internet is a challenge, and the
55
56 480 participants felt that having easy access to IT services may overcome this
57
58
59
60

issue. Although manuals are available, participants reported that these are complex, and hard to navigate and find the solution. Therapists wanted a simple laminated crib sheet or a quick starter guide explaining troubleshooting, screenshots of what has been shown in the manual, set-up, games, and calibration techniques.

“....the problem with the manuals and things are so long often that you can't find the thing that you're looking for..... having some crib sheet that quickly tells you this would help with this aspect....” (Participant-5)

From the equipment perspective, some need to be reset each time the children play a different game. Some children find it hard to grip the VR sets and need to readjust their handhold for each new game. For the complex VR sets, the children need someone around them to set up the programme, motivate them, and engage in the games for therapy. This is a challenge after their allocated time with the therapists, particularly over the weekend. This usually results in children not doing anything over the weekend. Most VR gadgets are made for typically developing children, and very little equipment is available for children with disabilities. When children with ABI tried to use the usual consoles, they found it challenging to manipulate them, and they tired quickly. Most children had prior experience of playing VR games with rich visual content, but rehabilitation-focused games are generally primitive; hence, children quickly lose interest.

“....kids these days will have high expectations from the point of view of graphics and action quality because of what they see in films and games.”
(Participant-2)

3.3.4 Time factor

Enseignement Supérieur (ABES) .
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

506 Lack of time is a common problem expressed by most of the therapists. They
507 acknowledged that they need dedicated time for training to use VR properly. Despite
508 the availability of the systems, therapists need more training time and clinical time to
509 use them appropriately.

510 *"....but often we have to reset it 3, 4, 5 times to make sure that (all) the Wi-Fi*
511 *connectors are actually connected. So, for one successful session, it's about 2*
512 *hours. So that means 100% more time than you (usually spend)...."*
513 (Participant-5)

514

515 3.3.5 Technology

516 Many participants reported that poor Wi-Fi connectivity in their hospital limited their
517 opportunity to use VR. They commented that the connection would not work or the
518 equipment would not pick up the signal. Having a reliable network connectivity is vital
519 if VR is to be used in the hospital setting.

520 *"I think the practical difficulties in the hospital is lack of time, space, Wi-Fi,*
521 *general resources (which) make it very difficult to be consistent in anything that*
522 *needs a set-up"* (Participant-4)

523

524 Participants believed that the technology constraints listed above often put them off. It
525 requires a lot more support if it is to be used effectively, which defeats the purpose of
526 why VR was introduced in the first instance.

527 *"....if the IT set up for this sort of thing was not easy to use and not reliable, that*
528 *would make me use other adjuncts instead.... I'll spend an hour setting it up*
529 *and then it won't work anyway"* (Participant-2)

530

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

531 3.3.6 Change management

532 Participants acknowledged that people take the path of least resistance and that they
533 generally do not like change. They require convincing and demonstrating the utility,
534 as people are not tech-savvy. They suggested example videos of VR in practice,
535 showing how children used it and the outcome achieved.

536 *"....it's a change in practice, so people don't like change. Particularly and will*
537 *only engage if the change is managed, almost like a change management*
538 *way....it's about saying, 'this could help; why don't you give a (try)'." (Participant-*
539 *1)*

541 3.3.7 Accessibility

542 Lack of space is an essential factor that limits the use of VR. Leaving the system close
543 to the bedside in some wards takes a lot of work. In some places, expensive units are
544 available, but therapists have yet to be trained to use them. Sometimes, more minor
545 issues, such as missing cables or sets needing to be charged more, caused
546 embarrassment when the session was disrupted. Many therapists felt excited initially
547 to use VR but became reluctant because of accessibility issues; therefore, sustaining
548 enthusiasm is difficult.

549 *"....it's also about how we've gotta make it accessible to everybody. In our*
550 *department, we find all the expensive kits that once were the pride and joy and*
551 *are now stuck at the back of a cupboard because nobody can remember how*
552 *to set up the Wii or nobody's got the key to get the search insert. So we've got*
553 *all these pieces of kit, but I've never seen them used." (Participant-6)*

555 3.4 Promotion & Consideration

3.4.1 Motivation for staff & children

The YPAG suggested different strategies to promote VR for rehabilitation purposes. These included age-appropriate games, offering taster sessions, encouraging games with parents or siblings, challenging others, or beating their previous scores. Watching other children enjoy VR may enthuse disinterested children. Promoting VR success stories online, getting people to talk about their experience, having competitions by playing against each other with a leader board, and playing interactively can give reassurance and motivation.

"....you should get other kids to try it out in front of kids that don't wanna do it yet...." (YPAG-3)

3.4.2 Reward/feedback

From the therapists' perspective, VR is not a universal treatment provision but an additional tool that can be used in conjunction with their usual treatment. Positive feedback is always a good motivational factor. This feedback can be the visual displays on the game, a certificate for reaching certain milestones, or purely verbal feedback.

"I definitely like that when they get the feedback as well, like party, pop sound effects or something just to kind of go well, you know, it's that big 'Well done! You've just done that level'." (Participant-5)

3.4.2 Perceived benefits

All the participants acknowledged that children love computer games and VR. They can spend hours playing as it is enjoyable, fun, engaging, and interactive. Therapists felt that VR helps to improve the child's engagement, which can facilitate movements,

581 increase movement control, and facilitate stability and function. The end products are
582 improved performance, grip strength, muscle length, satisfaction, spasticity
583 management, and function improvement underpinned by component skill
584 development. VR helps to distract the children, get them moving and enable them to
585 do their therapy without realizing that they are doing it.

*"Children love gaming, so I think it's a great way to distract them and actually
get them a bit more moving."* (YPAG-3)

589 VR can also be used for motivation instead of therapy, especially for fun if they do not
590 want to do anything in the ward. VR also has the potential to make hospitals more
591 interesting, turning a 'no' person into a 'yes' person by trusting their therapists. It
592 empowers parents to improve their independence.

*"He was smiling and laughing, and that was the first time I'd seen that.... you're
not going to do anything horrible and that just playing computer games and that
did have a knock-on effect as well to his positivity in other sessions."*
(Participant-5)

598 3.4.3 Confidence of Healthcare Professionals

599 Participants who had prior experience using VR were confident in turning on Bluetooth,
600 adjusting tables to use VR, deciding suitable VR games for different age groups,
601 selecting games' complexity, progressing to the next level if the child achieved the
602 target, using different movements and techniques to control VR and helped them to
603 use both hands. However, confidence varied:

"Do people feel trained & confident & competent?" (Participant-1)

"We don't have confidence...." (Participant-6)

606 *"I'd like you to guide how to do it..."* (Participant-8)

607

608 3.4.4 Prior experience

609 Some therapists had experience in using VR and noted that it was beneficial. It was
610 used to improve attention and concentration, engagement, movements, and
611 strength. They learned that a child does not need much movement to operate VR to
612 use it at a low level. However, the lack of Bluetooth connectivity limited the VR unit's
613 use for treatment and led to frustration. Children lost interest quickly when the system
614 did not work, and it compromised other aspects of therapy. Apart from connectivity,
615 they have noted other challenges, including children being competitive and making it
616 too difficult for themselves. Therapists had assumed that parents understood the
617 treatment goals and could follow what they were seeing, but this was only sometimes
618 the case.

619 *"I've had that a few times, and then they lose interest really quickly, then they*
620 *lose their faith in you knowing what you're doing because the whole system is*
621 *just not working. So, if you're not careful, that can rub off on other aspects of*
622 *your therapy, and you don't want that."* (Participant-4)

623

624 3.4.5 Deliberation

625 Participants from the YPAG group suggested that the commercially available VR
626 games they are playing are fast-moving, high-motion simulation, and have loud sound
627 effects. These games will be challenging to play for children with ABI due to their slow
628 reactions; the effects of flash and sound effects on eyes and ears, especially when
629 most already suffer from headaches. Therapists deliberated about whether they could
630 advocate screen time for the acute ABI children, when it would be appropriate to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

engage them in VR, and what games and equipment would kindle a child's interest and maintain their motivation to exercise and facilitate them taking responsibility. They would want to ensure that the child and their family perceived it as therapeutic and that barriers to its use for rehabilitation were minimised.

“...it's not good for an injured and healing brain to have a lot of screen time because of the way it fires up all your neurons....” (Participant-1)

3.4.6 Grading & Adaptation

Children with ABI experience various complex challenges, including slow reaction speed; therefore, adapting VR games is vital to make it inclusive for everyone. Adaptation of VR requires consideration of a child's age range, physical and cognitive ability, concentration and engagement, communication issues, sensory impairment, and hearing and visual impairment. In the acute phase, the children struggle to be in a busy ward environment where the noise and other distractions limit their focus in therapy sessions. These children will require both the games and their environment to be adapted. Instructions should be clear and concise for all parents to understand.

"In terms of understanding families and making sure that they can understand the instructions.... you need to adapt how you gave the instructions or how the interface worked....” (Participant-1)

3.5 Family factors

3.5.1 Family Needs

Digital illiteracy and digital poverty are essential factors, as some families and parents may need access to technology. During the acute rehabilitation phase, families

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

656 struggle to cope with sudden changes, and it was considered hard to expect them to
657 engage their children in VR programmes.

658 *"....thinking about high incidences of and younger parents, high incidences of*
659 *parents who might not have access to technology, high incidences of parents*
660 *who might have needs, learning disability, physical disabilities themselves.*
661 *How are we expecting them to carry over...."* (Participant-7)

663 3.5.2 Parenting culture and negotiation

664 Parenting covers a massive spectrum that is influenced by cultural and historical
665 context; any technology intervention should fit meaningfully around the family culture
666 and belief system. Having cultural sensitivity around parenting and boundaries is
667 required. One aspect of this that therapists observed was the varying approaches to
668 technology access and screen time. Parents must see the value and feel confident
669 that VR will support their child's rehabilitation. Therapists suggested negotiating a set-
670 up programme with the parents.

671 *"....different parents might have different parenting approaches relating to*
672 *access to technology and screen time, and one of the jobs we'd have to do to*
673 *set up a treatment program is negotiate that with them."* (Participant-6)

675 3.5.3 Goals

676 Engaging family and children around personalised goal setting is considered
677 paramount. The therapist's role is to decide whether VR helps achieve the agreed
678 goals.

679 *"....we don't want to set people up with a device that measures their failure*
680 *because that's really disheartening."* (Participant-3)

1
2
3 681
4
5 682 3.5.4 Structure to family

6
7
8 683 Families need to use VR for the well-being of their children, and they require a
9
10 684 structure around the use of VR. This includes clarity about how VR will be used, a
11
12 685 timetable for VR games, and limiting the play time and frequency when playing
13
14 686 complex games. Participants initially recommended playing for 10 minutes and
15
16
17 687 building up if the children were eager to play further. They have suggested playing
18
19 688 between 5 and 45 minutes, morning and evening, and having a break between
20
21 689 sessions.

22
23
24 690
25
26
27 691
28
29 692
30
31
32 693
33
34
35 694
36
37 695
38
39
40 696
41
42
43 697
44
45 698
46
47
48 699
49
50
51 700
52
53 701
54
55
56 702
57
58
59 703
60

Enseignement Supérieur (ABES).
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

4. Discussion

This scoping work is based on the perceptions of therapists, clinicians, YPAG members and people in an NHS setting when using VR for rehabilitation. Five major themes relating to issues around knowledge, training, and implementation emerged. Four key principles ('4i') were developed, which need consideration by professionals when using VR in upper limb rehabilitation (Figure 2). Those four principles structure the discussion.

4.1 Improving knowledge

Evidence-based intervention is the current principle that enables clinicians to use interventions with confidence[19]. To facilitate VR intervention, the NHS workforce requires extensive investment to improve awareness, knowledge, and convincing evidence base in VR use. Although VR technology has been used for rehabilitation for over two decades, there are not enough high-quality studies encouraging clinicians to utilise VR in routine practice[12]. Commercially available VR equipment has not been used by the ABI population and requires adaptation of VR equipment. It may not be possible to conduct research to the level of randomised controlled trials, especially under individual conditions; therefore, translating the knowledge gained from other conditions, such as adult stroke and children with CP should be considered within the paediatric ABI population.

4.2 Invest in training

Therapists have trained to be 'hands-on' and require considerable and extensive training to adapt technology for treatment. Therapists often feel uncomfortable trying

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

new technology as part of their practice due to their age perceptions[15]. Introducing new technology requires considerable training to change behaviour[11]. Various learning and training models and descriptors of available VR equipment, clinical indicators, and participants' experience need consideration when designing training packages. VR needs to be embedded in routine practice to facilitate behaviour change. Tact *et al.*, (2021) have suggested a digital adaptation framework for the NHS workforce, and it can be considered for technology-related training and education[20]. Technology-based intervention must be a core skill gained through graduate training so future therapists can become early adopters[21,22].

4.3 Infrastructure

Basic infrastructure provision will eliminate many barriers to using VR for rehabilitation[23]. Early adopters or technophobes within the workforce are at risk of abandoning VR if the required resources, such as uninterrupted Wi-Fi, updated software, and user-friendly hardware, are not readily available. Participants perceived that much of the available rehabilitation-focused VR equipment was heavy, came in universal/uniform sizes only, was hard to operate, unstable, and time-consuming. Finally, attrition of technology limits its use due to poor longevity, and a lot of equipment needs upgrading in healthcare settings due to ongoing cost factors associated with degradation over time[24]. Technology readiness is one of the main issues organisations face; the poor compatibility between the evolving technology and the existing electronic health records often limits this[25]. The evolving evidence base, lack of user guidelines[26], and uncertainty around personalised applications[27], alongside accessibility and cost factors, further limit the organisation's ability, willingness and capability to use technology[28]. The introduction of digital

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES).

technology, including VR, should take place alongside investment in the stability of the approach to rehabilitation, including the required infrastructure should be in place.

756

757 **4.4 Implementing change**

758 Our findings suggest that human factors play an essential role in digital adaptation.
759 Various factors related to affordability, digital poverty, and the family's readiness to try
760 technology as a tool significantly limit the uptake of VR for rehabilitation. The NHS
761 has a digital technology framework for allied health professionals[29] and a long-term
762 plan for digitally enabled care[30]. However, the embodied reality is far behind and in
763 relative infancy. If NHS organisations commit to embedding digital interventions as a
764 high priority, and investing heavily in their workforce and resources, technology-based
765 interventions such as VR will likely reach end users. Therapists are also responsible
766 for changing their rehabilitation strategies by moving away from traditional models and
767 adopting positive approaches of early adopters, which will lead to the evolution of
768 future leaders who will crystallise further embedding of change. For example, a group
769 of practitioners have used VR for ABI rehabilitation and noted the improvement and
770 engagement of children. A recent survey indicates therapists' higher confidence and
771 motivation when using digital technology[20]. This indicates either a step change in
772 therapists' age or a sea change in the UK's commitment to health services. Either
773 way, it is a positive sign.

774

775 Our findings enable us to understand factors that need addressing in the
776 implementation of VR for ABI rehabilitation. Implementation is a complex process that
777 requires managing barriers at various levels with multiple and holistic strategies.
778 Kouijzer et al. (2023) suggested using an implementation framework that facilitates

behavioural changes of all the stakeholders to integrate VR into healthcare practice [31]. VR intervention can be developed by addressing the issues identified in this work, and a future feasibility study is needed to see if VR can be used for upper limb rehabilitation in children with ABI, additionally focusing on the implementation issues from the beginning.

Limitations

This study did not report the parents' views of using VR, and this could add value to the themes developed. Triangulation of views was limited due to time availability of the clinicians, which could be strengthened with expert review, such as Delphi approaches but was not possible. A larger study would have encompassed more participants and a number of clinical service centres. Only two therapists and a play worker who took part in this project had some experience of VR, and the remaining therapists' views appear to be their perceptions rather than their experience of using VR. As a result, generalisability is not possible, but re-occurring themes could be made with some surety due to analytical saturation.

Conclusion

The use of VR technology for upper limb rehabilitation of children with ABI depends on investing in training for health professionals to improve their knowledge and thereby implement changes in rehabilitation techniques. We must invest in improving knowledge, repeated training, and positive behaviour change among health professionals to use VR technology for upper limb rehabilitation of children with ABI.

804 **Authors Contribution**

805 CR, WF, DR and RG: involved in study conceptualisation and preparing final
806 manuscript

807 CR and WF: responsible for study design

808 CR: responsible for data collection

809 CR, WF: responsible for data interpreting and coding

810 CR: guarantor of the study

811 All authors read, provided feedback and approved the final manuscript

812

813 **Acknowledgement**

814 The authors would like to thank Sharon Machin (Coordinator, Child Brain Injury Trust),
815 Ian Wood (Coordinator, YPAG, Birmingham Children's Hospital), YPAG participants
816 (Birmingham Children's Hospital), Children's Physiotherapists and Occupational
817 therapists (Birmingham Children's Hospital & Oxford Children's Hospital) for their help
818 and participation in interviews/focus groups. The authors also thank Ms Janet Peirson,
819 Retired Physiotherapist, Cambridge and Dr Vikram Mohan, Senior Lecturer,
820 Department of Rehabilitation and Sports Sciences, Bournemouth University, for their
821 constructive comments and preparation of this script.

822

823

824

825

826

827

828

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References:

[1] Dunford C, Cobbold C, Ray I, Wales L. The Information Gap for Children and Young People with Acquired Brain Injury. *Dev Neurorehabil* 2020;23:1–8. doi:10.1080/17518423.2018.1564394.

[2] Rathinam C, Mohan V, Bill P, Yates D, Gupta R, Peirson J. Impact of transcranial magnetic stimulation on motor function in children with acquired brain injury: a scoping review protocol. *BMJ Paediatr Open* 2023;7. doi:10.1136/bmjpo-2023-001885.

[3] Trefan L, Houston R, Pearson G, Edwards R, Hyde P, Maconochie I, et al. Epidemiology of children with head injury: a national overview. *Arch Dis Child* 2016;101:527–32. doi:10.1136/archdischild-2015-308424.

[4] Abgottspon S, Steiner L, Slavova N, Steinlin M, Grunt S, Everts R. Relationship between motor abilities and executive functions in patients after pediatric stroke. *Appl Neuropsychol Child* 2022;11:618–28. doi:10.1080/21622965.2021.1919111.

[5] Abdullahi A, Wong TWL, Ng SSM. Variation in the rate of recovery in motor function between the upper and lower limbs in patients with stroke: some proposed hypotheses and their implications for research and practice. *Front Neurol* 2023;14:1225924. doi:10.3389/fneur.2023.1225924.

[6] Rathinam C, Mohan V, Yates D, Bill P, Peirson J, Gupta R. Effect of non-invasive brain stimulation in children with acquired brain injury-a scoping review. *Front Neurol* 2024;15:1388718. doi:10.3389/fneur.2024.1388718.

[7] Kleim JA, Jones TA. Principles of experience-dependent neural plasticity: implications for rehabilitation after brain damage. *J Speech Lang Hear Res* 2008;51:S225-39. doi:10.1044/1092-4388(2008/018).

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

- 854 [8] Keetley R, Bennett E, Williams J, Stewart I, Whitehouse WP, Pilling P, et al.
855 Outcomes for children with acquired brain injury (ABI) admitted to acute
856 neurorehabilitation. *Dev Med Child Neurol* 2021. doi:10.1111/dmcn.14846.
- 857 [9] Rathinam C. Virtual reality for neurorehabilitation: hopes and challenges for
858 therapists. *Dev Med Child Neurol* 2021;63:370–1. doi:10.1111/dmcn.14791.
- 859 [10] Rathinam C, Mohan V, Peirson J, Skinner J, Nethaji KS, Kuhn I. Effectiveness
860 of virtual reality in the treatment of hand function in children with cerebral
861 palsy: A systematic review. *J Hand Ther* 2018. doi:10.1016/j.jht.2018.01.006.
- 862 [11] Choi JY, Yi S-H, Ao L, Tang X, Xu X, Shim D, et al. Virtual reality rehabilitation
863 in children with brain injury: a randomized controlled trial. *Dev Med Child*
864 *Neurol* 2021;63:480–7. doi:10.1111/dmcn.14762.
- 865 [12] Farr W, Green D. Lab to Rehab: Is virtual reality being adopted clinically for
866 children? 13th ICDVRAT with ITAG, 2020.
- 867 [13] Shen J, Johnson S, Chen C, Xiang H. Virtual Reality for Pediatric Traumatic
868 Brain Injury Rehabilitation: A Systematic Review. *Am J Lifestyle Med*
869 2020;14:6–15. doi:10.1177/1559827618756588.
- 870 [14] Burin-Chu S, Baillet H, Leconte P, Lejeune L, Thouvarecq R, Benguigui N.
871 Effectiveness of virtual reality interventions of the upper limb in children and
872 young adults with cerebral palsy: A systematic review with meta-analysis. *Clin*
873 *Rehabil* 2024;38:15–33. doi:10.1177/02692155231187858.
- 874 [15] Levac D, Glegg S, Colquhoun H, Miller P, Noubary F. Virtual Reality and
875 Active Videogame-Based Practice, Learning Needs, and Preferences: A
876 Cross-Canada Survey of Physical Therapists and Occupational Therapists.
877 *Games Health J* 2017;6:217–28. doi:10.1089/g4h.2016.0089.
- 878 [16] Ryan JM, Cassidy EE, Noorduyne SG, O'Connell NE. Exercise interventions for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

cerebral palsy. *Cochrane Database Syst Rev* 2017.

doi:10.1002/14651858.CD011660.pub2.

[17] O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for reporting qualitative research: a synthesis of recommendations. *Acad Med* 2014;89:1245–51. doi:10.1097/ACM.0000000000000388.

[18] Byrne D. A worked example of Braun and Clarke's approach to reflexive thematic analysis. *Qual Quant* 2022;56:1391–412. doi:10.1007/s11135-021-01182-y.

[19] Kumah EA, McSherry R, Bettany-Saltikov J, van Schaik P. Evidence-informed practice: simplifying and applying the concept for nursing students and academics. *Br J Nurs* 2022;31:322–30. doi:10.12968/bjon.2022.31.6.322.

[20] Tack C, Holdsworth L, Wilson A, McComiskie E, McCabe P, Wilkinson W, et al. Digital competency: a survey of UK allied health professionals. *Br J Healthc Manag* 2022;28:1–15. doi:10.12968/bjhc.2021.0123.

[21] Aungst TD, Patel R. Integrating Digital Health into the Curriculum- Considerations on the Current Landscape and Future Developments. *J Med Educ Curric Dev* 2020;7:2382120519901275. doi:10.1177/2382120519901275.

[22] Parsons D, MacCallum K, Schofield L, Johnstone A, Coulter S-K. Next-Generation Digital Curricula for Future Teaching and Learning BT - Emerging Technologies and Pedagogies in the Curriculum. In: Yu S, Ally M, Tsinakos A, editors., Singapore: Springer Singapore; 2020, p. 3–19. doi:10.1007/978-981-15-0618-5_1.

[23] Glegg SMN, Levac DE. Barriers, Facilitators and Interventions to Support Virtual Reality Implementation in Rehabilitation: A Scoping Review. *PM R*

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

- 2018;10:1237-1251.e1. doi:10.1016/j.pmrj.2018.07.004.
- [24] Williams MG, Stott R, Bromwich N, Oblak SK, Espie CA, Rose JB. Determinants of and barriers to adoption of digital therapeutics for mental health at scale in the NHS. *BMJ Innov* 2020;6:92 LP – 98. doi:10.1136/bmjinnov-2019-000384.
- [25] Iqbal AI, Aamir A, Hammad A, Hafsa H, Basit A, Oduoye MO, et al. Immersive Technologies in Healthcare: An In-Depth Exploration of Virtual Reality and Augmented Reality in Enhancing Patient Care, Medical Education, and Training Paradigms. *J Prim Care Community Health* 2024;15:21501319241293310. doi:10.1177/21501319241293311.
- [26] Nieto-Escamez F, Cortés-Pérez I, Obrero-Gaitán E, Fusco A. Virtual Reality Applications in Neurorehabilitation: Current Panorama and Challenges. *Brain Sci* 2023;13. doi:10.3390/brainsci13050819.
- [27] Cerritelli F, Chiera M, Abbro M, Megale V, Esteves J, Gallace A, et al. The Challenges and Perspectives of the Integration Between Virtual and Augmented Reality and Manual Therapies. *Front Neurol* 2021;12:700211. doi:10.3389/fneur.2021.700211.
- [28] Zanier ER, Zoerle T, Di Lernia D, Riva G. Virtual Reality for Traumatic Brain Injury. *Front Neurol* 2018;9:345. doi:10.3389/fneur.2018.00345.
- [29] England NHS. A digital framework for allied health professionals 2019.
- [30] Glasper A. A long-term plan for embracing digital healthcare technology. *Br J Nurs* 2019;28:204–5.
- [31] Kouijzer MMTE, Kip H, Bouman YHA, Kelders SM. Implementation of virtual reality in healthcare: a scoping review on the implementation process of virtual reality in various healthcare settings. *Implement Sci Commun* 2023;4:67.

doi:10.1186/s43058-023-00442-2.

Participant	Profession	Experience (years)	
		Professional	VR use
1	PT	32	4
2	PT	17	0
3	OT	6	0
4	OT	4	0
5	OT	20	0
6	OT	20	0
7	OT	10	6
8	PW	4	1

Table 1: Number of professional and virtual reality (VR) use experience among the participants in years (OT – Occupational Therapist; PT – Physiotherapist; PW - Play Worker)

Figure 1: Model of factors influencing virtual reality use for upper limb rehabilitation for children with acquired brain injury

Figure 2: 4i model to use implement VR for upper limb rehabilitation




Figure 1: Model of factors influencing virtual reality use for upper limb rehabilitation for children with acquired brain injury

107x103mm (300 x 300 DPI)




Figure 2: 4i model to use implement VR for upper limb rehabilitation

338x117mm (300 x 300 DPI)

Appendix 1: Factors Influencing Virtual Reality Use in Paediatric Acquired Brain Injury Upper Limb Rehabilitation - A qualitative study

Standards for Reporting Qualitative Research (SRQR) Checklist

O'Brien B.C., Harris, I.B., Beckman, T.J., Reed, D.A., & Cook, D.A. (2014). Standards for reporting qualitative research: a synthesis of recommendations. *Academic Medicine*, 89(9), 1245-1251.

No.	Topic	Item	Page
	Title and abstract		
S1	Title	Concise description of the nature and topic of the study identifying the study as qualitative or indicating the approach (e.g., ethnography, grounded theory) or data collection methods (e.g., interview, focus group) is recommended	Title page-1
S2	Abstract	Summary of key elements of the study using the abstract format of the intended publication; typically includes objective, methods, results, and conclusions	Title page – 3 & 4
	Introduction		
S3	Problem formulation	Description and significance of the problem/phenomenon studied; review of relevant theory and empirical work; problem statement	6 - 7
S4	Purpose or research question	Purpose of the study and specific objectives or questions	8
	Methods		
S5	Qualitative approach and research paradigm	Qualitative approach (e.g., ethnography, grounded theory, case study, phenomenology, narrative research) and guiding theory if appropriate; identifying the research paradigm (e.g., positivist, constructivist/interpretivist) is also recommended	9
S6	Researcher characteristics and reflexivity	Researchers' characteristics that may influence the research, including personal attributes, qualifications/experience, relationship with participants, assumptions, or presuppositions; potential or actual interaction between researchers' characteristics and the research questions, approach, methods, results, or transferability	10

S7	Context	Setting/site and salient contextual factors; rationale ^a	
S8	Sampling strategy	How and why research participants, documents, or events were selected; criteria for deciding when no further sampling was necessary (e.g., sampling saturation); rationale ^a	9 - 10
S9	Ethical issues pertaining to human subjects	Documentation of approval by an appropriate ethics review board and participant consent, or explanation for lack thereof; other confidentiality and data security issues	9
S10	Data collection methods	Types of data collected; details of data collection procedures including (as appropriate) start and stop dates of data collection and analysis, iterative process, triangulation of sources/methods, and modification of procedures in response to evolving study findings; rationale ^a	10 - 11
S11	Data collection instruments and technologies	Description of instruments (e.g., interview guides, questionnaires) and devices (e.g., audio recorders) used for data collection; if/how the instrument(s) changed over the course of the study	10 - 11
S12	Units of study	Number and relevant characteristics of participants, documents, or events included in the study; level of participation (could be reported in results)	9 - 10; 12 Table 1
S13	Data processing	Methods for processing data prior to and during analysis, including transcription, data entry, data management and security, verification of data integrity, data coding, and anonymization/deidentification of excerpts	10
S14	Data analysis	Process by which inferences, themes, etc., were identified and developed, including researchers involved in data analysis; usually references a specific paradigm or approach; rationale ^a	10
S15	Techniques to enhance trustworthiness	Techniques to enhance trustworthiness and credibility of data analysis (e.g., member checking, audit trail, triangulation); rationale ^a	11
Results/Findings			
S16	Synthesis and interpretation	Main findings (e.g., interpretations, inferences, and themes); might include development of a theory or model, or integration with prior research or theory	12 - 28
S17	Links to empirical data	Evidence (e.g., quotes, field notes, text excerpts, photographs) to substantiate analytic findings	12 - 28
Discussion			
S18	Integration with prior work, implications, transferability, and field	Short summary of main findings; explanation of how findings and conclusions connect to, support, elaborate on, or challenge conclusions of earlier scholarship; discussion of scope of application/generalizability; identification of unique contribution(s) to scholarship in a discipline or field	29 - 32

contribution(s) to the field		
S19 Limitations	Trustworthiness and limitations of findings	32
Other		
S20 Conflicts of interest	Potential sources of influence or perceived influence on study conduct and conclusions; how these were managed	n/a
S21 Funding	Sources of funding and other support; role of funders in data collection, interpretation, and reporting	Title page - 2

^aThe rationale should briefly discuss the justification for choosing that theory, approach, method, or technique rather than other options available, the assumptions and limitations implicit in those choices, and how those choices influence study conclusions and transferability. As appropriate, the rationale for several items might be discussed together.

1
2
3 **Appendix 2: Interview guide**
4
5
6
7

8 **Professionals**
9

- 10
- 11 • What is your understanding of virtual reality (VR)?
 - 12
 - 13 • Would Virtual reality-based video games be useful in treating children with
 - 14 brain injuries? What is the reason?
 - 15
 - 16
 - 17 • What motivates or stops you from using VR in clinical practice?
 - 18
 - 19
 - 20 • What is the support or training required?
 - 21
 - 22
 - 23 • How can we deliver VR treatment in clinical practice?
 - 24
 - 25 • What factors from the patient’s perspective should be considered when trying
 - 26 VR?
 - 27
 - 28
 - 29 • What factors from the staff and resource perspective do we need to consider
 - 30 when using VR?
 - 31
 - 32
 - 33
 - 34
 - 35

36 **YPAG group**
37

- 38
- 39 • What is your understanding of VR? What is your own experience of VR?
 - 40
 - 41 • Would Virtual reality-based video games be useful in treating children with
 - 42 brain injuries? What is the reason?
 - 43
 - 44
 - 45 • What motivates children to use VR? How can we encourage them to use it as
 - 46 a treatment tool?
 - 47
 - 48
 - 49
 - 50 • What type of games do you suggest and avoid?
 - 51
 - 52
 - 53 • When is the best time to use VR?
 - 54
 - 55 • What support do you need to use VR?
 - 56
 - 57 • What stops you from considering VR as a treatment tool?
 - 58
 - 59
 - 60