BMJ Open School nurse-led educational interventions for sexual and reproductive health promotion in adolescents in high-income countries: a mixed-methods systematic review protocol

Yu Mi Choi,¹ Suyeon Noh,² Hyun-Ju Seo ⁽¹⁾,³ Jaehee Yoon⁴

To cite: Choi YM, Noh S, Seo H-J, *et al.* School nurseled educational interventions for sexual and reproductive health promotion in adolescents in high-income countries: a mixed-methods systematic review protocol. *BMJ Open* 2025;**15**:e087528. doi:10.1136/ bmjopen-2024-087528

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (https://doi.org/10.1136/ bmjopen-2024-087528).

YMC and SN contributed equally.

Received 12 April 2024 Accepted 28 January 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to Professor Hyun-Ju Seo; shj5th@korea.ac.kr

ABSTRACT

Introduction As educators and health professionals, school nurses are in an optimal position to improve and advocate for adolescent reproductive and sexual health. This report outlines a protocol for a systematic review to synthesize evidence on the effects of school nurse-led education interventions and barriers and facilitators to implementing the interventions to improve students' knowledge, attitudes and behaviours related to sexual and reproductive health in high-income countries.

Methods and analysis We will develop a protocol to systematically review school nurse-led education interventions aimed at promoting adolescent sexual and reproductive health. This protocol will be based on the methodology of the Cochrane Handbook for Systematic Reviews of Interventions. The search will be conducted in Ovid-MEDLINE, CINAHL, Cochrane Library, Ovid-Embase, PsycINFO, Koreamed and ScienceON using relevant Medical Subject Headings and text words to identify the literature on different types of studies examining school nurse-led sexual and reproductive health education interventions in April 2024. Two independent reviewers will select relevant studies and extract data using a predefined template. We will assess methodological quality using the risk-of-bias tools appropriate for study designs and will resolve discrepancies through discussion with the review team. Where appropriate, we will conduct meta-analyses to estimate the effectiveness of school nurse-led sexual and reproductive health education interventions. Additionally, a qualitative evidence synthesis will be performed for the qualitative research included in the review. Finally, both a quantitative synthesis and a qualitative synthesis will be combined into a secondary synthesis addressing the facilitators and barriers of sexual and reproductive health educational interventions provided by school nurses to adolescents.

Ethics and dissemination This review will synthesise publicly available resources and does not require ethical approval. The findings will provide insights into how school nurses can improve students' sexual and reproductive health. The results will be disseminated through peerreviewed publications, reports and academic conferences.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This mixed-methods systematic review will adhere to the Joanna Briggs Institute (JBI) mixed-methods systematic review manual.
- ⇒ Methodological quality assessment will be conducted using the Cochrane Risk of Bias V.2 for randomised controlled trials and cluster-randomised trials, the Risk of Bias Assessment Tool for Nonrandomized Studies V.2 for nonrandomised studies and the JBI critical appraisal checklist for qualitative research.
- ⇒ Applying a mixed-methods systematic review to identify the effects of school nurse-led sexual and reproductive health interventions, along with information on facilitators and barriers to implementing these interventions in schools, will help maximise the review's findings to inform practice and policy.
- ⇒ Because this review focuses on studies conducted in high-income countries, the findings may not be generalised to low-income and middle-income countries with different school health systems and sociocultural contexts.

PROSPERO registration number CRD42022347625.

INTRODUCTION

Adolescence aged 10–19 years is a unique and critical period in an individual's life.¹ Moreover, it is characterised by rapid physical, cognitive and psychosocial changes, which might be exposed to a variety of sexual and reproductive risks.² Reproductive health is not simply the absence of disease or infirmity in all matters related to the reproductive system but a state of complete physical, mental and social well-being, which is very important in adolescence and has implications for the next generation's health.³

According to the literature, adolescents might face problems such as early pregnancy

data mining, AI training, and similar

Protected by copyright, including for uses related to text and



and birth, abortion, violence, unintended pregnancy, genital tract infections (RTIs) and sexually transmitted infections (STIs).² Therefore, age-specific sexual and reproductive health interventions are important to improve the knowledge and skills necessary for healthy sexual health across the lifespan.⁴ It is also an important time for young people to learn how to think critically about sex, make informed decisions and maintain healthy sexual relationships.⁵

Based on the 2020 report by the Centers for Disease Control and Prevention,⁶ only a small percentage of secondary schools across the USA provide services related to sexual health. While most schools provide medication administration for students with chronic diseases (86.1%), emergency management and medication provision (75.0%) and case management for students with chronic diseases (74.5%), the percentage of schools providing services related to HIV (2.2%), sexually transmitted diseases (STDs) (2.9%) and pregnancy management (2.7%) is notably low. Similarly, since September 2020, the UK has required all schools to implement relationships and sex education, addressing topics like puberty, STIs, healthy relationships and consent, tailored to students' age and maturity.⁷⁸ However, limited resources, teachers' expertise and parental opposition hindered the effective implementation of sexual and reproductive health education in schools.⁹ A safe and healthy school environment improves students' quality of life and learning effectiveness.¹⁰ Health education is important in school as it greatly influences students' future habits and attitudes.¹⁰ Given the school environment, school nurses can provide crucial support for accessible gender and reproductive health education interventions and counselling for adolescents.^{11 12}

School nurses can work with other experts to develop and implement programmes to provide appropriate information based on sexual health, contraception, STIs, safe sex relationships and adolescents' age.^{13 14} According to prior works, when students perceived high closeness to their teachers (in the context of HIV and pregnancy prevention), they valued what they learnt.¹⁵ Also, effective sexual health education in schools largely depends on the abilities of the teachers responsible for delivering it.¹⁶ In addition, the attitudes and expertise of educators play a significant role in the effectiveness of this education.¹⁶ Research showed that most teachers do not feel ready to manage and teach adolescent sexual issues and, in most cases, consider sexuality only from a biological point of view and not in a social context.¹⁷ School nurses could play a crucial role in providing sexual and reproductive health education as health professionals with clinical expertise.^{18 19}

Previous systematic literature reviews focused on parental and family-based adolescent sexual health interventions²⁰; the effectiveness of comprehensive sex education²¹; sexual health issues and mental health²²; pregnancy and STDs²³; contraception²⁴ and substance abuse and sexual violence.²⁵ The effects and implementing factors

<page-header><text><text><text><text><section-header><section-header><section-header><section-header><text><text>

on providing clinical health services³² were excluded, as they do not align with the study's objective of evaluating educational interventions provided by school nurses.

Comparisons

The comparison is defined as groups receiving sexual and reproductive health education provided by research nurses or teachers instead of school nurses or groups that did not receive any educational intervention.

Outcomes of interest

Primary outcomes

Sexual and reproductive health knowledge: the impact of sexual and reproductive health education interventions provided by school nurses on adolescents' knowledge (eg. STIs and AIDS/HIV; contraceptive methods; comprehensive knowledge of condom use, abstinence, pregnancy, unsafe abortion, sexual violence and sexual abuse; gender; sexual orientation and gender identity/expression; gender equality and adult rights).

Attitudes towards sexual and reproductive health: whether students' sexual attitudes change after receiving an educational intervention from school nurses to improve knowledge about sexual and reproductive health (eg, perceptions and practices related to contraception, attitudes towards condoms, abstinence, delaying sexual intercourse, attitudes and intentions towards HPV vaccination).

Sexual and reproductive health behaviours: these include contraceptive use (eg, condom use), age at sexual initiation, abstinence, number of sexual partners and sexual risk behaviours (eg, unprotected sexual activity).³³

Health and social outcomes: these refer to the consequences of such sexual behaviours, including pregnancy rates, incidence of STIs and other related indicators.³⁴

Secondary outcomes

Facilitators and obstacles to implementing sexual and reproductive health education interventions led by school nurses.

Study design

This review will include all primary research study designs, including quantitative, qualitative and mixed-methods research. Only studies published in English or Korean will be included.

Setting

To include the studies conducted in HICs, we will adhere to HICs' lists based on the World Bank per capita Gross National Income in 2024.³⁵

Search strategy

The following databases will be searched to identify relevant studies: the Cochrane Library, Ovid-Embase, PsycINFO, CINAHL, Ovid-Medline, Koreamed and ScienceON. Medical Subject Headings and text words related to school nurse-led sexual and reproductive health will be used in the search process. We will also

use the search filters to exclude articles published in low-income and middle-income countries.³⁶ The search will cover studies from each database's inception to April 2024. The Ovid-MEDLINE search strategies are demonstrated online supplemental appendix 1.

Study selection

After performing the initial search, the search results go through a two-step screening process. Two independent reviewers (YMC and SN) evaluate the title and abstract of the study separately to determine whether they meet the predefined inclusion criteria and excludes studies that do not meet these criteria. Disagreement will be resolved through discussion or 8 consultation with a third reviewer. The study selection process will be visualised using the PRISMA 2020 flowchart.³⁷ The screening and selection will be carried out using Covidence, a systematic review software developed by Veritas Health Innovation in Melbourne, Australia (http://www.covidence.org).

Risk-of-bias assessment

Risk-of-bias assessment After screening their full text, two reviewers will inde-pendently evaluate the risk of bias included in the review. The risk of bias in the included studies will be applied according to study designs.³⁸ The Cochrane Collaboration Risk-of-Bias 2 (RoB 2.0) tool for đ randomised controlled trials (RCTs) will be used to e assess the risk of bias in the included studies.³⁹ The tool is structured into five domains through which bias can be introduced. The domains cover all types of trends that may affect the results of randomised a trials, namely: (1) bias arising from the randomisa- \exists tion process, (2) bias due to deviations from intended interventions, (3) bias due to missing outcome data, (4) bias in measurement of the outcome and (5) bias in selection of the reported result. The answer options for the questions are 'yes' (Y), 'probably yes' (PY), 'probably no' (PN), 'no' (N) and 'no information' (NI). Responses to questions provide the basis for domain-level judgments about the risk of bias, and then, these domain-level judgments provide the basis for a general risk-of-bias judgement for the outcome of the study being evaluated. The possible judgments of overall risk of bias are (1) low risk of bias, (2) some concerns and (3) high risk of bias.

For the risk-of-bias assessment in cluster RCTs, the RoB 2.0 tool for cluster-randomised trials (RoB 2.0 8 CRTs) developed by the Cochrane Collaboration will be applied.⁴⁰ The RoB 2.0 CRT tool evaluates six domains of potential bias arising during the randomisation process as follows: (1) bias arising from the randomization process (domain 1a: risk of bias arising from the randomisation process), (2) bias arising from the timing of identification or recruitment of participants in a CRT (domain 1b: risk of bias arising from the timing of identification or recruitment of

participants), (3) bias due to deviations from intended intervention (effect of assignment to intervention/ effect of adhering to intervention), (4) risk of bias due to missing outcome data, (5) risk of bias in measurement of the outcome and (6) risk of bias in selection of the reported result. The answer options for the questions are 'yes' (Y), 'probably yes' (PY), 'probably no' (PN), 'no' (N) and 'no information' (NI). The overall risk of bias for each domain will be decided as (1) low risk of bias, (2) some concerns and (3) high risk of bias.

For non-random studies (cohort studies, case–control studies and before-and-after studies), a Revised Risk of Bias Assessment Tool for Nonrandomized Studies of Interventions V.2 (RoBANS 2) will be used.⁴¹ The RoBANS 2 tool includes comparability of the target group, target group selection, confounders, measurement of intervention/exposure, binding of assessors, outcome assessment, incomplete outcome data and selective outcome reporting. Based on the RoBANS 2 guidance, we will judge the overall risk of bias in each study as 'high', 'unclear' or 'low'.

For qualitative studies, the JBI critical appraisal checklist for qualitative research will be used.⁴² The JBI tool consists of ten questions with possible answers of yes, no, unclear or not applicable. A comprehensive evaluation will be conducted based on the JBI assessment guide to determine whether the study meets adequate methodological quality standards.⁴²

Data extraction

Two independent reviewers (YMC and SN) will be conducted to extract data using a prespecified template form after pilot testing. The data extracted will include specific details about the participants, intervention, context, study methods, outcome data and study findings relevant to the review questions.⁴³ Cross-checking will be conducted for the accuracy and completeness of the extracted data, and the discrepancy will be resolved through discussion and consultation with a third reviewer.

Statistical analysis

For quantitative data, meta-analysis will be employed to generate pooled estimates to assess the effects of school nurse-led sexual and reproductive health education. Relevant indicators will use the standardised mean difference for continuous variables and relative risk or OR for categorical variables.⁴⁴ We will use a random-effects model as the interventions and populations are likely to be heterogeneous across included studies. The heterogeneity of intervention effects across studies using the I² and the Q-statistic will be assessed.⁴⁵ A funnel plot will be used to detect any publication bias if more than 10 studies are included in this analysis.⁴⁶ Additionally, subgroup analyses will be performed according to study designs, such as randomised or cluster-controlled trials versus BMJ Open: first published as 10.1136/bmjopen-2024-087528 on 18 February 2025. Downloaded from http://bmjopen.bmj.com/ on June 7, 2025 at Agence Bibliographique de l Enseignement Superieur (ABES) . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

non-randomised studies or outcomes of interest, such as knowledge, attitudes and behaviours, or the cultural context of countries such as Asian HICs versus Western HICs. If a meta-analysis is inadequate to conduct because of the heterogeneity of studies included in the review, we will summarise and synthesise the results using a Synthesis Without Meta-analysis methodology, such as harvest or albatross plots.⁴⁷

Regarding qualitative evidence synthesis, we will adopt thematic synthesis using 'best fit' framework approach.⁴⁸ Qualitative studies will be coded sentence by sentence based on units of meaning, applied to both first-order and second-order constructs demonstrated in the results and discussion sections of articles, to capture both primary participants' opinions and authors' interpretations.⁴⁹ Finally, we will combine both quantitative evidence synthesis and qualitative evidence synthesis using convergent segregated approaches from qualitative evidence synthesis to inform the findings of quantitative evidence synthesis.^{50 51}

Patient and public involvement

Patients and/or the public were not involved in this research's design, conduct, or dissemination plans.

Ethics and dissemination

This review will synthesize publicly available resources and does not require ethical approval. The findings will provide insights into how school nurses can improve students' sexual and reproductive health. The results will be disseminated through peer-reviewed publications, reports and academic conferences.

Author affiliations

¹Graduate School of Chungnam National University, Daejeon, Korea (the Republic of)
²College of Nursing, Ewha Womans University, Seoul, Korea (the Republic of)
³College of Nursing, Chungnam National University, Daejeon, Korea (the Republic of)
⁴Seoul Wolchon Elementary School, Seoul, Korea (the Republic of)

Acknowledgements We acknowledge the BK21 FOUR project at the College of Nursing of Chungnam NationalUniversity for their support.

Contributors H-JS contributed to the study's conceptualisation and design. YMC and SN wrote the initial draft of the manuscript. H-JS is the guarantor of the review. JY reviewed the initial and revised manuscript. All authors read, provided feedback and approved the final manuscript.

Funding This study was supported by the Basic Science Research Program of the National Research Foundation of Korea, funded by the Ministry of Science, ICT and Future Planning (grant number NRF-2021R1A2C1093100).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not required

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content

includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iD

Hyun-Ju Seo http://orcid.org/0000-0001-9019-1135

REFERENCES

- Liang M, Simelane S, Fortuny Fillo G, et al. The State of Adolescent Sexual and Reproductive Health. J Adolesc Health 2019;65:S3-15.
- 2 World Health Organization. WHO recommendations on adolescent sexual and reproductive health and rights. Geneva: World Health Organization; 2018. Available: https://apps.who.int/iris/bitstream/han dle/10665/275374/9789241514606-eng.pdf?ua=1
- Bernstein S, Hansen CJ. Public choices, private decisions: sexual 3 and reproductive health and the millennium development goals. New York: United Nations Millennium Project; 2006. Available: https://www.themimu.info/sites/themimu.info/files/documents/ RefDoc_Health_PublicChoices-PrivateDecisions-SRHandMDGs_ UNMillenniumProject_2006.pdf
- Mantula F, Chauraya Y, Danda G, et al. Perspective chapter: sexual 4 health interventions for adolescents. In: Sterling R, ed. Sexual education around the world - past, present and future issues. London: IntechOpen, 2023.
- 5 Planned Parenthood. What are the goals of sex education for youth. 2020. Available: https://www.plannedparenthood.org/learn/foreducators/what-are-goals-sex-education-youth
- Centers for Disease Control and Prevention. School health profiles 6 2020. Atlanta (GA): Centers for Disease Control and Prevention; 2020. Available: https://www.cdc.gov/school-health-profiles/media/ pdf/cdc-profiles-2020.pdf
- Department for Education. Relationships and sex education (RSE) 7 and health education: statutory guidance on relationships education, relationships and sex education (RSE), and health education. London: Department for Education; 2019. Available: https://www.gov. uk/government/publications/relationships-education-relationshipsand-sex-education-rse-and-health-education
- Ponsford R, Meiksin R, Melendez-Torres GJ, et al. The 8 implementation of compulsory relationships and sex education in English secondary schools: qualitative research in the 2022-23 school year. Sex Educ 2024;1-17.
- 9 Aslan F. School-Based Sexual Health Education for Adolescents in Turkey: A Systematic Review. Community Health Equity Res Policy 2022;42:135-43.
- Korean Educational Development Institute. Status of school health 10 in the United States. 2020. Available: https://edpolicy.kedi.re.kr/frt/ boardView.do?strCurMenuId=10091&nTbBoardArticleSeq=826412
- 11 Bundy DAP, Schultz L, Sarr B, et al. The school as a platform for addressing health in middle childhood and adolescence. In: Bundy DAP, Silva ND, Horton S, et al, eds. Child and adolescent health and development. 3rd edn. Washington (DC): The International Bank for Reconstruction and Development / The World Bank, 2017.
- 12 Pavelová Ľ, Archalousová A, Slezáková Z, et al. The Need for Nurse Interventions in Sex Education in Adolescents. Int J Environ Res Public Health 2021:18:492.
- Silivri M, Wirf T, Hodges EA, et al. Conversations About Sexual and 13 Reproductive Health and Rights-From a School Nurse Perspective. J Sch Nurs 2023;39:406-14.
- Smith S, Platt JM, Clifford D, et al. A State-Level Examination of 14 School Nurses' Perceptions of Condom Availability Accompanied by Sex Education. J Sch Nurs 2020;36:386-93.
- 15 Dickson E, Parshall M, Brindis CD. Isolated Voices: Perspectives of Teachers, School Nurses, and Administrators Regarding Implementation of Sexual Health Education Policy. J Sch Health 2020:90:88-98
- 16 Lodge A, Duffy M, Feeney M. 'I think it depends on who you have, I was lucky I had a teacher who felt comfortable telling all this stuff'. Teacher comfortability: key to high-quality sexuality education? Ir Educ Stud 2024;43:171-88.

- Beech S, Sayer L. Analysis of sexual healthcare provided by school 17 nurses. Prim Health Care Res Dev 2018;19:288-300.
- 18 Borawski EA, Tufts KA, Trapl ES, et al. Effectiveness of health education teachers and school nurses teaching sexually transmitted infections/human immunodeficiency virus prevention knowledge and skills in high school. J Sch Health 2015;85:189–96.
- 19 Brewin D, Koren A, Morgan B, et al. Behind closed doors: school nurses and sexual education. J Sch Nurs 2014;30:31-41.
- 20 Santhya KG, Jejeebhoy SJ. Sexual and reproductive health and rights of adolescent girls: evidence from low- and middle-income countries. Glob Public Health 2015;10:189-221.
- 21 Goldfarb ES, Lieberman LD. Three Decades of Research: The Case for Comprehensive Sex Education. J Adolesc Health 2021;68:13-27.
- Mori C, Temple JR, Browne D, et al. Association of Sexting With 22 Sexual Behaviors and Mental Health Among Adolescents. JAMA Pediatr 2019;173:770.
- Mason-Jones AJ, Sinclair D, Mathews C, et al. School-based 23 interventions for preventing HIV, sexually transmitted infections, and pregnancy in adolescents. Cochrane Database Syst Rev 2016;11.
- Munakampe MN, Zulu JM, Michelo C. Contraception and abortion 24 knowledge, attitudes and practices among adolescents from low and middle-income countries: a systematic review. BMC Health Serv Res 2018:18:909
- 25 Piolanti A, Foran HM. Efficacy of Interventions to Prevent Physical and Sexual Dating Violence Among Adolescents: A Systematic Review and Meta-analysis. JAMA Pediatr 2022;176:142-9.
- 26 Shackleton N, Jamal F, Viner R, et al. Systematic review of reviews of observational studies of school-level effects on sexual health, violence and substance use. Health Place 2016;39:168–76.
- 27 Desrosiers A, Betancourt T, Kergoat Y, et al. A systematic review of sexual and reproductive health interventions for young people in humanitarian and lower-and-middle-income country settings. BMC Public Health 2020;20:666.
- 28 Higgins JP, Thomas J, Chandler J, eds. Cochrane handbook for systematic reviews of interventions. Version 6.2. Cochrane. 2021. Available: https://training.cochrane.org/handbook
- Stern C, Lizarondo L, Carrier J, et al. Methodological guidance for 29 the conduct of mixed methods systematic reviews. JBI Evidence Synthesis 2020:18:2108–18
- Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for 30 systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Syst Rev 2015;4:1.
- UNESCO. International technical guidance on sexuality education: an evidence-informed approach. Revised edition. Paris: United Nations Educational, Scientific and Cultural Organization; 2018. Available: https://www.unaids.org/sites/default/files/media_asset/ ITGSE_en.pdf
- Kjolhede C, Lee AC, Council on School Health. School-Based Health 32 Centers and Pediatric Practice. Pediatrics 2021;148:e2021053758.
- 33 Kann L, McManus T, Harris WA, et al. Youth Risk Behavior Surveillance - United States, 2015. MMWR Surveill Summ 2016:65:1-174.
- World Health Organization. Developing sexual health programmes: 34 a framework for action. Geneva: World Health Organization; 2010. Available: https://www.who.int/publications/i/item/WHO-RHR-HRP-10.22
- 35 World Bank Open Data. High income. Washington (DC): World Bank; 2024. Available: https://data.worldbank.org/income-level/highincome
- 36 Salam RA, Faqqah A, Sajjad N, et al. Improving Adolescent Sexual and Reproductive Health: A Systematic Review of Potential Interventions. J Adolesc Health 2016;59:S11-28.
- 37 Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. Syst Rev 2021;10:89.
- Seo H-J, Kim SY, Lee YJ, et al. A newly developed tool for 38 classifying study designs in systematic reviews of interventions and exposures showed substantial reliability and validity. J Clin Epidemiol 2016:70:200-5.
- Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for 39 assessing risk of bias in randomised trials. BMJ 2019;366.
- 40 Eldridge S, Campbell M, Dahota A. Revised cochrane risk of bias tool for randomized trials (RoB 2.0): additional considerations for cluster-randomized trials. Cochrane Methods Cochrane Database Svst Rev 2016:10.
- Seo HJ, Kim SY, Lee YJ, et al. RoBANS 2: A Revised Risk of Bias 41 Assessment Tool for Nonrandomized Studies of Interventions. Korean J Fam Med 2023:44:249-60.
- 42 Lockwood C, Munn Z, Porritt K. Qualitative research synthesis: methodological guidance for systematic reviewers utilizing metaaggregation. Int J Evid Based Healthc 2015;13:179-87.

and similar technologies

Open access

- 43 Munn Z, Tufanaru C, Aromataris E. JBI's systematic reviews: data extraction and synthesis. *Am J Nurs* 2014;114:49–54.
- 44 Cuijpers P, Weitz E, Cristea IA, *et al.* Pre-post effect sizes should be avoided in meta-analyses. *Epidemiol Psychiatr Sci* 2017;26:364–8.
- 45 Higgins JPT, Thomas J, Chandler J, et al, eds. Cochrane handbook for systematic reviews of interventions. Version 6.4. Cochrane, 2023. Available: http://www.training.cochrane.org/ handbook
- 46 Shea BJ, Grimshaw JM, Wells GA, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. BMC Med Res Methodol 2007;7:10.
- 47 Deeks JJ, Higgins JPT, Altman DG, *et al.* Chapter 10: analysing data and undertaking meta-analyses. In: Higgins JPT, Thomas J, Chandler J, et al, eds. *Cochrane handbook for systematic reviews of*

interventions. Version 6.5. Cochrane, 2024. Available: https://training. cochrane.org/handbook

- 48 Booth A, Carroll C. How to build up the actionable knowledge base: the role of 'best fit' framework synthesis for studies of improvement in healthcare. *BMJ Qual Saf* 2015;24:700–8.
- 49 Salmi EM, Basile FW, Khan FA, *et al*. Facilitators and barriers affecting the implementation of e-health for chronic respiratory diseases in remote settings: a qualitative evidence synthesis. *BMC Health Serv Res* 2025;25:19.
- 50 Pearson A, White H, Bath-Hextall F, *et al*. A mixed-methods approach to systematic reviews. *Int J Evid Based Healthc* 2015;13:121–31.
- 51 Hong QN, Rees R, Sutcliffe K, et al. Variations of mixed methods reviews approaches: A case study. *Res Synth Methods* 2020;11:795–811.