BMJ Open What are the symptom trajectories of self-regulatory fatigue among family caregivers of stroke survivors? A protocol of mixed-methods study in **Chinese rehabilitation settings**

Chao-Yue Xu (10),¹ Ping Zou (10),² Xi Chen (10),³ Shu-Lin Li,¹ Jia-Chun You,¹ Zhi-Qing He (10),¹ Yan-Jin Huang¹

ABSTRACT

To cite: Xu C-Y. Zou P. Chen X. et al. What are the symptom trajectories of self-regulatory fatigue among family caregivers of stroke survivors? A protocol of mixed-methods study in Chinese rehabilitation settings. BMJ Open 2025;15:e092724. doi:10.1136/ bmjopen-2024-092724

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (https://doi.org/10.1136/ bmjopen-2024-092724).

Received 21 August 2024 Accepted 12 February 2025

Check for updates

C Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

¹School of Nursing, University of South China Hengyang Medical School, Hengyang, Hunan, China ²School of Nursing, Nipissing University, North Bay, Ontario, Canada

³School of Nursing, The Hong Kong Polytechnic University School of Nursing, Hong Kong, China

Correspondence to Dr Yan-Jin Huang: 2019000050@usc.edu.cn

Introduction Stroke presents a considerable burden not only to patients but also to their families and society at large. In many instances, stroke patients opt for home rehabilitation, relying on family caregivers for daily assistance. This dynamic significantly influences the physiological, psychological and social well-being of these caregivers. Despite its importance, the phenomenon of self-regulatory fatigue (SRF) among family caregivers has received insufficient attention in the literature. Therefore, the objective of this study is to investigate the levels of SRF, the characteristics of associated symptoms and the trajectories of symptom change experienced by family caregivers of stroke patients.

Methods and analysis This research employs a mixed-methods approach, combining a cross-sectional study with a prospective longitudinal quantitative and gualitative design. The Chinese version of the SRF Scale and the Chinese version of Patient-Reported Outcomes Measurement Information System profile-29 are used to assess SRF, psychological and physiological symptoms, and related functional outcomes among family caregivers of stroke patients. Latent class growth analysis will be employed to model the heterogeneous developmental trajectories of SRF-related symptoms among family caregivers of stroke patients. Reflexive thematic analysis will be employed to analyse, organise and summarise qualitative data, to identify the experiences and management needs related to SRF among family caregivers during home care. Through this comprehensive mixed-methods approach, the study aims to: investigate the levels of SRF experienced by family caregivers of stroke patients, identify patterns and trajectories of related symptoms. The integration of cross-sectional and longitudinal data allows for a thorough examination of both immediate and long-term aspects of caregiver experiences, providing valuable insights into the complex dynamics of SRF in this population.

Ethics and dissemination The study protocol was approved by the Medical Ethics Committee of the University of South China (approval number 2023-NHHL-051). Data collection was authorised by the ethics committees of the First Affiliated Hospital. Second

 Suppose the property of the pro

not been effectively controlled. In recent years, China has experienced explosive growth in stroke incidence, with a concerning trend towards younger patients. The 'Annual Report on Cardiovascular Health and Diseases in China (2022 Edition)' highlights the substantial economic impact of stroke, reporting average annual growth rates of 16.81% and 12.79% in total hospitalisation costs for cerebral infarction and cerebral haemorrhage, respectively, since 2004.³ These statistics underscore the urgent need for improved stroke prevention and management strategies in China.

The role of family caregivers in poststroke rehabilitation is crucial and deserves significant attention. Research indicates that home caregivers are associated with a reduced risk of hospitalisation for stroke patients compared with those without such support.⁴ van Dijk and Buijck emphasise that informal caregivers are often considered the ideal source of care for stroke survivors.⁵ In China, cultural traditions and family values influence care decisions, with nearly 90% of stroke patients opting for home-based rehabilitation after hospital discharge. This trend underscores the importance of integrating caregivers into the rehabilitation process as essential health partners for patients. Family caregivers often endure prolonged physical and emotional exhaustion, which adversely affects their overall physical and mental health.^{7 8} According to the results of a cross-sectional survey, 43.9% of family caregivers have anxiety symptoms, and 53.9% of family caregivers have varying degrees of depression symptoms. About 68.4% of family caregivers bear a moderate to severe burden from caregiving.⁹ Two studies emphasise that caregivers often exhibit signs of stress, isolation and depression.¹⁰¹¹ According to a metaanalysis, approximately 40% of caregivers of stroke survivors experienced depressive symptoms at different time points during the assessment process.¹² Moreover, 18% of caregivers reported such symptoms within the first 4 months of caring for patients.¹³ The decline in the wellbeing of family caregivers is closely associated with the deteriorating health outcomes of the patients they care for. Em et al found that the physical and mental health of caregivers correlates with the severity of physical dysfunction in stroke patients.¹⁴ Furthermore, Pucciarelli et al suggest that improving caregivers' health can positively impact the functional recovery and daily living activities of stroke survivors.¹⁵ Given these findings, healthcare professionals should prioritise interventions that support the physical and mental health of caregivers. Such interventions can indirectly enhance patients' functional status and overall recovery. Currently, there is a paucity of personalised care plans that adequately address the heterogeneity among family caregivers of stroke patients.¹⁶ Additionally, there is a limited body of literature exploring the incorporation of self-regulatory fatigue (SRF) as a more advanced indicator for assessing the symptoms, functional capacities and burdens experienced by family caregivers during the poststroke recovery process.¹⁷ This gap highlights the need for further research in understanding and fulfilling

<page-header><text><text><text><text>

not widely used.^{32 34 35} This study will employ the Chinese version of the SRF Scale (SRF-S) and the Patient-Reported Outcomes Measurement Information System (PROMIS). These tools will enable us to accurately identify SRF and related symptoms in family caregivers. This, in turn, will allow us to provide guidance and references for future research on symptom improvement and enhance caregivers' understanding of disease processes and health management.

Therefore, this study will investigate the aetiology and categorise the manifestations of SRF-related symptoms among family caregivers of stroke patients. We will examine the phenomenology of SRF-related symptoms experienced by primary family caregivers throughout various stages of the caregiving continuum. A prospective longitudinal analysis will be conducted to assess the temporal dynamics of SRF-related symptoms among these caregivers. The findings from this research will provide a reference for the development of a comprehensive management plan addressing SRF for family caregivers during the continuity of care process for stroke patients in the future.

Research objectives

- 1. To investigate the aetiology and categorise the manifestations of SRF-related symptoms among family caregivers of stroke patients.
- 2. To examine the phenomenology of SRF-related symptoms experienced by primary family caregivers of stroke patients across various stages of the caregiving continuum.
- 3. To conduct a prospective longitudinal analysis of the temporal dynamics of SRF-related symptoms among family caregivers.

METHODOLOGY

Study design

This study adheres rigorously to the Good Reporting of A Mixed Methods Study checklist (online supplemental material 1), ensuring comprehensive and transparent reporting of our mixed-methods approach.³⁶ The qualitative research section of this article will adhere to the Consolidated Criteria for Reporting Qualitative Studies checklist (online supplemental material 2) designed to enhance the quality and transparency of qualitative research, particularly in the context of interviews and focus groups.³⁷ The focus of this study is to analyse the temporal dynamics and influencing factors of SRF-related symptoms among family caregivers of stroke patients through longitudinal observation. To enhance this analysis, longitudinal qualitative research will simultaneously capture the emotional experiences of family caregivers at various stages of caregiving. Consequently, an embedded design was selected for this study, allowing for the integration of qualitative insights within the predominantly quantitative framework.³⁸ This mixed-methods study employs a concurrent embedded design, comprising

two primary components.³⁹ Quantitative component: (a) cross-sectional survey-to elucidate the typology of SRFrelated symptoms among primary caregivers of stroke patients. (b) Prospective longitudinal study: to delineate the trajectory of SRF-related symptoms among primary caregivers over time. Qualitative component: longitudinal phenomenological inquiry⁴⁰—to explore the lived experiences of SRF symptoms among family caregivers of stroke patients with long-term functional impairments and self-care deficits. The study will employ a **p** multiphase data collection strategy. T0 (baseline): crosssectional survey administration and first qualitative interview within 1 month of the postcerebrovascular event. T1 (3 month follow-up): quantitative assessments and **Z** second qualitative interview. T2 (6month follow-up): guantitative assessments—final quantitative assessments and third qualitative interview. The entire study will last from November 2024 to August 2025. The diagram of the study is presented in figure 1.

Qualitative data will be used to contextualise and expand on quantitative findings, particularly in interpreting symptom scores and extreme values. This integra-SRF among family caregivers that neither method could g tion aims to provide a comprehensive understanding of achieve independently. The findings from this study will inform the development of evidence-based interventions lated to text and for managing SRF among family caregivers of stroke patients in subsequent research.

Quantitative research section

Sample size

data When using Bayesian Information Criteria as the primary consideration for model selection, a minimum sample size of 200 cases is recommended.⁴¹ The robustness of latent class models for analysing variable trajectory categories is also influenced by sample size, with larger sample ٩ sizes (up to 500 cases) providing more reliable results. training Therefore, considering both statistical requirements and the practical feasibility of data collection, this study has determined a minimum sample size of 200 cases. , and simila

Settings and sample recruitment

A cluster sampling method will be employed to recruit participants. The study will include all family caregivers of stroke patients who meet the inclusion and exclusion criteria from the rehabilitation and neurology departments of three tertiary hospitals in Hengyang City, Hunan Province. Preliminary estimates suggest that, on average, the neurology department at each hospital admits approximately 4000 patients annually, while the rehabilitation department admits about 400 patients per year. This patient volume provides a robust foundation for our recruitment efforts. The participants will be initially recruited in the ward, and subsequent follow-ups will be expected to be completed in rehabilitation settings. This approach ensures a representative sample and facilitates the practical implementation of the study.



Figure 1 Research diagram. PROMIS-29, Patient-Reported Outcomes Measurement Information System profile-29; SRF-S, Self-Regulatory Fatigue Scale.

Eligibility criteria for samples

The conditions that stroke patients meet include: (1) with first-ever stroke; (2) impaired self-care ability, defined as a Modified Barthel Index score <60 (26), assessed weekly during hospitalisation; (3) residence within Hengyang city limits; (4) absence of comorbid major diseases, malignancies, end-stage conditions or need for dialysis and (5) no history of intermittent psychiatric disorders. Family caregivers inclusion criteria:

- 1. Primary caregiver status, defined as the individual providing the most time-intensive care and assuming the majority of caregiving responsibilities.
- 2. Minimum age of 18 years.
- 3. Cognitive capacity to comprehend questionnaire content and provide accurate responses.
- 4. Provision of written informed consent and willingness to participate in the study protocol.

Family caregivers exclusion criteria:

- 1. Remunerated caregiving status.
- 2. Formal training in professional caregiving.
- 3. Personal history of psychiatric illness according to medical records review or assessment of the Chinese Health Questionnaire (CHQ).⁴²
- 4. Presence of severe medical conditions that compromise caregiving capacity.
- 5. Recent exposure to significant familial stressors.
- Family caregivers discontinuation criteria:
- 1. Development of severe illness or mortality during the follow-up period, resulting in cessation of caregiving ability.
- 2. Refusal of follow-up assessments or voluntary withdrawal from the study.

Data collection method

The collection of quantitative data will be conducted at three-time points between November 2024 and June 2025: T0 (1 month postcerebrovascular event), T1 (3 months postcerebrovascular event) and T2 (6 months postcerebrovascular event). After obtaining permission from the University of South China and the three tertiary hospitals, three postgraduate students will undergo research assistant (RA) training conducted by the first author (C-YX) and corresponding author (Y-JH). This training will encompass all research procedures, including participant recruitment based on the established inclusion and exclusion criteria, management of informed consent forms and techniques for administering questionnaires. 1 week following the admission of the stroke patient, the responsible nurse, along with the head nurse, will assess the eligibility of the family caregiver. They will provide a comprehensive explanation of the research objectives and procedures to the caregiver and subsequently obtain their written informed consent. The survey questionnaire will be administered on-site by RA (J-CY, S-LL and Z-QH) to ensure consistency and minimise bias. RA will provide standardised instructions on how to complete the questionnaire, ensuring uniformity in the data collection process across different researchers. All participants will be required to complete the questionnaire independently. In cases where participants are unable to read or write, RA will offer oral assistance to facilitate the completion of the questionnaire. If participants have any questions during the process, RA will provide immediate clarification. On completion, RA will review the questionnaires on-site to ensure they are fully completed. If any items are missing, participants will be asked to fill them in immediately. The questionnaires will be rechecked for completeness before being collected.

Research instruments

Patient Demographic and Clinical Characteristics Questionnaire

This instrument will collect comprehensive sociodemographic and clinical data, including: (a) sociodemographic variables: age, gender, anthropometric measures (height and weight), residential status, marital status, educational attainment, occupational status and

<page-header><page-header><text><section-header><section-header><section-header><section-header><section-header><section-header><section-header>

following an evaluation conducted in patients with aortic dissection, with a Cronbach's alpha greater than 0.90. Additionally, the research findings by Sikorskii *et al*⁴⁸ suggest that PROMIS-29 and traditional legacy tools yield comparable conclusions regarding the effects of interventions on pain, fatigue, sleep disturbances, anxiety, as well as physical and social functioning. Furthermore, due to its brevity and minimal response burden, the PROMIS 4-item form is an excellent choice for assessing patientreported outcomes while both measurement systems are valuable for assessing caregiver outcomes.

All items are scored using a 5-point Likert scale, with variations in the descriptions of the response options for each item. Each domain is scored separately, resulting in scores across three primary domains. Furthermore, scores can be converted into standardised T-scores, where the mean is set at 50 with a SD of 10. Higher T-scores indicate a greater degree of the measured construct. For instance, a T-score of 70 in social role functioning suggests that the individual's ability to engage socially is above the average for the general population, reflecting better social functioning. Conversely, a T-score of 70 in depression indicates a higher level of depressive symptoms compared with the general population. The PROMIS tools have been translated into over 40 languages and are widely used in various healthcare settings.

The 6-month period following a stroke is recognised as the optimal timeframe for patient recovery. Concurrently, this period is critical for family caregivers, who may experience substantial impacts on their quality of life during this time.⁴⁹ By including caregivers in this assessment, we gain a holistic view of the healthcare experience.

Chinese version of the SRF-S

The SRF-S was developed by Nes et al.⁵⁰ and consists of 18 items specifically designed to assess SRF in individuals with chronic diseases. The scale has demonstrated good reliability and validity, effectively distinguishing SRF from self-control and physical fatigue, thereby providing a more direct and accurate assessment of an individual's state of self-depletion.

The Chinese version of the SRF-S was translated and validated in young adults by Wang et al.⁵¹ Also validated in elderly maintenance haemodialysis patients and caregivers.¹⁷ It comprises three dimensions: cognitive control, emotional control and behavioural control, with a total of 16 items. Each item is scored using a 5-point Likert scale, ranging from 'strongly disagree' (1) to 'strongly agree' (5), yielding a total score between 16 and 80. Higher scores indicate greater severity of SRF.

Data analysis

Statistical analyses will be conducted using IBM SPSS V.26.0 and R statistical software V. 4.2.1. SPSS 26.0 provides a user-friendly interface for data management and basic statistical analyses, ideal for exploratory and descriptive statistics, while R 4.2.1 excels in handling large datasets and advanced statistical methods. Using both software

to text

ĩ

packages leverages their strengths, enhancing analytical efficiency and allowing for cross-validation of results to improve the robustness of findings. Quantitative data that conforms to a normal distribution will be summarised using mean±SD, while data that do not follow a normal distribution will be described using median and IQR. Categorical data will be presented as frequency and percentage to characterise the general demographic features of the study participants.

To compare demographic differences between study subjects who were lost to follow-up and those who remained, independent sample t-tests and χ^2 tests will be employed. Latent class analysis will be performed to ŝ classify family caregivers into distinct latent groups based on the varying characteristics of symptoms associated with SRF. For intergroup comparisons, t-tests, χ^2 tests and analysis of variance will be used, while logistic regression will be applied to identify independent influencing factors. including Additionally, latent class growth analysis (LCGA) will be employed to model the heterogeneous developmental trajectories of SRF-related symptoms among family caregivers of stroke patients. for uses related

Qualitative research section

Sample size

The sample size for qualitative research is determined by the informational power of the sample size, meaning that the more information the sample has that is relevant to the actual research, the fewer participants are required.⁵² The size of a sample with adequate informational capacity is contingent on the research objectives, the specificity of the sample, the application of established theories, the quality of dialogues and the analytical strategies employed. To ensure the collection of rich and relevant \blacksquare information, we will select participants who can provide valuable insights into the research topic. The anticipated sample size is approximately 10-15 individuals.

Settings and sample recruitment

ning, Al training, In this study, a purposive sampling method will be used to select family caregivers of stroke patients as interview participants. For caregivers of hospitalised stroke patients, qualitative interviews will be conducted in conference rooms within the ward. This setting is chosen to accommodate the significant caregiving burden that many family caregivers experience, providing a supportive environment for the interviews. For caregivers of discharged stroke patients, interviews will take place in their homes or 🗳 in community conference rooms, allowing for a comfort- **8** able and familiar atmosphere that may encourage open and candid discussions.

Data collection

Between November 2024 and June 2025, qualitative interviews will be conducted at three-time points for each stroke patient: T0 (1 month postcerebrovascular event), T1 (3 months postcerebrovascular event) and T2 (6 months postcerebrovascular event). All interviews will be scheduled with the participants at a time and location convenient for them, following the completion of quantitative data collection. All interviews will be conducted on a one-on-one basis, with only the family caregiver and the researcher present. A postgraduate student (C-YX) trained in qualitative interviewing will be responsible for face-to-face data collection. Prior to the interviews, the researcher will prepare all necessary materials and equipment, including informed consent forms, interview outlines, recording devices, notebooks, pens and thank you gifts. Before commencing the interview, the researcher will explain the purpose and significance of the study to the family caregiver. The interviewee will be informed that recording equipment will be used to ensure the completeness of the information gathered during the interview process. It will be emphasised that all content collected is for research purposes only, and the interviewee has the right to withdraw at any time if they experience any discomfort. The researcher will seek the interviewee's understanding and consent regarding these procedures.

Each interview is expected to last between 30 and 60 min. During the interview, the researcher will focus not only on capturing the verbal responses of the interviewee but also on carefully observing and recording non-verbal behaviours, such as facial expressions, tone of voice and body movements, which may reflect psychological changes. The researcher will conduct appropriate follow-ups on valuable information, providing timely responses and employing interview techniques such as repetition and probing questions to facilitate a deeper exploration of the information shared by the interviewee.

Interview content

The interview will focus on open-ended questions designed to elicit in-depth responses from participants. The questions will include:

Understanding of stroke:

"Can you talk about your understanding of stroke? What changes has the patient's illness brought to your life, and how do you respond to these changes?"

Patient care dynamics:

"What is the patient's response to your care while assisting with their daily activities and recovery? How did you feel during the caregiving process, and what changes has the caregiving responsibility brought to you?"

Health and well-being:

"How is your current health condition? How do you feel while caring for the patient? How do you balance the patient's health needs with your own? What is your perspective on the relationship between the patient's health and your health?"

Changes in personal time:

"Has your alone time changed since the patient's illness? How do you feel about these changes?"

Role of the family caregiver:

"How do you understand the role of a family caregiver? What changes do you believe you, as a family member, will bring to the patient's recovery?"

Challenges in caregiving:

"What are the current difficulties you face in caring for the patient and assisting in their rehabilitation process? What kind of support do you hope to receive?"

Definition and recognition of SRF:

"How would you define self-regulatory fatigue in your own words?"

"Have you noticed changes in your ability to manage stress or maintain routines since becoming a caregiver?"

Triggers and symptoms:

Š "What specific situations or tasks contribute most to your feelings of difficulty in maintaining emotional and physiological stability?" "Can you describe any physical or emotional symp-toms you experience that you believe are related to this

Including difficulty?"

Data analysis

All qualitative data analysis will be conducted using NVivo 10 software. A diachronic approach will be adopted for uses r data analysis, meaning that analysis will occur after the completion of all data collection. This approach allows for comprehensive consideration of all data waves, providing a deeper perspective on the changes over time that contribute to understanding the experiences and ç processes of family caregivers.

This structured approach ensures a rigorous analysis process while fostering collaboration among authors at each stage of reflexive thematic analysis (RTA) implementation. The recorded interview content will be transcribed verbatim into written documents, and an archive file will be established by C-YX and double-checked by Y-JH. This file will include essential information such ≥ as material number, basic information about the interviewee, and details regarding the method and location of data collection. RTA will be employed to analyse, organise and summarise the data, to identify the experiences and management needs related to SRF among family caregivers during home care.⁵³ The entire process was conducted under the guidance of three nursing PhD authors-PZ, XC and Y-JH-who possess extensive qualitative research experience in RTA methods for analysing qualitative data. All authors have completed at least 4 months of both theoretical and practical training related to RTA methodologies. In the event of disagreements arising before or during any stage of the process, **g** the corresponding author, Y-JH, will organise a meeting to facilitate discussion on the key challenges and complexities encountered by the authors.

Each specific step will be carried out according to the following arrangement:

Familiarisation with the data (step 1): C-YX, J-CY, S-LL and Z-QH will engage in a thorough review of the interviewrelated documents to ensure comprehensive familiarity with the data. Each author will highlight sections that

Protected

align with the qualitative research objectives and will document relevant notes. Consensus will be achieved through discussions in subsequent meetings. Generating initial codes (step 2): C-YX, J-CY, S-LL and Z-QH will work in pairs to code the materials, take pertinent notes and export the codes from NVivo software in the form of charts. Consensus regarding the coding will be established during a collaborative meeting. Developing themes (step 3): following consensus on the initial coding, C-YX, J-CY, S-LL and Z-QH will collaboratively create an initial thematic map based on the codes to facilitate theme development through discussion. Reviewing themes (step 4): the identified themes will undergo a thorough review to ensure they accurately reflect the dataset. This review will be conducted collectively by all four authors. Defining and naming themes (step 5): during a dedicated meeting, all authors will engage in discussions to finalise the definitions and names of the themes. Writing the report (step 6): C-YX will be responsible for drafting the final section of the qualitative research report.

Patient and public involvement

Patients and the public were not involved in the development of this research proposal or the conduct of the study. The research question and study design were informed by the review of existing evidence and discussions with clinical experts. However, we acknowledge the limitations of not incorporating patient input, particularly in ensuring the relevance and acceptability of the qualitative data collection. In future studies, we will seek patient involvement in refining the research question, selecting appropriate outcome measures and optimising the study design.

ETHICS AND DISSEMINATION

The study protocol was approved by the Medical Ethics Committee of the University of South China (Approval number 2023-NHHL-051). Data collection was authorised by the ethics committees of the First Affiliated Hospital, Second Affiliated Hospital and Nanhua Affiliated Hospital of the University of South China. The results of this study will be disseminated through publication in pertinent peer-reviewed journals, presentation at local and international conferences, and communication with all relevant stakeholders.

DISCUSSION

This study is expected to explore the patterns of SRFrelated symptoms among family caregivers, establishing a foundation for developing management plans addressing maladjustment in family caregivers at different stages of the poststroke recovery. The quantitative component of the study uses self-report data collection, which encourages the active engagement of family caregivers and fosters their participation in maintaining the health of stroke patients. By focusing on individual experiences

BMJ Open: first published as 10.1136/bmjopen-2024-092724 on 6 March Ense http://bmjopen.bmj.com/ on June 8, 2025 at Agence Bibliographique de l

and authentic emotions, this study fosters a deeper understanding among family caregivers regarding their own feelings. This awareness encourages them to take proactive steps in seeking appropriate assistance, ultimately enhancing the social support available for stroke patients. Additionally, the combination of reliable selfreport tools and one-on-one face-to-face interviews allows for a comprehensive and nuanced understanding of the experiences and self-care situations faced by family caregivers following cerebrovascular events.

givers following cerebrovascular events. However, potential limitations of the proposed study include the relatively short tracking period of 6 months. Future research could benefit from longer-term tracking Š to provide a more comprehensive understanding of the long-term SRF experienced by caregivers of stroke copyright, including for patients. Furthermore, the primary indicators used in the quantitative research section are largely self-reported by participants, which may lack the support of objective measures.

CONCLUSION

This protocol outlines a comprehensive mixed-methods study designed to investigate the symptom trajectory of This protocol outlines a comprehensive mixed-methods study designed to investigate the symptom trajectory of SRF-related symptoms in family caregivers of stroke survivors. By combining cross-sectional investigation, quantitative longitudinal surveys with qualitative in-depth interviews, this study aims to provide a nuanced understanding of how SRF develops, progresses and impacts caregivers over time. The proposed research addresses a significant gap in the literature by focusing specifically on SRF, a critical yet understudied aspect of caregiver burden. By mapping the trajectory of SRF-related symptoms, we anticipate identification points and contributing factors that

fying key inflection points and contributing factors that *a* could inform targeted interventions. The mixed-methods ≥ approach will allow for a rich, multifaceted exploration of caregivers' experiences, capturing both the measur-able changes in fatigue levels and the lived experiences of **g** and similar technologies managing this challenge.

Acknowledgements We would like to express our sincere gratitude to Dr. Qiao Xiao for her invaluable assistance in the statistical analysis of this study. E-mail: xo. hnit@gmail.com. Her expertise in mathematicmathematics (statistics)/epidemiology and biostatistics will greatly enhance the quality of our research. We extend our heartfelt appreciation to all the research participants who generously volunteered their time and shared their experiences. Your contributions were invaluable and form the cornerstone of this research.

Contributors C-YX: conceived and designed the study, developed the theoretical framework and wrote the initial draft of the manuscript, PZ, XC, S-LL, J-CY and Z-QH: assisted in the design of the methodology, contributed to drafting sections of the manuscript and contributed to the critical revision of the manuscript for important intellectual content. Y-JH: acted as guarantor. Assisted in the design of the methodology, and reviewed and provided feedback on all drafts of the manuscript, ensuring clarity and coherence.

Funding This work was supported by the Social Science Achievement Review Committee of Hunan Province under Grant number: XSP25YBZ054; the Natural Science Foundation of Hunan Province under Grant number: 2023JJ40558; and the Health Commission of Hunan Province under Grant number: B202314057822.

Competing interests None declared.

for uses related to text and data mining, Al training,

and similar technologies

Protected by copyright, including

Patient and public involvement Patients and/or the public were not involved in the design or conduct or reporting or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Chao-Yue Xu http://orcid.org/0009-0001-4545-8643 Ping Zou http://orcid.org/0000-0002-3973-7830 Xi Chen http://orcid.org/0000-0001-9392-6096 Zhi-Qing He http://orcid.org/0009-0002-7694-3817

REFERENCES

- 1 Yanyan H. Ischemic cerebrovascular disease and regulation of blood pressure. *Chinese Journal of Geriatric Heart Brain and Vessel Diseases* 2012;12:670–2.
- 2 Statistics NBo. Bulletin of the seventh national population census (no. 2). 2021.
- 3 National Center for Cardiovascular Diseases C. Annual report on cardiovascular health and diseases in China. 2022.
- 4 Tyagi S, Koh GC-H, Nan L, *et al.* Healthcare utilization and cost trajectories post-stroke: role of caregiver and stroke factors. *BMC Health Serv Res* 2018;18:881.
- 5 van Dijk M, Buijck BI. What is desirable care in the opinion of formal and informal caregivers in nursing-home care for patients with dementia? *Nurs Open* 2018;5:139–48.
- 6 Wenzhi W. Epidemiological trends and the necessity of community based prevention and control for stroke in China. *Chinese Journal of Health Management* 2017;11:393–6.
- 7 Wan Ab Rahman WMA, Mazlan M. The Prevalence of Depression and Its Correlates Among Informal Caregivers of Stroke Patients in an Urban Community. *Cureus* 2024;16:e51486.
- 8 Blanton S, Clark PC, Cotsonis G, *et al.* Factors associated with depressive symptoms of carepartners of stroke survivors after discharge from rehabilitation therapy. *Top Stroke Rehabil* 2020;27:590–600.
- 9 Hu P, Yang Q, Kong L, et al. Relationship between the anxiety/ depression and care burden of the major caregiver of stroke patients. *Medicine (Baltimore)* 2018;97:e12638.
- 10 Greenwood N, Habibi R, Mackenzie A. Respite: carers' experiences and perceptions of respite at home. *BMC Geriatr* 2012;12:42.
- 11 Haley WE, Roth DL, Hovater M, et al. Long-term impact of stroke on family caregiver well-being: a population-based case-control study. *Neurology (ECronicon)* 2015;84:1323–9.
- 12 Loh AZ, Tan JS, Zhang MW, *et al.* The Global Prevalence of Anxiety and Depressive Symptoms Among Caregivers of Stroke Survivors. *J* Am Med Dir Assoc 2017;18:111–6.
- 13 Byun E, Evans L, Sommers M, et al. Depressive symptoms in caregivers immediately after stroke. *Top Stroke Rehabil* 2019;26:187–94.
- 14 Em S, Bozkurt M, Caglayan M, et al. Psychological health of caregivers and association with functional status of stroke patients. *Top Stroke Rehabil* 2017;24:323–9.
- 15 Pucciarelli G, Vellone E, Savini S, *et al.* Roles of Changing Physical Function and Caregiver Burden on Quality of Life in Stroke: A Longitudinal Dyadic Analysis. *Stroke* 2017;48:733–9.
- 16 Creasy KR, Lutz BJ, Young ME, et al. Clinical Implications of Family-Centered Care in Stroke Rehabilitation. *Rehabil Nurs* 2015;40:349–59.

17 Chai Xiaopei LS, Xuan X, Xing H. Self-regulatory Fatigue on the Quality of Life Among Elderly Maintenance Hemodialysis Patients and Their Caregivers. *Military Nursing* 2023;40:77–80.

Open access

- 18 Ling Qian JS, Wangling S, Liujun L. Analysis of Self-regulatory Fatigue Status and Influencing Factors in Parents of Children with Asthma. J Clin Nurs 2024;23:6–10.
- 19 Xiang Jing CC. Analysis of the current status and influencing factors of self-regulatory fatigue in parents of premature infants. *Modern Nurse* 2023;30:120–3.
- Baumeister RF, Bratslavsky E, Muraven M, *et al.* Ego depletion: is the active self a limited resource? *J Pers Soc Psychol* 1998;74:1252–65.
 Sethager J, Oursla M, Tang W, Ool JD, 200 (1998);74:1252–65.
- 21 Schlaegel C, Gunkel M, Taras V. COVID-19 and individual performance in global virtual teams: The role of self-regulation and individual cultural value orientations. *J Organ Behav* 2023;44:102–31.
- 22 Pandey A, Hale D, Das S, et al. Effectiveness of Universal Selfregulation-Based Interventions in Children and Adolescents: A Systematic Review and Meta-analysis. JAMA Pediatr 2018;172:566–75.
- 23 Shields GS, Moons WG, Slavich GM. Inflammation, Self-Regulation, and Health: An Immunologic Model of Self-Regulatory Failure. *Perspect Psychol Sci* 2017;12:588–612.
- 24 Cameron D, Webb T. Self-regulatory fatigue. In: Gellman MD, Turner JR, eds. *Encyclopedia of Behavioral Medicine*. New York, NY: Springer, 2013: 1760–2.
- 25 Eriku GA, Bekele G, Yitayal MM, *et al.* Depressive Symptoms and Its Associated Factors Among Primary Caregivers of Stroke Survivors at Amhara Regional State Tertiary Hospitals: Multicenter Study. *Neuropsychiatr Dis Treat* 2023;19:1675–84.
- 26 Kumar A, Yadav AK, Singh VK, et al. Caregiver Burden in Caregivers of Stroke Survivors: A Hospital-Based Study. Ann Indian Acad Neurol 2022;25:1092–8.
- 27 Ou PR, Wu MH, Tsai ST, et al. The Relationship Between Social Support and Stress in Family Caregivers of Stroke Patients. J Neurosci Nurs 2024;56:209–13.
- 28 Tyagi S, Luo N, Tan CS, et al. Support system diversity among family caregivers of stroke survivors: a qualitative study exploring Asian perspectives. BMC Geriatr 2021;21:594.
- 29 Evans DR, Boggero IA, Segerstrom SC. The Nature of Self-Regulatory Fatigue and "Ego Depletion": Lessons From Physical Fatigue. *Pers Soc Psychol Rev* 2016;20:291–310.
- 30 Tao Y, Liu T, Li P, et al. Self-management experiences of haemodialysis patients with self-regulatory fatigue: A phenomenological study. J Adv Nurs 2023;79:2250–8.
- 31 Karimirad MR, Seyedfatemi N, Mirsepassi Z, et al. Barriers to Self-Care Planning for Family Caregivers of Patients with Severe Mental Illness. J Patient Exp 2022;9:23743735221092630.
- 32 Lin C, Zhang F, Yang F, et al. Factors influencing self-regulatory fatigue in patients undergoing chemotherapy for gynecologic cancer: a cross-sectional study. Front Psychiatry 2024;15:1273151.
- 33 Lu F, Wong CKH, Ng APP, et al. Effectiveness of a 5-year health empowerment programme on promoting cardiovascular health for adults from low-income families in Hong Kong. *Patient Educ Couns* 2024;124:108240.
- 34 Cui Y, Yang T, Gao H, *et al.* The relationship between ego depletion and work alienation in Chinese nurses: A network analysis. *Front Psychol* 2022;13:915959.
- 35 Ying J, Ren L, Zhang J, *et al.* The network structure of ego depletion in Chinese male young adults. *Front Psychol* 2023;14:1102624.
- 36 O'cathain A, Murphy E, Nicholl J. The Quality of Mixed Methods Studies in Health Services Research. J Health Serv Res Policy 2008;13:92–8.
- 37 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- 38 Schoonenboom J, Johnson RB. How to Construct a Mixed Methods Research Design. Kolner Z Soz Sozpsychol 2017;69:107–31.
- 39 Wang Q, Xu C, He Z, *et al.* An exploration of proactive health oriented symptom patterns in patients undergoing percutaneous coronary intervention with stent implantation: A mixed-methods study protocol. *PLoS ONE* 2023;18:e0292285.
- 40 Lu Q, Mårtensson J, Zhao Y, *et al.* Living on the edge: Family caregivers' experiences of caring for post-stroke family members in China: A qualitative study. *Int J Nurs Stud* 2019;94:1–8.
- 41 Shi D, DiStefano C, Zheng X, et al. Fitting Latent Growth Models with Small Sample Sizes and Non-normal Missing Data. Int J Behav Dev 2021;45:179–92.
- 42 Cheng TA, Wu JT, Chong MY, *et al.* Internal consistency and factor structure of the Chinese Health Questionnaire. *Acta Psychiatr Scand* 1990;82:304–8.
- 43 Goldberg DP, Hillier VF. A scaled version of the General Health Questionnaire. *Psychol Med* 1979;9:139–45.

Open access

- 44 Yang T, Huang L, Wu Z. The application of Chinese health questionnaire for mental disorder screening in community settings in China's mainland. *Chinese Journal of Epidemiology* 2003;23:769–73.
- 45 Evon DM, Amador J, Stewart P, et al. Psychometric properties of the PROMIS short form measures in a U.S. cohort of 961 patients with chronic hepatitis C prescribed direct acting antiviral therapy. *Aliment Pharmacol Ther* 2018;47:1001–11.
- 46 Krohe M, Tang DH, Klooster B, et al. Content validity of the National Comprehensive Cancer Network - Functional Assessment of Cancer Therapy - Breast Cancer Symptom Index (NFBSI-16) and Patient-Reported Outcomes Measurement Information System (PROMIS) Physical Function Short Form with advanced breast cancer patients. *Health Qual Life Outcomes* 2019;17:92.
- 47 Huang W, Wu Q, Zhang Y, et al. Preliminary evaluation of the Chinese version of the patient-reported outcomes measurement information system 29-item profile in patients with aortic dissection. *Health Qual Life Outcomes* 2022;20:94.
- 48 Sikorskii A, Victorson D, O'Connor P, et al. PROMIS and legacy measures compared in a supportive care intervention for breast

cancer patients and caregivers: Experience from a randomized trial. *Psychooncology* 2018;27:2265–73.

- 49 Lee KB, Lim SH, Kim KH, et al. Six-month functional recovery of stroke patients: a multi-time-point study. Int J Rehabil Res 2015;38:173–80.
- 50 Nes LS, Ehlers SL, Whipple MO, *et al.* Self-regulatory fatigue in chronic multisymptom illnesses: scale development, fatigue, and self-control. *J Pain Res* 2013;6:181–8.
- 51 Wang L, Zhang J, Wang J, *et al.* Validity and reliability of the Chinese version of the Self-Regulatory Fatigue Scale in young adults. *Chinese Mental Health Journal* 2015;29:290–4.
- 52 Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview Studies: Guided by Information Power. *Qual Health Res* 2016;26:1753–60.
- 53 Braun V, Clarke V. Supporting best practice in reflexive thematic analysis reporting in *Palliative Medicine*: A review of published research and introduction to the *Reflexive Thematic Analysis Reporting Guidelines* (RTARG). *Palliat Med* 2024;38:608–16.