



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Can culturally safe general practice telehealth overcome barriers to care for Aboriginal and Torres Strait Islander Australians? A Qualitative Study

Journal:	BMJ Open
Manuscript ID	bmjopen-2024-089436
Article Type:	Original research
Date Submitted by the Author:	30-May-2024
Complete List of Authors:	Woodall, Hannah; Griffith University Griffith Health, ; Rural Medical Education Australia, Evans, Rebecca; James Cook University, College of Medicine and Dentistry and Anton Breinl Research Centre for Health Systems Strengthening McArthur, Lawrie; University of Adelaide Sen Gupta, Tarun; James Cook Univ Ward, Raelene; University of Southern Queensland Brumpton, Kay; Gold Coast Campus, School of Medicine and Dentistry; James Cook University, College of Medicine and Dentistry
Keywords:	Primary Care < Primary Health Care, Telemedicine < BIOTECHNOLOGY & BIOINFORMATICS, Safety

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Can culturally safe general practice telehealth overcome barriers to care for Aboriginal and Torres Strait Islander Australians? A Qualitative Study

Woodall, H*.^{1,2}, Evans, R.³, McArthur, L.⁴, Sen Gupta, T.³, Ward, R.,⁵ Brumpton, K.^{1,2}

¹Griffith University, Gold Coast, Queensland, Australia

²Rural Medical Education Australia, Toowoomba, Queensland, Australia

³James Cook University, Townsville, Queensland, Australia

⁴Division of General Practice, University of Adelaide, South Australia, Australia

⁵University of Southern Queensland, Toowoomba, Queensland, Australia

*Corresponding author: Hannah Woodall, h.woodall@griffith.edu.au, 190 Hume Street, Toowoomba, QLD, Australia

Word count: 2994 words

Keywords: Cultural safety, primary care, telehealth, Aboriginal and Torres Strait Islander

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives: To explore Aboriginal and Torres Strait Islander Australians’ perceptions of telehealth general practice consultations and elements required for a culturally safe telehealth consultation.

Design: Qualitative study

Setting: Primary care telehealth in three centres in regional and remote Australia.

Participants: Seventeen Aboriginal or Torres Strait Islander individuals participated in semi-structured interviews exploring experiences of telehealth in general practice settings. Participants were eligible for inclusion if they were Aboriginal or Torres Strait Islander, over 18 years of age and had experienced at least one telehealth appointment with their general practitioner in the preceding 12 months. Data was collected in the form of a short survey and semi-structured interview. Data was analysed using thematic and content analysis techniques.

Results

Participants had experienced teleconference (88%) and videoconference appointments (12%). Reasons for choosing telehealth included being unable to attend due to respiratory symptoms and/or COVID-19 restrictions on in-person consultations (reflecting the study period) and issues of access (e.g. availability of doctor, convenience of hours). Participants described benefits of telehealth around reduced barriers to care but also described practical and communication challenges experienced during telehealth. Elements of culturally safe telehealth identified included: consultation skills, pre-existing relationship, doctor-patient relationship, cultural and community knowledge.

Conclusion

This study demonstrates the benefits of telehealth and its ability to reduce barriers to care for Aboriginal and Torres Strait Islander Australians. However, the identified disadvantages demonstrate that this modality should be considered an addition to, rather than replacement for, face-to-face consultations. The elements identified interact as part of a complex interplay of factors contributing to cultural safety in the telehealth context. These elements provide useful recommendations for practice and policy.

Strengths and Limitations:

- This study sought Aboriginal and Torres Strait Islander individuals' experiences and perspectives to explore how cultural safety can be enhanced within telehealth.
- In depth data was obtained from participants (n=17) in both regional and remote Australia
- Study participants had predominantly experienced teleconference (rather than videoconference or other telehealth modalities). Cultural safety within teleconference consultations has not been significantly explored in the previous telehealth literature
- Participants were predominantly women (n=15) which may the influence of a female interviewer.

Introduction

Cultural safety is recognised as a vital aspect of medical care, improving access and reducing health inequities. The term “cultural safety” was coined in the 1990s¹ but the elements required for a culturally safe consultation remain difficult to define.² Identifying elements of cultural safety is complicated by the diversity of terms and definitions in use, as well as the intrinsic differences between what individuals, communities and countries may consider to be culturally safe.^{2, 3}

The medical system, and society more broadly, experienced significant shifts from 2020 due to the COVID-19 pandemic. One such change was the widespread introduction of telehealth, including in primary care.⁴ In Australia, telehealth was relatively rare within general practice prior to COVID-19.^{4, 5} Australian healthcare is funded under Medicare; a government-funded universal health insurance scheme that subsidises medical consultations, investigations and procedures.⁶ Prior to COVID-19, funding of telehealth was limited to specialist or Royal Flying Doctor Service consultations and only in restricted situations (e.g., significant geographical distance to services).^{4, 7, 8} As a result, telehealth accounted for only 0.1% of all government-funded consultations.⁵ While the informal use of telehealth was relatively common (e.g. in follow up of results), the lack of funding meant that telehealth was not economically feasible in private general practice based on fee-for-service.⁷ However, expanding telehealth funding to include general practitioners (GPs) in response to the COVID-19 pandemic altered this picture significantly.⁴

Telehealth usage has continued into the post-COVID era. In 2023, 33.5 million telehealth consultations were conducted in Australia, constituting 17% of total Medicare-funded services. Within general practice, telehealth represented 20% of funded services (27.8 million consultations).⁹ Significantly, the majority of telehealth consultations conducted in Australia

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

were conducted by teleconference (87%), with videoconference making up the remaining 13%.⁹

Medical professional bodies have released best-practice telehealth guidelines in response to the increased popularity of this form of healthcare.^{10, 11} However, these guidelines do not explore how cultural safety can be achieved in the telehealth setting.

Practitioner and environmental attributes contributing to cultural safety of telehealth for Indigenous peoples have been explored previously in the literature.¹² Practitioner attributes for culturally safe care included their community and cultural knowledge, building and maintaining of clinician-patient relationships, and communication skills.¹² Environmental factors included technology, the availability of support staff and the telehealth setting (e.g. soundproofing, ensuring privacy and confidentiality).¹² Importantly, few studies have explored the cultural safety of telehealth in primary care, with most studies situated within a specialist mental health context. In addition, most studies explored videoconferencing or store-and-forward consultations (where data is transmitted to a remote clinician who replies with a plan).¹² Thus, current literature has not investigated cultural safety in telehealth relevant to the Australian general practice context.

As this research is situated within the Australian setting, cultural safety will be defined according to the Aboriginal and Torres Strait Islander Health Strategy Group for the Australian Health Practitioner Regulation Agency (AHPRA):

“Cultural safety is determined by Aboriginal and Torres Strait Islander individuals, families, and communities. Culturally safe practise (sic) is the ongoing critical reflection of health practitioner knowledge, skills, attitudes, practising behaviours and power differentials in delivering safe, accessible, and responsive healthcare free of racism.”¹³

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Thus, the “knowledge, skills, attitudes, practising behaviours, and power differentials”¹³ required in the telehealth setting should be determined by Aboriginal and Torres Strait Islander people. The aim of this project is to explore the experiences of Aboriginal and Torres Strait Islander people with telehealth in primary care and understand perspectives on what makes a telehealth consultation culturally safe in this setting.

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Methods

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

This study utilised a constructivist framework¹⁴ to allow exploration of patient preferences and experiences of telehealth with their GPs through a narrative qualitative approach. The concept arose from a concurrent study exploring cultural safety within face-to-face consultations, derived from experiences of staff at a participating Aboriginal Community-Controlled Health Organisation (ACCHO).¹⁵ The onset of COVID-19 prompted the need to explore cultural safety within telehealth.

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Participants were required to be Aboriginal and/or Torres Strait Islander, over 18, and have had at least one experience of telehealth in the preceding 12 months. Inclusion criteria did not set a minimum number of telehealth experiences to avoid excluding participants who may have had a negative experience and thus not proceeded with further telehealth appointments. Participants were recruited within participating ACCHOs, identified through partnerships developed in the research team’s previous work. The study was conducted from June 2022 to August 2023.

49

50

51

52

53

54

55

56

57

58

59

60

Participants completed a short survey including demographic information and questions about identity, followed by a semi-structured interview exploring preferences and experiences of telehealth consultations and cultural safety in this context. The interview also explored the importance of factors identified within cultural safety literature (e.g. use of traditional language, including family in consultations, or the practitioner’s knowledge of

Australian history). The full interview guide can be found in Appendix 1. This was based on the protocol utilised in the study by Brumpton et al. in their study exploring cultural safety in face-to-face general practice consultations.¹⁵ All participants provided informed consent to participate in this study. Member checking of transcripts was offered to all participants.

Demographic data was analysed using descriptive statistics. Interview data was transcribed using Sonix™. Thematic and content analysis was conducted by two independent researchers, facilitated by NVivo™ and informed by Braun and Clark's methodology.¹⁶

HW, TSG and LM are academic GPs with clinical and research experience in Aboriginal and Torres Strait Islander health. KB is a clinician researcher and senior GP within a participating ACCHO. RE is a health services researcher with experience in qualitative and mixed research methods. RW is an Aboriginal health academic from Kunja Nations.

An advisory group of local Aboriginal leaders and community members provided oversight and review of the project and associated processes. This study received ethical approval through the James Cook University Human Research Ethics Committee (H8296).

Results

Seventeen participants were recruited from three ACCHOs within southern Queensland (Table 1). Participating ACCHOs were spread over a large geographical area (750km between the most distant sites). The data was felt to have reached thematic saturation after seventeen participants and thus data collection was ceased.

Reasons for telehealth

Participant reason/s for utilising telehealth consultations are outlined in Figure 1. Some participants used telehealth because of the nature of their presenting complaint. This included being unable to attend due to respiratory symptoms and/or restrictions on in-person

consultations due to COVID-19 public health recommendations. Some participants also indicated they would preferentially select telehealth if they only required prescriptions.

Telehealth was also selected due to convenience (accessing an appointment from work or outside of business hours) or due to doctor availability (e.g. a GP was not available locally, making telehealth the only local option). All patients who reported using telehealth due to GP unavailability were in a remote location, while all participants citing convenience were from a regional location.

Benefits of telehealth

The benefits of telehealth reported by participants centred around the potential for telehealth to improve access to care. This included reducing practical barriers such as transportation and time efficiency.

“The majority of the time I haven't got transport.” (1714)

“It's just a lot easier for that script from your doctor's appointment over the phone to get sent to the chemist. It cuts out a lot of time.” (7966)

Telehealth (specifically teleconference) also allowed people to access care without their physical appearance being visible, which was felt to reduce fear of judgement or concerns about their image.

“I would rather not have them see what I look like sometimes... It avoids them assuming things about my personal life.” (5014)

“What you're wearing, your appearance, all of that thing, all of that feels a bit more relaxed on the phone.” (7966)

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Finally, some participants described feeling that a telehealth consultation was less stressful.

"I've actually found it better talking over the phone, to be honest...Maybe because I'm a bit more relaxed." (6892)

"Because I don't have to come into the building and sit around a lot of people. It's just over the phone, so it makes it a lot easier." (9095)

Disadvantages of telehealth

Participants also described disadvantages of telehealth. Practical constraints of telehealth included the lack of physical examination and potential technological challenges.

"I'd like to get my blood pressure checked. All of that kind of stuff. I think the weight checked all of that, sugar tested. Yeah. So that's the huge disadvantage." (7966)

"I'm not very, computer wise." (8609)

Participants also described communication challenges including the lack of translators and the challenges of communication without non-verbal cues.

"There might not be a translator that can translate directly if the doctor's working from home." (4226)

"If you're not seeing someone's facial expressions...you don't know what they're really thinking." (8840)

Some participants described a lack of control in the telehealth setting. Participants felt a lack of control around being able to choose their preferred GP or feared that their concerns may

not be heard or actioned. In some cases, this was manifested in the practical fear that the promised script would not arrive.

“I suppose with telehealth you just get a random doctor too.” (4226)

“You can't see my expression but can you understand my concern?... I'll just go down there because I probably won't get the outcome I'm expecting (by telehealth).” (3218)

“You're getting that script sent to my chemist. It hasn't arrived. Oh, I shoulda just went to the doctors and got my script in my hand.” (7966)

Finally, participants also described feeling overall that telehealth was different. This related to communication, but also to the challenge of the doctor-patient relationship in this setting. Participants described feeling that the experience of seeing a doctor in person was more “honest” than a telehealth consultation. The difference experienced in a face-to-face consultation was partially attributed to the presence of non-verbal communication. However, some participants did not seem able to fully express why a telehealth consultation felt different, only that it felt less real or “true”.

“There's something about when you're sitting with a doctor, it's a lot more honest and true.” (4226)

“I always feel weird when I'm not sitting in front of someone talking to them....it just feels like you're not really talking to someone.” (5014)

Culturally safe telehealth

Four elements were identified as important for culturally safe telehealth (see Figure 2).

Consultation skills

Consultation skills including communication skills, respect and holistic care were valued by participants. Participants gave specific suggestions about appropriate communication for culturally safe telehealth.

Ask them some indirect questions before you start the direct questions.” (4226)

“Don't speak above us. Speak at our level where we understand.” (8278)

Mutual respect was also valued and expected.

“I treat them with respect and that's how they treat me.” (1714)

“My advice is we're all human. We all get treated the same. It doesn't matter what, just respect is all anyone asks for, not just Aboriginal.” (7599)

Participants also preferred holistic and personalised care, considering the priorities and challenges of the individual person.

“Don't make it just like it's got to be a 15 minute consult. If the consult turns into an hour, it does, because blackfellas like to talk.” (4226)

“Understanding that, okay, wow, she's not feeling this good right now. Let's get her in [arrange for an in-person consultation] and let's get her to see somebody or, you know, a specialist or something like that.” (6892)

Relationship

Participants also highlighted the importance of a pre-existing relationship with the doctor. This relationship contributed to the acceptability and safety of the consultation, with the reverse being true when the doctor was not known to the patient.

“We can't trust him because we don't know him.” (7674)

1
2
3 *“If you've got a doctor that's just knows you just straight off the bat, then you're*
4 *comfortable with that doctor.” (6892)*
5
6

7
8 *Cultural knowledge*
9

10
11 Participants identified the importance of cultural knowledge, including understanding the
12 context and culture of the individual patient. An understanding of culture was felt to be
13 important to the patients’ health and for culturally safe care.
14
15

16
17 *“Understand their culture, where they're coming from, their background. If you have*
18 *to sit and yarn for a little bit then, sit and yarn for a little bit. And take the time to*
19 *listen to them, really listen because there's a lot of them are hurting. And that's all*
20 *they want is that doctor that will listen, take note, understand their culture, be*
21 *culturally aware.” (6892)*
22
23

24 *“People should definitely be briefed on our culture because it's very important when*
25 *it comes to health.” (5014)*
26
27

28
29 *Community knowledge*
30

31
32 Finally, community knowledge was vital, as expressed through the importance of the GP
33 having visited the community to demonstrate commitment to the community and interest in
34 local culture.
35
36

37
38 *“I reckon they should come here... they should sit with the people and even part of the*
39 *community with the Aboriginal people and, and sit with them. And then that's the only*
40 *way they'll know.” (7674)*
41
42

43 *“If they're going to do the telehealth or even in person, they've got to want to, want to*
44 *come and do it and they want to mingle with Indigenous culture.” (6307)*
45
46

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Enseignement Supérieur (ABES)

Discussion

This study explored Aboriginal and Torres Strait Islander people's experiences and preferences around telehealth and the features which contribute to a culturally safe telehealth consultation.

Participant preferences

Participant rurality impacted on reasons for accessing telehealth. Remote participants identified having used telehealth because a GP was not physically available in their town. Conversely, regional participants cited accessibility issues related to convenience of hours (rather than the absence of any doctor). This reflects the general maldistribution of the GP workforce in Australia, i.e. decreasing workforce with increasing rurality.^{17, 18} The most common reason for choosing telehealth was the presence of respiratory symptoms and/or COVID-19 restrictions, in keeping with the timing of this study.

The impact on barriers to care demonstrates telehealth's value in primary care provision. However, the challenges of telehealth demonstrate the need for ongoing availability of in-person appointments for those who do not want telehealth or where telehealth is not appropriate. This is a particular challenge in remote areas, where telehealth may at times be the only option available.

The majority (88%) of participants in this study had experienced teleconference appointments (rather than videoconference or other modalities), consistent with Australian national data.¹⁹ Participants described the lack of video contact as both a potential disadvantage and benefit in this study. These findings are particularly useful in the Australian context where teleconference is common. Experiences may vary in regions where videoconference is the more common telehealth modality.

Culturally safe telehealth

A key aim of this study was to identify features contributing to culturally safe telehealth for Aboriginal and Torres Strait Islander people. Several of the identified features are not unique to the telehealth setting. Many of the consultation skills described (e.g., respect, communication and holistic care) mirror the consultation skills taught more generally as best practice.^{20, 21} Similarly, the importance of the doctor-patient relationship has been described elsewhere, both by patients and healthcare professionals.²²⁻²⁴ The importance of community visits and cultural knowledge was highlighted in a recent literature review exploring cultural safety for Indigenous peoples in telehealth globally.¹²

Further, there is likely to be significant overlap between the four features identified as important for culturally safe telehealth. For instance, community visits potentially improve cultural as well as community knowledge. Similarly, a pre-existing relationship is likely to impact communication and both cultural and community knowledge. Thus, these factors should not be seen as separate items on a checklist, but rather as a complex interplay of factors contributing to cultural safety within telehealth.

In addition, incorporating identified elements of culturally safe telehealth may reduce disadvantages of telehealth. For example, knowledge of the person's computer literacy and context (e.g. technology access) may help to reduce and overcome difficulties. Similarly, time constraints may be mitigated by an understanding of the patient, based on pre-existing relationship. Finally, telehealth with a known GP, with good communication, local and cultural knowledge, may decrease any perceived lack of control in telehealth by assisting the practitioner to understand the patient's priorities and context.

Finally, many of the features described in this study are not unique to the Aboriginal and Torres Strait Islander population. While specific cultural features were identified in this

study, the importance of relationship with a GP, good communication skills, holistic care, and even community visits are likely to be relevant to the wider population.

Limitations

This study was conducted in an Australian context within rural and regional settings. It is possible that urban contexts may exhibit some differences, particularly in the reasons for using telehealth. In addition, the female predominance (15 of 17 participants) should be noted. This may be related to the use of a female interviewer in this study. While the findings are likely to be transferable, future studies may consider purposively recruiting male participants for balance amongst participants. Finally, this study focussed on teleconference appointments, as this was the modality experienced by most participants. Similar studies exploring other modalities of telehealth (such as videoconference or asynchronous telehealth such as store and forward) would be useful.

Recommendations for practice:

1. Telehealth is a valuable tool with potential to reduce barriers to care for Aboriginal and Torres Strait Islander patients.
2. Clinicians should remain aware of potential disadvantages to telehealth, particularly those which can be mitigated, such as community challenges and the patient's potential lack of control.
3. Culturally safe telehealth is a complex interplay of a number of factors:
 - a. Good consultation skills of the practitioner,
 - b. A practitioner with whom the patient has a pre-existing therapeutic relationship, and
 - c. Knowledge of the culture and community with which the patient identifies.

Areas for future development and research

Future study exploring whether these findings translate to the urban setting, and to the wider population would be of value. More study focussed on cultural safety in other forms of telehealth would be valuable (e.g. videoconferencing, asynchronous options).

Implementation of these findings into clinical guidelines and/or telehealth training would be a valuable addition to encourage the cultural safety of telehealth consultations and assist clinicians’ understanding of the benefits and challenges of telehealth for their patients.

Acknowledgements

The authors wish to acknowledge the work of Dr Sumudu Wickramasinghe who assisted in data analysis for this project.

Declaration of Conflicting Interests

The Authors declare that there is no conflict of interest.

Funding

The authors wish to thank the Australian College of Rural and Remote Medicine for funding this research through an Education Research Grant.

Data Availability Statement

The datasets analysed during the current study are not publicly available due to participants’ potential identifiability because of the small dataset. Data is available from the corresponding author on reasonable request.

Author Contributions

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

1
2
3 HW and KB designed the study with advice and input from all authors. HW was responsible
4
5 for data collection, and analysis, and development of the manuscript. KB, RE, LM, TSG and
6
7
8 RW were involved in making sense of the data and reviewing the developed manuscript.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Curtis E, Jones R, Tipene-Leach D, et al. Why cultural safety rather than cultural competency is required to achieve health equity: a literature review and recommended definition. *Int J Equity Health* 2019; 18: 174. DOI: 10.1186/s12939-019-1082-3.

2. Brumpton K, Evans R, Ward R, et al. A consistent definition of cultural safety within Australian health professional education: a scoping review. *AlterNative* 2022; 18: 436-444. DOI: 10.1177/11771801221118950.

3. Australian Institute of Health and Welfare. Cultural Safety in health care for Indigenous Australians: monitoring framework, <https://www.aihw.gov.au/reports/indigenous-australians/cultural-safety-health-care-framework/contents/about> (2023, accessed 19th January 2024).

4. Snoswell CL, Caffery LJ, Haydon HM, et al. Telehealth uptake in general practice as a result of the coronavirus (COVID-19) pandemic. *Aust Health Rev* 2020; 44: 737-740. DOI: 10.1071/AH20183.

5. Pearce C, McLeod A, Supple J, et al. Responding to COVID-19 with real-time general practice data in Australia. *Int J Med Inform* 2022; 157. DOI: 10.1016/j.ijmedinf.2021.104624.

6. Medical Sciences Advisory Committee. What is the MBS and Medicare?, <http://www.msac.gov.au/internet/msac/publishing.nsf/Content/Factsheet-03> (2016, accessed 17th November 2023).

7. Fisher K, Davey A and Magin P. Telehealth for Australian general practice: The present and the future. *Aust J Gen Pract* 2022; 51: 626-629. 2022/08/01. DOI: 10.31128/ajgp-11-21-6229.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

8. Bradford NK, Caffery LJ and Smith AC. Telehealth services in rural and remote Australia: a systematic review of models of care and factors influencing success and sustainability. *Rural Remote Health* 2016; 16: 3808. 2016/10/18.
9. Snoswell C, Caffery, LJ., Taylor, ML, Haydon, HM., Thomas, E., Smith, AC. Centre for Online Health, The University of Queensland. Telehealth and coronavirus: Medicare Benefits Schedule (MBS) activity in Australia., <https://coh.centre.uq.edu.au/telehealth-and-coronavirus-medicare-benefits-schedule-mbs-activity-australia> (2024, accessed 16th May 2024).
10. Australian College of Rural and Remote Medicine. ACCRM Framework and Guidelines for Telehealth Services. Brisbane, QLD: ACCRM, 2020.
11. The Royal Australian College of General Practitioners. Guide to providing telephone and video consultations in general practice. East Melbourne, Victoria: RACGP, 2020.
12. Terrill K, Woodall H, Evans R, et al. Cultural safety in telehealth consultations with Indigenous people: A scoping review of global literature. *J Telemed Telecare* 2023. DOI: 10.1177/1357633X231203874.
13. Australian Health Practitioner Regulation Agency. *The National Scheme's Aboriginal and Torres Strait Islander Health and Cultural Safety Strategy 2020-2025*. 2020. Canberra: AHPRA.
14. Creswell JW and Plano Clark VL. *Designing and conducting mixed methods research*. Third edition. ed. Thousand Oaks, California: SAGE, 2018.
15. Brumpton K, Ward R, Evans R, et al. Assessing cultural safety in general practice consultations for Indigenous patients: protocol for a mixed methods sequential embedded design study. *BMC Med Educ* 2023; 23: 306. DOI: 10.1186/s12909-023-04249-6.
16. Braun V and Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006; 3: 77-101. DOI: 10.1191/1478088706qp063oa.

17. Wilkinson D. Selected demographic, social and work characteristics of the Australian general medical practitioner workforce: Comparing capital cities with regional areas. *Aust J Rural Health* 2000; 8: 327-334. DOI: 10.1046/j.1440-1584.2000.00316.x.

18. Walters LK, McGrail MR, Carson DB, et al. Where to next for rural general practice policy and research in Australia? *Med J Aust* 2017; 207: 56-58. DOI: <https://doi.org/10.5694/mja17.00216>.

19. Snoswell C, Caffery L, Taylor M, et al. Telehealth and coronavirus: Medicare Benefits Schedule (MBS) activity in Australia., <https://coh.centre.uq.edu.au/telehealth-and-coronavirus-medicare-benefits-schedule-mbs-activity-australia> (2023, accessed 12th October 2023).

20. Denness C. What are consultation models for? *InnovAiT* 2013; 6: 592-599. DOI: 10.1177/1755738013475436.

21. Murtagh J. John Murtagh's general practice. 4th ed. ed. North Ryde, N.S.W. :: McGraw-Hill Australia, 2007.

22. Mathew S, Fitts MS, Liddle Z, et al. Telehealth in remote Australia: a supplementary tool or an alternative model of care replacing face-to-face consultations? *BMC Health Serv Res* 2023; 23: 341. DOI: 10.1186/s12913-023-09265-2.

23. Andreadis K, Muellers K, Ancker JS, et al. Telemedicine Impact on the Patient-Provider Relationship in Primary Care During the COVID-19 Pandemic. *Med Care* 2023; 61: S83-s88. 2023/03/10. DOI: 10.1097/mlr.0000000000001808.

24. Ekegren CL, Clark-Ash M, Callaway L, et al. Perspectives of telehealth access and implementation in people recovering from serious transport injury, health care providers and compensation system staff during the COVID-19 pandemic in Australia. *Injury* 2023; 54: 110987. DOI: <https://doi.org/10.1016/j.injury.2023.110987>.

For peer review only

Table 1: Participant Demographics

		Number of participants (n)	Percentage (%)
Gender	Male	2	11.76%
	Female	15	88.24%
	Other / prefer not to say	0	0%
Age	18-24	3	17.65%
	25-34	3	17.65%
	35-44	2	11.76%
	45-54	3	17.65%
	55-64	4	23.53%
	65+	2	11.76%
Rurality	Modified Monash 2 (regional centre)	1	5.88%
	Modified Monash 3 (large rural town)	6	35.29%
	Modified Monash 7 (remote centre)	10	58.82%
Telehealth modality	Teleconference	15	88.24%
	Videoconference	2	11.76%




Figure 1: Reasons for participant selection of telehealth as a modality for consultation.




Figure 2: Elements of Culturally Safe Telehealth

Appendix 1: Semi-structured interview guide

For these questions, we want to learn specifically about your experience with telehealth appointments. These may have been by phone or by video-call.

1. We would like to know, in your opinion, what makes a good doctor?
 Prompt: Can you think a GP that you like? Why do you like them?
2. Still thinking about a good doctor, in your opinion, how would you describe their attitude?
 Prompt: For example, are they patient, positive, funny?
3. How many times have you been seen a GP by telehealth over the past 12 months?
 Prompts: Have they all been with the same GP?
 Have they been with your usual GP?
 What has made you choose telehealth?
 What has made you avoid telehealth?
3. When seeing your GP by telehealth, what are the things you like them to do to make you feel like you are being treated well?
4. I am interested in learning about what it means to be respectful. What should a doctor do to show they respect you in a telehealth visit?
 Prompt: Can you tell me about some examples of ways in which you have been treated with respect by your doctor during a telehealth visit?
 Can you tell me about some examples of ways in which you have been treated disrespectfully by your doctor during a telehealth visit?
5. When a doctor asks you where you are from, where is your country or your mob, what do you think?
 How does this make you feel?
6. As an Aboriginal and/or Torres Strait Islander person, what makes you feel safe in a telehealth consultation?
7. How is culture important to you, particularly, when seeking healthcare from a GP via telehealth?
8. Unfortunately, sometimes people have bad experiences with a doctor. Can you think of a time when you had a telehealth appointment, and this happened to you?
 What was this doctor like? How would you describe their attitude?
9. Sadly, we also know racism and discrimination can occur during GP consultations. Have you ever had an experience during a telehealth appointment when you felt they were being racist or discriminating against you or your family?
 Prompt: Sometimes patients will describe feeling controlled or punished by their GP. Have you had experiences like this?

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

10. I am now going to read you a series of statements. For each statement, I would like you to tell me, on a scale of 1-5, how important these are to you in a telehealth visit and why you have chosen this rating.

How important is:

- Your Aboriginality to you and your identity
- Your Connection to culture
- Your Connection to land
- The consideration of spirituality by your GP
- Including family, Elders, or significant others in your consultations
- The use of silence in consultations
- GPs to use your traditional language (Aboriginal and/or Torres Strait Islander) words during a consultation
- Your knowledge of Australia before and after colonisation (Prompt: before and after white people arrived)
- Your GP's knowledge of Australia pre-colonisation
- Your GP's knowledge of the experiences of Aboriginal and/or Torres Strait Islander people after colonisation
- GP's medical/clinical knowledge and skills
- GP's knowledge of programs designed to improve Aboriginal and Torres Strait Islander health (such as Closing the Gap for subsidised scripts, 715 health assessments)

13. Thank-you for sharing your experiences. We are wanting to improve the way medical students and doctors deliver care to Aboriginal and/or Torres Strait Islander patients.

What tips/tricks/words of wisdom would you like to give to medical students or doctors to:

- improve the way they consult by telehealth with you and other Aboriginal and/or Torres Strait Islander patients
- or
- avoid others having similar experiences to you?

14. Do you have anything else you want to add?




Figure 1: Reasons for participant selection of telehealth as a modality for consultation.




Figure 2: Elements of Culturally Safe Telehealth

BMJ Open

Can culturally safe general practice telehealth overcome barriers to care for Aboriginal and Torres Strait Islander Australians? A Qualitative Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2024-089436.R1
Article Type:	Original research
Date Submitted by the Author:	14-Jan-2025
Complete List of Authors:	Woodall, Hannah; Griffith University Griffith Health, ; Rural Medical Education Australia, Evans, Rebecca; James Cook University, College of Medicine and Dentistry and Anton Breinl Research Centre for Health Systems Strengthening McArthur, Lawrie; University of Adelaide Sen Gupta, Tarun; James Cook Univ Ward, Raelene; University of Southern Queensland Brumpton, Kay; Gold Coast Campus, School of Medicine and Dentistry; James Cook University, College of Medicine and Dentistry
Primary Subject Heading:	General practice / Family practice
Secondary Subject Heading:	Patient-centred medicine
Keywords:	Primary Care < Primary Health Care, Telemedicine < BIOTECHNOLOGY & BIOINFORMATICS, Safety, Australian Aboriginal and Torres Strait Islander Peoples

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Can culturally safe general practice telehealth overcome barriers to care for Aboriginal and Torres Strait Islander Australians? A Qualitative Study

Woodall, H*.^{1,2}, Evans, R.³, McArthur, L.⁴, Sen Gupta, T.³, Ward, R.,⁵ Brumpton, K.^{1,3}

¹Griffith University, Gold Coast, Queensland, Australia

²Rural Medical Education Australia, Toowoomba, Queensland, Australia

³James Cook University, Townsville, Queensland, Australia

⁴Division of General Practice, University of Adelaide, South Australia, Australia

⁵University of Southern Queensland, Toowoomba, Queensland, Australia

*Corresponding author: Hannah Woodall, h.woodall@griffith.edu.au, 190 Hume Street, Toowoomba, QLD, Australia

Word count: 3573 words

Keywords: Cultural safety, primary care, telehealth, Aboriginal and Torres Strait Islander

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives: To explore Aboriginal and Torres Strait Islander Australians’ perceptions of telehealth general practice consultations and elements required for a culturally safe telehealth consultation.

Design: Qualitative study

Setting: Primary care telehealth in three centres in regional and remote Australia.

Participants: Seventeen Aboriginal or Torres Strait Islander individuals participated in semi-structured interviews exploring experiences of telehealth in general practice settings. Participants were eligible for inclusion if they were Aboriginal or Torres Strait Islander, over 18 years of age and had experienced at least one telehealth appointment with their general practitioner in the preceding 12 months. Data was collected in the form of a short survey and semi-structured interview. Data collection occurred between June 2022 and August 2023. Data was analysed using thematic and content analysis techniques.

Results

Participants had experienced telephone (88%) and videoconference appointments (12%). Reasons for choosing telehealth included being unable to attend due to respiratory symptoms and/or COVID-19 restrictions on in-person consultations (reflecting the study period) and issues of access (e.g. availability of doctor, convenience of hours). Participants described benefits of telehealth around reduced barriers to care but also described practical and communication challenges experienced during telehealth. Elements of culturally safe telehealth identified included: consultation skills, a pre-existing doctor-patient relationship and local knowledge (including knowledge of the local cultural and community context).

Conclusion

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

This study demonstrates the benefits of telehealth and its ability to reduce barriers to care for Aboriginal and Torres Strait Islander Australians. However, the identified disadvantages demonstrate that this modality should be considered an addition to, rather than replacement for, face-to-face consultations. The elements identified interact as part of a complex interplay of factors contributing to cultural safety in the telehealth context. These elements provide useful recommendations for practice and policy.

Strengths and Limitations:

- This study sought Aboriginal and Torres Strait Islander individuals' experiences and perspectives to explore how cultural safety can be enhanced within telehealth.
- In depth data was obtained from participants (n=17) in both regional and remote Australia
- Study participants had predominantly experienced telephone consultations (rather than videoconference or other telehealth modalities). Cultural safety within telephone consultations has not been significantly explored in the previous telehealth literature
- Participants were predominantly women (n=15) which may be the influence of a female interviewer.

Introduction

The medical system, and society more broadly, experienced significant shifts from 2020 due to the COVID-19 pandemic. One such change was the widespread introduction of telehealth, including in primary care.¹ In Australia, telehealth was relatively rare within general practice prior to COVID-19.^{1, 2} However, expanding telehealth funding to include general practitioners (GPs) in response to the COVID-19 pandemic altered this picture significantly.¹

Australian healthcare is funded under Medicare; a government-funded universal health insurance scheme that subsidises medical consultations, investigations and procedures.³ Prior to COVID-19, Medicare funding of telehealth was limited to specialist or Royal Flying Doctor Service consultations and only in restricted situations (e.g., significant geographical distance to services).^{1, 4, 5} While the informal use of telehealth was relatively common (e.g. in follow up of results), the lack of funding meant that telehealth was not economically feasible in private general practice based on fee-for-service.⁴ As a result, telehealth accounted for only 0.1% of all government-funded consultations.²

Expansion of funding due to COVID-19 caused a rapid uptake of telehealth in general practice. This increase in telehealth usage continued into the post-COVID era. In 2023, 33.5 million telehealth consultations were conducted in Australia, constituting 17% of total Medicare-funded services. Within general practice, telehealth represented 20% of funded services (27.8 million consultations).⁶ Significantly, the majority of telehealth consultations conducted in Australia were conducted by telephone (87%), with videoconference making up the remaining 13%.⁶

Medical professional bodies have released best-practice telehealth guidelines in response to the increased popularity of this form of healthcare.^{7, 8} However, these guidelines do not explore how cultural safety can be achieved in the telehealth setting.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

What is cultural safety?

Cultural safety is recognised as a vital aspect of medical care, improving healthcare access and reducing health inequities. The health of Aboriginal and Torres Strait Islander Australians continues to be impacted by the ongoing effects of colonisation and experiences of racism and inequality.⁹ Thus, ensuring culturally safe health services is one way to improve access to care and improve health outcomes.

The term “cultural safety” was coined in the 1990s¹⁰ but the elements required for a culturally safe consultation remain difficult to define.¹¹ Identifying elements of cultural safety is complicated by the diversity of terms and definitions in use, as well as the intrinsic differences between what individuals, communities and countries may consider to be culturally safe.^{11, 12}

The term “cultural safety” is one of many which have been used in literature and practice. Other terms in use include cultural sensitivity, cultural competency, cultural respect, cultural humility, cultural security and cultural appropriateness, amongst many others¹³⁻¹⁷ However, the distinctions between these terms are not always clear or consistent.^{13, 18}

For example, the authors of one review proposed that cultural competency was predominantly related to building cultural knowledge and developing an awareness of one’s own background.¹⁹ Cultural safety was seen as higher level of skill which included both practical skills and knowledge, as well as considering patient-defined outcomes of care.¹⁹

Other authors have proposed that cultural awareness, safety and security reflect a sequence of skills which build upon one another.¹⁶ Through this lens, cultural awareness is seen as knowledge-based (e.g. understanding an aspect of culture). Cultural safety is then viewed as the application of this knowledge into practice. Cultural security is considered the highest

level of skill, integrating individual knowledge and actions and incorporating appropriate policies and procedures.¹⁶

By contrast, some authors have identified cultural humility (rather than security) as the highest level of skill. Cultural humility is defined as a transformative process by which a person’s perspective is changed, becoming aware of power differentials and acting with humility at all times.²⁰ These conflicting views on terminology and definitions can create uncertainty and contribute to a lack of clarity in this field.

However, since this research is situated within an Australian context, we will use the definition developed by the Aboriginal and Torres Strait Islander Health Strategy Group for the Australian Health Practitioner Regulation Agency (AHPRA). This definition, included below, was developed in consultation with community and uses the term “cultural safety”. Thus, the term cultural safety was used in this study and is defined as below.

“Cultural safety is determined by Aboriginal and Torres Strait Islander individuals, families, and communities. Culturally safe practise (sic) is the ongoing critical reflection of health practitioner knowledge, skills, attitudes, practising behaviours and power differentials in delivering safe, accessible, and responsive healthcare free of racism.”²¹

This definition recognises firstly the importance of cultural safety being determined by Aboriginal and Torres Strait Islander individuals and communities. It also identifies elements that contribute to cultural safety, e.g. knowledge, skills and attitudes. This definition has been recognised by medical regulators and colleges, including the Australian Medical Council and the Royal Australian College of General Practitioners^{22, 23}

Culturally safe telehealth

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

Practitioner and environmental attributes contributing to culturally safe telehealth for Indigenous peoples have been explored previously.²⁴ Practitioner attributes for culturally safe care included their community and cultural knowledge, building and maintaining of clinician-patient relationships, and communication skills.²⁴ Environmental factors included technology, the availability of support staff and the telehealth setting (e.g. soundproofing, ensuring privacy and confidentiality).²⁴ Importantly, few studies have explored the cultural safety of telehealth in primary care, with most studies situated within a specialist mental health context. In addition, most studies explored videoconferencing or store-and-forward consultations (where data is transmitted to a remote clinician who replies with a plan).²⁴ Thus, current literature has not investigated cultural safety in telehealth relevant to the Australian general practice context.

The aim of this project is to explore the experiences of Aboriginal and Torres Strait Islander people with telehealth in primary care and understand perspectives on what makes a telehealth consultation culturally safe in this setting.

Methods

This study utilised a constructivist framework²⁵ to allow exploration of patient preferences and experiences of telehealth with their GPs through a narrative qualitative approach. The concept arose from a concurrent study exploring cultural safety within face-to-face consultations, derived from experiences of staff at a participating Aboriginal Community-Controlled Health Organisation (ACCHO).²⁶ The onset of COVID-19 prompted the need to explore cultural safety within telehealth.

Participants were required to be Aboriginal and/or Torres Strait Islander, over 18, and have had at least one experience of telehealth in the preceding 12 months. Inclusion criteria did not set a minimum number of telehealth experiences to avoid excluding participants who may

have had a negative experience and thus not proceeded with further telehealth appointments. Participants were recruited within participating ACCHOs, identified through partnerships developed in the research team’s previous work. The study was conducted from June 2022 to August 2023.

Participants completed a short survey including demographic information and questions about identity, followed by a semi-structured interview exploring preferences and experiences of telehealth consultations and cultural safety in this context. The interview also explored the importance of factors identified within cultural safety literature (e.g. use of traditional language, including family in consultations, or the practitioner’s knowledge of Australian history). The full interview guide can be found in Appendix 1. This was based on the protocol utilised in the study by Brumpton et al. exploring cultural safety in face-to-face general practice consultations.²⁶

All participants provided written informed consent to participate in this study. Member checking of transcripts was offered to all participants. Seven participants requested a copy of their transcript, which was sent by email or registered mail depending on participant preference. None of these participants made any changes to their transcript.

The interview was conducted in person for 16 participants and via telephone for one participant. The choice of in-person versus telephone was dependent on participant preference. Interviews were conducted by HW, who is not Aboriginal or Torres Strait Islander. The advisory group determined that HW was an appropriate person to conduct the interview.

Demographic data was analysed using descriptive statistics. When exploring the reason for telehealth consultation/s, any reason provided by the participant was coded. More than one reason could be provided by a participant. Interview data was transcribed using Sonix™.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Thematic and content analysis was conducted by two independent researchers, facilitated by NVivo™ and informed by Braun and Clark's methodology.²⁷

HW, TSG and LM are academic GPs with clinical and research experience in Aboriginal and Torres Strait Islander health. KB is a clinician researcher and senior GP within a participating ACCHO. RE is a health services researcher with experience in qualitative and mixed research methods. RW is an Aboriginal health academic from Kunja Nations.

This study received ethical approval through the James Cook University Human Research Ethics Committee (H8296).

Patient and public involvement statement

This project was developed from another which explored cultural safety in face-to-face general practice consultations. This initial project was developed in partnership with a local ACCHO. The COVID pandemic began during this initial study, causing the question of telehealth to be raised.

Through this study, an advisory group of local leaders and community members provided oversight and review of the project and associated processes. The advisory group was formed within the ACCHO with whom the study was designed. The group included patients, local leaders and health professionals, all of whom identified as Aboriginal and/or Torres Strait Islander.

Results

Seventeen participants were recruited from three ACCHOs within southern Queensland (Table 1). Participating ACCHOs were spread over a large geographical area (750km between the most distant sites). The data was felt to have reached thematic saturation after seventeen participants and thus data collection was ceased.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Reasons for telehealth

Participant reason/s for utilising telehealth consultations are outlined in Figure 1. Some participants used telehealth because of the nature of their presenting complaint. This included being unable to attend due to respiratory symptoms and restrictions on in-person consultations due to COVID-19 public health recommendations. Some participants also indicated they would preferentially select telehealth if they only required prescriptions.

Telehealth was also selected due to convenience (accessing an appointment from work or outside of business hours) or due to doctor availability (e.g. a GP was not available locally, making telehealth the only local option). All patients who reported using telehealth due to GP unavailability were in a remote location, while all participants citing convenience were from a regional location.

Benefits of telehealth

The benefits of telehealth reported by participants centred around the potential for telehealth to improve access to care. This included reducing practical barriers such as transportation and time efficiency.

“The majority of the time I haven't got transport.” (1714)
“It's just a lot easier for that script from your doctor's appointment over the phone to get sent to the chemist. It cuts out a lot of time.” (7966)

Telehealth (specifically telephone consultations) also allowed people to access care without their physical appearance being visible, which was felt to reduce fear of judgement or concerns about their image.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

1
2
3 *"I would rather not have them see what I look like sometimes... It avoids*
4 *them assuming things about my personal life."* (5014)

5
6
7
8 *"What you're wearing, your appearance, all of that thing, all of that feels a bit*
9 *more relaxed on the phone."* (7966)

10
11
12
13
14
15 Finally, some participants described feeling that a telehealth consultation was less stressful.

16
17
18 *"I've actually found it better talking over the phone, to be honest...Maybe because I'm*
19 *a bit more relaxed."* (6892)

20
21
22
23 *"Because I don't have to come into the building and sit around a lot of people. It's*
24 *just over the phone, so it makes it a lot easier."* (9095)

25 26 27 28 29 30 ***Disadvantages of telehealth***

31
32
33 Participants also described disadvantages of telehealth. Practical constraints of telehealth
34 included the lack of physical examination and potential technological challenges.

35
36
37
38
39 *"I'd like to get my blood pressure checked. All of that kind of stuff. I think the weight*
40 *checked all of that, sugar tested. Yeah. So that's the huge disadvantage."* (7966)

41
42
43 *"I'm not very, computer wise."* (8609)

44
45
46
47
48 Participants also described communication challenges including the lack of translators and
49 the challenges of communication without non-verbal cues.

50
51
52
53
54 *"There might not be a translator that can translate directly if the doctor's working*
55 *from home."* (4226)

1
2
3 *“If you're not seeing someone's facial expressions...you don't know what they're*
4 *really thinking.” (8840)*
5
6
7
8
9

10
11 Some participants described a lack of control in the telehealth setting. Participants felt a lack
12 of control around being able to choose their preferred GP or feared that their concerns may
13 not be heard or actioned. In some cases, this was manifested in the practical fear that the
14 promised script would not arrive.
15
16
17
18

19
20 *“I suppose with telehealth you just get a random doctor too.” (4226)*
21
22 *“You can't see my expression but can you understand my concern?... I'll just go down*
23 *there because I probably won't get the outcome I'm expecting (by telehealth).” (3218)*
24
25
26
27 *“You're getting that script sent to my chemist. It hasn't arrived. Oh, I shoulda just*
28 *went to the doctors and got my script in my hand.” (7966)*
29
30
31

32 Finally, participants also described feeling overall that telehealth was different. This related
33 to communication, but also to the challenge of the doctor-patient relationship in this setting.
34
35 Participants described feeling that the experience of seeing a doctor in person was more
36
37 “honest” than a telehealth consultation. The difference experienced in a face-to-face
38
39 consultation was partially attributed to the presence of non-verbal communication. However,
40
41 some participants did not seem able to fully express why a telehealth consultation felt
42
43 different, only that it felt less real or “true”.
44
45
46
47
48

49 *“There's something about when you're sitting with a doctor, it's a lot more honest*
50 *and true.” (4226)*
51
52
53 *“I always feel weird when I'm not sitting in front of someone talking to them....it just*
54 *feels like you're not really talking to someone.” (5014)*
55
56
57
58
59
60

Culturally safe telehealth

Four elements were identified as important for culturally safe telehealth (see Figure 2).

Consultation skills

Consultation skills including communication skills, respect and holistic care were valued by participants. Participants gave specific suggestions about appropriate communication for culturally safe telehealth.

Ask them some indirect questions before you start the direct questions.” (4226)

“Don't speak above us. Speak at our level where we understand.” (8278)

Mutual respect was also valued and expected.

“I treat them with respect and that's how they treat me.” (1714)

“My advice is we're all human. We all get treated the same. It doesn't matter what, just respect is all anyone asks for, not just Aboriginal.” (7599)

Participants also preferred holistic and personalised care, considering the priorities and challenges of the individual person.

“Don't make it just like it's got to be a 15 minute consult. If the consult turns into an hour, it does, because blackfellas like to talk.” (4226)

“Understanding that, okay, wow, she's not feeling this good right now. Let's get her in [arrange for an in-person consultation] and let's get her to see somebody or, you know, a specialist or something like that.” (6892)

Relationship

Participants also highlighted the importance of a pre-existing relationship with the doctor. This relationship contributed to the acceptability and safety of the consultation, with the reverse being true when the doctor was not known to the patient.

“We can't trust him because we don't know him.” (7674)

“If you've got a doctor that's just knows you just straight off the bat, then you're comfortable with that doctor.” (6892)

Cultural knowledge

Participants identified the importance of cultural knowledge, including understanding the context and culture of the individual patient. An understanding of culture was felt to be important to the patients’ health and for culturally safe care.

“Understand their culture, where they're coming from, their background. If you have to sit and yarn for a little bit then, sit and yarn for a little bit. And take the time to listen to them, really listen because there's a lot of them are hurting. And that's all they want is that doctor that will listen, take note, understand their culture, be culturally aware.” (6892)

“People should definitely be briefed on our culture because it's very important when it comes to health.” (5014)

Community knowledge

Finally, community knowledge was vital, as expressed through the importance of the GP having visited the community to demonstrate commitment to the community and interest in local culture.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

1
2
3 *"I reckon they should come here... they should sit with the people and even part of the*
4 *community with the Aboriginal people and, and sit with them. And then that's the only*
5 *way they'll know."* (7674)
6
7

8
9
10
11 *"If they're going to do the telehealth or even in person, they've got to want to, want to*
12 *come and do it and they want to mingle with Indigenous culture."* (6307)
13
14

15 16 **Discussion**

17
18
19 This study explored Aboriginal and Torres Strait Islander people's experiences and
20 preferences around telehealth and the features which contribute to a culturally safe telehealth
21 consultation.
22
23

24 25 26 ***Participant preferences***

27
28
29 Participant rurality impacted on reasons for accessing telehealth. Remote participants
30 identified having used telehealth because a GP was not physically available in their town.
31
32 Conversely, regional participants cited accessibility issues related to convenience of hours
33 (rather than the absence of any doctor). This reflects the general maldistribution of the GP
34 workforce in Australia, i.e. decreasing workforce with increasing rurality.^{28, 29} The most
35 common reason for choosing telehealth was the presence of respiratory symptoms and/or
36 COVID-19 restrictions, in keeping with the timing of this study.
37
38

39
40
41 The impact on barriers to care demonstrates telehealth's value in primary care provision.
42
43 However, the challenges of telehealth demonstrate the need for ongoing availability of in-
44 person appointments for those who do not want telehealth or where telehealth is not
45 appropriate. This is a particular challenge in remote areas, where telehealth may at times be
46 the only option available.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The majority (88%) of participants in this study had experienced telephone appointments (rather than videoconference or other modalities), consistent with Australian national data.³⁰ Participants described the lack of video contact as both a potential disadvantage and benefit in this study. These findings are particularly useful in the Australian context where telephone consultations are common. Experiences may vary in regions where videoconference is the more common telehealth modality.

Culturally safe telehealth

A key aim of this study was to identify features contributing to culturally safe telehealth for Aboriginal and Torres Strait Islander people. Several of the identified features are not unique to the telehealth setting. Many of the consultation skills described (e.g., respect, communication and holistic care) mirror the consultation skills taught more generally as best practice.^{31, 32} Similarly, the importance of the doctor-patient relationship has been described elsewhere, both by patients and healthcare professionals.³³⁻³⁵ The importance of community visits and cultural knowledge was highlighted in a recent literature review exploring cultural safety for Indigenous peoples in telehealth globally.²⁴

Further, there is likely to be significant overlap between the four features identified as important for culturally safe telehealth. For instance, community visits potentially improve cultural as well as community knowledge. Similarly, a pre-existing relationship is likely to impact communication and both cultural and community knowledge. Thus, these factors should not be seen as separate items on a checklist, but rather as a complex interplay of factors contributing to cultural safety within telehealth.

In addition, incorporating identified elements of culturally safe telehealth may reduce disadvantages of telehealth. For example, knowledge of the person’s computer literacy and context (e.g. technology access) may help to reduce and overcome difficulties. Similarly,

time constraints may be mitigated by an understanding of the patient, based on pre-existing relationship. Finally, telehealth with a known GP, with good communication, local and cultural knowledge, may decrease any perceived lack of control in telehealth by assisting the practitioner to understand the patient's priorities and context.

Finally, many of the features described in this study are not unique to the Aboriginal and Torres Strait Islander population. While specific cultural features were identified in this study, the importance of relationship with a GP, good communication skills, holistic care, and even community visits are likely to be relevant to the wider population.

Limitations

This study was conducted in an Australian context within rural and regional settings. It is possible that urban contexts may exhibit some differences, particularly in the reasons for using telehealth. In addition, the female predominance (15 of 17 participants) should be noted. This may be related to the use of a female interviewer in this study. While the findings are likely to be transferable, future studies may consider purposively recruiting male participants for balance amongst participants. Finally, this study focussed on telephone appointments, as this was the modality experienced by most participants. Similar studies exploring other modalities of telehealth (such as videoconference or asynchronous telehealth such as store and forward) would be useful.

Recommendations for practice:

1. Rural Aboriginal and Torres Strait Islander patients in this study found telehealth to be a vital healthcare resource.

Recommendations for Doctors: Doctors should recognise the role of telehealth in healthcare and attempt to offer it where desired / possible

Recommendations for Health Systems: Medicare should expand funding to meet the growing needs of historically marginalized communities that may otherwise be unable to access healthcare.

2. Although telehealth may increase opportunities for access to care, this study identified disadvantages that must be considered.

Recommendations for Doctors: Doctors should familiarise themselves with the disadvantages of telehealth such as limitations in access to support (e.g. translators) and the perceived lack of control over the encounter, and work with health systems to mitigate these disadvantages. Doctors should also recognise that some patients will not prefer telehealth and provide alternatives where possible.

Recommendations for Health Systems: Institutions which offer telehealth should develop robust systems to provide education and support to patients to mitigate disadvantages and ensure that non-telehealth appointments remain available.

3. Culturally safe telehealth is a complex interplay of factors including consultation skills, pre-existing therapeutic relationship/s, and local knowledge of culture and community

Recommendations for Doctors: Doctors should be aware of the factors which make a telehealth consultation culturally safe and strive to put them into place within their daily practice. This may include trying to deliver holistic, person-centred care, communicating clearly without jargon, intentionally developing the doctor-patient relationship and/or seeking to learn about the local community and cultural context.

Recommendations for Health Systems: Health systems should strive to engage GPs who have the necessary consultation skills to provide culturally safe telehealth. Health systems should work to reduce GP turnover in primary care to enhance the development of therapeutic relationships which enhance cultural safe in telehealth.

Conclusion

Our study identified that Aboriginal and Torres Strait Islander people perceived culturally safe telehealth to include appropriate communication and consultation skills, the building of relationship between doctor and patient, and local knowledge (including both cultural and community knowledge).

Future study exploring whether these findings translate to the urban setting, and to the wider population would be of value. More study focussed on cultural safety in other forms of telehealth would be valuable (e.g. videoconferencing, asynchronous options).

Implementation of these findings into clinical guidelines and/or telehealth training would be a valuable addition to encourage the cultural safety of telehealth consultations and assist clinicians' understanding of the benefits and challenges of telehealth for their patients.

Acknowledgements

The authors wish to acknowledge the work of Dr Sumudu Wickramasinghe who assisted in data analysis for this project.

Declaration of Conflicting Interests

The Authors declare that there is no conflict of interest.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Funding

The authors wish to thank the Australian College of Rural and Remote Medicine for funding this research through an Education Research Grant.

Data Availability Statement

The datasets analysed during the current study are not publicly available due to participants’ potential identifiability because of the small dataset. Data is available from the corresponding author on reasonable request.

Author Contributions

HW and KB designed the study with advice and input from all authors. HW was responsible for data collection, and analysis, and development of the manuscript. KB, RE, LM, TSG and RW were involved in interpreting data and reviewing the developed manuscript. HW is the guarantor.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

References

1. Snoswell CL, Caffery LJ, Haydon HM, et al. Telehealth uptake in general practice as a result of the coronavirus (COVID-19) pandemic. *Aust Health Rev* 2020; 44: 737-740. DOI: 10.1071/AH20183.
2. Pearce C, McLeod A, Supple J, et al. Responding to COVID-19 with real-time general practice data in Australia. *Int J Med Inform* 2022; 157. DOI: 10.1016/j.ijmedinf.2021.104624.
3. Medical Sciences Advisory Committee. What is the MBS and Medicare?, <http://www.msac.gov.au/internet/msac/publishing.nsf/Content/Factsheet-03> (2016, accessed 17th November 2023).
4. Fisher K, Davey A and Magin P. Telehealth for Australian general practice: The present and the future. *Aust J Gen Pract* 2022; 51: 626-629. 2022/08/01. DOI: 10.31128/ajgp-11-21-6229.
5. Bradford NK, Caffery LJ and Smith AC. Telehealth services in rural and remote Australia: a systematic review of models of care and factors influencing success and sustainability. *Rural Remote Health* 2016; 16: 3808. 2016/10/18.
6. Snoswell C, Caffery, LJ., Taylor, ML, Haydon, HM., Thomas, E., Smith, AC. Centre for Online Health, The University of Queensland. Telehealth and coronavirus: Medicare Benefits Schedule (MBS) activity in Australia., <https://coh.centre.uq.edu.au/telehealth-and-coronavirus-medicare-benefits-schedule-mbs-activity-australia> (2024, accessed 16th May 2024).
7. Australian College of Rural and Remote Medicine. ACCRM Framework and Guidelines for Telehealth Services. Brisbane, QLD: ACCRM, 2020.
8. The Royal Australian College of General Practitioners. Guide to providing telephone and video consultations in general practice. East Melbourne, Victoria: RACGP, 2020.

9. Productivity Commisison. *Closing the Gap Annual Data Compilation Report July 2024*. 2024. Canberra.

10. Curtis E, Jones R, Tipene-Leach D, et al. Why cultural safety rather than cultural competency is required to achieve health equity: a literature review and recommended definition. *Int J Equity Health* 2019; 18: 174. DOI: 10.1186/s12939-019-1082-3.

11. Brumpton K, Evans R, Ward R, et al. A consistent definition of cultural safety within Australian health professional education: a scoping review. *AlterNative* 2022; 18: 436-444. DOI: 10.1177/11771801221118950.

12. Australian Institute of Health and Welfare. Cultural Safety in health care for Indigenous Australians: monitoring framework, <https://www.aihw.gov.au/reports/indigenous-australians/cultural-safety-health-care-framework/contents/about> (2023, accessed 19th January 2024).

13. Curtis E, Jones R, Tipene-Leach D, et al. Why cultural safety rather than cultural competency is required to achieve health equity: a literature review and recommended definition. *International Journal for Equity in Health* 2019; 18: 174. DOI: 10.1186/s12939-019-1082-3.

14. Liaw ST and Wade V. Cultural respect in general practice: a cluster randomised controlled trial. *Med J Aust* 2019; 211: 43-43.e41. Letter. DOI: 10.5694/mja2.50214.

15. Chang ES, Simon M and Dong X. Integrating cultural humility into health care professional education and training. *Advances in Health Sciences Education* 2012; 17: 269-278. Article. DOI: 10.1007/s10459-010-9264-1.

16. Coffin J. Rising to the Challenge in Aboriginal Health by Creating Cultural Security. *Aboriginal and Islander Health Worker Journal* 2007; 31: 22-24. Journal Article.

17. Amanda R, Rana K, Saunders P, et al. Evaluation of the usability, content, readability and cultural appropriateness of online alcohol and other drugs resources for Aboriginal and

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

Torres Strait Islander Peoples in New South Wales, Australia. *BMJ Open* 2023; 13. Article.

DOI: 10.1136/bmjopen-2022-069756.

18. Brumpton K, Evans R, Ward R, et al. A consistent definition of cultural safety within Australian health professional education: a scoping review. *AlterNative: An International Journal of Indigenous Peoples* 2022; 18: 436-444. DOI: 10.1177/11771801221118950.

19. Evans N, Meñaca A, Koffman J, et al. Cultural Competence in End-of-Life Care: Terms, Definitions, and Conceptual Models from the British Literature. *Journal of Palliative Medicine* 2012; 15: 812-820. DOI: 10.1089/jpm.2011.0526.

20. Foronda C, Baptiste D-L, Reinholdt MM, et al. Cultural Humility: A Concept Analysis. *Journal of Transcultural Nursing* 2016; 27: 210-217. DOI: 10.1177/1043659615592677.

21. Australian Health Practitioner Regulation Agency. *The National Scheme's Aboriginal and Torres Strait Islander Health and Cultural Safety Strategy 2020-2025*. 2020. Canberra: AHPRA.

22. Australian Medical Council Limited. Standards for Assessment and Accreditation of Primary Medical Programs. 2023.

23. Royal Australian College of General Practitioners. Cultural Safety, <https://www.racgp.org.au/cultural-and-health-training-framework/framework/cultural-safety/> (2024, accessed 19th December 2024).

24. Terrill K, Woodall H, Evans R, et al. Cultural safety in telehealth consultations with Indigenous people: A scoping review of global literature. *J Telemed Telecare* 2023. DOI: 10.1177/1357633X231203874.

25. Creswell JW and Plano Clark VL. *Designing and conducting mixed methods research*. Third edition. ed. Thousand Oaks, California: SAGE, 2018.

26. Brumpton K, Ward R, Evans R, et al. Assessing cultural safety in general practice consultations for Indigenous patients: protocol for a mixed methods sequential embedded design study. *BMC Med Educ* 2023; 23: 306. DOI: 10.1186/s12909-023-04249-6.

27. Braun V and Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006; 3: 77-101. DOI: 10.1191/1478088706qp063oa.

28. Wilkinson D. Selected demographic, social and work characteristics of the Australian general medical practitioner workforce: Comparing capital cities with regional areas. *Aust J Rural Health* 2000; 8: 327-334. DOI: 10.1046/j.1440-1584.2000.00316.x.

29. Walters LK, McGrail MR, Carson DB, et al. Where to next for rural general practice policy and research in Australia? *Med J Aust* 2017; 207: 56-58. DOI: <https://doi.org/10.5694/mja17.00216>.

30. Snoswell C, Caffery L, Taylor M, et al. Telehealth and coronavirus: Medicare Benefits Schedule (MBS) activity in Australia., <https://coh.centre.uq.edu.au/telehealth-and-coronavirus-medicare-benefits-schedule-mbs-activity-australia> (2023, accessed 12th October 2023).

31. Denness C. What are consultation models for? *InnovAiT* 2013; 6: 592-599. DOI: 10.1177/1755738013475436.

32. Murtagh J. John Murtagh's general practice. 4th ed. ed. North Ryde, N.S.W. :: McGraw-Hill Australia, 2007.

33. Mathew S, Fitts MS, Liddle Z, et al. Telehealth in remote Australia: a supplementary tool or an alternative model of care replacing face-to-face consultations? *BMC Health Serv Res* 2023; 23: 341. DOI: 10.1186/s12913-023-09265-2.

34. Andreadis K, Muellers K, Ancker JS, et al. Telemedicine Impact on the Patient-Provider Relationship in Primary Care During the COVID-19 Pandemic. *Med Care* 2023; 61: S83-s88. 2023/03/10. DOI: 10.1097/mlr.0000000000001808.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

- 1
2
3 35. Ekegren CL, Clark-Ash M, Callaway L, et al. Perspectives of telehealth access and
4 implementation in people recovering from serious transport injury, health care providers and
5 compensation system staff during the COVID-19 pandemic in Australia. *Injury* 2023; 54:
6
7
8 110987. DOI: <https://doi.org/10.1016/j.injury.2023.110987>.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1: Participant Demographics

		Number of participants (n)	Percentage (%)
Gender	Male	2	12%
	Female	15	88%
	Other / prefer not to say	0	0%
Age	18-24	3	18%
	25-34	3	18%
	35-44	2	12%
	45-54	3	18%
	55-64	4	22%
	65+	2	12%
Rurality	Modified Monash 2 (regional centre)	1	6%
	Modified Monash 3 (large rural town)	6	35%
	Modified Monash 7 (remote centre)	10	59%
Telehealth modality	Teleconference	15	88%
	Videoconference	2	12%

Figure 1: Reasons for participant selection of telehealth as a modality for consultation.

Figure 2: Elements of Culturally Safe Telehealth




Figure 1: Reasons for participant selection of telehealth as a modality for consultation.

501x237mm (130 x 130 DPI)




Figure 2: Elements of culturally safe telehealth

398x224mm (38 x 38 DPI)

Appendix 1: Semi-structured interview guide

For these questions, we want to learn specifically about your experience with telehealth appointments. These may have been by phone or by video-call.

1. We would like to know, in your opinion, what makes a good doctor?

Prompt: Can you think a GP that you like? Why do you like them?

2. Still thinking about a good doctor, in your opinion, how would you describe their attitude?

Prompt: For example, are they patient, positive, funny?

3. How many times have you been seen a GP by telehealth over the past 12 months?

Prompts: Have they all been with the same GP?

Have they been with your usual GP?

What has made you choose telehealth?

What has made you avoid telehealth?

3. When seeing your GP by telehealth, what are the things you like them to do to make you feel like you are being treated well?

4. I am interested in learning about what it means to be respectful. What should a doctor do to show they respect you in a telehealth visit?

Prompt: Can you tell me about some examples of ways in which you have been treated with respect by your doctor during a telehealth visit?

Can you tell me about some examples of ways in which you have been treated disrespectfully by your doctor during a telehealth visit?

5. When a doctor asks you where you are from, where is your country or your mob, what do you think?

How does this make you feel?

6. As an Aboriginal and/or Torres Strait Islander person, what makes you feel safe in a telehealth consultation?

7. How is culture important to you, particularly, when seeking healthcare from a GP via telehealth?

8. Unfortunately, sometimes people have bad experiences with a doctor. Can you think of a time when you had a telehealth appointment, and this happened to you?

What was this doctor like? How would you describe their attitude?

9. Sadly, we also know racism and discrimination can occur during GP consultations. Have you ever had an experience during a telehealth appointment when you felt they were being racist or discriminating against you or your family?

Prompt: Sometimes patients will describe feeling controlled or punished by their GP.

Have you had experiences like this?

10.I am now going to read you a series of statements. For each statement, I would like you to tell me, on a scale of 1-5, how important these are to you in a telehealth visit and why you have chosen this rating.

How important is:

- Your Aboriginality to you and your identity
- Your Connection to culture
- Your Connection to land
- The consideration of spirituality by your GP
- Including family, Elders, or significant others in your consultations
- The use of silence in consultations
- GPs to use your traditional language (Aboriginal and/or Torres Strait Islander) words during a consultation
- Your knowledge of Australia before and after colonisation (Prompt: before and after white people arrived)
- Your GP’s knowledge of Australia pre-colonisation
- Your GP’s knowledge of the experiences of Aboriginal and/or Torres Strait Islander people after colonisation
- GP’s medical/clinical knowledge and skills
- GP’s knowledge of programs designed to improve Aboriginal and Torres Strait Islander health(such as Closing the Gap for subsidised scripts, 715 health assessments)

13. Thank-you for sharing your experiences. We are wanting to improve the way medical students and doctors deliver care to Aboriginal and/or Torres Strait Islander patients.

What tips/tricks/words of wisdom would you like to give to medical students or doctors to:

- improve the way they consult by telehealth with you and other Aboriginal and/or Torres Strait Islander patients
- or
- avoid others having similar experiences to you?

14.Do you have anything else you want to add?

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).

BMJ Open

Can culturally safe general practice telehealth overcome barriers to care for Aboriginal and Torres Strait Islander Australians? A Qualitative Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2024-089436.R2
Article Type:	Original research
Date Submitted by the Author:	27-Feb-2025
Complete List of Authors:	Woodall, Hannah; Griffith University Griffith Health, ; Rural Medical Education Australia, Evans, Rebecca; James Cook University, College of Medicine and Dentistry and Anton Breinl Research Centre for Health Systems Strengthening McArthur, Lawrie; University of Adelaide Sen Gupta, Tarun; James Cook Univ Ward, Raelene; University of Southern Queensland Brumpton, Kay; Gold Coast Campus, School of Medicine and Dentistry; James Cook University, College of Medicine and Dentistry
Primary Subject Heading:	General practice / Family practice
Secondary Subject Heading:	Patient-centred medicine
Keywords:	Primary Care < Primary Health Care, Telemedicine < BIOTECHNOLOGY & BIOINFORMATICS, Safety, Australian Aboriginal and Torres Strait Islander Peoples

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

Can culturally safe general practice telehealth overcome barriers to care for Aboriginal and Torres Strait Islander Australians? A Qualitative Study

Woodall, H*.^{1,2}, Evans, R.³, McArthur, L.⁴, Sen Gupta, T.³, Ward, R.,⁵ Brumpton, K.^{1,3}

¹Griffith University, Gold Coast, Queensland, Australia

²Rural Medical Education Australia, Toowoomba, Queensland, Australia

³James Cook University, Townsville, Queensland, Australia

⁴Division of General Practice, University of Adelaide, South Australia, Australia

⁵University of Southern Queensland, Toowoomba, Queensland, Australia

*Corresponding author: Hannah Woodall, h.woodall@griffith.edu.au, 190 Hume Street, Toowoomba, QLD, Australia

Word count: 3573 words

Keywords: Cultural safety, primary care, telehealth, Aboriginal and Torres Strait Islander

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives: To explore Aboriginal and Torres Strait Islander Australians’ perceptions of telehealth general practice consultations and elements required for a culturally safe telehealth consultation.

Design: Qualitative study

Setting: Primary care telehealth in three centres in regional and remote Australia.

Participants: Seventeen Aboriginal or Torres Strait Islander individuals participated in semi-structured interviews exploring experiences of telehealth in general practice settings.

Participants were eligible for inclusion if they were Aboriginal or Torres Strait Islander, over 18 years of age and had experienced at least one telehealth appointment with their general practitioner in the preceding 12 months. Data was collected in the form of a short survey and semi-structured interview. Data collection occurred between June 2022 and August 2023. Data was analysed using thematic and content analysis techniques.

Results

Participants had experienced telephone (88%) and videoconference appointments (12%). Reasons for choosing telehealth included being unable to attend due to respiratory symptoms and/or COVID-19 restrictions on in-person consultations (reflecting the study period) and issues of access (e.g. availability of doctor, convenience of hours). Participants described benefits of telehealth around reduced barriers to care but also described practical and communication challenges experienced during telehealth. Elements of culturally safe telehealth identified included: consultation skills, a pre-existing doctor-patient relationship and local knowledge (including knowledge of the local cultural and community context).

Conclusion

This study demonstrates the benefits of telehealth and its ability to reduce barriers to care for Aboriginal and Torres Strait Islander Australians. However, the identified disadvantages demonstrate that this modality should be considered an addition to, rather than replacement for, face-to-face consultations. The elements identified interact as part of a complex interplay of factors contributing to cultural safety in the telehealth context. These elements provide useful recommendations for practice and policy.

Strengths and Limitations:

- This study sought Aboriginal and Torres Strait Islander individuals' experiences and perspectives to explore how cultural safety can be enhanced within telehealth.
- In depth data was obtained from participants (n=17) in both regional and remote Australia
- Study participants had predominantly experienced telephone consultations (rather than videoconference or other telehealth modalities). Cultural safety within telephone consultations has not been significantly explored in the previous telehealth literature
- Participants were predominantly women (n=15) which may be the influence of a female interviewer.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Introduction

2 The medical system, and society more broadly, experienced significant shifts from 2020 due
3 to the COVID-19 pandemic. One such change was the widespread introduction of telehealth,
4 including in primary care.¹ In Australia, telehealth was relatively rare within general practice
5 prior to COVID-19.^{1, 2} However, expanding telehealth funding to include general
6 practitioners (GPs) in response to the COVID-19 pandemic altered this picture significantly.¹
7 Australian healthcare is funded under Medicare; a government-funded universal health
8 insurance scheme that subsidises medical consultations, investigations and procedures.³ Prior
9 to COVID-19, Medicare funding of telehealth was limited to specialist or Royal Flying
10 Doctor Service consultations and only in restricted situations (e.g., significant geographical
11 distance to services).^{1, 4, 5} While the informal use of telehealth was relatively common (e.g. in
12 follow up of results), the lack of funding meant that telehealth was not economically feasible
13 in private general practice based on fee-for-service.⁴ As a result, telehealth accounted for only
14 0.1% of all government-funded consultations.²
15 Expansion of funding due to COVID-19 caused a rapid uptake of telehealth in general
16 practice. This increase in telehealth usage continued into the post-COVID era. In 2023, 33.5
17 million telehealth consultations were conducted in Australia, constituting 17% of total
18 Medicare-funded services. Within general practice, telehealth represented 20% of funded
19 services (27.8 million consultations).⁶ Significantly, the majority of telehealth consultations
20 conducted in Australia were conducted by telephone (87%), with videoconference making up
21 the remaining 13%.⁶
22 Medical professional bodies have released best-practice telehealth guidelines in response to
23 the increased popularity of this form of healthcare.^{7, 8} However, these guidelines do not
24 explore how cultural safety can be achieved in the telehealth setting.

1

2 ***What is cultural safety?***

3 Cultural safety is recognised as a vital aspect of medical care, improving healthcare access
4 and reducing health inequities. The health of Aboriginal and Torres Strait Islander
5 Australians continues to be impacted by the ongoing effects of colonisation and experiences
6 of racism and inequality.⁹ Thus, ensuring culturally safe health services is one way to
7 improve access to care and improve health outcomes.

8 The term “cultural safety” was coined in the 1990s¹⁰ but the elements required for a culturally
9 safe consultation remain difficult to define.¹¹ Identifying elements of cultural safety is
10 complicated by the diversity of terms and definitions in use, as well as the intrinsic
11 differences between what individuals, communities and countries may consider to be
12 culturally safe.^{11, 12}

13 The term “cultural safety” is one of many which have been used in literature and practice.
14 Other terms in use include cultural sensitivity, cultural competency, cultural respect, cultural
15 humility, cultural security and cultural appropriateness, amongst many others¹³⁻¹⁷ However,
16 the distinctions between these terms are not always clear or consistent.^{13, 18}

17 For example, the authors of one review proposed that cultural competency was
18 predominantly related to building cultural knowledge and developing an awareness of one’s
19 own background.¹⁹ Cultural safety was seen as higher level of skill which included both
20 practical skills and knowledge, as well as considering patient-defined outcomes of care.¹⁹

21 Other authors have proposed that cultural awareness, safety and security reflect a sequence of
22 skills which build upon one another.¹⁶ Through this lens, cultural awareness is seen as
23 knowledge-based (e.g. understanding an aspect of culture). Cultural safety is then viewed as
24 the application of this knowledge into practice. Cultural security is considered the highest

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 level of skill, integrating individual knowledge and actions and incorporating appropriate
2 policies and procedures.¹⁶

3 By contrast, some authors have identified cultural humility (rather than security) as the
4 highest level of skill. Cultural humility is defined as a transformative process by which a
5 person’s perspective is changed, becoming aware of power differentials and acting with
6 humility at all times.²⁰ These conflicting views on terminology and definitions can create
7 uncertainty and contribute to a lack of clarity in this field.

8 However, since this research is situated within an Australian context, we will use the
9 definition developed by the Aboriginal and Torres Strait Islander Health Strategy Group for
10 the Australian Health Practitioner Regulation Agency (AHPRA). This definition, included
11 below, was developed in consultation with community and uses the term “cultural safety”.
12 Thus, the term cultural safety was used in this study and is defined as below.

13 *“Cultural safety is determined by Aboriginal and Torres Strait Islander individuals,*
14 *families, and communities. Culturally safe practise (sic) is the ongoing critical*
15 *reflection of health practitioner knowledge, skills, attitudes, practising behaviours*
16 *and power differentials in delivering safe, accessible, and responsive healthcare free*
17 *of racism.”²¹*

18 This definition recognises firstly the importance of cultural safety being determined by
19 Aboriginal and Torres Strait Islander individuals and communities. It also identifies elements
20 that contribute to cultural safety, e.g. knowledge, skills and attitudes. This definition has
21 been recognised by medical regulators and colleges, including the Australian Medical
22 Council and the Royal Australian College of General Practitioners^{22, 23}

23 ***Culturally safe telehealth***

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Ensignment Supérieur (ABES) .

Practitioner and environmental attributes contributing to culturally safe telehealth for Indigenous peoples have been explored previously.²⁴ Practitioner attributes for culturally safe care included their community and cultural knowledge, building and maintaining of clinician-patient relationships, and communication skills.²⁴ Environmental factors included technology, the availability of support staff and the telehealth setting (e.g. soundproofing, ensuring privacy and confidentiality).²⁴ Importantly, few studies have explored the cultural safety of telehealth in primary care, with most studies situated within a specialist mental health context. In addition, most studies explored videoconferencing or store-and-forward consultations (where data is transmitted to a remote clinician who replies with a plan).²⁴ Thus, current literature has not investigated cultural safety in telehealth relevant to the Australian general practice context.

The aim of this project is to explore the experiences of Aboriginal and Torres Strait Islander people with telehealth in primary care and understand perspectives on what makes a telehealth consultation culturally safe in this setting.

Methods

This study utilised a constructivist framework²⁵ to allow exploration of patient preferences and experiences of telehealth with their GPs through a narrative qualitative approach. The concept arose from a concurrent study exploring cultural safety within face-to-face consultations, derived from experiences of staff at a participating Aboriginal Community-Controlled Health Organisation (ACCHO).²⁶ The onset of COVID-19 prompted the need to explore cultural safety within telehealth.

Participants were required to be Aboriginal and/or Torres Strait Islander, over 18, and have had at least one experience of telehealth in the preceding 12 months. Inclusion criteria did not set a minimum number of telehealth experiences to avoid excluding participants who may

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 have had a negative experience and thus not proceeded with further telehealth appointments.

2 Participants were recruited within participating ACCHOs, identified through partnerships

3 developed in the research team’s previous work. The study was conducted from June 2022 to

4 August 2023.

5 Participants completed a short survey including demographic information and questions

6 about identity, followed by a semi-structured interview exploring preferences and

7 experiences of telehealth consultations and cultural safety in this context. The interview also

8 explored the importance of factors identified within cultural safety literature (e.g. use of

9 traditional language, including family in consultations, or the practitioner’s knowledge of

10 Australian history). The full interview guide can be found in Appendix 1. This was based on

11 the protocol utilised in the study by Brumpton et al. exploring cultural safety in face-to-face

12 general practice consultations.²⁶

13 All participants provided written informed consent to participate in this study. Member

14 checking of transcripts was offered to all participants. Seven participants requested a copy of

15 their transcript, which was sent by email or registered mail depending on participant

16 preference. None of these participants made any changes to their transcript.

17 The interview was conducted in person for 16 participants and via telephone for one

18 participant. The choice of in-person versus telephone was dependent on participant

19 preference. Interviews were conducted by HW, who is not Aboriginal or Torres Strait

20 Islander. The advisory group determined that HW was an appropriate person to conduct the

21 interview.

22 Demographic data was analysed using descriptive statistics. When exploring the reason for

23 telehealth consultation/s, any reason provided by the participant was coded. More than one

24 reason could be provided by a participant. Interview data was transcribed using Sonix™.

Thematic and content analysis was conducted by two independent researchers, facilitated by NVivo™ and informed by Braun and Clark's methodology.²⁷

HW, TSG and LM are academic GPs with clinical and research experience in Aboriginal and Torres Strait Islander health. KB is a clinician researcher and senior GP within a participating ACCHO. RE is a health services researcher with experience in qualitative and mixed research methods. RW is an Aboriginal health academic from Kunja Nations.

This study received ethical approval through the James Cook University Human Research Ethics Committee (H8296).

Patient and public involvement statement

This project was developed from another which explored cultural safety in face-to-face general practice consultations. This initial project was developed in partnership with a local ACCHO. The COVID pandemic began during this initial study, causing the question of telehealth to be raised.

Through this study, an advisory group of local leaders and community members provided oversight and review of the project and associated processes. The advisory group was formed within the ACCHO with whom the study was designed. The group included patients, local leaders and health professionals, all of whom identified as Aboriginal and/or Torres Strait Islander.

Results

Seventeen participants were recruited from three ACCHOs within southern Queensland (Table 1). Participating ACCHOs were spread over a large geographical area (750km between the most distant sites). The data was felt to have reached thematic saturation after seventeen participants and thus data collection was ceased.

Reasons for telehealth

Participant reason/s for utilising telehealth consultations are outlined in Figure 1. Some participants used telehealth because of the nature of their presenting complaint. This included being unable to attend due to respiratory symptoms and restrictions on in-person consultations due to COVID-19 public health recommendations. Some participants also indicated they would preferentially select telehealth if they only required prescriptions. Telehealth was also selected due to convenience (accessing an appointment from work or outside of business hours) or due to doctor availability (e.g. a GP was not available locally, making telehealth the only local option). All patients who reported using telehealth due to GP unavailability were in a remote location, while all participants citing convenience were from a regional location.

Benefits of telehealth

The benefits of telehealth reported by participants centred around the potential for telehealth to improve access to care. This included reducing practical barriers such as transportation and time efficiency.

“The majority of the time I haven't got transport.” (1714)
“It's just a lot easier for that script from your doctor's appointment over the phone to get sent to the chemist. It cuts out a lot of time.” (7966)

Telehealth (specifically telephone consultations) also allowed people to access care without their physical appearance being visible, which was felt to reduce fear of judgement or concerns about their image.

1
2
3 1 *"I would rather not have them see what I look like sometimes... It avoids*
4
5 2 *them assuming things about my personal life."* (5014)
6

7
8 3 *"What you're wearing, your appearance, all of that thing, all of that feels a bit*
9
10 4 *more relaxed on the phone."* (7966)
11
12 5

13
14
15 6 Finally, some participants described feeling that a telehealth consultation was less stressful.
16

17
18 7 *"I've actually found it better talking over the phone, to be honest...Maybe because I'm*
19
20 8 *a bit more relaxed."* (6892)
21

22
23 9 *"Because I don't have to come into the building and sit around a lot of people. It's*
24
25 10 *just over the phone, so it makes it a lot easier."* (9095)
26
27 11

28 29 30 12 ***Disadvantages of telehealth***

31
32
33 13 Participants also described disadvantages of telehealth. Practical constraints of telehealth
34
35 14 included the lack of physical examination and potential technological challenges.
36

37
38
39 15 *"I'd like to get my blood pressure checked. All of that kind of stuff. I think the weight*
40
41 16 *checked all of that, sugar tested. Yeah. So that's the huge disadvantage."* (7966)
42

43 17 *"I'm not very, computer wise."* (8609)
44
45 18

46
47
48 19 Participants also described communication challenges including the lack of translators and
49
50 20 the challenges of communication without non-verbal cues.
51

52
53
54 21 *"There might not be a translator that can translate directly if the doctor's working*
55
56 22 *from home."* (4226)
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 *“If you're not seeing someone's facial expressions...you don't know what they're*
2 *really thinking.” (8840)*

4 Some participants described a lack of control in the telehealth setting. Participants felt a lack
5 of control around being able to choose their preferred GP or feared that their concerns may
6 not be heard or actioned. In some cases, this was manifested in the practical fear that the
7 promised script would not arrive.

8 *“I suppose with telehealth you just get a random doctor too.” (4226)*
9 *“You can't see my expression but can you understand my concern?... I'll just go down*
10 *there because I probably won't get the outcome I'm expecting (by telehealth).” (3218)*
11 *“You’re getting that script sent to my chemist. It hasn’t arrived. Oh, I shoulda just*
12 *went to the doctors and got my script in my hand.” (7966)*

13 Finally, participants also described feeling overall that telehealth was different. This related
14 to communication, but also to the challenge of the doctor-patient relationship in this setting.
15 Participants described feeling that the experience of seeing a doctor in person was more
16 “honest” than a telehealth consultation. The difference experienced in a face-to-face
17 consultation was partially attributed to the presence of non-verbal communication. However,
18 some participants did not seem able to fully express why a telehealth consultation felt
19 different, only that it felt less real or “true”.

20 *“There’s something about when you’re sitting with a doctor, it’s a lot more honest*
21 *and true.” (4226)*
22 *“I always feel weird when I'm not sitting in front of someone talking to them....it just*
23 *feels like you're not really talking to someone.” (5014)*

1 **Culturally safe telehealth**

2 Four elements were identified as important for culturally safe telehealth (see Figure 2).

3 *Consultation skills*

4 Consultation skills including communication skills, respect and holistic care were valued by
5 participants. Participants gave specific suggestions about appropriate communication for
6 culturally safe telehealth.

7 *Ask them some indirect questions before you start the direct questions.” (4226)*

8 *“Don't speak above us. Speak at our level where we understand.” (8278)*

9 Mutual respect was also valued and expected.

10 *“I treat them with respect and that's how they treat me.” (1714)*

11 *“My advice is we're all human. We all get treated the same. It doesn't matter what,
12 just respect is all anyone asks for, not just Aboriginal.” (7599)*

13 Participants also preferred holistic and personalised care, considering the priorities and
14 challenges of the individual person.

15 *“Don't make it just like it's got to be a 15 minute consult. If the consult turns into an
16 hour, it does, because blackfellas like to talk.” (4226)*

17 *“Understanding that, okay, wow, she's not feeling this good right now. Let's get her in
18 [arrange for an in-person consultation] and let's get her to see somebody or, you
19 know, a specialist or something like that.” (6892)*

20 *Relationship*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Participants also highlighted the importance of a pre-existing relationship with the doctor.
2 This relationship contributed to the acceptability and safety of the consultation, with the
3 reverse being true when the doctor was not known to the patient.

4 *“We can't trust him because we don't know him.” (7674)*

5 *“If you've got a doctor that's just knows you just straight off the bat, then you're*
6 *comfortable with that doctor.” (6892)*

7 *Cultural knowledge*

8 Participants identified the importance of cultural knowledge, including understanding the
9 context and culture of the individual patient. An understanding of culture was felt to be
10 important to the patients' health and for culturally safe care.

11 *“Understand their culture, where they're coming from, their background. If you have*
12 *to sit and yarn for a little bit then, sit and yarn for a little bit. And take the time to*
13 *listen to them, really listen because there's a lot of them are hurting. And that's all*
14 *they want is that doctor that will listen, take note, understand their culture, be*
15 *culturally aware.” (6892)*

16 *“People should definitely be briefed on our culture because it's very important when*
17 *it comes to health.” (5014)*

18
19 *Community knowledge*

20 Finally, community knowledge was vital, as expressed through the importance of the GP
21 having visited the community to demonstrate commitment to the community and interest in
22 local culture.

1
2
3 1 *"I reckon they should come here... they should sit with the people and even part of the*
4
5 2 *community with the Aboriginal people and, and sit with them. And then that's the only*
6
7 3 *way they'll know."* (7674)
8
9

10
11 4 *"If they're going to do the telehealth or even in person, they've got to want to, want to*
12
13 5 *come and do it and they want to mingle with Indigenous culture."* (6307)
14
15

16 **Discussion**

17
18
19 7 This study explored Aboriginal and Torres Strait Islander people's experiences and
20
21 8 preferences around telehealth and the features which contribute to a culturally safe telehealth
22
23 9 consultation.
24
25

26 ***Participant preferences***

27
28
29 11 Participant rurality impacted on reasons for accessing telehealth. Remote participants
30
31 12 identified having used telehealth because a GP was not physically available in their town.
32
33 13 Conversely, regional participants cited accessibility issues related to convenience of hours
34
35 14 (rather than the absence of any doctor). This reflects the general maldistribution of the GP
36
37 15 workforce in Australia, i.e. decreasing workforce with increasing rurality.^{28, 29} The most
38
39 16 common reason for choosing telehealth was the presence of respiratory symptoms and/or
40
41 17 COVID-19 restrictions, in keeping with the timing of this study.
42
43
44
45

46 18 The impact on barriers to care demonstrates telehealth's value in primary care provision.
47
48 19 However, the challenges of telehealth demonstrate the need for ongoing availability of in-
49
50 20 person appointments for those who do not want telehealth or where telehealth is not
51
52 21 appropriate. This is a particular challenge in remote areas, where telehealth may at times be
53
54 22 the only option available.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The majority (88%) of participants in this study had experienced telephone appointments
2 (rather than videoconference or other modalities), consistent with Australian national data.³⁰
3 Participants described the lack of video contact as both a potential disadvantage and benefit
4 in this study. These findings are particularly useful in the Australian context where telephone
5 consultations are common. Experiences may vary in regions where videoconference is the
6 more common telehealth modality.

7 ***Culturally safe telehealth***

8 A key aim of this study was to identify features contributing to culturally safe telehealth for
9 Aboriginal and Torres Strait Islander people. Several of the identified features are not unique
10 to the telehealth setting. Many of the consultation skills described (e.g., respect,
11 communication and holistic care) mirror the consultation skills taught more generally as best
12 practice.^{31, 32} Similarly, the importance of the doctor-patient relationship has been described
13 elsewhere, both by patients and healthcare professionals.³³⁻³⁵ The importance of community
14 visits and cultural knowledge was highlighted in a recent literature review exploring cultural
15 safety for Indigenous peoples in telehealth globally.²⁴

16 Further, there is likely to be significant overlap between the four features identified as
17 important for culturally safe telehealth. For instance, community visits potentially improve
18 cultural as well as community knowledge. Similarly, a pre-existing relationship is likely to
19 impact communication and both cultural and community knowledge. Thus, these factors
20 should not be seen as separate items on a checklist, but rather as a complex interplay of
21 factors contributing to cultural safety within telehealth.

22 In addition, incorporating identified elements of culturally safe telehealth may reduce
23 disadvantages of telehealth. For example, knowledge of the person’s computer literacy and
24 context (e.g. technology access) may help to reduce and overcome difficulties. Similarly,

time constraints may be mitigated by an understanding of the patient, based on pre-existing relationship. Finally, telehealth with a known GP, with good communication, local and cultural knowledge, may decrease any perceived lack of control in telehealth by assisting the practitioner to understand the patient's priorities and context.

Finally, many of the features described in this study are not unique to the Aboriginal and Torres Strait Islander population. While specific cultural features were identified in this study, the importance of relationship with a GP, good communication skills, holistic care, and even community visits are likely to be relevant to the wider population.

Limitations

This study was conducted in an Australian context within rural and regional settings. It is possible that urban contexts may exhibit some differences, particularly in the reasons for using telehealth. In addition, the female predominance (15 of 17 participants) should be noted. This may be related to the use of a female interviewer in this study. While the findings are likely to be transferable, future studies may consider purposively recruiting male participants for balance amongst participants. Data was not collected on whether families or caregivers were present in consultations. In addition, data was not collected on multiple consultations and thus reasons for choosing telehealth were coded only once per patient. Finally, this study focussed on telephone appointments, as this was the modality experienced by most participants. This focus on telephone interactions may have impacted participants' attitudes to remote care modalities. Similar studies exploring other modalities of telehealth (such as videoconference or asynchronous telehealth such as store and forward) would be useful.

Recommendations for practice:

1. Rural Aboriginal and Torres Strait Islander patients in this study found telehealth to be a vital healthcare resource.

Recommendations for Doctors: Doctors should recognise the role of telehealth in healthcare and attempt to offer it where desired / possible

Recommendations for Health Systems: Medicare should expand funding to meet the growing needs of historically marginalized communities that may otherwise be unable to access healthcare.

2. Although telehealth may increase opportunities for access to care, this study identified disadvantages that must be considered.

Recommendations for Doctors: Doctors should familiarise themselves with the disadvantages of telehealth such as limitations in access to support (e.g. interpreters) and the perceived lack of control over the encounter, and work with health systems to mitigate these disadvantages. Doctors should also recognise that some patients will not prefer telehealth and provide alternatives where possible.

Recommendations for Health Systems: Institutions which offer telehealth should develop robust systems to provide education and support to clinicians to mitigate disadvantages and ensure that non-telehealth appointments remain available.

3. Culturally safe telehealth is a complex interplay of factors including consultation skills, pre-existing therapeutic relationship/s, and local knowledge of culture and community

Recommendations for Doctors: Doctors should be aware of the factors which make a telehealth consultation culturally safe and strive to put them into place

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignement Supérieur (ABES).

1 within their daily practice. This may include trying to deliver holistic, person-
2 centred care, communicating clearly without jargon, intentionally developing
3 the doctor-patient relationship and/or seeking to learn about the local
4 community and cultural context.

5 ***Recommendations for Health Systems:*** Health systems should strive to
6 engage GPs who have the necessary consultation skills to provide culturally
7 safe telehealth. Health systems should work to reduce GP turnover in primary
8 care to enhance the development of therapeutic relationships which promote
9 culturally safe telehealth. Ongoing clinician education should be provided to
10 foster development of skills which can improve cultural safety, such as
11 telehealth-specific communication training or education regarding cultural and
12 community knowledge.

13 ***Conclusion***

14 Our study identified that Aboriginal and Torres Strait Islander people perceived culturally
15 safe telehealth to include appropriate communication and consultation skills, the building of
16 relationship between doctor and patient, and local knowledge (including both cultural and
17 community knowledge).

18 Future studies exploring whether these findings translate to the urban setting, and to the wider
19 population would be of value. More study focussed on cultural safety in other forms of
20 telehealth would be valuable (e.g. videoconferencing, asynchronous options).

21 Implementation of these findings into clinical guidelines and/or telehealth training would be a
22 valuable addition to encourage the cultural safety of telehealth consultations and assist
23 clinicians' understanding of the benefits and challenges of telehealth for their patients.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

The authors wish to acknowledge the work of Dr Sumudu Wickramasinghe who assisted in data analysis for this project.

Declaration of Conflicting Interests

The Authors declare that there is no conflict of interest.

Funding

The authors wish to thank the Australian College of Rural and Remote Medicine for funding this research through an Education Research Grant.

Data Availability Statement

The datasets analysed during the current study are not publicly available due to participants' potential identifiability because of the small dataset. Data is available from the corresponding author on reasonable request.

Author Contributions

HW and KB designed the study with advice and input from all authors. HW was responsible for data collection, and analysis, and development of the manuscript. KB, RE, LM, TSG and RW were involved in interpreting data and reviewing the developed manuscript. HW is the guarantor.

References

1. Snoswell CL, Caffery LJ, Haydon HM, et al. Telehealth uptake in general practice as a result of the coronavirus (COVID-19) pandemic. *Aust Health Rev* 2020; 44: 737-740. DOI: 10.1071/AH20183.
2. Pearce C, McLeod A, Supple J, et al. Responding to COVID-19 with real-time general practice data in Australia. *Int J Med Inform* 2022; 157. DOI: 10.1016/j.ijmedinf.2021.104624.
3. Medical Sciences Advisory Committee. What is the MBS and Medicare?, <http://www.msac.gov.au/internet/msac/publishing.nsf/Content/Factsheet-03> (2016, accessed 17th November 2023).
4. Fisher K, Davey A and Magin P. Telehealth for Australian general practice: The present and the future. *Aust J Gen Pract* 2022; 51: 626-629. 2022/08/01. DOI: 10.31128/ajgp-11-21-6229.
5. Bradford NK, Caffery LJ and Smith AC. Telehealth services in rural and remote Australia: a systematic review of models of care and factors influencing success and sustainability. *Rural Remote Health* 2016; 16: 3808. 2016/10/18.
6. Snoswell C, Caffery, LJ., Taylor, ML, Haydon, HM., Thomas, E., Smith, AC. Centre for Online Health, The University of Queensland. Telehealth and coronavirus: Medicare Benefits Schedule (MBS) activity in Australia., <https://coh.centre.uq.edu.au/telehealth-and-coronavirus-medicare-benefits-schedule-mbs-activity-australia> (2024, accessed 16th May 2024).
7. Australian College of Rural and Remote Medicine. ACCRM Framework and Guidelines for Telehealth Services. Brisbane, QLD: ACCRM, 2020.
8. The Royal Australian College of General Practitioners. Guide to providing telephone and video consultations in general practice. East Melbourne, Victoria: RACGP, 2020.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

9. Productivity Commisison. *Closing the Gap Annual Data Compilation Report July 2024*. 2024. Canberra.

10. Curtis E, Jones R, Tipene-Leach D, et al. Why cultural safety rather than cultural competency is required to achieve health equity: a literature review and recommended definition. *Int J Equity Health* 2019; 18: 174. DOI: 10.1186/s12939-019-1082-3.

11. Brumpton K, Evans R, Ward R, et al. A consistent definition of cultural safety within Australian health professional education: a scoping review. *AlterNative* 2022; 18: 436-444. DOI: 10.1177/11771801221118950.

12. Australian Institute of Health and Welfare. Cultural Safety in health care for Indigenous Australians: monitoring framework, <https://www.aihw.gov.au/reports/indigenous-australians/cultural-safety-health-care-framework/contents/about> (2023, accessed 19th January 2024).

13. Curtis E, Jones R, Tipene-Leach D, et al. Why cultural safety rather than cultural competency is required to achieve health equity: a literature review and recommended definition. *International Journal for Equity in Health* 2019; 18: 174. DOI: 10.1186/s12939-019-1082-3.

14. Liaw ST and Wade V. Cultural respect in general practice: a cluster randomised controlled trial. *Med J Aust* 2019; 211: 43-43.e41. Letter. DOI: 10.5694/mja2.50214.

15. Chang ES, Simon M and Dong X. Integrating cultural humility into health care professional education and training. *Advances in Health Sciences Education* 2012; 17: 269-278. Article. DOI: 10.1007/s10459-010-9264-1.

16. Coffin J. Rising to the Challenge in Aboriginal Health by Creating Cultural Security. *Aboriginal and Islander Health Worker Journal* 2007; 31: 22-24. Journal Article.

17. Amanda R, Rana K, Saunders P, et al. Evaluation of the usability, content, readability and cultural appropriateness of online alcohol and other drugs resources for Aboriginal and

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Enseignement Supérieur (ABES).

- 1
2
3 1 Torres Strait Islander Peoples in New South Wales, Australia. *BMJ Open* 2023; 13. Article.
4
5 2 DOI: 10.1136/bmjopen-2022-069756.
6
7
8 3 18. Brumpton K, Evans R, Ward R, et al. A consistent definition of cultural safety within
9
10 4 Australian health professional education: a scoping review. *AlterNative: An International*
11
12 5 *Journal of Indigenous Peoples* 2022; 18: 436-444. DOI: 10.1177/11771801221118950.
13
14
15 6 19. Evans N, Meñaca A, Koffman J, et al. Cultural Competence in End-of-Life Care:
16
17 7 Terms, Definitions, and Conceptual Models from the British Literature. *Journal of Palliative*
18
19 8 *Medicine* 2012; 15: 812-820. DOI: 10.1089/jpm.2011.0526.
20
21
22 9 20. Foronda C, Baptiste D-L, Reinholdt MM, et al. Cultural Humility: A Concept
23
24 10 Analysis. *Journal of Transcultural Nursing* 2016; 27: 210-217. DOI:
25
26 11 10.1177/1043659615592677.
27
28
29 12 21. Australian Health Practitioner Regulation Agency. *The National Scheme's Aboriginal*
30
31 13 *and Torres Strait Islander Health and Cultural Safety Strategy 2020-2025*. 2020. Canberra:
32
33 14 AHPRA.
34
35
36 15 22. Australian Medical Council Limited. Standards for Assessment and Accreditation of
37
38 16 Primary Medical Programs. 2023.
39
40
41 17 23. Royal Australian College of General Practitioners. Cultural Safety,
42
43 18 <https://www.racgp.org.au/cultural-and-health-training-framework/framework/cultural-safety/>
44
45 19 (2024, accessed 19th December 2024).
46
47
48 20 24. Terrill K, Woodall H, Evans R, et al. Cultural safety in telehealth consultations with
49
50 21 Indigenous people: A scoping review of global literature. *J Telemed Telecare* 2023. DOI:
51
52 22 10.1177/1357633X231203874.
53
54
55 23 25. Creswell JW and Plano Clark VL. *Designing and conducting mixed methods*
56
57 24 *research*. Third edition. ed. Thousand Oaks, California: SAGE, 2018.
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

26. Brumpton K, Ward R, Evans R, et al. Assessing cultural safety in general practice consultations for Indigenous patients: protocol for a mixed methods sequential embedded design study. *BMC Med Educ* 2023; 23: 306. DOI: 10.1186/s12909-023-04249-6.

27. Braun V and Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006; 3: 77-101. DOI: 10.1191/1478088706qp063oa.

28. Wilkinson D. Selected demographic, social and work characteristics of the Australian general medical practitioner workforce: Comparing capital cities with regional areas. *Aust J Rural Health* 2000; 8: 327-334. DOI: 10.1046/j.1440-1584.2000.00316.x.

29. Walters LK, McGrail MR, Carson DB, et al. Where to next for rural general practice policy and research in Australia? *Med J Aust* 2017; 207: 56-58. DOI: <https://doi.org/10.5694/mja17.00216>.

30. Snoswell C, Caffery L, Taylor M, et al. Telehealth and coronavirus: Medicare Benefits Schedule (MBS) activity in Australia., <https://coh.centre.uq.edu.au/telehealth-and-coronavirus-medicare-benefits-schedule-mbs-activity-australia> (2023, accessed 12th October 2023).

31. Denness C. What are consultation models for? *InnovAiT* 2013; 6: 592-599. DOI: 10.1177/1755738013475436.

32. Murtagh J. John Murtagh's general practice. 4th ed. ed. North Ryde, N.S.W. :: McGraw-Hill Australia, 2007.

33. Mathew S, Fitts MS, Liddle Z, et al. Telehealth in remote Australia: a supplementary tool or an alternative model of care replacing face-to-face consultations? *BMC Health Serv Res* 2023; 23: 341. DOI: 10.1186/s12913-023-09265-2.

34. Andreadis K, Muellers K, Ancker JS, et al. Telemedicine Impact on the Patient-Provider Relationship in Primary Care During the COVID-19 Pandemic. *Med Care* 2023; 61: S83-s88. 2023/03/10. DOI: 10.1097/mlr.0000000000001808.

- 1 35. Ekegren CL, Clark-Ash M, Callaway L, et al. Perspectives of telehealth access and
2 implementation in people recovering from serious transport injury, health care providers and
3 compensation system staff during the COVID-19 pandemic in Australia. *Injury* 2023; 54:
4 110987. DOI: <https://doi.org/10.1016/j.injury.2023.110987>.

For peer review only

1 *Table 1: Participant Demographics*

		Number of participants (n)	Percentage (%)
Gender	Male	2	12%
	Female	15	88%
	Other / prefer not to say	0	0%
Age	18-24	3	18%
	25-34	3	18%
	35-44	2	12%
	45-54	3	18%
	55-64	4	22%
	65+	2	12%
Rurality	Modified Monash 2 (regional centre)	1	6%
	Modified Monash 3 (large rural town)	6	35%
	Modified Monash 7 (remote centre)	10	59%
Telehealth modality	Teleconference	15	88%
	Videoconference	2	12%

Figure 1: Reasons for participant selection of telehealth as a modality for consultation.

Figure 2: Elements of Culturally Safe Telehealth




Figure 1: Reasons for participant selection of telehealth as a modality for consultation.

501x237mm (130 x 130 DPI)




Figure 2: Elements of culturally safe telehealth

398x224mm (38 x 38 DPI)

Appendix 1: Semi-structured interview guide

For these questions, we want to learn specifically about your experience with telehealth appointments. These may have been by phone or by video-call.

1. We would like to know, in your opinion, what makes a good doctor?

Prompt: Can you think a GP that you like? Why do you like them?

2. Still thinking about a good doctor, in your opinion, how would you describe their attitude?

Prompt: For example, are they patient, positive, funny?

3. How many times have you been seen a GP by telehealth over the past 12 months?

Prompts: Have they all been with the same GP?

Have they been with your usual GP?

What has made you choose telehealth?

What has made you avoid telehealth?

3. When seeing your GP by telehealth, what are the things you like them to do to make you feel like you are being treated well?

4. I am interested in learning about what it means to be respectful. What should a doctor do to show they respect you in a telehealth visit?

Prompt: Can you tell me about some examples of ways in which you have been treated with respect by your doctor during a telehealth visit?

Can you tell me about some examples of ways in which you have been treated disrespectfully by your doctor during a telehealth visit?

5. When a doctor asks you where you are from, where is your country or your mob, what do you think?

How does this make you feel?

6. As an Aboriginal and/or Torres Strait Islander person, what makes you feel safe in a telehealth consultation?

7. How is culture important to you, particularly, when seeking healthcare from a GP via telehealth?

8. Unfortunately, sometimes people have bad experiences with a doctor. Can you think of a time when you had a telehealth appointment, and this happened to you?

What was this doctor like? How would you describe their attitude?

9. Sadly, we also know racism and discrimination can occur during GP consultations. Have you ever had an experience during a telehealth appointment when you felt they were being racist or discriminating against you or your family?

Prompt: Sometimes patients will describe feeling controlled or punished by their GP.

Have you had experiences like this?

10.I am now going to read you a series of statements. For each statement, I would like you to tell me, on a scale of 1-5, how important these are to you in a telehealth visit and why you have chosen this rating.

How important is:

- Your Aboriginality to you and your identity
- Your Connection to culture
- Your Connection to land
- The consideration of spirituality by your GP
- Including family, Elders, or significant others in your consultations
- The use of silence in consultations
- GPs to use your traditional language (Aboriginal and/or Torres Strait Islander) words during a consultation
- Your knowledge of Australia before and after colonisation (Prompt: before and after white people arrived)
- Your GP's knowledge of Australia pre-colonisation
- Your GP's knowledge of the experiences of Aboriginal and/or Torres Strait Islander people after colonisation
- GP's medical/clinical knowledge and skills
- GP's knowledge of programs designed to improve Aboriginal and Torres Strait Islander health(such as Closing the Gap for subsidised scripts, 715 health assessments)

13. Thank-you for sharing your experiences. We are wanting to improve the way medical students and doctors deliver care to Aboriginal and/or Torres Strait Islander patients. What tips/tricks/words of wisdom would you like to give to medical students or doctors to:

- improve the way they consult by telehealth with you and other Aboriginal and/or Torres Strait Islander patients
- or
- avoid others having similar experiences to you?

14.Do you have anything else you want to add?

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Ensignment Supérieur (ABES).