

BMJ Open Parental involvement in infection prevention and control in low- and middle-income country neonatal units: a scoping review protocol

Lydia Davidson ^{1,2}, Chikomborero Kitikiti,³ Hannah Blencowe ¹,
Felicity Fitzgerald ⁴, Sarah Moxon ¹, Gwendoline Chimhini,⁵ Rudo Chingono²

To cite: Davidson L, Kitikiti C, Blencowe H, *et al*. Parental involvement in infection prevention and control in low- and middle-income country neonatal units: a scoping review protocol. *BMJ Open* 2025;**15**:e093967. doi:10.1136/bmjopen-2024-093967

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-093967>).

Received 24 September 2024
Accepted 28 March 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY. Published by BMJ Group.

¹MARCH Centre, London School of Hygiene & Tropical Medicine, London, UK

²The Health Research Institute, Biomedical Research and Training Institute, Harare, Zimbabwe

³Division of Paediatrics, Harare Central Hospital, Harare, Zimbabwe

⁴Department of Infectious Diseases, Imperial College London, London, UK

⁵Department of Child Adolescent and Women's Health, University of Zimbabwe Faculty of Medicine and Health Sciences, Harare, Zimbabwe

Correspondence to

Lydia Davidson;
Lydia.Davidson@LSHTM.ac.uk

ABSTRACT

Introduction Neonatal sepsis is a key contributor to neonatal mortality worldwide, and low- and middle-income countries (LMIC) are disproportionately affected. With antimicrobial resistance challenging effective treatment of neonatal sepsis, it is increasingly urgent to improve infection prevention and control (IPC) in LMIC neonatal units (NNU) and reduce transmission of infections. One pathway to improvement which merits further exploration is the collaboration with families to build an IPC intervention.

Families are constantly present on neonatal units, and much of the hands-on care for their newborns is given by them. For IPC to be effective, families must adhere to IPC standards within the NNU, but furthermore, any IPC intervention implemented must be feasible and acceptable for families as well as the hospital staff as this will increase uptake and effectiveness of the intervention. This scoping review aims to provide an overview of parental involvement in infection prevention and control in low- and middle-income setting neonatal units.

Methods and analysis This protocol was developed in line with the Joanna Briggs Institute recommendations. Searches will be carried out on six databases (Medline, CINAHL, Global Health, EMBASE, Web of Science and Global Index Medicus), and reference searching will be carried out on included studies. The search will be carried out from 2000 to present (end date 28/02/2024), and included languages will be English, French, Spanish and Portuguese. Screening and data extraction will be performed independently by two reviewers, with a third reviewer to resolve conflicts. Results will be reported by narrative synthesis of each sub-question in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews guidelines.

Ethics and dissemination This study will be carried out using already published data exclusively and therefore does not require further ethical approval. Results will be disseminated through peer-reviewed publications and conference presentations and through engagement with peers and relevant stakeholders.

Trial registration number Registered with Open Science Framework - https://osf.io/snc7a/?view_only=8ffc39d837594b4388c7394a838c3a9e

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Despite the constant presence of parents within neonatal units, there is limited information on parental involvement in infection prevention and control (IPC) in low- and middle-income country neonatal units. This scoping review provides an overview of existing evidence and consideration of further areas of exploration.
- ⇒ The search strategy is comprehensive including broad search terms to capture as much information as possible and covers six databases, four languages and hand-searching references.
- ⇒ A scoping review is more likely to identify the breadth of evidence around parental involvement in IPC, as both interventions and study methodologies are likely to be heterogeneous and the strict criteria necessary for a systematic review may limit which studies are identified.
- ⇒ There may be publications in languages other than those searched here.
- ⇒ This study focuses on the neonatal unit only, due to the unique nature of parental involvement in this setting but acknowledges that infections can be contracted within the labour ward, at the time of delivery or in the community.

INTRODUCTION

Neonatal sepsis is a key contributor to neonatal mortality worldwide, and low- and middle-income countries (LMIC) are disproportionately affected.¹ Although neonatal sepsis is in principle a treatable illness, and the treatment is well known, there are multiple barriers to preventing, identifying and managing sepsis in LMIC. Due to these challenges, neonatal sepsis in LMIC frequently results in significant mortality, morbidity and healthcare costs.²

Within healthcare settings in LMIC, it is challenging to identify sepsis as clinical signs are often non-specific and subtle, reliant on close observation to identify.³ Additionally, there are often high patient-to-staff ratios,

meaning that close observation is extremely challenging.⁴ Blood cultures are recognised as the current gold standard required to confirm the diagnosis of sepsis and identify the causative pathogen—in LMIC, blood culture access is limited and inconsistent. This is in part due to a lack of resources but also due to challenges with the supply chain of consumables, availability of staff with the specialist skills necessary and relative cost of processing a blood culture within the healthcare system.^{5 6} Where blood cultures are available, there is often a significant delay in results or high rates of contamination, both impacting the clinical use and challenging interpretation.

Although neonatal infections are in principle treatable, increasing antimicrobial resistance is making treatment of sepsis and other bacterial infections increasingly difficult in LMIC. Indeed, the WHO recommended first-line treatment (a combination of Ampicillin and Gentamicin) is not effective in an increasing number of cases.⁷ A recent global review identified that 62.9% of positive blood cultures were Gram-negative organisms commonly resistant to WHO-recommended regimens.⁸ A meta-analysis reviewing the aetiology of invasive bacterial infections in neonates in sub-Saharan Africa indicated that in this region, Gram negative bacteria are the predominant cause of early onset sepsis, and combined with increasing antimicrobial resistance, this leads to a limited effectiveness of WHO regimens.⁷ This has, in part, been driven by antimicrobial use. For example, in Malawi, rates of resistance of Gram-negative pathogens to all first-line antimicrobials have increased dramatically since the introduction of ceftriaxone in the early 2000s.⁹

The pathogen *Klebsiella pneumoniae* is increasingly being recognised as a significant threat to global health, and in some settings carries a 70% case fatality rate.¹⁰ Reported resistance levels vary across regions and settings. Other pathogens are predominant in different areas, making the first-line choice of antibiotic challenging for clinicians, as the recommended first-line antibiotics are not effective against the predominant pathogens in the region.⁷ *Klebsiella* has traditionally been associated with late-onset sepsis (sepsis >72 hours after delivery) and is associated with healthcare-associated infections. However, it is now increasingly being identified in the first 24 hours after birth, indicating a change in the epidemiology of the timing of the onset of infection with healthcare-associated infections being transmitted in the immediate hours after birth in labour and postnatal wards.¹¹ Indeed, Zaidi *et al* have recommended that neonatal sepsis following any facility-based delivery be classified as a healthcare-associated infection, regardless of the time of onset, with this being recognised in Strengthening the Reporting of Observational Studies in Epidemiology for Newborn Infection guidelines by moving away from a binary classification of early versus late to documenting the time of onset of any infection postdelivery.^{12 13}

It is increasingly urgent to improve infection prevention and control (IPC) in LMIC neonatal units (NNU) and reduce transmission of infections. One pathway to

improvement that has yet to be underexplored is the collaboration with families to build an IPC intervention. Families are constantly present on neonatal units, and much of the hands-on care for their newborns is given by them. This provision of care, however, is informal with limited education for families in how to safely provide care for their newborns.^{4 14} Care practices that require hygiene measures (eg, handwashing, feeding, changing diapers) are often informally adopted through peer learning among mothers of newborns.

A scoping review on bundles of care addressing healthcare-associated infections (HCAI) in LMIC NNUs both highlights some strategies that may be used to involve families in IPC and the very limited involvement of parents thus far. Identified IPC interventions were analysed by each part of an intervention, called an 'element'. In total, four separate elements were found from 44 papers that related to parental involvement. Two elements related to feeding, one to skin-to-skin contact and one to empowering mothers to engage in routine care.¹⁵

For IPC to be effective, families must adhere to IPC standards within the NNU, but furthermore, any IPC intervention implemented must be feasible and acceptable for families as well as the hospital staff as this will increase uptake of and adherence to the intervention.

Research questions

At present, there is scarce literature on how parents have been involved in IPC in LMIC NNUs. This review aims to review the scope of the evidence around this topic by addressing three sub-questions.

1. How have parents been involved in designing interventions in LMIC NNUs?
2. How have parents been involved in the delivery of IPC interventions?
3. What is the parental experience of hygiene and care in the NNU in LMIC?

Rationale for a scoping review

A scoping review was chosen because at present there is limited information on parental involvement in IPC on LMIC NNUs, insufficient information to form a specific and answerable question for a systematic review, and identified methods and interventions are likely to be heterogeneous and unsuited for a systematic review. There are systematic reviews of connected concepts, for example, effective interventions to prevent infections, gaps in IPC in LMIC NNUs, parental roles in hand hygiene in paediatrics or antimicrobial resistance, but none directly about parents and IPC in LMIC.^{7 11 16 17}

Objectives

To evaluate published literature on parental involvement in the designing and implementation of IPC interventions in LMIC NNUs.

To evaluate published literature on parental experience of IPC interventions (design, delivery and impact) in LMIC NNUs.

METHODS

This review protocol has been written in compliance with the Joanna Briggs Institute manual and will be registered with the Open Science Framework database.^{18 19} The proposed review will follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist.²⁰ Research objectives, inclusion and exclusion criteria and search methods are described and defined prior to study commencement. The end date for searches is 28/02/2024.

Participants

The target population of this review are guardians (parents, families or non-medical caregivers) of both term and preterm neonates in hospitals in low- and middle-income countries.

Concept

The key concept of this review is infection prevention and control interventions in LMIC neonatal units.

Context

The context of this review is inpatient neonatal units in LMIC.

INCLUSION AND EXCLUSION CRITERIA

Criteria	Inclusion	Exclusion
Care context	Studies carried out in the neonatal unit.	Studies carried out in the community and in paediatric wards.
Publication date	Studies published from 2000 to current.	Studies published prior to 2000.
Language	English, French, Portuguese, Spanish.	Literature not available in English, French, Portuguese or Spanish.
Participants	Studies that report on parents' involvement in the design, implementation or experience of IPC interventions in NNU.	Studies with nil evidence or documentation of parental participation in design, implementation or experience of IPC interventions in NNU.
Setting	Studies must be carried out in LMIC.	Studies not carried out in LMIC.

Rationale

Study type: The concept of study type will include qualitative studies to identify studies that report on family experience of IPC, and cohort studies and programme evaluation will capture intervention development and implementation related to parents in the NNU.

Care context: Paediatric wards frequently have a different layout, staffing pattern and patient type compared with neonatal wards—this will not provide the information needed to improve IPC in NNUs. Studies aimed at care in the community will not provide information on parental involvement with IPC in LMIC NNUs.

Publication date: This time frame will allow for gathering as much evidence as possible while ensuring that

Table 1 Keywords and synonyms

Parents	IPC	Neonatal unit
Exp Parents/ Exp Family/ Mother Father Carer Caregiver Guardian Parent Family Families	exp Infection Control/ exp Hygiene/ exp Cross Infection/ *Neonatal Sepsis/ WASH HAI IPC Hygiene Neonatal infection Neonatal sepsis Infection adj2 (prevention or control)	exp Infant, Newborn/ exp Intensive Care Units, Neonatal/ NBU NNU NICU or SCBU or NNU or Special care baby unit Newborn* Neonat*
***indicates truncation of a search term.		

it remains relevant. Key understanding and evidence on neonatal infection have been gathered since 2000.

Language: The majority of studies are published in English, including French; this will ensure that French-speaking African countries are included, and the inclusion of Portuguese and Spanish will ensure that LMIC in Latin America are included.

Setting: The context of neonatal care is very different in high-income country settings. Availability of resources, such as power, water and antibiotics, makes the kinds of IPC interventions likely to be different. Although IPC principles remain the same, the challenges faced and therefore interventions needed are not likely to be transferable across contexts. Therefore, this study will include only studies carried out in LMIC. The Medline filter for low- and middle-income countries will be applied and adapted to other databases. The filter has been created using countries in the World Bank lending groups of low- and middle-income countries for 2021.

Search strategy

The search strategy including the concepts of parents, IPC and neonatal unit was created to capture all three study sub-questions.

A test search has been created in Ovid Medline and reviewed by a librarian (online supplemental file 1). The keywords and synonyms are below in [table 1](#).

Titles and abstracts will be screened for inclusion criteria concepts and keywords. Subsequent full-text screening will then be carried out based on inclusion criteria.

Filters applied

- Medline all countries LMIC
- Study type—qualitative to capture experience, interventional to capture IPC interventions from https://libguides.sph.uth.tmc.edu/search_filters/ovid_medline_filters

Limits applied

- 2000 to present

- ▶ English, French, Portuguese and Spanish

Databases

- ▶ Medline
- ▶ CINAHL
- ▶ Global Health
- ▶ EMBASE
- ▶ Web of Science
- ▶ Global Index Medicus

The search strategy was created in Medline and will be applied to EMBASE and Global Health and adapted for CINAHL, Web of Science and Global Index Medicus. This study will be carried out by two independent reviewers, with a third to discuss disagreements. All results will be imported to EndNote 21 and duplicates removed. Titles and abstracts will then be searched against the selection criteria on Rayyan. A full text review for inclusion will then be carried out. Reference searching of identified articles will be carried out. Grey literature from reference searching will be included if inclusion criteria are met; however, no formal grey literature search will be carried out.

Charting

General data points to be extracted

- ▶ Author(s)
- ▶ Year of publication
- ▶ Country where the study was conducted
- ▶ Study type
- ▶ Outcome measures

Data for studies relevant to Q1

- ▶ Stage of research parents were involved in
- ▶ What form did parental involvement take, for example, survey, focus group discussions, in-depth interviews or workshops

Data for studies relevant to Q2

- ▶ The parental role in delivering the intervention, for example, carrying out chlorhexidine washes, skin care, environmental cleaning.

Data for studies relevant to Q3

- ▶ Themes identified in parental experience related to IPC

Qualitative content analysis will be descriptive and allow an understanding of the reported experience of parents in LMIC NNU and identify common themes across LMIC, as well as the range of experience reported.

Strengths and limitations

This study will provide an overview of the involvement of parents in low- and middle-income country neonatal units. At present, although parents are frequently providing hands-on care to newborns in these settings, there is little evidence on how they are involved with infection prevention and control. This protocol will enable a broad range of evidence to be collected, as the search terms are broad, six databases are included and four languages are used. Additionally, a scoping review allows the inclusion of

heterogeneous studies so the breadth of the evidence can be understood.

Limitations of this study are that it does not include a wider range of languages—there may be papers in languages other than English, French, Spanish and Portuguese. Additionally, this scoping review focuses solely on the neonatal unit. The aim of this is to develop a review that will underpin research and policy on IPC within the NNU. However, there may be evidence on parental involvement in hygiene and infection control at home or in other settings that may be related and informative.

Dissemination

The findings of this review will be disseminated to key stakeholders through publication and presentations. The findings will also be used to inform future policy and research aimed at improving infection prevention and control in low- and middle-income neonatal units.

Patient and public involvement

We plan to utilise the results of this study to underpin future collaborative research with families in neonatal units, as well as to engage patient representatives for communication or dissemination of results.

X Hannah Blencowe @HannahBlencowe

Acknowledgements The authors would like to thank Kathleen Perris (librarian at the London School of Hygiene and Tropical Medicine Library) for her advice and guidance when developing the search strategy for this review.

Contributors LD and CK are joint first authors. LD conceived the idea for the review, developed the research questions and study methods and drafted the protocol. RC, HB, SM, FF and GC supervised the protocol design and continue to supervise the review process. LD and CK piloted the protocol and will screen records for inclusion. GC will act as a third party in case of disagreement. All authors discussed the protocol design and critically edited and approved the final manuscript. LD is the guarantor.

Funding This work was carried out as part of LD PhD research, funded by the Wellcome Trust [318536/Z/24/Z] under the CREATE PhD programme.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Patients and the public were not consulted on the scope of this review. However, the review aims to understand current literature on parental involvement in aspects of neonatal care and aims to underpin future collaborative research with families in neonatal units.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Lydia Davidson <http://orcid.org/0009-0009-3150-2234>
Hannah Blencowe <http://orcid.org/0000-0003-1556-3159>
Felicity Fitzgerald <http://orcid.org/0000-0001-9594-3228>
Sarah Moxon <http://orcid.org/0000-0001-6600-6147>

REFERENCES

- 1 Fleischmann C, Reichert F, Cassini A, *et al.* Global incidence and mortality of neonatal sepsis: a systematic review and meta-analysis. *Arch Dis Child* 2021;106:745–52.
- 2 Milton R, Gillespie D, Dyer C, *et al.* Neonatal sepsis and mortality in low-income and middle-income countries from a facility-based birth cohort: an international multisite prospective observational study. *Lancet Glob Health* 2022;10:e661–72.
- 3 Seale AC, Blencowe H, Manu AA, *et al.* Estimates of possible severe bacterial infection in neonates in sub-Saharan Africa, south Asia, and Latin America for 2012: a systematic review and meta-analysis. *Lancet Infect Dis* 2014;14:731–41.
- 4 Oluoch D, Hinton L, English M, *et al.* Mothers' involvement in providing care for their hospitalised sick newborns in Kenya: a focused ethnographic account. *BMC Pregnancy Childbirth* 2023;23:389.
- 5 Chimhini G, Chimhuya S, Madzudzo L, *et al.* Auditing use of antibiotics in Zimbabwean neonates. *Infect Prev Pract* 2020;2:100046.
- 6 Murless-Collins S, Kawaza K, Salim N, *et al.* Blood culture versus antibiotic use for neonatal inpatients in 61 hospitals implementing with the NEST360 Alliance in Kenya, Malawi, Nigeria, and Tanzania: a cross-sectional study. *BMC Pediatr* 2023;23:568.
- 7 Okomo U, Akpalu ENK, Le Doare K, *et al.* Aetiology of invasive bacterial infection and antimicrobial resistance in neonates in sub-Saharan Africa: a systematic review and meta-analysis in line with the STROBE-NI reporting guidelines. *Lancet Infect Dis* 2019;19:1219–34.
- 8 Russell NJ, Stöhr W, Plakkal N, *et al.* Patterns of antibiotic use, pathogens, and prediction of mortality in hospitalized neonates and young infants with sepsis: A global neonatal sepsis observational cohort study (NeoOBS). *PLoS Med* 2023;20:e1004179.
- 9 Iroh Tam P-Y, Musicha P, Kawaza K, *et al.* Emerging Resistance to Empiric Antimicrobial Regimens for Pediatric Bloodstream Infections in Malawi (1998–2017). *Clin Infect Dis* 2019;69:61–8.
- 10 Chimhini G, Olaru ID, Fitzgerald F, *et al.* Evaluation of a Novel Culture System for Rapid Pathogen Identification and Detection of Cephalosporin Resistance in Neonatal Gram-negative Sepsis at a Tertiary Referral Unit in Harare, Zimbabwe. *Pediatr Infect Dis J* 2021;40:785–91.
- 11 Fitzgerald FC, Zingg W, Chimhini G, *et al.* The Impact of Interventions to Prevent Neonatal Healthcare-associated Infections in Low- and Middle-income Countries: A Systematic Review. *Pediatr Infect Dis J* 2022;41:S26–35.
- 12 Zaidi AKM, Huskins WC, Thaver D, *et al.* Hospital-acquired neonatal infections in developing countries. *Lancet* 2005;365:1175–88.
- 13 Fitchett EJA, Seale AC, Vergnano S, *et al.* Strengthening the Reporting of Observational Studies in Epidemiology for Newborn Infection (STROBE-NI): an extension of the STROBE statement for neonatal infection research. *Lancet Infect Dis* 2016;16:e202–13.
- 14 Sunkwa-Mills G, Rawal L, Enweronu-Laryea C, *et al.* Perspectives and practices of healthcare providers and caregivers on healthcare-associated infections in the neonatal intensive care units of two hospitals in Ghana. *Health Policy Plan* 2020;35:i38–50.
- 15 Molina García A, Cross JH, Fitchett EJA, *et al.* Infection prevention and care bundles addressing health care-associated infections in neonatal care in low-middle income countries: a scoping review. *EClinicalMedicine* 2022;44:101259.
- 16 Bellissimo-Rodrigues F, Pires D, Zingg W, *et al.* Role of parents in the promotion of hand hygiene in the paediatric setting: a systematic literature review. *J Hosp Infect* 2016;93:159–63.
- 17 Yee D, Osuka H, Weiss J, *et al.* Identifying the priority infection prevention and control gaps contributing to neonatal healthcare-associated infections in low-and middle-income countries: results from a modified Delphi process. *J Glob Health Rep* 2021;5.
- 18 Peters MD, Godfrey C, McInerney P, *et al.* Chapter 11: scoping reviews. In: *JBI manual for evidence synthesis*. 2020.
- 19 Available: <https://doi.org/10.17605/OSF.IO/NMK9P>
- 20 Tricco AC, Lillie E, Zarin W, *et al.* PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018;169:467–73.