

The relationship between employment and social participation among Australians with a disabling chronic health condition: a cross-sectional analysis

Emily Callander, Deborah J Schofield

To cite: Callander E, Schofield DJ. The relationship between employment and social participation among Australians with a disabling chronic health condition: a cross-sectional analysis. *BMJ Open* 2013;**3**:e002054. doi:10.1136/bmjopen-2012-002054

► Prepublication history for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-002054>).

Received 4 September 2012
Accepted 21 December 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

The Sydney Medical School, NHMRC Clinical Trials Centre, University of Sydney, Camperdown, New South Wales, Australia

Correspondence to

Dr Emily Callander;
emily.callander@ctc.usyd.edu.au

ABSTRACT

Objectives: Social interaction may be particularly important for people with chronic health conditions, due to the numerous benefits to an individual's health. This paper aims to determine if labour force participation is a factor that influences individuals with chronic health conditions partaking in social or cultural events.

Design and setting: The study undertakes a cross-sectional analysis of the *2009 Survey of Disability, Ageing and Carers*, a nationally representative survey of the Australian population.

Participants: 33 376 records of persons aged 25–64 years.

Outcome measures: Participation in social and community activities.

Results: It was found that after controlling for age, sex, level of highest education, income unit type and severity of disability, people with a chronic health condition that were in the labour force were more than twice as likely to be participating in social or community events (OR 2.54, 95% CI 1.95 to 3.29, $p<0.0001$), and in cultural events (OR 2.57, 95% CI 2.21 to 3.00, $p<0.0001$) as their counterparts who were out of the labour force. The results were then repeated, with the addition of income as a confounding variable. People with a chronic health condition that were in the labour force were still a little more than twice as likely to be participating in social or community events (OR 2.25, 95% CI 1.69 to 3.00, $p<0.0001$), and to be participating in cultural events (OR 2.08, 95% CI 1.76 to 2.45, $p<0.0001$) as their counterparts who were out of the labour force.

Conclusions: Participating in the labour force may be an important driver of social participation among those with chronic health conditions, independent of income. People with chronic health conditions who are not in the labour force and do not participate in social or cultural activities may have a compounding disadvantage.

INTRODUCTION

Participating in social events and having access to social networks are known to

ARTICLE SUMMARY

Article focus

- Does labour force participation influence the likelihood of an individual with poor health engaging in social events?

Key messages

- People with a chronic health condition in the labour force were 2.3 times more likely to be participating in social events.
- People with a chronic health condition in the labour force were 2.1 times more likely to be participating in cultural events.

Strengths and limitations of this study

- A limitation of this study is that it was conducted using cross-sectional data.

produce numerous health benefits. Some studies have documented improved survival rates associated with social activities,^{1–4} with one study finding that social activities had similar survival advantages as exercise activities.² Similarly, social participation has been associated with lower chronic disease rates,⁵ positive outcomes relating to cardiovascular, endocrine and immune systems,⁶ improved quality of life and better overall mental well-being.^{7–8} The potential health benefits of social participation also extend to those who already have chronic health conditions.^{9–11}

The explanations for these health benefits are complex and diverse, ranging from social support being linked with a lower likelihood of chronic disease risk factors such as smoking and lack of exercise to tangible benefits such as help in times of need, to biochemical explanations such as reductions in the levels of fibrinogen and catecholamine, to social contact reducing stress by enhancing cellular and humoral immune responses.^{5–6}

Regardless of the explanation, this past research has highlighted the importance of social networks and socialising for better health.

It is known that labour force participation has been linked with higher rates of social inclusion.^{12–14} Therefore, participating in the labour force may be an important factor in facilitating participation in social activities. Given the high chance people with chronic health conditions have of exiting the labour force—with one Australian study finding that over half of the 45-year-old to 64-year-old Australians with a chronic health condition were not in the labour force¹⁵—it may be likely that people who have had to retire early due to their ill health may also be excluded from social activities and may be missing out on the associated health benefits.

This paper will explore the relationship between participation in the labour force and participation in social, community or cultural activities among Australians aged 25–64 years with a chronic health condition. We will examine whether people with a chronic health condition who are in the labour force are more likely to participate in social and community activities than those who are out of the labour force.

It is possible that those in the labour force may have higher rates of social participation as their employment income may give them the financial capacity to do so. It has been found that in a number of European countries those who experience long-term unemployment have less chance of meeting with friends and participating in social clubs or organisations, and the authors theorised that this could be due to financial difficulties in maintaining relationships.¹⁴ In Australia, Saunders¹⁶ reported that 16% of households could not afford to participate in social activities, and Berry has also speculated on the restrictions that the lack of financial resources plays on social and community participation.¹⁷ As such, labour force participation could facilitate social and community participation as it merely provides the financial capability. Owing to this possibility, the results will be controlled for the influence of income unit income.

METHODS

This study was conducted using the 2009 *Survey of Disability, Ageing and Carers* (SDAC), a nationally representative survey of health and disability conducted by the Australian Bureau of Statistics.¹⁸ The 2009 SDAC had a response rate of 89.9%, with 28 474 households participating, and 64 123 persons fully responding. The data are weighted such that the weight assigned to each survey record provides an estimate of the number of similar individuals in the Australian population,¹⁸ which allowed nationally representative results to be produced.

Measure of social participation

Respondents in the 2009 SDAC were asked about their participation in social or community activities away from

home in the 3 months prior to the interview and their culture or leisure activities in the 12 months prior to the interview. Social or community participation away from home consisted of either visiting relatives or friends or a restaurant or club, or participating in church, a voluntary performing arts groups, art or craft group or other special interest group activities or other activities not specified. Cultural or leisure activities consisted of either visiting a museum, art gallery, library, animal or marine park, or botanic gardens, or attending the theatre, a concert, cinema or sports event as a spectator, or taking part in a sport or recreation activity.

The study was limited to people aged 25–64 years and identified people who had a chronic health condition and indicated that their condition was disabling. On the 2009 SDAC, all people with a disability had a long-term health condition, but not all people with a long-term health condition had a disability. Within the 2009 SDAC, a disability is defined as ‘a limitation, restriction or impairment, which has lasted, or is likely to last, for at least six months and restricts everyday activities’.¹⁸ Previous studies have shown that there is considerable variation in the functional outcomes of people with different health conditions.^{15–19} As the authors were concerned with identifying people with poor health and whose capabilities were limited by their health condition, those with a long-term health condition (chronic health condition) that was disabling were selected to be the focus of this paper. In this paper, the comparator group—those who did not have a disabling chronic health condition—consists of those with no chronic health conditions and those with a chronic health condition who stated that they did not have a disability.

Statistical analysis

Initial descriptive analysis was undertaken to compare the rates of labour force participation and social interaction between those with and without a disabling chronic health condition. The study compared individuals who were in and out of the labour force, with ‘in the labour force’ being defined as being employed full time, employed part time or looking for full-time or part-time work.

Logistic regression models were then constructed to look at the OR of participating in social or community activities and cultural or leisure activities for those with a disabling chronic health condition and who were in the labour force compared with those out of the labour force. The models were adjusted for age, sex, level of highest education, severity of disability and income unit type (single, couple, single parent with dependent children or couple with dependent children). Disability severity classified people as having either a disability and not limited in core activities or restricted in schooling or employment; a disability and not limited in core activities but restricted in schooling or employment; a disability and mildly limited in core activities; a disability and moderately limited in core activities; a disability and

severely limited in core activities or a disability and profoundly limited in core activities. This severity of disability variable has been shown to correlate with self-assessed health status,²⁰ but was considered to provide more detailed information on health and functional limitation.

A sensitivity analysis was conducted to control for the impact of income unit income. The logistic regression models were repeated with the inclusion of the income unit income decile. The analyses were undertaken using SAS V.9.1 (SAS Institute Inc, Cary, North Carolina, USA). All statistical tests were two sided with the significance level set at 5%.

RESULTS

In the 2009 SDAC, there were 33 376 records of people aged 25–64 years who were members of an income unit. Once weighted, this represented 11 350 900 individuals in the Australian population in 2009. There were 5704 records of persons aged 25–64 years who had a chronic health condition and stated that their condition was disabling (representing 1 896 400 individuals in the Australian population in 2009).

Of the people who had a chronic health condition that was disabling, 55% were in the labour force (1 035 000 individuals) and 45% were out of the labour force (861 400 individuals). Of those who did not have a disabling chronic health condition, 86% were in the labour force (8 141 000 individuals) and 14% were out of the labour force (1 313 500 individuals).

Table 1 shows the proportion of people that were participating in various social activities among those with a chronic health condition that was disabling and those without a disabling chronic health condition who were in and out of the labour force. Almost 100% of the people who did *not* have a disabling chronic health condition participated in social or community activities, regardless of the labour force participation status. Similarly, almost all (96%) of the people with a chronic health condition who were in the labour force participated in a social or community activity, whereas only 88% of those with a chronic health condition that were not in the labour force did so.

Almost all people *without* a disabling chronic health condition participated in cultural or leisure activities, regardless of whether they were in or out of the labour force (99% and 96%, respectively). A higher proportion of people with a disabling chronic health condition that were in the labour force participated in cultural or leisure activities (85%) than people with a disabling chronic health condition that were not in the labour force (60%). Over one-third of those out of the labour force with a disabling chronic health condition had *not* participated in a cultural or leisure activity in the past 12 months in 2009.

The analysis was then limited to those with a disabling chronic health condition. After adjusting for age, sex, level of highest education attainment, severity of disability and income unit type, there were found to be significant differences in the experience of social contact between those in the labour force with a chronic health condition and those out of the labour force with a chronic health condition (table 2). Those in the labour force with a chronic health condition were more than twice as likely to be participating in social or community activities away from home compared with those not in the labour force with a chronic health condition (OR 2.54, 95% CI 1.95 to 3.29). Those in the labour force with a chronic health condition were 2.57 times more likely to have participated in cultural or leisure activities away from home than people with a chronic health condition that were out of the labour force (OR 2.57, 95% CI 2.21 to 3.00).

Sensitivity analysis

After adjusting for income unit income as well as age, sex, level of highest education attainment, severity of disability and income unit type, there was still a significant difference in the experience of social contact between those in the labour force with a chronic health condition and those out of the labour force with a chronic health condition (table 3). After also adjusting for income, those in the labour force with a chronic health condition were still more than twice as likely to have participated in social or community activities away from the home compared with those not in the labour force with a chronic health condition (OR 2.25, 95% CI 1.69 to

Table 1 Proportion of individuals, in and out of the labour force, participating in activities away from home, 2009

	Participating in social or community activities away from home		Participating in cultural or leisure activities away from home		Total
	N	%	N	%	
Did not have a disabling chronic health condition					
In the labour force	8 126 300	100	8 074 300	99	8 141 000
Not in the labour force	1 302 100	99	1 266 600	96	1 313 500
Had a disabling chronic health condition					
In the labour force	994 800	96	880 400	85	1 035 000
Not in the labour force	759 500	88	525 200	60	861 500

Table 2 Likelihood of participating in social or community activities away from home or participating in cultural or leisure activities away from home, adjusted for age, sex, education, severity of disability and income unit type, Australians aged 25–64 years, 2009

	Participating in social or community activities away from home			Participating in cultural or leisure activities away from home		
	OR	95% CI	p Value	OR	95% CI	p Value
In the labour force	Reference			Reference		
Not in the labour force	2.54	1.95 to 3.29	<0.0001	2.57	2.21 to 3.00	<0.0001

3.00). Those in the labour force with a chronic health condition were now 2.08 times more likely to have participated in social or community activities away from the home than people with a chronic health condition that were out of the labour force (OR 2.08, 95% CI 1.76 to 2.45).

DISCUSSION

A higher proportion of people who did *not* have a disabling chronic health condition participated in social and cultural activities, regardless of labour force participation status, than people who had a disabling chronic health condition. Among people with a disabling chronic health condition, people who were in the labour force were significantly more likely to participate in social and cultural activities. In this study, the authors controlled for the effects of age, sex, level of education attainment, severity and type of disability, income unit type and level of income, indicating that labour force participation may have unique benefits that foster participation in social activities.

The economic benefit of labour force participation for people with a health condition has been well established. It is known that those who retire early due to ill health have a lower income, lower amounts of accumulated wealth and will reach the age of 65 years with lower amounts of savings to support their retirement than those who remain in the labour force.^{20–25} However, this study has shown that there are additional social benefits of remaining in the workforce that are independent from the economic benefits.

Causality

A limitation of this study is that it was conducted using cross-sectional data. As a result of this, only the

relationship between labour force participation and participation in social and cultural activities at one point in time could be assessed. While participating in the labour force may facilitate social and cultural participation among those with a chronic health condition, it is also possible that participating in social and cultural activities may facilitate labour force participation among those with a disabling chronic health condition.

It has been theorised that being part of a social network gives members access to ‘social leverage’—a form of social capital that is generated by a network of people. Social leverage helps members of a network to access information to improve their social position, such as details of employment opportunities or job referrals. However, access to this information is usually limited to members of the social network.²⁶ Developing a chronic health condition may result in individuals developing limitations that may restrict the type of tasks they can perform. This may require people to change jobs or industries to find employment that provides suitable working conditions. As such, being part of a social network may increase the chances of labour force participation among those with poor health. However, despite theories that link social networks and finding employment, a longitudinal study of European Union countries reported no link between social networks and finding employment.²⁷ Thus, it is likely that labour force participation does facilitate social participation, rather than the other way around.

Health implications

Regardless of causality, those with a disabling chronic health condition who are out of the labour force have lower rates of social and community participation, and so may be forgoing the health benefits that social

Table 3 Sensitivity analysis: likelihood of participating in social or community activities away from home or participating in cultural or leisure activities away from home, adjusted for age, sex, education, severity of disability, income unit type and income unit income, Australians aged 25–64 years, 2009

	Participating in social or community activities away from home			Participating in cultural or leisure activities away from home		
	OR	95% CI	p Value	OR	95% CI	p Value
In the labour force	Reference			Reference		
Not in the labour force	2.25	1.69 to 3.00	<0.0001	2.08	1.76 to 2.45	<0.0001

participation provides. Social capital—which includes, but is not limited to, social participation—is linked to health status.^{28–29} Berry and Welsh³⁰ have looked at the relationship between social capital and health in Australia and found that informal social connections and civic engagement were linked to better health outcomes. Informal social connections and civic engagement broadly align with the variables of social participation utilised in this study. Social exclusion has also been linked with poor health outcomes,³¹ and some studies have documented an improvement in health that can be gained by those with chronic health conditions participating in social activities and through engaging in social networks.^{9–11}

This study has shown that people with a disabling chronic health condition who are not in the labour force are less likely to be participating in social and cultural activities. The study controlled for a number of variables that may have confounded this relationship—such as income, income unit type (being in a relationship or single, with or without children) and severity and type of disability. This indicates that participating in the labour force in itself may be an important driver of social participation and an important opportunity for social contact among those with chronic health conditions. Employment may provide people with chronic health conditions the opportunity to create social contacts and networks, as well as also the opportunity to engage with them.

Contributors EC conceived the study, carried out the data analysis and drafted the manuscript. DJS provided expert advice for the design of the study and the interpretation of the results, and contributed to the final manuscript.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None.

Ethics approval The use of this data was approved by the Australian Bureau of Statistics.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

REFERENCES

- Bygnen LO, Konlaan BB, Johansson SE. Attendance at cultural events, reading books or periodicals, and making music or singing in a choir as determinants for survival: Swedish interview survey of living standards. *BMJ* 1996;313:1577–80.
- Glass TA, Mendes de Leon C, Martotoli RA, et al. Population based study of social and productive activities as predictors of survival among elderly Americans. *BMJ* 1999;319:478–83.
- House JS, Robbins C, Metzger HL. The association of social relationships and activities with mortality: prospective evidence from the Tecumseh Community Health Study. *Am J Epidemiol* 1982;116:123–40.
- Kaplan GA, Salonen JT, Cohen RD, et al. Social connections and mortality from all causes and from CVD: prospective evidence from eastern Finland. *Am J Epidemiol* 1988;128:370–80.
- Orth-Gomer K, Rosengren A, Wilhelmsen L. Lack of social support and incidence of coronary heart disease in middle-aged Swedish men. *Psychosom Med* 1993;55:37–43.
- Uchino BN, Cacioppo JT, Kiecolt-Glaser JK. The relationship between social support and physiological processes: a review with emphasis on underlying mechanisms and implications for health. *Psychol Bull* 1996;119:488–531.
- Bloom JR, Stewart SL, Johnston M, et al. Sources of support and the physical and mental well-being of young women with breast cancer. *Soc Sci Med* 2001;53:1513–24.
- Lynch EB, Butt Z, Heinemann A, et al. A qualitative study of quality of life after stroke: the importance of social relationships. *J Rehabil Med* 2008;40:518–23.
- Gallant MP. The influence of social support on chronic illness self-management: a review and directions for research. *Health Educ Behav* 2003;30:170–95.
- Bennett SJ, Perkins SM, Lane KA, et al. Social support and health-related quality of life in chronic heart failure patients. *Qual Life Res* 2001;10:671–82.
- Cohen S. Social relationships and health. *Am Psychol* 2004;59:676–84.
- Paugam S. The spiral of precariousness: a multidimensional approach to the process of social disqualification in France. In: Room G. *Beyond the threshold: the measurement and analysis of social exclusion*. Bristol: Policy Press, 1995.
- Twena M, Aaheim HA. *Social exclusion and unemployment in the European Union: current and future trends*. Oslo: Center for International Climate and Environmental Research, 2005.
- Furaker B. On the social consequences of unemployment. In: Policy Department EaSP, European Parliament. Brussels: European Parliament, 2010.
- Schofield D, Shrestha R, Passey M, et al. Chronic disease and labour force participation among older Australians. *Med J Aust* 2008;189:447–50.
- Saunders P. *Can social exclusion provide a new framework for measuring poverty? Discussion paper 127*. Sydney: Social Policy Research Centre (SPRC), 2003.
- Berry HL. Social capital elite, excluded participants, busy working parents and aging, participating less: types of community participants and their mental health. *Soc Psychiatry Psychiatr Epidemiol* 2008;43:527–37.
- Australian Bureau of Statistics. *Disability, ageing and carers, Australia: user guide ABS catalogue No. 4431.0.55.001*. Canberra: Australian Bureau of Statistics, 2011.
- Callander E, Schofield D, Shrestha R. Freedom poverty: a new tool to identify the multiple disadvantages affecting those with CVD. *Int J Cardiol* 2011; (Epub ahead of print).
- Callander EJ, Schofield DJ, Shrestha RN, et al. Health and functional capacity-capturing capability limitations in measures of health. *Health Soc Rev* 2012;21:322–31.
- Schofield D, Kelly S, Shrestha R, et al. Long term financial impacts of CVD: living standards in retirement. *Int J Cardiol* 2010;155:406–8.
- Schofield D, Percival R, Passey M, et al. The financial vulnerability of individuals with diabetes. *Br J Diab Vasc Dis* 2010;10:300–4.
- Schofield D, Shrestha R, Percival R, et al. Early retirement and the financial assets of individuals with back problems. *Eur Spine J* 2011;20:731–6.
- Schofield D, Shrestha R, Percival R, et al. Quantifying the effect of early retirement on the wealth of individuals with depression or other mental illness. *Br J Psychiatry* 2011;198:123–8.
- Schofield D, Shrestha R, Callander E, et al. Modelling the cost of ill health in Health&WealthMOD (Version II): lost labour force participation, income and taxation, and the impact of disease prevention. *Int J Microsimul* 2011;4:32–6.
- Carpiano RM. Towards a neighbourhood resource-based theory of social capital for health: can Bourdieu and sociology help? *Soc Sci Med* 2006;62:165–75.
- Gallie D, Paugam S, Jacobs S. Unemployment, poverty and social isolation: is there a vicious circle of social exclusion? *Eur Soc* 2003;5:1–32.
- Lochner KA, Kawachi I, Brennan RT, et al. Social capital and neighbourhood mortality rates in Chicago. *Soc Sci Med* 2003;56:1797–805.
- De Silva MJ, McKenzie K, Haroman T, et al. Social capital and mental illness: a systematic review. *J Epidemiol Commun Capital* 2005;59:619–27.
- Berry HL, Welsh JA. Social capital and health in Australia: an overview from the household, income and labour dynamics in Australia survey. *Soc Sci Med* 2010;70:588–96.
- Shaw M, Doorling D, Smith GD. Poverty, social exclusion, and minorities. In: Marmont M, Wilkinson R, eds. *Social determinants of health*. Online: Oxford Scholarship Online, 2005.