

BMJ Open

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-005292
Article Type:	Research
Date Submitted by the Author:	20-Mar-2014
Complete List of Authors:	Ford, John; University of East Anglia, Public Health Shyangdan, Deepson; University of Warwick, Warwick Evidence, Warwick Medical School Uthman, Olalekan; Warwick Medical School, Lois, Noemi; Queen's University, Centre for Vision and Vascular Science, Waugh, Norman; University of Warwick, Warwick Evidence
Primary Subject Heading:	Ophthalmology
Secondary Subject Heading:	Health policy
Keywords:	Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Medical retina < OPHTHALMOLOGY, Medical ophthalmology < OPHTHALMOLOGY

SCHOLARONE™
Manuscripts

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

John A Ford, Deepson Shyangdan, Olalekan A. Uthman, Noemi Lois, Norman Waugh

John A Ford, Academic Clinical Fellow in Public Health, Department of Population Health and Primary Care, University of East Anglia
Deepson Shyangdan, Research Fellow, Warwick Evidence, University of Warwick
Olalekan Uthman, Assistant Professor in Applied Research, University of Warwick
Noemi Lois, Professor of Ophthalmology, Centre for Vision and Vascular Science, Queen's University
Norman Waugh, Professor of Health Technology Assessment, University of Warwick

Corresponding author
Dr. John Ford
Norwich Medical School
University of East Anglia
Chancellor Drive
Norwich
NR4 7TJ
John.ford@uea.ac.uk
Tel: 01603 591269

Protocol – no published protocol exists for this study.

Ethics approval – not required

Funding – no external funding required

Data sharing: no additional data available

Word count: 3,331 words

What is already known on this subject

Anti-VEGF drugs (ranibizumab, bevacizumab and aflibercept) and corticosteroids (dexamethasone and triamcinolone), given intravitreally, have all been shown to be effective compared to placebo for the treatment of macular oedema secondary to central retinal vein occlusion.

There are no head-to-head trials.

What this study adds

There was no evidence of a difference in the effectiveness of aflibercept, ranibizumab, bevacizumab and triamcinolone for improving vision.

Clinicians may prefer aflibercept because steroids are associated with cataract formation and ranibizumab might require more frequent injections.

Abstract

Objective: To indirectly compare aflibercept, bevacizumab, dexamethasone, ranibizumab and triamcinolone for treatment of macular oedema secondary to central retinal vein occlusion using a network meta-analysis.

Design: Network meta-analysis

Data sources: The following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE; CDSR, DARE, HTA, NHSEED, CENTRAL; Science Citation Index and Conference Proceedings Citation Index-Science

Eligibility criteria for selecting studies: Only randomized controlled trials assessing patients with macular oedema secondary to central retinal vein occlusion were included. Studies had to report either proportions of patients gaining more than or equal to 3 lines, losing more than or equal to 3 lines, or mean change in best corrected visual acuity. Two authors screened titles and abstracts, extracted data and undertook risk of bias assessment. Bayesian network meta-analysis was used to compare the different interventions.

Results: Seven studies, assessing five drugs, were judged to be sufficiently comparable for inclusion in the NMA. For the proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2mg had a higher probability of being more effective than sham and dexamethasone. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision compared to those treated with sham. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab

1.25mg and aflibercept 2mg had a higher probability of improvement in mean best correct visual acuity compared to those treated with sham injections.

Conclusions: We found no evidence of differences between ranibizumab, aflibercept, bevacizumab and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections

Systematic review registration – Not registered

Strengths and limitations of this study

- Important topic area, with significant policy implications
- Robust method used to identify studies
- Network meta-analysis are based on a number of assumptions
- Network meta-analysis is the best method to compare interventions in the absence of head to head trials

Introduction

Central retinal vein occlusion (CRVO) dramatically reduces an individual's functioning and quality of life.[1] It is estimated that the 15 year cumulative incidence of central retinal vein occlusion is 0.5%.[2] Visual loss is caused by thrombosis of the central retinal vein which leads to a rise in venous pressure and an increase in vascular endothelial growth factor (VEGF), consequently causing an increase in vascular permeability. Macular oedema subsequently ensues with varying degrees of ischaemia and neo-vascularisation. Although CRVO is generally classified as ischaemic or non-ischaemic, ischaemia should be regarded as a spectrum.[3] Cases with ischaemia carry a considerably worse prognosis as in around a third of them, neovascular glaucoma may develop; the most devastating complication of CRVO.[4]

CRVO is more common in older people with risk factors such as diabetes, hypertension or hyperlipidaemia, but can occur in young people with inflammatory disorders. Hayreh and colleagues in a 27-year cohort study found that only 13% of people with CRVO were under 45 years of age.[3] In 95% of cases CRVO affects only one eye.[3] However visual loss in this already co-morbid patient group significantly compounds their already impaired functioning and quality of life. Patients can lose confidence, struggle with daily activities and become increasingly dependent on friends and family.[1]

For many years, laser photocoagulation was the only effective therapeutic strategy that could be used in the management of patients with CRVO. It was only useful for reducing the risk of neovascular glaucoma, but not effective for the treatment of macular oedema in CRVO.[5] Over the past decade a number of drugs to treat macular oedema have been introduced, including the steroids, triamcinolone and dexamethasone, and the anti-VEGFs, ranibizumab, bevacizumab, pegaptanib and aflibercept. Dexamethasone,

ranibizumab and aflibercept have been assessed in large commercially funded trials.[6-13] Bevacizumab was originally developed as an anti-cancer drug and has been found to be effective in treating macular oedema secondary to age-related macular degeneration,[14] diabetic macular oedema, [15] branch retinal vein occlusion[16] and central retinal vein occlusion.[17] Like triamcinolone, bevacizumab is used off licence in the eye. Ranibizumab is a derived from the same parent molecule of the bevacizumab monoclonal antibody and was developed and commercially marketed specifically for use in the eye.

In the United Kingdom, the National Institute of Health and Care Excellence (NICE) has recommended the use of dexamethasone, ranibizumab and aflibercept for the treatment of macular oedema secondary to CRVO in separate appraisals[18-20] Therefore clinicians have three NICE-recommended treatments for CRVO without head-to-head trials or clear guidance on which one may be best for their patients. On this basis, the aim of this study was to indirectly compare in a network meta-analysis the clinical effectiveness of aflibercept, ranibizumab, bevacizumab, dexamethasone and triamcinolone for the treatment of macular oedema secondary to CRVO.

Methods

Information sources and search strategy

To identify suitable studies, initially for a systematic review of treatment of macular oedema after CRVO (submitted for publication) the following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE (all via OVID); CDSR, DARE, HTA, NHSEED, CENTRAL (all via The Cochrane Library); Science Citation Index and Conference Proceedings Citation Index-Science (via Web of Knowledge). The MEDLINE search strategy is shown in appendix 1. This search strategy was modified for other databases. In addition to the bibliographic database searching, supplementary searches were undertaken to look for recent and unpublished studies in the WHO International Clinical Trials Registry Platform and ophthalmology conference websites (American Academy of Ophthalmology, Association for Research in Vision and Ophthalmology from 2010 to 2012).

Study selection

Only randomised controlled trials which included patients with macular oedema secondary to central retinal vein occlusion were included. It was acceptable for a study to include both branch retinal vein occlusion and central retinal vein occlusion provided that the central retinal vein occlusion group was reported separately. The following drugs were included: dexamethasone, triamcinolone, ranibizumab, bevacizumab and aflibercept. Pegaptanib was not included because it is not used routinely in clinical practice. Only doses which are used in clinical practice were included. Studies had to report at least one of the following outcomes: proportions of patients gaining more than or equal to 3 lines from baseline to six months, proportions of patients losing more than or equal to 3 lines from baseline to six months and mean change in best corrected visual acuity (BCVA) from baseline to six months

Risk of bias assessment

The Cochrane Collaboration's tool for assessing risk of bias was used.[21] The trials were graded (unclear, high or low risk of bias) based on: (i) sequence generation, (ii) allocation concealment, (iii) blinding of outcome assessor, (iv) incomplete outcome data, and (v) selective outcome reporting.

Study selection and data abstraction

Two authors independently assessed the eligibility and methodological quality of the studies identified during the literature search. Two authors extracted and compared the data. For each study identified that met the selection criteria, details on study design, study population characteristics, intervention, outcome measures, and study quality were extracted. Discrepancies were resolved by consensus through discussion. Studies were assessed for comparability based on the populations included, trial arms, outcome measures and duration of follow-up. Common comparators were identified from the trials and a network diagram was created.

Summary measures

The primary measures of treatment effects were relative risk (RR) for the proportions of patients gaining more than or equal to 3 lines of vision, proportions of patients losing more than or equal to 3 lines of vision and weighted mean difference (WMD) for mean change BCVA. We used the following methods to calculate standard deviations, when incompletely reported: (1) contact with the corresponding author; or (2) estimation of the standard deviation on the basis of the sample size, median, and range as suggested by Hozo and colleagues[22] or on the basis of the sample size and P value.

In one trial (SCORE),[23-36] six month data was not available because patients were followed up every four months. For the dichotomous outcomes i.e. proportions of patients gaining and losing ≥ 3 lines, we averaged four and eight month data to get the six months follow-up data. For the third outcome i.e. mean change BCVA, again data from two time-points were used. Weighted mean and SDs for each treatment arm was calculated using mean and SDs of two time-points.

Data synthesis and model implementation

Bayesian network meta-analysis [37 38] (NMA) was used to compare the different interventions. Network meta-analysis is a generalization of meta-analysis methods because they allow comparisons of agents not addressed within individual primary trials. Bayesian statistical inference provides probability distributions for treatment effect parameters (RR and WMD), with 95% credible intervals (95% CrI), rather than 95% confidence intervals (95% CI). A 95% credible interval can be interpreted as there being a 95% probability that the parameter takes a value in the specified range.[37 38]

All analyses were conducted using a Bayesian Markov Chain Monte Carlo (MCMC) method and fitted in the freely available Bayesian software, WinBUGS 1.4.3.[39] Two Markov chains were run simultaneously using different initial values. Convergence to a stable solution was checked by viewing plots of the sampled simulations and using the Brooks-Gelman-Rubin diagnostic tool.[40] Convergence was found to be adequate after running 20 000 samples for both chains. These samples were then discarded and a further 70 000 sampled simulation was then run, on which the results were based. We also calculated the probability of treatment being the most effective (first best), the second best, the third best, and so on, and presented the results graphically with rankograms.[41]

Like standard meta-analysis comparison, a NMA can be either a fixed- or a random-effect models. We used the Bayesian Deviation Information Criterion (DIC) to compare fixed and random effect models. The most appropriate NMA model can be identified as the one with the lowest DIC. The DIC measures the fit of the model while penalizing it for the number of effective parameters. The fixed - effect model was chosen because of the small number of trials available for each comparison and difficulty in estimating between studies variance if random-effect model was implemented and the difference in DIC is less than 5.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Study selection and characteristics

The literature search identified 945 articles, as shown in Figure 1. Seven studies were judged to be sufficiently comparable to be included in the network meta-analysis. Tables 1 and 2 present the characteristics and results of the included trials. Two studies [11-13] compared aflibercept 2 mg against sham; two identical studies [6-8] compared dexamethasone 0.7 mg (Ozurdex) against sham; one study [9 10] compared ranibizumab 0.5 mg against sham; one study [42-44] compared bevacizumab 1.25 mg against sham, and finally one study [23-36] compared triamcinolone 4 mg against observation. Sham or observation were used as the common comparator. The number of included participants varied from 60 [42-44] to 437 [6-8]. Most studies required patients to be treatment naive and have macular oedema with retinal thickness measuring at least 250 or 300 µm on optical coherence tomography. Sham injection was undertaken by placing a needleless syringe onto the eye. All studies, except for Epstein and colleagues 2012[42-44], were multi-centre, international studies. Most studies had an extension phase after the primary outcome, but this was not included in the network meta-analysis.

The sufficiently comparable studies were combined into a network analysis based on a common comparator. The network for the proportions of patients gaining more than or equal to 3 lines is shown in Figure 2. This network is the same for the other two outcomes, but without dexamethasone because the trial did not report these outcomes.

Risk of bias of included trials

Risk of bias is shown in Table 3. Included studies were generally of high quality, with all studies being judged to be of low or unclear bias for all criteria. The non-commercially

funded bevacizumab trial had fewer patients and inevitably results had wider confidence intervals.[42-44] In no study does it appear that patients were asked at the end of the trial what arm they thought they had been assigned. It is unclear how many could distinguish injections (intervention arm) from punctureless pressure (sham arm).

Effects of interventions on proportions of patients gaining ≥ 3 lines

Figure 3 displays a forest plot of the risk ratio and 95% credible interval in proportions of patients gaining more than or equal to 3 lines for all the possible pairwise comparisons. In terms of proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of being more effective than a sham and dexamethasone (Figure 4). There was no difference in the proportions of patients gaining more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg.

Effects of interventions on proportions of patients losing ≥ 3 lines

Figure 5 displays forest plot of the risk ratio and 95% credible interval of proportions of patients losing more than or equal to 3 lines for all the possible pairwise comparisons. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision than those treated with sham. There was no difference in the proportions of patients losing more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25 mg and aflibercept 2mg. Figure 6 shows ranking for efficacy in terms of proportions of patients losing ≥ 3 lines.

Effects of interventions on mean change in BCVA

Figure 7 displays a forest plot of the mean changes and 95% credible intervals of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

improvement in BCVA for all the possible pairwise comparisons. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of improvement in BCVA compared to those treated with sham injections. Patients treated with aflibercept 2mg had a higher probability of improvement in BCVA compared with those treated with triamcinolone 4mg (Figure 8). There was no difference in mean change in BCVA from baseline between patients treated with ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2 mg.

For peer review only

Discussion

Statement of principal findings

Our results show no evidence of a difference in effectiveness between aflibercept, ranibizumab and triamcinolone. Bevacizumab was similar to these drugs in terms of letters gained and mean change in BCVA. Dexamethasone was less effective compared to these drugs.

Strengths and limitations

This is the first study providing an indirect comparison of drugs to treat macular oedema secondary to CRVO. A robust search strategy, screening process and data extraction was used, and this analysis drew on a systematic review. The studies included had, in general, a low risk of bias. Safety was not considered in this study but is described in detail elsewhere.[45] Five different drugs were suitable for network meta-analysis. Unpublished data was obtained from one author. [42-44] Bayesian methods were used for the NMA. There was good model fit and convergence within the analysis.

However pre-specified outcomes were not reported in all studies and the sample size varied considerably. For example Epstein 2012, assessing bevacizumab, only included 30 participants in each arm.[42-44] This resulted in wide credible intervals from the network meta-analysis which may lead to a type 1 error especially with regards to the proportions of patients losing more than or equal to 3 lines. The SCORE study compared triamcinolone to observation.[23-36] The NMA assumes a [11] similar effect of sham and observation and this may result in a small degree of bias. Only six months of data was included, and the long term effects are not known. Using a six-month follow-up period may disadvantage dexamethasone because peak effect in the GENEVA trials was seen at 90 days, and by six months, benefits had been largely lost.[6-8]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

As with most network meta-analyses, methodological heterogeneity was present. There were some differences amongst the trials. For example CRUISE[9 10], assessing ranibizumab, did not include as many patients with ischaemic CRVO as the aflibercept trials.[12 13] There were also some small differences in the chronicity of macular oedema and the mean BCVA at baseline.

Meaning of the study: possible explanations and implications for clinicians and policymakers

No head-to-head trials comparing aflibercept, bevacizumab, ranibizumab, triamcinolone or dexamethasone have been published in central retinal vein occlusion. Part of the reason for this is that the Food and Drug Administration require proof of the safety and effectiveness of a drug.[46] The easiest and quickest method for pharmaceutical companies to produce this is through placebo controlled trials. Trials comparing new medications to current best treatment would be considerably more useful to clinicians and patients.

Head-to-head trials comparing some of these drugs are available in other conditions. For example a comparison of ranibizumab and bevacizumab was undertaken in age related macular degeneration in the Comparison of Age-related macular degeneration Treatment Trials (CATT)[47] and alternative treatments to Inhibit VEGF in patients with Age-related choroidal Neovascularisation (IVAN)[48] trials. Both of these trials found no difference in effectiveness between ranibizumab and bevacizumab. Furthermore an indirect comparison of ranibizumab and bevacizumab found no evidence of a difference between these drugs.[49] Thus, it is highly probable that this may also apply in CRVO. The difference seen in our results regarding bevacizumab may be due to the low number of patients included in Epstein 2012.[42-44] In the CATT trial, more patients were

hospitalized in the bevacizumab arm, but the authors did not believe that this was explained by a direct effect of bevacizumab.[47] The 2-year results from the IVAN showed little difference in cardiovascular events, with the number being insignificantly lower with bevacizumab.[50] Ranibizumab and aflibercept were directly compared in two similarly designed trials, VEGF Trap-eye: investigation of Efficacy and safety in Wet age-related macular degeneration (VIEW 1 and 2).[51] Similar efficacy and safety was found in both drugs.

From the included trials it is clear that intraocular steroids are associated with complications, including increased intra-ocular pressure and cataract formation.[6-8 23-36] These are substantial drawbacks for using steroids to treat macular oedema in CRVO. However, many affected patients may be already pseudophakic and, on these, the use of intraocular steroids may be reasonable. Steroids may have a place in the treatment pathway of patients who have failed on anti-VEGF therapy, but this has yet to be tested. The anti-VEFG drugs have a good safety profile and do not cause cataract formation.[9-13 42-44] For this reason are likely to be more favoured by clinicians than steroids.

Aflibercept, compared with ranibizumab and bevacizumab, targets a wider range of cytokines and may have a stronger binding affinity.[52] Initial results suggested that aflibercept would require fewer injections than ranibizumab.[51] Heier and colleagues compared aflibercept and ranibizumab in two similarly designed randomised controlled trials in age related macular degeneration. They found that 2 mg aflibercept administered every eight weeks produced similar effects at 96 weeks to 0.5 mg ranibizumab every four weeks.[51] This was reflected in the FDA Dermatologic and Ophthalmic Drugs Advisory Committee recommendation that aflibercept should be given every two months following three initial monthly doses in age related macular

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

oedema.[53] This may be because aflibercept also appears to last longer in the eye than ranibizumab.[54] Age related macular degeneration is a more aggressive condition than central retinal vein occlusion and so it is unlikely that more frequent dosing would be needed. Therefore aflibercept may be preferred because it would reduce pressure on out-patient clinics. Furthermore there is some evidence from patients with age-related macular degeneration that aflibercept may be effective in patients who have not responded to ranibizumab.[55 56] This may be due to the higher affinity and wider number of cytokines that are targeted. There is no reason to suspect that these effects be any different for the macular oedema caused by central retinal vein occlusion. However we have as yet no evidence as to whether ranibizumab would be effective after aflibercept has failed.

The National Institute of Health and Care Excellence has recommended dexamethasone and ranibizumab,[18 19] and is currently appraising aflibercept. Until these technologies are reviewed together and compared with each other, clinicians may be left with three recommended drugs for macular oedema secondary to central retinal vein occlusion. It should be noted that during the appraisal of ranibizumab the evidence review group found that in the cost-effectiveness analysis dexamethasone was extendedly dominated by ranibizumab (an intervention is judged not be cost-effective because it has an ICER that is greater than that that of a more effective intervention). The committee appraising ranibizumab did not re-consider the previous appraisal decision on dexamethasone.

Our results show that dexamethasone was not as effective as ranibizumab or aflibercept, at six months follow-up and with the dosing regimens in the trials. However these results do not assess quality of life or cost effectiveness. Bevacizumab is likely to prove more cost effective than both aflibercept and ranibizumab because it costs substantially

less.[57] However the National Institute for Health and Care Excellence has not issued guidance on bevacizumab because it does not have a license for use in the eye.

Unanswered questions and future research

Not all patients benefit from the use of anti-VEGF drugs; only about 60% gain 15 or more letters. It is not clear why some patients benefit more than others. Future research should focus on identifying subgroups of patients who are likely to benefit. Only a few of these trials included ischaemic patients, and in these trials only a few patients with ischaemia were included.[11-13] More research assessing the effectiveness of these drugs in severely ischaemic patients is needed.

Head-to-head trials comparing ranibizumab, aflibercept, bevacizumab and triamcinolone are needed. These should include assessment of cost effectiveness. To assist this, a better measure of quality of life is needed for patients with eye conditions. The widely-used EQ5D may not be sensitive enough to measure changes which are important to patients, such as the ability to drive.

In conclusion, we have found no evidence of difference between ranibizumab, bevacizumab, aflibercept and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

We thank Christine Clar, Sian Thomas and Rachel Court for assisting with searches, screening and data extraction for the systematic review which precede this study. We thank the authors of the Epstein 2012 trial for providing addition data.

Copyright statement

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above."

Declaration of competing interests

"All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work."

Contribution statement

NW conceived the idea. All authors contributed to the design of the study. DS and OU undertook the statistical analysis. JF, DS and OU wrote the first draft of the manuscript. All authors redrafted and agreed the final article. JF is the guarantor.

Transparency statement

JF affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

Table 1: Baseline characteristics and results of all included studies

Study	Participants	Intervention / Outcomes
DEXAMETHASONE		
GENEVA 2010[6-8] International Setting: multicentre (167 centres in 24 countries, so a mean of 2.6 patients per centre) Design: 2 identical double-blind, sham-controlled RCTs, phase 3 Follow-up: primary endpoint for the masked trial: 6 months; primary endpoint for the open-label extension: 12 months	N: CRVO – 437 eyes of 437 patients randomised; 94% follow-up at 6 months Participants: adults with visual acuity reduced because of macular oedema due to CRVO or BRVO	1. Dexamethasone 0.7 mg (n=136) Single dose 2. Dexamethasone 0.35 mg (n=154) Single dose 3. Sham (n=147) Single dose - a needleless applicator was placed against the conjunctiva to simulate the placement of study medication. Primary end point: gain of ≥ 15 ETDRS letters; for the open-label extension: safety

TRIAMCINOLONE		
SCORE 2009[23-36] USA Setting: multicentre Design: RCT Follow-up: primary end point 12 months, FU planned up to 36 months	N: 271 eyes of 271 patients randomised; 83% (observation) and 90% (triamcinolone) completed 12 months Participants: centre-involved macular oedema secondary to CRVO	1. Triamcinolone 1 mg (n=92) Every 4 months depending on retreatment regimen (ave 2.2 injections at 12 months) 2. Triamcinolone 4 mg (n=91) Every 4 months depending on retreatment regimen (ave 2.0 injections at 12 months) (The form of triamcinolone used was Trivaris, no longer available. It was made by the manufacturer of Ozurdex (Allergan)) 3. Observation (n=88) Primary end point: gain of ≥15 ETDRS letters
AFLIBERCEPT		
COPERNICUS 2012[12 13] International Setting: multicentre, 70 sites in North and South America, India and Israel. Mean 2.7 patients per centre. Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU 2 years	N: 189 eyes of 189 patients randomised; 95.7% (aflibercept) and 81.1% (sham) completed 24 weeks; 93% (aflibercept) and 77% (sham) completed 52 weeks Participants: adult patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=114) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=73) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥15 ETDRS letters

GALILEO 2012[11] International Setting: multicentre, 10 countries in Europe and Asia; 63 centres in total Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU up to 12 months, planned up to 76 weeks	N: 177 eyes of 177 patients randomised; 90.6% (aflibercept) and 78.9% (sham) completed 24 weeks Participants: treatment-naïve patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=103) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=71) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥ 15 ETDRS letters
--	--	--

RANIBIZUMAB		
CRUISE 2010 ^[9 10] USA Setting: multicentre Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 6 months, FU up to 12 months	N: 392 eyes of 392 patients randomised; 97.7% (ranibizumab 0.3 mg), 91.5% (ranibizumab 0.5 mg), and 88.5% (sham) completed 6 months Participants: patients with foveal centre-involved macular oedema secondary to CRVO diagnosed within 12 months	1. Ranibizumab 0.3 mg (n=132) Every 4 weeks for 6 months (ave number not available) 2. Ranibizumab 0.5 mg (n=130) Every 4 weeks for 6 months (ave number not available) 3. Sham (n=130) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to the injection site) Primary end point: mean change from baseline BCVA
BEVACIZUMAB		
Epstein 2012 ^[42-44] Sweden Setting: Single centre; St. Eriks Eye Hospital Stockholm Design: sham-injection controlled, double masked RCT Follow-up: primary end-point 6 months; open label extension up to 12 months	N: 60 eyes of 60 patients randomised; 93% completed open label extension Participants: patients with CRVO of ≤6 months	1. Bevacizumab 1.25 mg (n=30) Every 6 weeks for 6 months (ave number not available) 2. Sham (n=30) Every 6 weeks for 6 months (ave number not available) (syringe without needle pressed to the globe) Primary end point: gain of ≥15 ETDRS letters

FU= follow-up, RCT = randomised controlled trial, N = number, CRVO = central retinal vein occlusion, ETDRS = Early Treatment Diabetic Retinopathy Study, BRVO = branch retinal vein occlusion

Table 2: Baseline characteristics and results of included trials

	COPERNICUS[12 13]	GALILEO[11]	CRUISE[9 10]	GENEVA[6- 8]	Epstein et al (2012)[42- 44]	SCORE[23-36]
BASELINE SIMILARITIES						
Number (%) of patients						
	Aflib 2 mg: 114 Sham: 73	Aflib 2 mg: 103 Sham: 68	Rani 0.5 mg: 130 Sham: 130	Dexa0.7 mg: 136 Sham: 147	Beva 1.25 mg: 30 Sham: 30	Triam 4 mg: 91 Obser: 88
Age (years)						
	Aflib 2 mg: 65.5 SD13.6 Sham: 67.5 SD14.3	Aflib 2 mg: 59.9 SD12.4 Sham: 63.8 SD13.3	Rani 0.5 mg: 67.6 SD12.4 Sham: 65.4 SD13.1	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 70.6 SD 12.6 Sham: 70.4 SD 10.4	Triam 4 mg: 67.5 SD 12.0 Obser: 69.2 SD 12.8
BCVA at baseline (SD)						
	Aflib 2 mg: 50.7 SD13.90 Sham: 48.9 SD14.42	Aflib 2 mg: 53.6 SD15.8 Sham: 50.9 SD15.4	Rani 0.5 mg: 48.1 SD14.6 Sham: 49.2 SD14.7	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 44.4 SD 15.3 Sham: 43.6 SD 16.0	Triam 4 mg: 51.0 SD 14.4 Obser: 52.1 SD 13.1
Duration of MO from diagnosis to screening						
	Aflib 2 mg: 2.73 SD3.09(in months) Sham: 1.88 SD2.19 (in months)	Aflib 2 mg: 50.9 SD15.4)(in days) Sham: 87.6 SD79.1 (in days)	Rani 0.5 mg: - Sham: -	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: NR Sham: NR	Triam 4 mg: 4.2 SD 3.6 (in months) Obser: 4.2 SD 3.1 (in months)
RESULTS						
Number (%) of patients gaining ≥15 letters improvement from baseline to 6 months						
	Aflib 2 mg: 64 (56.1)	Aflib 2 mg: 62 (60.2)	Rani 0.5 mg: 62 (47.7)	Dexa 0.7 mg: 25 (18)	Beva 1.25 mg: 18 (60%)	Triam 4 mg: 18 (19.5%) (avg of 4 and 8 mths)

	Sham: 9 (12.3)	Sham: 15 (22.1)	Sham: 22 (16.9)	Sham: 18 (12)	Sham: 6 (20%)	Obser: 3 (4%) (avg of 4 and 8 mths)
Number (%) of patients losing ≥ 15 letters of BCVA from baseline to 6 months						
	Aflib 2 mg: 2 (1.8)	Aflib 2 mg: 8 (7.8)	Rani 0.5 mg: 2 (1.5)	Dexa 0.7 mg: NR	Beva 1.25 mg: 2 (6.7%)	Triam 4 mg: 19 (20.5%) (avg of 4 and 8 mths)
	Sham: 20 (27.4)	Sham: 15 (22.1)	Sham: 20 (15.4)	Sham: NR	Sham: 7 (23.3%)	Obser: 31 (35.5%) (avg of 4 and 8 mths)
Mean change (SD) from baseline in BCVA						
	Aflib 2 mg: 17.3 (12.8)	Aflib 2 mg: 18.0 (12.2)	Rani 0.5 mg: 14.9 (13.2)	Dexa 0.7 mg: 0.1 (NR)	Beva 1.25 mg: 14.1 SD 18.7	Triam 4 mg: -0.15 SD20.67 (n=85) (weight mean and SD of 4 and 8 months)
	Sham: -4 (18)	Sham: 3.3 (14.1)	Sham: 0.8 (16.2)	Sham: -1.8 (NR)	Sham: -2.0 SD 20.5	Obser: -9.66 SD18.04 (n=75) (weighted mean and SD of 4 and 8 months)

NR = not reported, Aflib = aflibercept, Rani = ranibizumab, Dexa = dexamethasone, Triam = triamcinolone, Obser = observation, SD = standard deviation, avg = average

Table 3: Risk of bias

Study (author and year)	Adequate sequence generation	Allocation concealment	Masking	Incomplete outcome data addressed	Free of selective reporting	Free of other bias (e.g. similarity at baseline, power assessment)	Funder
GENEVA 2010[6-8]	Low	Low	Partial: patients and assessors of efficacy variables	Low: ITT analysis, 94% FU at 6 months	Low	Power: 81% power to detect difference in primary outcome with n=495 for each trial Similarity at baseline: yes	Allergan Inc.
SCORE 2009[23-36]	Low	Unclear	Partial (physicians and patients masked to dose but not triamcinolone versus observation)	Low: ITT analysis, 83 to 90% FU at 12 months	Low	Power: 80% power to detect difference in primary outcome with n=486 (but only 271 randomised) Similarity at baseline: yes	National Eye Institute grants, Allergan
COPERNICUS 2012[12 13]	Low	Unclear	Low: double-blind	Low: ITT analysis, 89.9% assessed at primary end point	Low	Power: 90% power to detect difference in primary outcome with n=165 Similarity at baseline: yes	Bayer HealthCare, Regeneron Pharmaceuticals
GALILEO 2012[11]	Unclear	Unclear	Low: double-blind	Low: ITT analysis, 86% assessed at primary end point	Low	Power: 90% power to detect difference in primary outcome with n=150 Similarity at baseline: yes	Bayer HealthCare, Regeneron Pharmaceuticals

CRUISE 2010[9 10]	Low	Unclear	Low: patients and evaluating examiners, injecting physicians masked to dose	Low: ITT analysis, 88.5 to 97.7% completed 6 months	Low	<i>Power: not reported</i> <i>Similarity at baseline: yes</i>	Genentech Inc.
Epstein 2012[42-44]	Unclear	Low	Low: patients, outcome assessors	Low: ITT analysis; missing data for 2 patients (primary endpoint)	Low	<i>Power: 80% power to detect difference in primary outcome with n=24 per group</i> <i>Similarity at baseline: yes</i>	Unclear; authors are consultants for Allergan, Novartis, Alcon, Bayer

ITT= intention to treat, FU = follow-up

Figure 1: study selection flow diagram




Figure 2: Network of randomized controlled trials comparing different treatments for proportions of gaining 3 or more lines of vision




Figure 3: Proportions of patients gaining 3 lines or more from baseline to six months




Figure 4: Rankogram for gaining ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions





Figure 5: Proportions of patients losing 3 lines or more from baseline to six months



Confidential review only

Figure 6: Rankogram for losing ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions




Figure 7: Mean BCVA change from baseline to 6 months.





Figure 8: Rankogram for mean change in BCVA - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions



Appendix: MEDLINE search strategy

Ovid MEDLINE(R) 1946 to March Week 1 2013, searched on 20 March 2013

- 1 CRVO.mp.
- 2 Retinal Vein Occlusion/
- 3 retinal vein occlusion.mp.
- 4 retinal vein obstruction.mp.
- 5 retinal venous occlusion.mp.
- 6 retinal venous obstruction.mp.
- 7 retina*.mp.
- 8 ("central vein occlusion" or "central vein obstruction" or "central venous occlusion" or "central venous obstruction").mp.
- 9 7 and 8
- 10 1 or 2 or 3 or 4 or 5 or 6 or 9
- 11 randomized controlled trial.pt.
- 12 (random* or "controlled trial*" or "clinical trial*" or rct).tw.
- 13 11 or 12
- 14 (metaanalys* or "meta analys*" or "meta-analys*").tw.
- 15 "systematic review*".tw.
- 16 meta analysis.pt.
- 17 14 or 15 or 16
- 18 10 and 13
- 19 10 and 17
- 20 18 or 19
- 21 limit 20 to yr="2005 -Current"

References

1. Deramo VA, Cox TA, Syed AB, Lee PP, Fekrat S. Vision-related quality of life in people with central retinal vein occlusion using the 25-item National Eye Institute Visual Function Questionnaire. Arch.Ophthalmol. 2003;**121**(9):1297-302 doi: 10.1001/archophth.121.9.1297 [doi];121/9/1297 [pii][published Online First: Epub Date]].
2. Klein R, Moss SE, Meuer SM, Klein BE. The 15-year cumulative incidence of retinal vein occlusion: the Beaver Dam Eye Study. Archives of ophthalmology 2008;**126**(4):513-8 doi: 10.1001/archophth.126.4.513[published Online First: Epub Date]].
3. Hayreh SS, Podhajsky PA, Zimmerman MB. Natural history of visual outcome in central retinal vein occlusion. Ophthalmology 2011;**118**(1):119-33 doi: S0161-6420(10)00447-1 [pii];10.1016/j.ophtha.2010.04.019 [doi][published Online First: Epub Date]].
4. McIntosh RL, Rogers SL, Lim L, et al. Natural history of central retinal vein occlusion: an evidence-based systematic review. Ophthalmology 2010;**117**(6):1113-23 doi: S0161-6420(10)00133-8 [pii];10.1016/j.ophtha.2010.01.060 [doi][published Online First: Epub Date]].
5. The Royal College of Ophthalmologists. Interim Guidelines for Management of Retinal Vein Occlusion. Secondary Interim Guidelines for Management of Retinal Vein Occlusion 2010. http://www.rcophth.ac.uk/core/core_picker/download.asp?id=728&filetitle=Interim+Guidelines+for+Management+of+Retinal+Vein+Occlusion+2010.

6. Haller JA, Bandello F, Belfort R, Jr., et al. Randomized, sham-controlled trial of dexamethasone intravitreal implant in patients with macular edema due to retinal vein occlusion. *Ophthalmology* 2010;**117**(6):1134-46

7. Haller JA, Bandello F, Belfort R, Jr., et al. Dexamethasone intravitreal implant in patients with macular edema related to branch or central retinal vein occlusion twelve-month study results. *Ophthalmology* 2011;**118**(12):2453-60

8. Yeh WS, Haller JA, Lanzetta P, et al. Effect of the duration of macular edema on clinical outcomes in retinal vein occlusion treated with dexamethasone intravitreal implant. *Ophthalmology* 2012;**119**(6):1190-98

9. Brown DM, Campochiaro PA, Singh RP, et al. Ranibizumab for macular edema following central retinal vein occlusion: six-month primary end point results of a phase III study. *Ophthalmology* 2010;**117**(6):1124-33

10. Campochiaro PA, Brown DM, Awh CC, et al. Sustained benefits from ranibizumab for macular edema following central retinal vein occlusion: twelve-month outcomes of a phase III study. *Ophthalmology* 2011;**118**(10):2041-49

11. Holz FG, Roeder J, Ogura Y, et al. VEGF Trap-Eye for macular oedema secondary to central retinal vein occlusion: 6-month results of the phase III GALILEO study. *British Journal of Ophthalmology* 2013;**97**(3):278-84

12. Boyer D, Heier J, Brown DM, et al. Vascular endothelial growth factor Trap-Eye for macular edema secondary to central retinal vein occlusion: six-month results of the phase 3 COPENICUS study. *Ophthalmology* 2012;**119**(5):1024-32

13. Brown DM, Heier JS, Clark WL, et al. Intravitreal Aflibercept Injection for Macular Edema Secondary to Central Retinal Vein Occlusion: 1-Year Results From the Phase 3 COPENICUS Study. *American Journal of Ophthalmology* 2013;**155**(3):429-37
14. Michels S, Rosenfeld PJ, Puliafito CA, Marcus EN, Venkatraman AS. Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration twelve-week results of an uncontrolled open-label clinical study. *Ophthalmology* 2005;**112**(6):1035-47 doi: 10.1016/j.ophtha.2005.02.007[published Online First: Epub Date]].
15. Arevalo JF, Fromow-Guerra J, Quiroz-Mercado H, et al. Primary intravitreal bevacizumab (Avastin) for diabetic macular edema: results from the Pan-American Collaborative Retina Study Group at 6-month follow-up. *Ophthalmology* 2007;**114**(4):743-50 doi: 10.1016/j.ophtha.2006.12.028[published Online First: Epub Date]].
16. Rabena MD, Pieramici DJ, Castellarin AA, Nasir MA, Avery RL. Intravitreal bevacizumab (Avastin) in the treatment of macular edema secondary to branch retinal vein occlusion. *Retina (Philadelphia, Pa.)* 2007;**27**(4):419-25 doi: 10.1097/IAE.0b013e318030e77e[published Online First: Epub Date]].
17. Algvere PV, Epstein D, von Wendt G, Seregard S, Kvanta A. Intravitreal bevacizumab in central retinal vein occlusion: 18-month results of a prospective clinical trial. *European journal of ophthalmology* 2011;**21**(6):789-95 doi: 10.5301/ejo.2011.6522[published Online First: Epub Date]].

18. National Institute for Health and Care Excellence. Ranibizumab for treating visual impairment caused by macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 283. 2013.
<http://guidance.nice.org.uk/TA283/Guidance/pdf/English>.

19. National Institute for Health and Clinical Excellence. Dexamethasone intravitreal implant for the treatment of macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 229. 2011.
<http://guidance.nice.org.uk/TA229/Guidance/pdf/English>.

20. National Institute for Health and Care Excellence. Aflibercept for treating visual impairment caused by macular oedema secondary to central retinal vein occlusion. NICE technology appraisal guidance 305. 2014.
<http://guidance.nice.org.uk/TA305>.

21. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. BMJ (Clinical research ed.) 2011;**343**:d5928 doi: 10.1136/bmj.d5928[published Online First: Epub Date]].

22. Hozo S, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. BMC Medical Research Methodology 2005;**5**(1):13

23. Bhavsar AR, Ip MS, Glassman AR, Groups DatSS. The risk of endophthalmitis following intravitreal triamcinolone injection in the DRCRnet and SCORE clinical trials. American Journal of Ophthalmology 2007;**144**(3):454-56

24. Blodi BA, Domalpally A, Scott IU, et al. Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study system for evaluation of

- stereoscopic color fundus photographs and fluorescein angiograms:
SCORE Study Report 9. Archives of ophthalmology 2010;**128**(9):1140-45
25. Chan CK, Ip MS, VanVeldhuisen PC, et al. SCORE Study report #11: incidences
of neovascular events in eyes with retinal vein occlusion. Ophthalmology
2011;**118**(7):1364-72
26. Ip M, Oden N, VanVeldhuisen P, Scott I, Blodi B. The Standard Care vs.
Corticosteroid for Retinal Vein Occlusion Study: Design and Baseline
Characteristics. American Academy of Ophthalmology 2008:260
27. Ip MS, Oden NL, Scott IU, et al. SCORE Study report 3: study design and
baseline characteristics. Ophthalmology 2009;**116**(9):1770-77
28. Ip MS, Scott IU, VanVeldhuisen PC, et al. A randomized trial comparing the
efficacy and safety of intravitreal triamcinolone with observation to treat
vision loss associated with macular edema secondary to central retinal
vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein
Occlusion (SCORE) study report 5. Archives of ophthalmology
2009;**127**(9):1101-14
29. Myers D, Blodi B, Ip M, Scott I, Warren K. Reading Center Evaluation of OCT
Images From Patients Enrolled in the Standard Care vs. Corticosteroid for
Retinal Vein Occlusion (SCORE) Study. Iovs. 2006;**47**:ARVO
30. Oden NL, Veldhuisen PC, Scott IU, Ip MS, Blodi BA. Temporal Variability of
OCT in Retinal Vein Occlusion Participants in the SCORE Study. Iovs.
2007;**48**:ARVO
31. Scott IU, Blodi BA, Ip MS, et al. SCORE Study Report 2: Interobserver
agreement between investigator and reading center classification of
retinal vein occlusion type. Ophthalmology 2009;**116**(4):756-61

32. Scott IU, Oden NL, VanVeldhuisen PC, et al. SCORE Study Report 7: incidence of intravitreal silicone oil droplets associated with staked-on vs luer cone syringe design. American Journal of Ophthalmology 2009;**148**(5):725-32
33. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline predictors of visual acuity and retinal thickness outcomes in patients with retinal vein occlusion: Standard Care Versus Corticosteroid for Retinal Vein Occlusion Study report 10. Ophthalmology 2011;**118**(2):345-52
34. Scott IU, VanVeldhuisen PC, Oden NL, et al. SCORE Study report 1: baseline associations between central retinal thickness and visual acuity in patients with retinal vein occlusion. Ophthalmology 2009;**116**(3):504-12
35. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline Characteristics and Response to Treatment of Participants With Hemiretinal Compared With Branch Retinal or Central Retinal Vein Occlusion in the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study: SCORE Study Report 14. Archives of ophthalmology 2012;**130**(12):1517-24
36. Warren K, Blodi BA, Oden N, Veldhuisen P, Scott IU, Ip M. Reading Center Evaluation of Baseline Retinal Images in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. Iovs. 2008:ARVO
37. Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. BMJ (Clinical research ed.) 2005;**331**(7521):897-900 doi: 10.1136/bmj.331.7521.897[published Online First: Epub Date]].
38. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. Statistics in medicine 2004;**23**(20):3105-24 doi: 10.1002/sim.1875[published Online First: Epub Date]].

39. Spiegelhalter D, Thomas A, Best N, Lunn D. *WinBUGS User Manual: Version 1.4*. Cambridge, Mass: MRC Biostatistics Unit, 2003.
40. Brooks SP, Gelman A. General Methods for Monitoring Convergence of Iterative Simulations. *Journal of Computational and Graphical Statistics* 1998;**7**(4):434-55 doi: 10.1080/10618600.1998.10474787[published Online First: Epub Date]].
41. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;**64**(2):163-71 doi: 10.1016/j.jclinepi.2010.03.016[published Online First: Epub Date]].
42. Epstein D, Algvere P, Von WG, Seregard S, Kvanta A. Long-term benefit from bevacizumab for macular edema in central retinal vein occlusion: 12-month results of a prospective study. *Acta Ophthalmologica* 2012;**90**:48 doi: 10.1111/j.1755-3768.2012.02549.x[published Online First: Epub Date]].
43. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Benefit from bevacizumab for macular edema in central retinal vein occlusion: twelve-month results of a prospective, randomized study. *Ophthalmology* 2012;**119**(12):2587-91
44. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Bevacizumab for macular edema in central retinal vein occlusion: a prospective, randomized, double-masked clinical study. *Ophthalmology* 2012;**119**(6):1184-89

45. Ford JA, Clar C, Lois N, et al. Treatments for macular oedema following central retinal vein occlusion: systematic review. BMJ open 2014;**4**(2):e004120 doi: 10.1136/bmjopen-2013-004120[published Online First: Epub Date]].
46. FDA. Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators. Secondary Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators 2011. <http://www.fda.gov/RegulatoryInformation/Guidances/ucm126501.htm>.
47. Martin DF, Maguire MG, Fine SL, et al. Ranibizumab and bevacizumab for treatment of neovascular age-related macular degeneration: two-year results. Ophthalmology 2012;**119**(7):1388-98 doi: 10.1016/j.ophtha.2012.03.053[published Online First: Epub Date]].
48. Chakravarthy U, Harding SP, Rogers CA, et al. Ranibizumab versus bevacizumab to treat neovascular age-related macular degeneration: one-year findings from the IVAN randomized trial. Ophthalmology 2012;**119**(7):1399-411 doi: S0161-6420(12)00358-2 [pii];10.1016/j.ophtha.2012.04.015 [doi][published Online First: Epub Date]].
49. Ford JA, Elders A, Shyangdan D, Royle P, Waugh N. The relative clinical effectiveness of ranibizumab and bevacizumab in diabetic macular oedema: an indirect comparison in a systematic review. BMJ (Clinical research ed.) 2012;**345**:e5182

50. Chakravarthy U, Harding SP, Rogers CA, et al. Alternative treatments to inhibit VEGF in age-related choroidal neovascularisation: 2-year findings of the IVAN randomised controlled trial. *Lancet* 2013 doi: 10.1016/s0140-6736(13)61501-9[published Online First: Epub Date]].
51. Heier JS, Brown DM, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology* 2012;**119**(12):2537-48 doi: 10.1016/j.ophtha.2012.09.006[published Online First: Epub Date]].
52. Papadopoulos N, Martin J, Ruan Q, et al. Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* 2012;**15**(2):171-85 doi: 10.1007/s10456-011-9249-6[published Online First: Epub Date]].
53. FDA. VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee. Secondary VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee 2011. <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/DermatologicandOphthalmicDrugsAdvisoryCommittee/UCM259143.pdf>.
54. Stewart MW, Rosenfeld PJ. Predicted biological activity of intravitreal VEGF Trap. *The British journal of ophthalmology* 2008;**92**(5):667-8 doi: 10.1136/bjo.2007.134874[published Online First: Epub Date]].
55. Bakall B, Folk JC, Boldt HC, et al. Aflibercept therapy for exudative age-related macular degeneration resistant to bevacizumab and ranibizumab. *Am J Ophthalmol* 2013;**156**(1):15-22.e1 doi: 10.1016/j.ajo.2013.02.017[published Online First: Epub Date]].

56. Cho H, Shah CP, Weber M, Heier JS. Aflibercept for exudative AMD with persistent fluid on ranibizumab and/or bevacizumab. The British journal of ophthalmology 2013;**97**(8):1032-5 doi: 10.1136/bjophthalmol-2013-303344[published Online First: Epub Date]].

57. Raftery J, Clegg A, Jones J, Tan SC, Lotery A. Ranibizumab (Lucentis) versus bevacizumab (Avastin): modelling cost effectiveness. The British journal of ophthalmology 2007;**91**(9):1244-6 doi: 10.1136/bjo.2007.116616[published Online First: Epub Date]].

Confidential peer review only



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	3-4
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5-6
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5-6
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	1
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	7
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	41
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	7-8
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7-8
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	7-8
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	7-8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	8-9
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2 for each meta-analysis).	8-9



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	8-9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	8-9
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11+27
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	20-22
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	25-26
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	23-24
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	29-31
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	11-13
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	NA
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	14-18
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	14-18
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	14-18
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	1

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

John A Ford, Deepson Shyangdan, Olalekan A. Uthman, Noemi Lois, Norman Waugh

John A Ford, Academic Clinical Fellow in Public Health, Department of Population Health and Primary Care, University of East Anglia
Deepson Shyangdan, Research Fellow, Warwick Evidence, University of Warwick
Olalekan Uthman, Assistant Professor in Applied Research, University of Warwick
Noemi Lois, Professor of Ophthalmology, Centre for Vision and Vascular Science, Queen's University
Norman Waugh, Professor of Health Technology Assessment, University of Warwick

Corresponding author
Dr. John Ford
Norwich Medical School
University of East Anglia
Chancellor Drive
Norwich
NR4 7TJ
John.ford@uea.ac.uk
Tel: 01603 591269

Protocol – no published protocol exists for this study.

Ethics approval – not required

Funding – no external funding required

Data sharing: no additional data available

Word count: 3,331 words

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

What is already known on this subject

Anti-VEGF drugs (ranibizumab, bevacizumab and aflibercept) and corticosteroids (dexamethasone and triamcinolone), given intravitreally, have all been shown to be effective compared to placebo for the treatment of macular oedema secondary to central retinal vein occlusion.

There are no head-to-head trials.

What this study adds

There was no evidence of a difference in the effectiveness of aflibercept, ranibizumab, bevacizumab and triamcinolone for improving vision.

Clinicians may prefer aflibercept because steroids are associated with cataract formation and ranibizumab might require more frequent injections.

For peer review only

BMJ Open: first published as 10.1136/bmjopen-2014-005292 on 23 July 2014. Downloaded from <http://bmjopen.bmj.com/> on June 9, 2025 at Agence Bibliographique de l'Enseignement Supérieur (ABES).
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Abstract

Objective: To indirectly compare aflibercept, bevacizumab, dexamethasone, ranibizumab and triamcinolone for treatment of macular oedema secondary to central retinal vein occlusion using a network meta-analysis.

Design: Network meta-analysis

Data sources: The following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE; CDSR, DARE, HTA, NHSEED, CENTRAL; Science Citation Index and Conference Proceedings Citation Index-Science

Eligibility criteria for selecting studies: Only randomized controlled trials assessing patients with macular oedema secondary to central retinal vein occlusion were included. Studies had to report either proportions of patients gaining more than or equal to 3 lines, losing more than or equal to 3 lines, or mean change in best corrected visual acuity. Two authors screened titles and abstracts, extracted data and undertook risk of bias assessment. Bayesian network meta-analysis was used to compare the different interventions.

Results: Seven studies, assessing five drugs, were judged to be sufficiently comparable for inclusion in the NMA. For the proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2mg had a higher probability of being more effective than sham and dexamethasone. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision compared to those treated with sham. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1.25mg and aflibercept 2mg had a higher probability of improvement in mean best correct visual acuity compared to those treated with sham injections.

Conclusions: We found no evidence of differences between ranibizumab, aflibercept, bevacizumab and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections

Systematic review registration – Not registered

Strengths and limitations of this study

- Important topic area, with significant policy implications
- Robust method used to identify studies
- Network meta-analysis are based on a number of assumptions
- Network meta-analysis is the best method to compare interventions in the absence of head to head trials

Introduction

Central retinal vein occlusion (CRVO) dramatically reduces an individual's functioning and quality of life.[1] It is estimated that the 15 year cumulative incidence of central retinal vein occlusion is 0.5%.[2] Visual loss is caused by thrombosis of the central retinal vein which leads to a rise in venous pressure and an increase in vascular endothelial growth factor (VEGF), consequently causing an increase in vascular permeability. Macular oedema subsequently ensues with varying degrees of ischaemia and neo-vascularisation. Although CRVO is generally classified as ischaemic or non-ischaemic, ischaemia should be regarded as a spectrum.[3] Cases with ischaemia carry a considerably worse prognosis as in around a third of them, neovascular glaucoma may develop; the most devastating complication of CRVO.[4]

CRVO is more common in older people with risk factors such as diabetes, hypertension or hyperlipidaemia, but can occur in young people with inflammatory disorders. Hayreh and colleagues in a 27-year cohort study found that only 13% of people with CRVO were under 45 years of age.[3] In 95% of cases CRVO affects only one eye.[3] However visual loss in this already co-morbid patient group significantly compounds their already impaired functioning and quality of life. Patients can lose confidence, struggle with daily activities and become increasingly dependent on friends and family.[1]

For many years, laser photocoagulation was the only effective therapeutic strategy that could be used in the management of patients with CRVO. It was only useful for reducing the risk of neovascular glaucoma, but not effective for the treatment of macular oedema in CRVO.[5] Over the past decade a number of drugs to treat macular oedema have been introduced, including the steroids, triamcinolone and dexamethasone, and the anti-VEGFs, ranibizumab, bevacizumab, pegaptanib and aflibercept. Dexamethasone,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ranibizumab and aflibercept have been assessed in large commercially funded trials.[6-13] Bevacizumab was originally developed as an anti-cancer drug and has been found to be effective in treating macular oedema secondary to age-related macular degeneration,[14] diabetic macular oedema, [15] branch retinal vein occlusion[16] and central retinal vein occlusion.[17] Like triamcinolone, bevacizumab is used off licence in the eye. Ranibizumab is a derived from the same parent molecule of the bevacizumab monoclonal antibody and was developed and commercially marketed specifically for use in the eye.

In the United Kingdom, the National Institute of Health and Care Excellence (NICE) has recommended the use of dexamethasone, ~~and~~ ranibizumab and aflibercept for the treatment of macular oedema secondary to CRVO in separate appraisals[18-20] ~~and it is currently evaluating aflibercept. If aflibercept is also endorsed and with no head-to-head trials comparing these drugs, Therefore~~ clinicians ~~will be in the position of~~ having three NICE-recommended treatments for CRVO without head-to-head trials or clear guidance on which one may be best for their patients. On this basis, the aim of this study was to indirectly compare in a network meta-analysis the clinical effectiveness of aflibercept, ranibizumab, bevacizumab, dexamethasone and triamcinolone for the treatment of macular oedema secondary to CRVO.

Methods

Information sources and search strategy

To identify suitable studies, initially for a systematic review of treatment of macular oedema after CRVO (submitted for publication) the following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE (all via OVID); CDSR, DARE, HTA, NHSEED, CENTRAL (all via The Cochrane Library); Science Citation Index and Conference Proceedings Citation Index-Science (via Web of Knowledge). The MEDLINE search strategy is shown in appendix 1. This search strategy was modified for other databases. In addition to the bibliographic database searching, supplementary searches were undertaken to look for recent and unpublished studies in the WHO International Clinical Trials Registry Platform and ophthalmology conference websites (American Academy of Ophthalmology, Association for Research in Vision and Ophthalmology from 2010 to 2012).

Study selection

Only randomised controlled trials which included patients with macular oedema secondary to central retinal vein occlusion were included. It was acceptable for a study to include both branch retinal vein occlusion and central retinal vein occlusion provided that the central retinal vein occlusion group was reported separately. The following drugs were included: dexamethasone, triamcinolone, ranibizumab, bevacizumab and aflibercept. Pegaptanib was not included because it is not used routinely in clinical practice. Only doses which are used in clinical practice were included. Studies had to report at least one of the following outcomes: proportions of patients gaining more than or equal to 3 lines from baseline to six months, proportions of patients losing more than or equal to 3 lines from baseline to six months and mean change in best corrected visual acuity (BCVA) from baseline to six months

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Risk of bias assessment

The Cochrane Collaboration’s tool for assessing risk of bias was used.[21] The trials were graded (unclear, high or low risk of bias) based on: (i) sequence generation, (ii) allocation concealment, (iii) blinding of outcome assessor, (iv) incomplete outcome data, and (v) selective outcome reporting.

Study selection and data abstraction

Two authors independently assessed the eligibility and methodological quality of the studies identified during the literature search. Two authors extracted and compared the data. For each study identified that met the selection criteria, details on study design, study population characteristics, intervention, outcome measures, and study quality were extracted. Discrepancies were resolved by consensus through discussion. Studies were assessed for comparability based on the populations included, trial arms, outcome measures and duration of follow-up. Common comparators were identified from the trials and a network diagram was created.

Summary measures

The primary measures of treatment effects were relative risk (RR) for the proportions of patients gaining more than or equal to 3 lines of vision, proportions of patients losing more than or equal to 3 lines of vision and weighted mean difference (WMD) for mean change BCVA. We used the following methods to calculate standard deviations, when incompletely reported: (1) contact with the corresponding author; or (2) estimation of the standard deviation on the basis of the sample size, median, and range as suggested by Hozo and colleagues[22] or on the basis of the sample size and P value.

In one trial (SCORE),[23-36] six month data was not available because patients were followed up every four months. For the dichotomous outcomes i.e. proportions of patients gaining and losing ≥ 3 lines, we averaged four and eight month data to get the six months follow-up data. For the third outcome i.e. mean change BCVA, again data from two time-points were used. Weighted mean and SDs for each treatment arm was calculated using mean and SDs of two time-points.

Data synthesis and model implementation

Bayesian network meta-analysis [37 38] (NMA) was used to compare the different interventions. Network meta-analysis is a generalization of meta-analysis methods because they allow comparisons of agents not addressed within individual primary trials. Bayesian statistical inference provides probability distributions for treatment effect parameters (RR and WMD), with 95% credible intervals (95% CrI), rather than 95% confidence intervals (95% CI). A 95% credible interval can be interpreted as there being a 95% probability that the parameter takes a value in the specified range.[37 38]

All analyses were conducted using a Bayesian Markov Chain Monte Carlo (MCMC) method and fitted in the freely available Bayesian software, WinBUGS 1.4.3.[39] Two Markov chains were run simultaneously using different initial values. Convergence to a stable solution was checked by viewing plots of the sampled simulations and using the Brooks-Gelman-Rubin diagnostic tool.[40] Convergence was found to be adequate after running 20 000 samples for both chains. These samples were then discarded and a further 70 000 sampled simulation was then run, on which the results were based. [We also calculated the probability of treatment being the most effective \(first best\), the second best, the third best, and so on, and presented the results graphically with rankograms. \(Salanti\)](#)[41]:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Like standard meta-analysis comparison, a NMA can be either a fixed- or a random-effect models. We used the Bayesian Deviation Information Criterion (DIC) to compare fixed and random effect models. The most appropriate NMA model can be identified as the one with the lowest DIC. The DIC measures the fit of the model while penalizing it for the number of effective parameters. The fixed - effect model was chosen because of the small number of trials available for each comparison and difficulty in estimating between studies variance if random-effect model was implemented and the difference in DIC is less than 5.

For peer review only

Results

Study selection and characteristics

The literature search identified 945 articles, as shown in Figure 1. Seven studies were judged to be sufficiently comparable to be included in the network meta-analysis. Tables 1 and 2 present the characteristics and results of the included trials. Two studies [11-13] compared aflibercept 2 mg against sham; two identical studies [6-8] compared dexamethasone 0.7 mg (Ozurdex) against sham; one study [9 10] compared ranibizumab 0.5 mg against sham; one study [42-44] compared bevacizumab 1.25 mg against sham, and finally one study [23-36] compared triamcinolone 4 mg against observation. Sham or observation were used as the common comparator. The number of included participants varied from 60 [42-44] to 437 [6-8]. Most studies required patients to be treatment naive and have macular oedema with retinal thickness measuring at least 250 or 300 μm on optical coherence tomography. Sham injection was undertaken by placing a needleless syringe onto the eye. All studies, except for Epstein and colleagues 2012[42-44], were multi-centre, international studies. Most studies had an extension phase after the primary outcome, but this was not included in the network meta-analysis.

The sufficiently comparable studies were combined into a network analysis based on a common comparator. The network for the proportions of patients gaining more than or equal to 3 lines is shown in Figure 2. This network is the same for the other two outcomes, but without dexamethasone because the trial did not report these outcomes.

Risk of bias of included trials

Risk of bias is shown in Table 3. Included studies were generally of high quality, with all studies being judged to be of low or unclear bias for all criteria. The non-commercially

funded bevacizumab trial had fewer patients and inevitably results had wider confidence intervals.[42-44] In no study does it appear that patients were asked at the end of the trial what arm they thought they had been assigned. It is unclear how many could distinguish injections (intervention arm) from punctureless pressure (sham arm).

Effects of interventions on proportions of patients gaining ≥ 3 lines

Figure 3 displays a forest plot of the risk ratio and 95% credible interval in proportions of patients gaining more than or equal to 3 lines for all the possible pairwise comparisons. In terms of proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of being more effective than a sham and dexamethasone (eFigure 41). There was no difference in the proportions of patients gaining more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg.

Effects of interventions on proportions of patients losing ≥ 3 lines

Figure 54 displays forest plot of the risk ratio and 95% credible interval of proportions of patients losing more than or equal to 3 lines for all the possible pairwise comparisons. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision than those treated with sham. There was no difference in the proportions of patients losing more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25 mg and aflibercept 2mg. eFigure 62 shows ranking for efficacy in terms of proportions of patients losing ≥ 3 lines.

Effects of interventions on mean change in BCVA

Figure 75 displays a forest plot of the mean changes and 95% credible intervals of

improvement in BCVA for all the possible pairwise comparisons. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of improvement in BCVA compared to those treated with sham injections. Patients treated with aflibercept 2mg had a higher probability of improvement in BCVA compared with those treated with triamcinolone 4mg (eFigure 83). There was no difference in mean change in BCVA from baseline between patients treated with ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2 mg.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

Statement of principal findings

Our results show no evidence of a difference in effectiveness between aflibercept, ranibizumab and triamcinolone. Bevacizumab was similar to these drugs in terms of letters gained and mean change in BCVA. Dexamethasone was less effective compared to these drugs.

Strengths and limitations

This is the first study providing an indirect comparison of drugs to treat macular oedema secondary to CRVO. A robust search strategy, screening process and data extraction was used, and this analysis drew on a systematic review. The studies included had, in general, a low risk of bias. Safety was not considered in this study but is described in detail elsewhere.[45] Five different drugs were suitable for network meta-analysis. Unpublished data was obtained from one author. [42-44] Bayesian methods were used for the NMA. There was good model fit and convergence within the analysis.

However pre-specified outcomes were not reported in all studies and the sample size varied considerably. For example Epstein 2012, assessing bevacizumab, only included 30 participants in each arm.[42-44] This resulted in wide credible intervals from the network meta-analysis which may lead to a type 1 error especially with regards to the proportions of patients losing more than or equal to 3 lines. The SCORE study compared triamcinolone to observation.[23-36] The NMA assumes a [11] similar effect of sham and observation and this may result in a small degree of bias. Only six months of data was included, and the long term effects are not known. Using a six-month follow-up period may disadvantage dexamethasone because peak effect in the GENEVA trials was seen at 90 days, and by six months, benefits had been largely lost.[6-8]

As with most network meta-analyses, methodological heterogeneity was present. There were some differences amongst the trials. For example CRUISE[9 10], assessing ranibizumab, did not include as many patients with ischaemic CRVO as the aflibercept trials.[12 13] There were also some small differences in the chronicity of macular oedema and the mean BCVA at baseline.

Meaning of the study: possible explanations and implications for clinicians and policymakers

No head-to-head trials comparing aflibercept, bevacizumab, ranibizumab, triamcinolone or dexamethasone have been published in central retinal vein occlusion. Part of the reason for this is that the Food and Drug Administration require proof of the safety and effectiveness of a drug.[46] The easiest and quickest method for pharmaceutical companies to produce this is through placebo controlled trials. Trials comparing new medications to current best treatment would be considerably more useful to clinicians and patients.

Head-to-head trials comparing some of these drugs are available in other conditions. For example a comparison of ranibizumab and bevacizumab was undertaken in age related macular degeneration in the Comparison of Age-related macular degeneration Treatment Trials (CATT)[47] and alternative treatments to Inhibit VEGF in patients with Age-related choroidal Neovascularisation (IVAN)[48] trials. Both of these trials found no difference in effectiveness between ranibizumab and bevacizumab. Furthermore an indirect comparison of ranibizumab and bevacizumab found no evidence of a difference between these drugs.[49] Thus, it is highly probable that this may also apply in CRVO. The difference seen in our results regarding bevacizumab may be due to the low number of patients included in Epstein 2012.[42-44] In the CATT trial, more patients were

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

hospitalized in the bevacizumab arm, but the authors did not believe that this was explained by a direct effect of bevacizumab.[47] The 2-year results from the IVAN showed little difference in cardiovascular events, with the number being insignificantly lower with bevacizumab.[50] Ranibizumab and aflibercept were directly compared in two similarly designed trials, VEGF Trap-eye: investigation of Efficacy and safety in Wet age-related macular degeneration (VIEW 1 and 2).[51] Similar efficacy and safety was found in both drugs.

From the included trials it is clear that intraocular steroids are associated with complications, including increased intra-ocular pressure and cataract formation.[6-8 23-36] These are substantial drawbacks for using steroids to treat macular oedema in CRVO. However, many affected patients may be already pseudophakic and, on these, the use of intraocular steroids may be reasonable. Steroids may have a place in the treatment pathway of patients who have failed on anti-VEGF therapy, but this has yet to be tested. The anti-VEFG drugs have a good safety profile and do not cause cataract formation.[9-13 42-44] For this reason are likely to be more favoured by clinicians than steroids.

Aflibercept, compared with ranibizumab and bevacizumab, targets a wider range of cytokines and may have a stronger binding affinity.[52] Initial results suggested that aflibercept would require fewer injections than ranibizumab.[51] Heier and colleagues compared aflibercept and ranibizumab in two similarly designed randomised controlled trials in age related macular degeneration. They found that 2 mg aflibercept administered every eight weeks produced similar effects at 96 weeks to 0.5 mg ranibizumab every four weeks.[51] This was reflected in the FDA Dermatologic and Ophthalmic Drugs Advisory Committee recommendation that aflibercept should be given every two months following three initial monthly doses in age related macular

oedema.[53] This may be because aflibercept also appears to last longer in the eye than ranibizumab.[54] Age related macular degeneration is a more aggressive condition than central retinal vein occlusion and so it is unlikely that more frequent dosing would be needed. Therefore aflibercept may be preferred because it would reduce pressure on out-patient clinics. Furthermore there is some evidence from patients with age-related macular degeneration that aflibercept may be effective in patients who have not responded to ranibizumab.[55 56] This may be due to the higher affinity and wider number of cytokines that are targeted. There is no reason to suspect that these effects be any different for the macular oedema caused by central retinal vein occlusion. However we have as yet no evidence as to whether ranibizumab would be effective after aflibercept has failed.

The National Institute of Health and Care Excellence has recommended dexamethasone and ranibizumab,[18 19] and is currently appraising aflibercept. Until these technologies are reviewed together and compared with each other, clinicians may be left with three recommended drugs for macular oedema secondary to central retinal vein occlusion. It should be noted that during the appraisal of ranibizumab the evidence review group found that in the cost-effectiveness analysis dexamethasone was extendedly dominated by ranibizumab (an intervention is judged not be cost-effective because it has an ICER that is greater than that of a more effective intervention). The committee appraising ranibizumab did not re-consider the previous appraisal decision on dexamethasone.

Our results show that dexamethasone was not as effective as ranibizumab or aflibercept, at six months follow-up and with the dosing regimens in the trials. However these results do not assess quality of life or cost effectiveness. Bevacizumab is likely to prove more cost effective than both aflibercept and ranibizumab because it costs substantially

less.[57] However the National Institute for Health and Care Excellence has not issued guidance on bevacizumab because it does not have a license for use in the eye.

Unanswered questions and future research

Not all patients benefit from the use of anti-VEGF drugs; only about 60% gain 15 or more letters. It is not clear why some patients benefit more than others. Future research should focus on identifying subgroups of patients who are likely to benefit. Only a few of these trials included ischaemic patients, and in these trials only a few patients with ischaemia were included.[11-13] More research assessing the effectiveness of these drugs in severely ischaemic patients is needed.

Head-to-head trials comparing ranibizumab, aflibercept, bevacizumab and triamcinolone are needed. These should include assessment of cost effectiveness. To assist this, a better measure of quality of life is needed for patients with eye conditions. The widely-used EQ5D may not be sensitive enough to measure changes which are important to patients, such as the ability to drive.

In conclusion, we have found no evidence of difference between ranibizumab, bevacizumab, aflibercept and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections.

Acknowledgements

We thank Christine Clar, Sian Thomas and Rachel Court for assisting with searches, screening and data extraction for the systematic review which precede this study. We thank the authors of the Epstein 2012 trial for providing additional data.

Copyright statement

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above."

Declaration of competing interests

"All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work."

Contribution statement

NW conceived the idea. All authors contributed to the design of the study. DS and OU undertook the statistical analysis. JF, DS and OU wrote the first draft of the manuscript. All authors redrafted and agreed the final article. JF is the guarantor.

Transparency statement

JF affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

Table 1: Baseline characteristics and results of all included studies

Study	Participants	Intervention / Outcomes
DEXAMETHASONE		
GENEVA 2010[6-8] International Setting: multicentre (167 centres in 24 countries, so a mean of 2.6 patients per centre) Design: 2 identical double-blind, sham-controlled RCTs, phase 3 Follow-up: primary endpoint for the masked trial: 6 months; primary endpoint for the open-label extension: 12 months	N: CRVO – 437 eyes of 437 patients randomised; 94% follow-up at 6 months Participants: adults with visual acuity reduced because of macular oedema due to CRVO or BRVO	1. Dexamethasone 0.7 mg (n=136) Single dose 2. Dexamethasone 0.35 mg (n=154) Single dose 3. Sham (n=147) Single dose - a needleless applicator was placed against the conjunctiva to simulate the placement of study medication. Primary end point: gain of ≥15 ETDRS letters; for the open-label extension: safety

TRIAMCINOLONE		
SCORE 2009[23-36] USA Setting: multicentre Design: RCT Follow-up: primary end point 12 months, FU planned up to 36 months	N: 271 eyes of 271 patients randomised; 83% (observation) and 90% (triamcinolone) completed 12 months Participants: centre-involved macular oedema secondary to CRVO	1. Triamcinolone 1 mg (n=92) Every 4 months depending on retreatment regimen (ave 2.2 injections at 12 months) 2. Triamcinolone 4 mg (n=91) Every 4 months depending on retreatment regimen (ave 2.0 injections at 12 months) (The form of triamcinolone used was Trivaris, no longer available. It was made by the manufacturer of Ozurdex (Allergan)) 3. Observation (n=88) Primary end point: gain of ≥ 15 ETDRS letters
AFLIBERCEPT		
COPERNICUS 2012[12 13] International Setting: multicentre, 70 sites in North and South America, India and Israel. Mean 2.7 patients per centre. Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU 2 years	N: 189 eyes of 189 patients randomised; 95.7% (aflibercept) and 81.1% (sham) completed 24 weeks; 93% (aflibercept) and 77% (sham) completed 52 weeks Participants: adult patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=114) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=73) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥ 15 ETDRS letters

GALILEO 2012[11] International Setting: multicentre, 10 countries in Europe and Asia; 63 centres in total Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU up to 12 months, planned up to 76 weeks	N: 177 eyes of 177 patients randomised; 90.6% (aflibercept) and 78.9% (sham) completed 24 weeks Participants: treatment-naïve patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=103) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=71) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥15 ETDRS letters
--	--	--

RANIBIZUMAB		
CRUISE 2010[9 10] USA Setting: multicentre Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 6 months, FU up to 12 months	N: 392 eyes of 392 patients randomised; 97.7% (ranibizumab 0.3 mg), 91.5% (ranibizumab 0.5 mg), and 88.5% (sham) completed 6 months Participants: patients with foveal centre-involved macular oedema secondary to CRVO diagnosed within 12 months	1. Ranibizumab 0.3 mg (n=132) Every 4 weeks for 6 months (ave number not available) 2. Ranibizumab 0.5 mg (n=130) Every 4 weeks for 6 months (ave number not available) 3. Sham (n=130) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to the injection site) Primary end point: mean change from baseline BCVA
BEVACIZUMAB		
Epstein 2012[42-44] Sweden Setting: Single centre; St. Eriks Eye Hospital Stockholm Design: sham-injection controlled, double masked RCT Follow-up: primary end-point 6 months; open label extension up to 12 months	N: 60 eyes of 60 patients randomised; 93% completed open label extension Participants: patients with CRVO of ≤6 months	1. Bevacizumab 1.25 mg (n=30) Every 6 weeks for 6 months (ave number not available) 2. Sham (n=30) Every 6 weeks for 6 months (ave number not available) (syringe without needle pressed to the globe) Primary end point: gain of ≥15 ETDRS letters

FU= follow-up, RCT = randomised controlled trial, N = number, CRVO = central retinal vein occlusion, ETDRS = Early Treatment Diabetic Retinopathy Study, BRVO = branch retinal vein occlusion

For peer review only

Confidential

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 2: Baseline characteristics and results of included trials

	COPERNICUS[12 13]	GALILEO[11]	CRUISE[9 10]	GENEVA[6-8]	Epstein et al (2012)[42-44]	SCORE[23-36]
BASELINE SIMILARITIES						
Number (%) of patients						
	Aflib 2 mg: 114 Sham: 73	Aflib 2 mg: 103 Sham: 68	Rani 0.5 mg: 130 Sham: 130	Dexa 0.7 mg: 136 Sham: 147	Beva 1.25 mg: 30 Sham: 30	Triam 4 mg: 91 Obser: 88
Age (years)						
	Aflib 2 mg: 65.5 SD13.6 Sham: 67.5 SD14.3	Aflib 2 mg: 59.9 SD12.4 Sham: 63.8 SD13.3	Rani 0.5 mg: 67.6 SD12.4 Sham: 65.4 SD13.1	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 70.6 SD 12.6 Sham: 70.4 SD 10.4	Triam 4 mg: 67.5 SD 12.0 Obser: 69.2 SD 12.8
BCVA at baseline (SD)						
	Aflib 2 mg: 50.7 SD13.90 Sham: 48.9 SD14.42	Aflib 2 mg: 53.6 SD15.8 Sham: 50.9 SD15.4	Rani 0.5 mg: 48.1 SD14.6 Sham: 49.2 SD14.7	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 44.4 SD 15.3 Sham: 43.6 SD 16.0	Triam 4 mg: 51.0 SD 14.4 Obser: 52.1 SD 13.1
Duration of MO from diagnosis to screening						
	Aflib 2 mg: 2.73 SD3.09(in months) Sham: 1.88 SD2.19 (in months)	Aflib 2 mg: 50.9 SD15.4)(in days) Sham: 87.6 SD79.1 (in days)	Rani 0.5 mg: - Sham: -	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: NR Sham: NR	Triam 4 mg: 4.2 SD 3.6 (in months) Obser: 4.2 SD 3.1 (in months)
RESULTS						
Number (%) of patients gaining ≥ 15 letters improvement from baseline to 6 months						
	Aflib 2 mg: 64 (56.1)	Aflib 2 mg: 62 (60.2)	Rani 0.5 mg: 62 (47.7)	Dexa 0.7 mg: 25 (18)	Beva 1.25 mg: 18 (60%)	Triam 4 mg: 18 (19.5%) (avg of 4 and 8 mths)

	Sham: 9 (12.3)	Sham: 15 (22.1)	Sham: 22 (16.9)	Sham: 18 (12)	Sham: 6 (20%)	Obser: 3 (4%) (avg of 4 and 8 mths)
Number (%) of patients losing ≥15 letters of BCVA from baseline to 6 months						
	Aflib 2 mg: 2 (1.8)	Aflib 2 mg: 8 (7.8)	Rani 0.5 mg: 2 (1.5)	Dexa 0.7 mg: NR	Beva 1.25 mg: 2 (6.7%)	Triam 4 mg: 19 (20.5%) (avg of 4 and 8 mths)
	Sham: 20 (27.4)	Sham: 15 (22.1)	Sham: 20 (15.4)	Sham: NR	Sham: 7 (23.3%)	Obser: 31 (35.5%) (avg of 4 and 8 mths)
Mean change (SD) from baseline in BCVA						
	Aflib 2 mg: 17.3 (12.8)	Aflib 2 mg: 18.0 (12.2)	Rani 0.5 mg: 14.9 (13.2)	Dexa 0.7 mg: 0.1 (NR)	Beva 1.25 mg: 14.1 SD 18.7	Triam 4 mg: -0.15 SD20.67 (n=85) (weight mean and SD of 4 and 8 months)
	Sham: -4 (18)	Sham: 3.3 (14.1)	Sham: 0.8 (16.2)	Sham: -1.8 (NR)	Sham: -2.0 SD 20.5	Obser: -9.66 SD18.04 (n=75) (weighted mean and SD of 4 and 8 months)

NR = not reported, Aflib = aflibercept, Rani = ranibizumab, Dexa = dexamethasone, Triam = triamcinolone, Obser = observation, SD = standard deviation, avg = average

Table 3: Risk of bias

Study (author and year)	Adequate sequence generation	Allocation concealment	Masking	Incomplete outcome data addressed	Free of selective reporting	Free of other bias (e.g. similarity at baseline, power assessment)	Funder
GENEVA 2010[6-8]	Low	Low	Partial: patients and assessors of efficacy variables	Low: ITT analysis, 94% FU at 6 months	Low	<i>Power</i> : 81% power to detect difference in primary outcome with n=495 for each trial <i>Similarity at baseline</i> : yes	Allergan Inc.
SCORE 2009[23-36]	Low	Unclear	Partial (physicians and patients masked to dose but not triamcinolone versus observation)	Low: ITT analysis, 83 to 90% FU at 12 months	Low	<i>Power</i> : 80% power to detect difference in primary outcome with n=486 (but only 271 randomised) <i>Similarity at baseline</i> : yes	National Eye Institute grants, Allergan
COPERNICUS 2012[12 13]	Low	Unclear	Low: double-blind	Low: ITT analysis, 89.9% assessed at primary end point	Low	<i>Power</i> : 90% power to detect difference in primary outcome with n=165 <i>Similarity at baseline</i> : yes	Bayer HealthCare, Regeneron Pharmaceuticals
GALILEO 2012[11]	Unclear	Unclear	Low: double-blind	Low: ITT analysis, 86% assessed at primary end point	Low	<i>Power</i> : 90% power to detect difference in primary outcome with n=150 <i>Similarity at baseline</i> : yes	Bayer HealthCare, Regeneron Pharmaceuticals

CRUISE 2010[9 10]	Low	Unclear	Low: patients and evaluating examiners, injecting physicians masked to dose	Low: ITT analysis, 88.5 to 97.7% completed 6 months	Low	<i>Power: not reported</i> <i>Similarity at baseline: yes</i>	Genentech Inc.
Epstein 2012[42-44]	Unclear	Low	Low: patients, outcome assessors	Low: ITT analysis; missing data for 2 patients (primary endpoint)	Low	<i>Power: 80% power to detect difference in primary outcome with n=24 per group</i> <i>Similarity at baseline: yes</i>	Unclear; authors are consultants for Allergan, Novartis, Alcon, Bayer

ITT= intention to treat, FU = follow-up

Figure 1: study selection flow diagram





Figure 2: Network of randomized controlled trials comparing different treatments for proportions of gaining 3 or more lines of vision



Copyright only

Figure 3: Proportions of patients gaining 3 lines or more from baseline to six months




Figure 4: Rankogram for gaining ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions






Figure 5: Proportions of patients losing 3 lines or more from baseline to six months



Content not for review only

Figure 6: Rankogram for losing ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions



Confidential
Peer Review only

Figure 7: Mean BCVA change from baseline to 6 months.





Figure 8: Rankogram for mean change in BCVA - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions



Appendix: MEDLINE search strategy

Ovid MEDLINE(R) 1946 to March Week 1 2013, searched on 20 March 2013

- 1 CRVO.mp.
- 2 Retinal Vein Occlusion/
- 3 retinal vein occlusion.mp.
- 4 retinal vein obstruction.mp.
- 5 retinal venous occlusion.mp.
- 6 retinal venous obstruction.mp.
- 7 retina*.mp.
- 8 ("central vein occlusion" or "central vein obstruction" or "central venous occlusion" or "central venous obstruction").mp.
- 9 7 and 8
- 10 1 or 2 or 3 or 4 or 5 or 6 or 9
- 11 randomized controlled trial.pt.
- 12 (random* or "controlled trial*" or "clinical trial*" or rct).tw.
- 13 11 or 12
- 14 (metaanalys* or "meta analys*" or "meta-analys*").tw.
- 15 "systematic review*".tw.
- 16 meta analysis.pt.
- 17 14 or 15 or 16
- 18 10 and 13
- 19 10 and 17
- 20 18 or 19
- 21 limit 20 to yr="2005 -Current"

References

1. Deramo VA, Cox TA, Syed AB, Lee PP, Fekrat S. Vision-related quality of life in people with central retinal vein occlusion using the 25-item National Eye Institute Visual Function Questionnaire. Arch.Ophthalmol. 2003;**121**(9):1297-302 doi: 10.1001/archophth.121.9.1297 [doi];121/9/1297 [pii][published Online First: Epub Date]].

2. Klein R, Moss SE, Meuer SM, Klein BE. The 15-year cumulative incidence of retinal vein occlusion: the Beaver Dam Eye Study. Archives of ophthalmology 2008;**126**(4):513-8 doi: 10.1001/archophth.126.4.513[published Online First: Epub Date]].

3. Hayreh SS, Podhajsky PA, Zimmerman MB. Natural history of visual outcome in central retinal vein occlusion. Ophthalmology 2011;**118**(1):119-33 doi: S0161-6420(10)00447-1 [pii];10.1016/j.ophtha.2010.04.019 [doi][published Online First: Epub Date]].

4. McIntosh RL, Rogers SL, Lim L, et al. Natural history of central retinal vein occlusion: an evidence-based systematic review. Ophthalmology 2010;**117**(6):1113-23 doi: S0161-6420(10)00133-8 [pii];10.1016/j.ophtha.2010.01.060 [doi][published Online First: Epub Date]].

5. The Royal College of Ophthalmologists. Interim Guidelines for Management of Retinal Vein Occlusion. Secondary Interim Guidelines for Management of Retinal Vein Occlusion 2010. http://www.rcophth.ac.uk/core/core_picker/download.asp?id=728&file_title=Interim+Guidelines+for+Management+of+Retinal+Vein+Occlusion+2010.

6. Haller JA, Bandello F, Belfort R, Jr., et al. Randomized, sham-controlled trial of dexamethasone intravitreal implant in patients with macular edema due to retinal vein occlusion. *Ophthalmology* 2010;**117**(6):1134-46
7. Haller JA, Bandello F, Belfort R, Jr., et al. Dexamethasone intravitreal implant in patients with macular edema related to branch or central retinal vein occlusion twelve-month study results. *Ophthalmology* 2011;**118**(12):2453-60
8. Yeh WS, Haller JA, Lanzetta P, et al. Effect of the duration of macular edema on clinical outcomes in retinal vein occlusion treated with dexamethasone intravitreal implant. *Ophthalmology* 2012;**119**(6):1190-98
9. Brown DM, Campochiaro PA, Singh RP, et al. Ranibizumab for macular edema following central retinal vein occlusion: six-month primary end point results of a phase III study. *Ophthalmology* 2010;**117**(6):1124-33
10. Campochiaro PA, Brown DM, Awh CC, et al. Sustained benefits from ranibizumab for macular edema following central retinal vein occlusion: twelve-month outcomes of a phase III study. *Ophthalmology* 2011;**118**(10):2041-49
11. Holz FG, Roeder J, Ogura Y, et al. VEGF Trap-Eye for macular oedema secondary to central retinal vein occlusion: 6-month results of the phase III GALILEO study. *British Journal of Ophthalmology* 2013;**97**(3):278-84
12. Boyer D, Heier J, Brown DM, et al. Vascular endothelial growth factor Trap-Eye for macular edema secondary to central retinal vein occlusion: six-month results of the phase 3 COPENICUS study. *Ophthalmology* 2012;**119**(5):1024-32

13. Brown DM, Heier JS, Clark WL, et al. Intravitreal Aflibercept Injection for Macular Edema Secondary to Central Retinal Vein Occlusion: 1-Year Results From the Phase 3 COPENICUS Study. American Journal of Ophthalmology 2013;**155**(3):429-37

14. Michels S, Rosenfeld PJ, Puliafito CA, Marcus EN, Venkatraman AS. Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration twelve-week results of an uncontrolled open-label clinical study. Ophthalmology 2005;**112**(6):1035-47 doi: 10.1016/j.ophtha.2005.02.007[published Online First: Epub Date]].

15. Arevalo JF, Fromow-Guerra J, Quiroz-Mercado H, et al. Primary intravitreal bevacizumab (Avastin) for diabetic macular edema: results from the Pan-American Collaborative Retina Study Group at 6-month follow-up. Ophthalmology 2007;**114**(4):743-50 doi: 10.1016/j.ophtha.2006.12.028[published Online First: Epub Date]].

16. Rabena MD, Pieramici DJ, Castellarin AA, Nasir MA, Avery RL. Intravitreal bevacizumab (Avastin) in the treatment of macular edema secondary to branch retinal vein occlusion. Retina (Philadelphia, Pa.) 2007;**27**(4):419-25 doi: 10.1097/IAE.0b013e318030e77e[published Online First: Epub Date]].

17. Algvere PV, Epstein D, von Wendt G, Seregard S, Kvanta A. Intravitreal bevacizumab in central retinal vein occlusion: 18-month results of a prospective clinical trial. European journal of ophthalmology 2011;**21**(6):789-95 doi: 10.5301/ejo.2011.6522[published Online First: Epub Date]].

18. National Institute for Health and Care Excellence. Ranibizumab for treating visual impairment caused by macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 283. 2013.
<http://guidance.nice.org.uk/TA283/Guidance/pdf/English>.
19. National Institute for Health and Clinical Excellence. Dexamethasone intravitreal implant for the treatment of macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 229. 2011.
<http://guidance.nice.org.uk/TA229/Guidance/pdf/English>.
20. National Institute for Health and Care Excellence. Aflibercept for treating visual impairment caused by macular oedema secondary to central retinal vein occlusion. NICE technology appraisal guidance 305. 2014.
<http://guidance.nice.org.uk/TA305>.
21. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. BMJ (Clinical research ed.) 2011;**343**:d5928 doi: 10.1136/bmj.d5928[published Online First: Epub Date]].
22. Hozo S, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. BMC Medical Research Methodology 2005;**5**(1):13
23. Bhavsar AR, Ip MS, Glassman AR, Groups DatSS. The risk of endophthalmitis following intravitreal triamcinolone injection in the DRCRnet and SCORE clinical trials. American Journal of Ophthalmology 2007;**144**(3):454-56
24. Blodi BA, Domalpally A, Scott IU, et al. Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study system for evaluation of

stereoscopic color fundus photographs and fluorescein angiograms:
SCORE Study Report 9. Archives of ophthalmology 2010;**128**(9):1140-45

25. Chan CK, Ip MS, VanVeldhuisen PC, et al. SCORE Study report #11: incidences
of neovascular events in eyes with retinal vein occlusion. Ophthalmology
2011;**118**(7):1364-72

26. Ip M, Oden N, VanVeldhuisen P, Scott I, Blodi B. The Standard Care vs.
Corticosteroid for Retinal Vein Occlusion Study: Design and Baseline
Characteristics. American Academy of Ophthalmology 2008:260

27. Ip MS, Oden NL, Scott IU, et al. SCORE Study report 3: study design and
baseline characteristics. Ophthalmology 2009;**116**(9):1770-77

28. Ip MS, Scott IU, VanVeldhuisen PC, et al. A randomized trial comparing the
efficacy and safety of intravitreal triamcinolone with observation to treat
vision loss associated with macular edema secondary to central retinal
vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein
Occlusion (SCORE) study report 5. Archives of ophthalmology
2009;**127**(9):1101-14

29. Myers D, Blodi B, Ip M, Scott I, Warren K. Reading Center Evaluation of OCT
Images From Patients Enrolled in the Standard Care vs. Corticosteroid for
Retinal Vein Occlusion (SCORE) Study. Iovs. 2006;**47**:ARVO

30. Oden NL, Veldhuisen PC, Scott IU, Ip MS, Blodi BA. Temporal Variability of
OCT in Retinal Vein Occlusion Participants in the SCORE Study. Iovs.
2007;**48**:ARVO

31. Scott IU, Blodi BA, Ip MS, et al. SCORE Study Report 2: Interobserver
agreement between investigator and reading center classification of
retinal vein occlusion type. Ophthalmology 2009;**116**(4):756-61

32. Scott IU, Oden NL, VanVeldhuisen PC, et al. SCORE Study Report 7: incidence of intravitreal silicone oil droplets associated with staked-on vs luer cone syringe design. American Journal of Ophthalmology 2009;**148**(5):725-32
33. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline predictors of visual acuity and retinal thickness outcomes in patients with retinal vein occlusion: Standard Care Versus Corticosteroid for Retinal Vein Occlusion Study report 10. Ophthalmology 2011;**118**(2):345-52
34. Scott IU, VanVeldhuisen PC, Oden NL, et al. SCORE Study report 1: baseline associations between central retinal thickness and visual acuity in patients with retinal vein occlusion. Ophthalmology 2009;**116**(3):504-12
35. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline Characteristics and Response to Treatment of Participants With Hemiretinal Compared With Branch Retinal or Central Retinal Vein Occlusion in the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study: SCORE Study Report 14. Archives of ophthalmology 2012;**130**(12):1517-24
36. Warren K, Blodi BA, Oden N, Veldhuisen P, Scott IU, Ip M. Reading Center Evaluation of Baseline Retinal Images in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. Iovs. 2008:ARVO
37. Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. BMJ (Clinical research ed.) 2005;**331**(7521):897-900 doi: 10.1136/bmj.331.7521.897[published Online First: Epub Date]].
38. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. Statistics in medicine 2004;**23**(20):3105-24 doi: 10.1002/sim.1875[published Online First: Epub Date]].

39. Spiegelhalter D, Thomas A, Best N, Lunn D. *WinBUGS User Manual: Version 1.4*. Cambridge, Mass: MRC Biostatistics Unit, 2003.
40. Brooks SP, Gelman A. General Methods for Monitoring Convergence of Iterative Simulations. *Journal of Computational and Graphical Statistics* 1998;**7**(4):434-55 doi: 10.1080/10618600.1998.10474787[published Online First: Epub Date]].
41. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;**64**(2):163-71 doi: 10.1016/j.jclinepi.2010.03.016[published Online First: Epub Date]].
42. Epstein D, Algvere P, Von WG, Seregard S, Kvanta A. Long-term benefit from bevacizumab for macular edema in central retinal vein occlusion: 12-month results of a prospective study. *Acta Ophthalmologica* 2012;**90**:48 doi: 10.1111/j.1755-3768.2012.02549.x[published Online First: Epub Date]].
43. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Benefit from bevacizumab for macular edema in central retinal vein occlusion: twelve-month results of a prospective, randomized study. *Ophthalmology* 2012;**119**(12):2587-91
44. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Bevacizumab for macular edema in central retinal vein occlusion: a prospective, randomized, double-masked clinical study. *Ophthalmology* 2012;**119**(6):1184-89

45. Ford JA, Clar C, Lois N, et al. Treatments for macular oedema following central retinal vein occlusion: systematic review. BMJ open 2014;**4**(2):e004120 doi: 10.1136/bmjopen-2013-004120[published Online First: Epub Date]].
46. FDA. Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators. Secondary Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators 2011.
<http://www.fda.gov/RegulatoryInformation/Guidances/ucm126501.htm>
47. Martin DF, Maguire MG, Fine SL, et al. Ranibizumab and bevacizumab for treatment of neovascular age-related macular degeneration: two-year results. Ophthalmology 2012;**119**(7):1388-98 doi: 10.1016/j.ophtha.2012.03.053[published Online First: Epub Date]].
48. Chakravarthy U, Harding SP, Rogers CA, et al. Ranibizumab versus bevacizumab to treat neovascular age-related macular degeneration: one-year findings from the IVAN randomized trial. Ophthalmology 2012;**119**(7):1399-411 doi: S0161-6420(12)00358-2 [pii];10.1016/j.ophtha.2012.04.015 [doi][published Online First: Epub Date]].
49. Ford JA, Elders A, Shyangdan D, Royle P, Waugh N. The relative clinical effectiveness of ranibizumab and bevacizumab in diabetic macular oedema: an indirect comparison in a systematic review. BMJ (Clinical research ed.) 2012;**345**:e5182

50. Chakravarthy U, Harding SP, Rogers CA, et al. Alternative treatments to inhibit VEGF in age-related choroidal neovascularisation: 2-year findings of the IVAN randomised controlled trial. *Lancet* 2013 doi: 10.1016/s0140-6736(13)61501-9[published Online First: Epub Date]].
51. Heier JS, Brown DM, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology* 2012;**119**(12):2537-48 doi: 10.1016/j.ophtla.2012.09.006[published Online First: Epub Date]].
52. Papadopoulos N, Martin J, Ruan Q, et al. Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* 2012;**15**(2):171-85 doi: 10.1007/s10456-011-9249-6[published Online First: Epub Date]].
53. FDA. VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee. Secondary VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee 2011. <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/DermatologicandOphthalmicDrugsAdvisoryCommittee/UCM259143.pdf>.
54. Stewart MW, Rosenfeld PJ. Predicted biological activity of intravitreal VEGF Trap. *The British journal of ophthalmology* 2008;**92**(5):667-8 doi: 10.1136/bjo.2007.134874[published Online First: Epub Date]].
55. Bakall B, Folk JC, Boldt HC, et al. Aflibercept therapy for exudative age-related macular degeneration resistant to bevacizumab and ranibizumab. *Am J Ophthalmol* 2013;**156**(1):15-22.e1 doi: 10.1016/j.ajo.2013.02.017[published Online First: Epub Date]].

56. Cho H, Shah CP, Weber M, Heier JS. Aflibercept for exudative AMD with persistent fluid on ranibizumab and/or bevacizumab. The British journal of ophthalmology 2013;**97**(8):1032-5 doi: 10.1136/bjophthalmol-2013-303344[published Online First: Epub Date]].
57. Raftery J, Clegg A, Jones J, Tan SC, Lotery A. Ranibizumab (Lucentis) versus bevacizumab (Avastin): modelling cost effectiveness. The British journal of ophthalmology 2007;**91**(9):1244-6 doi: 10.1136/bjo.2007.116616[published Online First: Epub Date]].

Confidential peer review only

BMJ Open

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-005292.R1
Article Type:	Research
Date Submitted by the Author:	23-May-2014
Complete List of Authors:	Ford, John; University of East Anglia, Public Health Shyangdan, Deepson; University of Warwick, Warwick Evidence, Warwick Medical School Uthman, Olalekan; Warwick Medical School, Lois, Noemi; Queen's University, Centre for Vision and Vascular Science, Waugh, Norman; University of Warwick, Warwick Evidence
Primary Subject Heading:	Ophthalmology
Secondary Subject Heading:	Health policy
Keywords:	Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Medical retina < OPHTHALMOLOGY, Medical ophthalmology < OPHTHALMOLOGY

SCHOLARONE™
Manuscripts

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

John A Ford, Deepson Shyangdan, Olalekan A. Uthman, Noemi Lois, Norman Waugh

John A Ford, Academic Clinical Fellow in Public Health, Department of Population Health and Primary Care, University of East Anglia
Deepson Shyangdan, Research Fellow, Warwick Evidence, University of Warwick
Olalekan Uthman, Assistant Professor in Applied Research, University of Warwick
Noemi Lois, Professor of Ophthalmology, Centre for Vision and Vascular Science, Queen's University
Norman Waugh, Professor of Health Technology Assessment, University of Warwick

Corresponding author
Dr. John Ford
Norwich Medical School
University of East Anglia
Chancellor Drive
Norwich
NR4 7TJ
John.ford@uea.ac.uk
Tel: 01603 591269

Protocol – no published protocol exists for this study.

Ethics approval – not required

Funding – no external funding required

Data sharing: no additional data available

Word count: 3,331 words

What is already known on this subject

Anti-VEGF drugs (ranibizumab, bevacizumab and aflibercept) and corticosteroids (dexamethasone and triamcinolone), given intravitreally, have all been shown to be effective compared to placebo for the treatment of macular oedema secondary to central retinal vein occlusion.

There are no head-to-head trials.

What this study adds

There was no evidence of a difference in the effectiveness of aflibercept, ranibizumab, bevacizumab and triamcinolone for improving vision.

Clinicians may prefer aflibercept because steroids are associated with cataract formation and ranibizumab might require more frequent injections.

Abstract

Objective: To indirectly compare aflibercept, bevacizumab, dexamethasone, ranibizumab and triamcinolone for treatment of macular oedema secondary to central retinal vein occlusion using a network meta-analysis.

Design: Network meta-analysis

Data sources: The following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE; CDSR, DARE, HTA, NHSEED, CENTRAL; Science Citation Index and Conference Proceedings Citation Index-Science

Eligibility criteria for selecting studies: Only randomized controlled trials assessing patients with macular oedema secondary to central retinal vein occlusion were included. Studies had to report either proportions of patients gaining more than or equal to 3 lines, losing more than or equal to 3 lines, or mean change in best corrected visual acuity. Two authors screened titles and abstracts, extracted data and undertook risk of bias assessment. Bayesian network meta-analysis was used to compare the different interventions.

Results: Seven studies, assessing five drugs, were judged to be sufficiently comparable for inclusion in the NMA. For the proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2mg had a higher probability of being more effective than sham and dexamethasone. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision compared to those treated with sham. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab

1.25mg and aflibercept 2mg had a higher probability of improvement in mean best correct visual acuity compared to those treated with sham injections.

Conclusions: We found no evidence of differences between ranibizumab, aflibercept, bevacizumab and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections

Systematic review registration – Not registered

Strengths and limitations of this study

- Important topic area, with significant policy implications
- Robust method used to identify studies
- Network meta-analysis are based on a number of assumptions
- Network meta-analysis is the best method to compare interventions in the absence of head to head trials

Introduction

Central retinal vein occlusion (CRVO) dramatically reduces an individual's functioning and quality of life.[1] It is estimated that the 15 year cumulative incidence of central retinal vein occlusion is 0.5%.[2] Visual loss is caused by thrombosis of the central retinal vein which leads to a rise in venous pressure and an increase in vascular endothelial growth factor (VEGF), consequently causing an increase in vascular permeability. Macular oedema subsequently ensues with varying degrees of ischaemia and neo-vascularisation. Although CRVO is generally classified as ischaemic or non-ischaemic, ischaemia should be regarded as a spectrum.[3] Cases with ischaemia carry a considerably worse prognosis as in around a third of them, neovascular glaucoma may develop; the most devastating complication of CRVO.[4]

CRVO is more common in older people with risk factors such as diabetes, hypertension or hyperlipidaemia, but can occur in young people with inflammatory disorders. Hayreh and colleagues in a 27-year cohort study found that only 13% of people with CRVO were under 45 years of age.[3] In 95% of cases CRVO affects only one eye.[3] However visual loss in this already co-morbid patient group significantly compounds their already impaired functioning and quality of life. Patients can lose confidence, struggle with daily activities and become increasingly dependent on friends and family.[1]

For many years, laser photocoagulation was the only effective therapeutic strategy that could be used in the management of patients with CRVO. It was only useful for reducing the risk of neovascular glaucoma, but not effective for the treatment of macular oedema in CRVO.[5] Over the past decade a number of drugs to treat macular oedema have been introduced, including the steroids, triamcinolone and dexamethasone, and the anti-VEGFs, ranibizumab, bevacizumab, pegaptanib and aflibercept. Dexamethasone,

ranibizumab and aflibercept have been assessed in large commercially funded trials.[6-13] Bevacizumab was originally developed as an anti-cancer drug and has been found to be effective in treating macular oedema secondary to age-related macular degeneration,[14] diabetic macular oedema, [15] branch retinal vein occlusion[16] and central retinal vein occlusion.[17] Like triamcinolone, bevacizumab is used off licence in the eye. Ranibizumab is a derived from the same parent molecule of the bevacizumab monoclonal antibody and was developed and commercially marketed specifically for use in the eye.

In the United Kingdom, the National Institute of Health and Care Excellence (NICE) has recommended the use of dexamethasone, ranibizumab and aflibercept for the treatment of macular oedema secondary to CRVO in separate appraisals[18-20] Therefore clinicians have three NICE-recommended treatments for CRVO without head-to-head trials or clear guidance on which one may be best for their patients. On this basis, the aim of this study was to indirectly compare in a network meta-analysis the clinical effectiveness of aflibercept, ranibizumab, bevacizumab, dexamethasone and triamcinolone for the treatment of macular oedema secondary to CRVO.

Methods

Information sources and search strategy

To identify suitable studies, initially for a systematic review of treatment of macular oedema after CRVO (submitted for publication) the following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE (all via OVID); CDSR, DARE, HTA, NHSEED, CENTRAL (all via The Cochrane Library); Science Citation Index and Conference Proceedings Citation Index-Science (via Web of Knowledge). The MEDLINE search strategy is shown in appendix 1. This search strategy was modified for other databases. In addition to the bibliographic database searching, supplementary searches were undertaken to look for recent and unpublished studies in the WHO International Clinical Trials Registry Platform and ophthalmology conference websites (American Academy of Ophthalmology, Association for Research in Vision and Ophthalmology from 2010 to 2012).

Study selection

Only randomised controlled trials which included patients with macular oedema secondary to central retinal vein occlusion were included. It was acceptable for a study to include both branch retinal vein occlusion and central retinal vein occlusion provided that the central retinal vein occlusion group was reported separately. The following drugs were included: dexamethasone, triamcinolone, ranibizumab, bevacizumab and aflibercept. Pegaptanib was not included because it is not used routinely in clinical practice. Only doses which are used in clinical practice were included. Studies had to report at least one of the following outcomes: proportions of patients gaining more than or equal to 3 lines from baseline to six months, proportions of patients losing more than or equal to 3 lines from baseline to six months and mean change in best corrected visual acuity (BCVA) from baseline to six months

Risk of bias assessment

The Cochrane Collaboration's tool for assessing risk of bias was used.[21] The trials were graded (unclear, high or low risk of bias) based on: (i) sequence generation, (ii) allocation concealment, (iii) blinding of outcome assessor, (iv) incomplete outcome data, and (v) selective outcome reporting.

Study selection and data abstraction

Two authors independently assessed the eligibility and methodological quality of the studies identified during the literature search. Two authors extracted and compared the data. For each study identified that met the selection criteria, details on study design, study population characteristics, intervention, outcome measures, and study quality were extracted. Discrepancies were resolved by consensus through discussion. Studies were assessed for comparability based on the populations included, trial arms, outcome measures and duration of follow-up. Common comparators were identified from the trials and a network diagram was created.

Summary measures

The primary measures of treatment effects were relative risk (RR) for the proportions of patients gaining more than or equal to 3 lines of vision, proportions of patients losing more than or equal to 3 lines of vision and weighted mean difference (WMD) for mean change BCVA. We used the following methods to calculate standard deviations, when incompletely reported: (1) contact with the corresponding author; or (2) estimation of the standard deviation on the basis of the sample size, median, and range as suggested by Hozo and colleagues[22] or on the basis of the sample size and P value.

In one trial (SCORE),[23-36] six month data was not available because patients were followed up every four months. For the dichotomous outcomes i.e. proportions of patients gaining and losing ≥ 3 lines, we averaged four and eight month data to get the six months follow-up data. For the third outcome i.e. mean change BCVA, again data from two time-points were used. Weighted mean and SDs for each treatment arm was calculated using mean and SDs of two time-points.

Data synthesis and model implementation

Bayesian network meta-analysis [37 38] (NMA) was used to compare the different interventions. Network meta-analysis is a generalization of meta-analysis methods because they allow comparisons of agents not addressed within individual primary trials. Bayesian statistical inference provides probability distributions for treatment effect parameters (RR and WMD), with 95% credible intervals (95% CrI), rather than 95% confidence intervals (95% CI). A 95% credible interval can be interpreted as there being a 95% probability that the parameter takes a value in the specified range.[37 38]

All analyses were conducted using a Bayesian Markov Chain Monte Carlo (MCMC) method and fitted in the freely available Bayesian software, WinBUGS 1.4.3.[39] Two Markov chains were run simultaneously using different initial values. Convergence to a stable solution was checked by viewing plots of the sampled simulations and using the Brooks-Gelman-Rubin diagnostic tool.[40] Convergence was found to be adequate after running 20 000 samples for both chains. These samples were then discarded and a further 70 000 sampled simulation was then run, on which the results were based. We also calculated the probability of treatment being the most effective (first best), the second best, the third best, and so on, and presented the results graphically with rankograms.[41]

Like standard meta-analysis comparison, a NMA can be either a fixed- or a random-effect models. We used the Bayesian Deviation Information Criterion (DIC) to compare fixed and random effect models. The most appropriate NMA model can be identified as the one with the lowest DIC. The DIC measures the fit of the model while penalizing it for the number of effective parameters. The fixed - effect model was chosen because of the small number of trials available for each comparison and difficulty in estimating between studies variance if random-effect model was implemented and the difference in DIC is less than 5.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Study selection and characteristics

The literature search identified 945 articles, as shown in Figure 1. Seven studies were judged to be sufficiently comparable to be included in the network meta-analysis. Tables 1 and 2 present the characteristics and results of the included trials. Two studies [11-13] compared aflibercept 2 mg against sham; two identical studies [6-8] compared dexamethasone 0.7 mg (Ozurdex) against sham; one study [9 10] compared ranibizumab 0.5 mg against sham; one study [42-44] compared bevacizumab 1.25 mg against sham, and finally one study [23-36] compared triamcinolone 4 mg against observation. Sham or observation were used as the common comparator. The number of included participants varied from 60 [42-44] to 437 [6-8]. Most studies required patients to be treatment naive and have macular oedema with retinal thickness measuring at least 250 or 300 µm on optical coherence tomography. Sham injection was undertaken by placing a needleless syringe onto the eye. All studies, except for Epstein and colleagues 2012[42-44], were multi-centre, international studies. Most studies had an extension phase after the primary outcome, but this was not included in the network meta-analysis.

The sufficiently comparable studies were combined into a network analysis based on a common comparator. The network for the proportions of patients gaining more than or equal to 3 lines is shown in Figure 2. This network is the same for the other two outcomes, but without dexamethasone because the trial did not report these outcomes.

Risk of bias of included trials

Risk of bias is shown in Table 3. Included studies were generally of high quality, with all studies being judged to be of low or unclear bias for all criteria. The non-commercially

funded bevacizumab trial had fewer patients and inevitably results had wider confidence intervals.[42-44] In no study does it appear that patients were asked at the end of the trial what arm they thought they had been assigned. It is unclear how many could distinguish injections (intervention arm) from punctureless pressure (sham arm).

Effects of interventions on proportions of patients gaining ≥ 3 lines

Figure 3 displays a forest plot of the risk ratio and 95% credible interval in proportions of patients gaining more than or equal to 3 lines for all the possible pairwise comparisons. In terms of proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of being more effective than a sham and dexamethasone (Figure 4). There was no difference in the proportions of patients gaining more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg.

Effects of interventions on proportions of patients losing ≥ 3 lines

Figure 5 displays forest plot of the risk ratio and 95% credible interval of proportions of patients losing more than or equal to 3 lines for all the possible pairwise comparisons. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision than those treated with sham. There was no difference in the proportions of patients losing more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25 mg and aflibercept 2mg. Figure 6 shows ranking for efficacy in terms of proportions of patients losing ≥ 3 lines.

Effects of interventions on mean change in BCVA

Figure 7 displays a forest plot of the mean changes and 95% credible intervals of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

improvement in BCVA for all the possible pairwise comparisons. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of improvement in BCVA compared to those treated with sham injections. Patients treated with aflibercept 2mg had a higher probability of improvement in BCVA compared with those treated with triamcinolone 4mg (Figure 8). There was no difference in mean change in BCVA from baseline between patients treated with ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2 mg.

For peer review only

Discussion

Statement of principal findings

Our results show no evidence of a difference in effectiveness between aflibercept, ranibizumab and triamcinolone. Bevacizumab was similar to these drugs in terms of letters gained and mean change in BCVA. Dexamethasone was less effective compared to these drugs.

Strengths and limitations

This is the first study providing an indirect comparison of drugs to treat macular oedema secondary to CRVO. A robust search strategy, screening process and data extraction was used, and this analysis drew on a systematic review. The studies included had, in general, a low risk of bias. Safety was not considered in this study but is described in detail elsewhere.[45] Five different drugs were suitable for network meta-analysis. Unpublished data was obtained from one author. [42-44] Bayesian methods were used for the NMA. There was good model fit and convergence within the analysis.

However pre-specified outcomes were not reported in all studies and the sample size varied considerably. For example Epstein 2012, assessing bevacizumab, only included 30 participants in each arm.[42-44] This resulted in wide credible intervals from the network meta-analysis which may lead to a type 1 error especially with regards to the proportions of patients losing more than or equal to 3 lines. The SCORE study compared triamcinolone to observation.[23-36] The NMA assumes a [11] similar effect of sham and observation and this may result in a small degree of bias. Only six months of data was included, and the long term effects are not known. Using a six-month follow-up period may disadvantage dexamethasone because peak effect in the GENEVA trials was seen at 90 days, and by six months, benefits had been largely lost.[6-8]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

As with most network meta-analyses, methodological heterogeneity was present. There were some differences amongst the trials. For example CRUISE[9 10], assessing ranibizumab, did not include as many patients with ischaemic CRVO as the aflibercept trials.[12 13] There were also some small differences in the chronicity of macular oedema and the mean BCVA at baseline.

Meaning of the study: possible explanations and implications for clinicians and policymakers

No head-to-head trials comparing aflibercept, bevacizumab, ranibizumab, triamcinolone or dexamethasone have been published in central retinal vein occlusion. Part of the reason for this is that the Food and Drug Administration require proof of the safety and effectiveness of a drug.[46] The easiest and quickest method for pharmaceutical companies to produce this is through placebo controlled trials. Trials comparing new medications to current best treatment would be considerably more useful to clinicians and patients.

Head-to-head trials comparing some of these drugs are available in other conditions. For example a comparison of ranibizumab and bevacizumab was undertaken in age related macular degeneration in the Comparison of Age-related macular degeneration Treatment Trials (CATT)[47] and alternative treatments to Inhibit VEGF in patients with Age-related choroidal Neovascularisation (IVAN)[48] trials. Both of these trials found no difference in effectiveness between ranibizumab and bevacizumab. Furthermore an indirect comparison of ranibizumab and bevacizumab found no evidence of a difference between these drugs.[49] Thus, it is highly probable that this may also apply in CRVO. The difference seen in our results regarding bevacizumab may be due to the low number of patients included in Epstein 2012.[42-44] In the CATT trial, more patients were

hospitalized in the bevacizumab arm, but the authors did not believe that this was explained by a direct effect of bevacizumab.[47] The 2-year results from the IVAN showed little difference in cardiovascular events, with the number being insignificantly lower with bevacizumab.[50] Ranibizumab and aflibercept were directly compared in two similarly designed trials, VEGF Trap-eye: investigation of Efficacy and safety in Wet age-related macular degeneration (VIEW 1 and 2).[51] Similar efficacy and safety was found in both drugs.

From the included trials it is clear that intraocular steroids are associated with complications, including increased intra-ocular pressure and cataract formation.[6-8 23-36] These are substantial drawbacks for using steroids to treat macular oedema in CRVO. However, many affected patients may be already pseudophakic and, on these, the use of intraocular steroids may be reasonable. Steroids may have a place in the treatment pathway of patients who have failed on anti-VEGF therapy, but this has yet to be tested. The anti-VEFG drugs have a good safety profile and do not cause cataract formation.[9-13 42-44] For this reason are likely to be more favoured by clinicians than steroids.

Aflibercept, compared with ranibizumab and bevacizumab, targets a wider range of cytokines and may have a stronger binding affinity.[52] Initial results suggested that aflibercept would require fewer injections than ranibizumab.[51] Heier and colleagues compared aflibercept and ranibizumab in two similarly designed randomised controlled trials in age related macular degeneration. They found that 2 mg aflibercept administered every eight weeks produced similar effects at 96 weeks to 0.5 mg ranibizumab every four weeks.[51] This was reflected in the FDA Dermatologic and Ophthalmic Drugs Advisory Committee recommendation that aflibercept should be given every two months following three initial monthly doses in age related macular

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

oedema.[53] This may be because aflibercept also appears to last longer in the eye than ranibizumab.[54] Age related macular degeneration is a more aggressive condition than central retinal vein occlusion and so it is unlikely that more frequent dosing would be needed. Therefore aflibercept may be preferred because it would reduce pressure on out-patient clinics. Furthermore there is some evidence from patients with age-related macular degeneration that aflibercept may be effective in patients who have not responded to ranibizumab.[55 56] This may be due to the higher affinity and wider number of cytokines that are targeted. There is no reason to suspect that these effects be any different for the macular oedema caused by central retinal vein occlusion. However we have as yet no evidence as to whether ranibizumab would be effective after aflibercept has failed.

The National Institute of Health and Care Excellence has recommended dexamethasone and ranibizumab,[18 19] and is currently appraising aflibercept. Until these technologies are reviewed together and compared with each other, clinicians may be left with three recommended drugs for macular oedema secondary to central retinal vein occlusion. It should be noted that during the appraisal of ranibizumab the evidence review group found that in the cost-effectiveness analysis dexamethasone was extendedly dominated by ranibizumab (an intervention is judged not be cost-effective because it has an ICER that is greater than that that of a more effective intervention). The committee appraising ranibizumab did not re-consider the previous appraisal decision on dexamethasone.

Our results show that dexamethasone was not as effective as ranibizumab or aflibercept, at six months follow-up and with the dosing regimens in the trials. However these results do not assess quality of life or cost effectiveness. Bevacizumab is likely to prove more cost effective than both aflibercept and ranibizumab because it costs substantially

less.[57] However the National Institute for Health and Care Excellence has not issued guidance on bevacizumab because it does not have a license for use in the eye.

Unanswered questions and future research

Not all patients benefit from the use of anti-VEGF drugs; only about 60% gain 15 or more letters. It is not clear why some patients benefit more than others. Future research should focus on identifying subgroups of patients who are likely to benefit. Only a few of these trials included ischaemic patients, and in these trials only a few patients with ischaemia were included.[11-13] More research assessing the effectiveness of these drugs in severely ischaemic patients is needed.

Head-to-head trials comparing ranibizumab, aflibercept, bevacizumab and triamcinolone are needed. These should include assessment of cost effectiveness. To assist this, a better measure of quality of life is needed for patients with eye conditions. The widely-used EQ5D may not be sensitive enough to measure changes which are important to patients, such as the ability to drive.

In conclusion, we have found no evidence of difference between ranibizumab, bevacizumab, aflibercept and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

We thank Christine Clar, Sian Thomas and Rachel Court for assisting with searches, screening and data extraction for the systematic review which precede this study. We thank the authors of the Epstein 2012 trial for providing addition data.

Copyright statement

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above."

Declaration of competing interests

"All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work."

Contribution statement

NW conceived the idea. All authors contributed to the design of the study. DS and OU undertook the statistical analysis. JF, DS and OU wrote the first draft of the manuscript. All authors redrafted and agreed the final article. JF is the guarantor.

Transparency statement

JF affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

Table 1: Baseline characteristics and results of all included studies

Study	Participants	Intervention / Outcomes
DEXAMETHASONE		
GENEVA 2010[6-8] International Setting: multicentre (167 centres in 24 countries, so a mean of 2.6 patients per centre) Design: 2 identical double-blind, sham-controlled RCTs, phase 3 Follow-up: primary endpoint for the masked trial: 6 months; primary endpoint for the open-label extension: 12 months	N: CRVO – 437 eyes of 437 patients randomised; 94% follow-up at 6 months Participants: adults with visual acuity reduced because of macular oedema due to CRVO or BRVO	1. Dexamethasone 0.7 mg (n=136) Single dose 2. Dexamethasone 0.35 mg (n=154) Single dose 3. Sham (n=147) Single dose - a needleless applicator was placed against the conjunctiva to simulate the placement of study medication. Primary end point: gain of ≥ 15 ETDRS letters; for the open-label extension: safety

TRIAMCINOLONE		
SCORE 2009[23-36] USA Setting: multicentre Design: RCT Follow-up: primary end point 12 months, FU planned up to 36 months	N: 271 eyes of 271 patients randomised; 83% (observation) and 90% (triamcinolone) completed 12 months Participants: centre-involved macular oedema secondary to CRVO	1. Triamcinolone 1 mg (n=92) Every 4 months depending on retreatment regimen (ave 2.2 injections at 12 months) 2. Triamcinolone 4 mg (n=91) Every 4 months depending on retreatment regimen (ave 2.0 injections at 12 months) (The form of triamcinolone used was Trivaris, no longer available. It was made by the manufacturer of Ozurdex (Allergan)) 3. Observation (n=88) Primary end point: gain of ≥15 ETDRS letters
AFLIBERCEPT		
COPERNICUS 2012[12 13] International Setting: multicentre, 70 sites in North and South America, India and Israel. Mean 2.7 patients per centre. Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU 2 years	N: 189 eyes of 189 patients randomised; 95.7% (aflibercept) and 81.1% (sham) completed 24 weeks; 93% (aflibercept) and 77% (sham) completed 52 weeks Participants: adult patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=114) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=73) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥15 ETDRS letters

GALILEO 2012[11] International Setting: multicentre, 10 countries in Europe and Asia; 63 centres in total Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU up to 12 months, planned up to 76 weeks	N: 177 eyes of 177 patients randomised; 90.6% (aflibercept) and 78.9% (sham) completed 24 weeks Participants: treatment-naïve patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=103) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=71) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥ 15 ETDRS letters
--	--	--

RANIBIZUMAB		
CRUISE 2010 ^[9 10] USA Setting: multicentre Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 6 months, FU up to 12 months	N: 392 eyes of 392 patients randomised; 97.7% (ranibizumab 0.3 mg), 91.5% (ranibizumab 0.5 mg), and 88.5% (sham) completed 6 months Participants: patients with foveal centre-involved macular oedema secondary to CRVO diagnosed within 12 months	1. Ranibizumab 0.3 mg (n=132) Every 4 weeks for 6 months (ave number not available) 2. Ranibizumab 0.5 mg (n=130) Every 4 weeks for 6 months (ave number not available) 3. Sham (n=130) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to the injection site) Primary end point: mean change from baseline BCVA
BEVACIZUMAB		
Epstein 2012 ^[42-44] Sweden Setting: Single centre; St. Eriks Eye Hospital Stockholm Design: sham-injection controlled, double masked RCT Follow-up: primary end-point 6 months; open label extension up to 12 months	N: 60 eyes of 60 patients randomised; 93% completed open label extension Participants: patients with CRVO of ≤6 months	1. Bevacizumab 1.25 mg (n=30) Every 6 weeks for 6 months (ave number not available) 2. Sham (n=30) Every 6 weeks for 6 months (ave number not available) (syringe without needle pressed to the globe) Primary end point: gain of ≥15 ETDRS letters

FU= follow-up, RCT = randomised controlled trial, N = number, CRVO = central retinal vein occlusion, ETDRS = Early Treatment Diabetic Retinopathy Study, BRVO = branch retinal vein occlusion

Table 2: Baseline characteristics and results of included trials

	COPERNICUS[12 13]	GALILEO[11]	CRUISE[9 10]	GENEVA[6-8]	Epstein et al (2012)[42-44]	SCORE[23-36]
BASELINE SIMILARITIES						
Number (%) of patients						
	Aflib 2 mg: 114 Sham: 73	Aflib 2 mg: 103 Sham: 68	Rani 0.5 mg: 130 Sham: 130	Dexa0.7 mg: 136 Sham: 147	Beva 1.25 mg: 30 Sham: 30	Triam 4 mg: 91 Obser: 88
Age (years)						
	Aflib 2 mg: 65.5 SD13.6 Sham: 67.5 SD14.3	Aflib 2 mg: 59.9 SD12.4 Sham: 63.8 SD13.3	Rani 0.5 mg: 67.6 SD12.4 Sham: 65.4 SD13.1	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 70.6 SD 12.6 Sham: 70.4 SD 10.4	Triam 4 mg: 67.5 SD 12.0 Obser: 69.2 SD 12.8
BCVA at baseline (SD)						
	Aflib 2 mg: 50.7 SD13.90 Sham: 48.9 SD14.42	Aflib 2 mg: 53.6 SD15.8 Sham: 50.9 SD15.4	Rani 0.5 mg: 48.1 SD14.6 Sham: 49.2 SD14.7	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 44.4 SD 15.3 Sham: 43.6 SD 16.0	Triam 4 mg: 51.0 SD 14.4 Obser: 52.1 SD 13.1
Duration of MO from diagnosis to screening						
	Aflib 2 mg: 2.73 SD3.09(in months) Sham: 1.88 SD2.19 (in months)	Aflib 2 mg: 50.9 SD15.4)(in days) Sham: 87.6 SD79.1 (in days)	Rani 0.5 mg: - Sham: -	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: NR Sham: NR	Triam 4 mg: 4.2 SD 3.6 (in months) Obser: 4.2 SD 3.1 (in months)
RESULTS						
Number (%) of patients gaining ≥15 letters improvement from baseline to 6 months						
	Aflib 2 mg: 64 (56.1)	Aflib 2 mg: 62 (60.2)	Rani 0.5 mg: 62 (47.7)	Dexa 0.7 mg: 25 (18)	Beva 1.25 mg: 18 (60%)	Triam 4 mg: 18 (19.5%) (avg of 4 and 8 mths)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

	Sham: 9 (12.3)	Sham: 15 (22.1)	Sham: 22 (16.9)	Sham: 18 (12)	Sham: 6 (20%)	Obser: 3 (4%) (avg of 4 and 8 mths)
Number (%) of patients losing ≥ 15 letters of BCVA from baseline to 6 months						
	Aflib 2 mg: 2 (1.8)	Aflib 2 mg: 8 (7.8)	Rani 0.5 mg: 2 (1.5)	Dexa 0.7 mg: NR	Beva 1.25 mg: 2 (6.7%)	Triam 4 mg: 19 (20.5%) (avg of 4 and 8 mths)
	Sham: 20 (27.4)	Sham: 15 (22.1)	Sham: 20 (15.4)	Sham: NR	Sham: 7 (23.3%)	Obser: 31 (35.5%) (avg of 4 and 8 mths)
Mean change (SD) from baseline in BCVA						
	Aflib 2 mg: 17.3 (12.8)	Aflib 2 mg: 18.0 (12.2)	Rani 0.5 mg: 14.9 (13.2)	Dexa 0.7 mg: 0.1 (NR)	Beva 1.25 mg: 14.1 SD 18.7	Triam 4 mg: -0.15 SD20.67 (n=85) (weight mean and SD of 4 and 8 months)
	Sham: -4 (18)	Sham: 3.3 (14.1)	Sham: 0.8 (16.2)	Sham: -1.8 (NR)	Sham: -2.0 SD 20.5	Obser: -9.66 SD18.04 (n=75) (weighted mean and SD of 4 and 8 months)

NR = not reported, Aflib = aflibercept, Rani = ranibizumab, Dexa = dexamethasone, Triam = triamcinolone, Obser = observation, SD = standard deviation, avg = average

Table 3: Risk of bias

Study (author and year)	Adequate sequence generation	Allocation concealment	Masking	Incomplete outcome data addressed	Free of selective reporting	Free of other bias (e.g. similarity at baseline, power assessment)	Funder
GENEVA 2010[6-8]	Low	Low	Partial: patients and assessors of efficacy variables	Low: ITT analysis, 94% FU at 6 months	Low	<i>Power:</i> 81% power to detect difference in primary outcome with n=495 for each trial <i>Similarity at baseline:</i> yes	Allergan Inc.
SCORE 2009[23-36]	Low	Unclear	Partial (physicians and patients masked to dose but not triamcinolone versus observation)	Low: ITT analysis, 83 to 90% FU at 12 months	Low	<i>Power:</i> 80% power to detect difference in primary outcome with n=486 (but only 271 randomised) <i>Similarity at baseline:</i> yes	National Eye Institute grants, Allergan
COPERNICUS 2012[12 13]	Low	Unclear	Low: double-blind	Low: ITT analysis, 89.9% assessed at primary end point	Low	<i>Power:</i> 90% power to detect difference in primary outcome with n=165 <i>Similarity at baseline:</i> yes	Bayer HealthCare, Regeneron Pharmaceuticals
GALILEO 2012[11]	Unclear	Unclear	Low: double-blind	Low: ITT analysis, 86% assessed at primary end point	Low	<i>Power:</i> 90% power to detect difference in primary outcome with n=150 <i>Similarity at baseline:</i> yes	Bayer HealthCare, Regeneron Pharmaceuticals

CRUISE 2010[9 10]	Low	Unclear	Low: patients and evaluating examiners, injecting physicians masked to dose	Low: ITT analysis, 88.5 to 97.7% completed 6 months	Low	<i>Power: not reported</i> <i>Similarity at baseline: yes</i>	Genentech Inc.
Epstein 2012[42-44]	Unclear	Low	Low: patients, outcome assessors	Low: ITT analysis; missing data for 2 patients (primary endpoint)	Low	<i>Power: 80% power to detect difference in primary outcome with n=24 per group</i> <i>Similarity at baseline: yes</i>	Unclear; authors are consultants for Allergan, Novartis, Alcon, Bayer

ITT= intention to treat, FU = follow-up

Figure 1: study selection flow diagram




Figure 2: Network of randomized controlled trials comparing different treatments for proportions of gaining 3 or more lines of vision




Figure 3: Proportions of patients gaining 3 lines or more from baseline to six months




Figure 4: Rankogram for gaining ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions






Figure 5: Proportions of patients losing 3 lines or more from baseline to six months



Confidential review only

Figure 6: Rankogram for losing ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions



Confidential

Figure 7: Mean BCVA change from baseline to 6 months.





Figure 8: Rankogram for mean change in BCVA - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions



Appendix: MEDLINE search strategy

Ovid MEDLINE(R) 1946 to March Week 1 2013, searched on 20 March 2013

- 1 CRVO.mp.
- 2 Retinal Vein Occlusion/
- 3 retinal vein occlusion.mp.
- 4 retinal vein obstruction.mp.
- 5 retinal venous occlusion.mp.
- 6 retinal venous obstruction.mp.
- 7 retina*.mp.
- 8 ("central vein occlusion" or "central vein obstruction" or "central venous occlusion" or "central venous obstruction").mp.
- 9 7 and 8
- 10 1 or 2 or 3 or 4 or 5 or 6 or 9
- 11 randomized controlled trial.pt.
- 12 (random* or "controlled trial*" or "clinical trial*" or rct).tw.
- 13 11 or 12
- 14 (metaanalys* or "meta analys*" or "meta-analys*").tw.
- 15 "systematic review*".tw.
- 16 meta analysis.pt.
- 17 14 or 15 or 16
- 18 10 and 13
- 19 10 and 17
- 20 18 or 19
- 21 limit 20 to yr="2005 -Current"

References

1. Deramo VA, Cox TA, Syed AB, Lee PP, Fekrat S. Vision-related quality of life in people with central retinal vein occlusion using the 25-item National Eye Institute Visual Function Questionnaire. Arch.Ophthalmol. 2003;**121**(9):1297-302 doi: 10.1001/archophth.121.9.1297 [doi];121/9/1297 [pii][published Online First: Epub Date]].
2. Klein R, Moss SE, Meuer SM, Klein BE. The 15-year cumulative incidence of retinal vein occlusion: the Beaver Dam Eye Study. Archives of ophthalmology 2008;**126**(4):513-8 doi: 10.1001/archophth.126.4.513[published Online First: Epub Date]].
3. Hayreh SS, Podhajsky PA, Zimmerman MB. Natural history of visual outcome in central retinal vein occlusion. Ophthalmology 2011;**118**(1):119-33 doi: S0161-6420(10)00447-1 [pii];10.1016/j.ophtha.2010.04.019 [doi][published Online First: Epub Date]].
4. McIntosh RL, Rogers SL, Lim L, et al. Natural history of central retinal vein occlusion: an evidence-based systematic review. Ophthalmology 2010;**117**(6):1113-23 doi: S0161-6420(10)00133-8 [pii];10.1016/j.ophtha.2010.01.060 [doi][published Online First: Epub Date]].
5. The Royal College of Ophthalmologists. Interim Guidelines for Management of Retinal Vein Occlusion. Secondary Interim Guidelines for Management of Retinal Vein Occlusion 2010. http://www.rcophth.ac.uk/core/core_picker/download.asp?id=728&filetitle=Interim+Guidelines+for+Management+of+Retinal+Vein+Occlusion+2010.

6. Haller JA, Bandello F, Belfort R, Jr., et al. Randomized, sham-controlled trial of dexamethasone intravitreal implant in patients with macular edema due to retinal vein occlusion. *Ophthalmology* 2010;**117**(6):1134-46

7. Haller JA, Bandello F, Belfort R, Jr., et al. Dexamethasone intravitreal implant in patients with macular edema related to branch or central retinal vein occlusion twelve-month study results. *Ophthalmology* 2011;**118**(12):2453-60

8. Yeh WS, Haller JA, Lanzetta P, et al. Effect of the duration of macular edema on clinical outcomes in retinal vein occlusion treated with dexamethasone intravitreal implant. *Ophthalmology* 2012;**119**(6):1190-98

9. Brown DM, Campochiaro PA, Singh RP, et al. Ranibizumab for macular edema following central retinal vein occlusion: six-month primary end point results of a phase III study. *Ophthalmology* 2010;**117**(6):1124-33

10. Campochiaro PA, Brown DM, Awh CC, et al. Sustained benefits from ranibizumab for macular edema following central retinal vein occlusion: twelve-month outcomes of a phase III study. *Ophthalmology* 2011;**118**(10):2041-49

11. Holz FG, Roeder J, Ogura Y, et al. VEGF Trap-Eye for macular oedema secondary to central retinal vein occlusion: 6-month results of the phase III GALILEO study. *British Journal of Ophthalmology* 2013;**97**(3):278-84

12. Boyer D, Heier J, Brown DM, et al. Vascular endothelial growth factor Trap-Eye for macular edema secondary to central retinal vein occlusion: six-month results of the phase 3 COPENICUS study. *Ophthalmology* 2012;**119**(5):1024-32

13. Brown DM, Heier JS, Clark WL, et al. Intravitreal Aflibercept Injection for Macular Edema Secondary to Central Retinal Vein Occlusion: 1-Year Results From the Phase 3 COPENICUS Study. *American Journal of Ophthalmology* 2013;**155**(3):429-37
14. Michels S, Rosenfeld PJ, Puliafito CA, Marcus EN, Venkatraman AS. Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration twelve-week results of an uncontrolled open-label clinical study. *Ophthalmology* 2005;**112**(6):1035-47 doi: 10.1016/j.ophtha.2005.02.007[published Online First: Epub Date]].
15. Arevalo JF, Fromow-Guerra J, Quiroz-Mercado H, et al. Primary intravitreal bevacizumab (Avastin) for diabetic macular edema: results from the Pan-American Collaborative Retina Study Group at 6-month follow-up. *Ophthalmology* 2007;**114**(4):743-50 doi: 10.1016/j.ophtha.2006.12.028[published Online First: Epub Date]].
16. Rabena MD, Pieramici DJ, Castellarin AA, Nasir MA, Avery RL. Intravitreal bevacizumab (Avastin) in the treatment of macular edema secondary to branch retinal vein occlusion. *Retina (Philadelphia, Pa.)* 2007;**27**(4):419-25 doi: 10.1097/IAE.0b013e318030e77e[published Online First: Epub Date]].
17. Algvere PV, Epstein D, von Wendt G, Seregard S, Kvanta A. Intravitreal bevacizumab in central retinal vein occlusion: 18-month results of a prospective clinical trial. *European journal of ophthalmology* 2011;**21**(6):789-95 doi: 10.5301/ejo.2011.6522[published Online First: Epub Date]].

18. National Institute for Health and Care Excellence. Ranibizumab for treating visual impairment caused by macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 283. 2013.
<http://guidance.nice.org.uk/TA283/Guidance/pdf/English>.

19. National Institute for Health and Clinical Excellence. Dexamethasone intravitreal implant for the treatment of macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 229. 2011.
<http://guidance.nice.org.uk/TA229/Guidance/pdf/English>.

20. National Institute for Health and Care Excellence. Aflibercept for treating visual impairment caused by macular oedema secondary to central retinal vein occlusion. NICE technology appraisal guidance 305. 2014.
<http://guidance.nice.org.uk/TA305>.

21. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. BMJ (Clinical research ed.) 2011;**343**:d5928 doi: 10.1136/bmj.d5928[published Online First: Epub Date]].

22. Hozo S, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. BMC Medical Research Methodology 2005;**5**(1):13

23. Bhavsar AR, Ip MS, Glassman AR, Groups DatSS. The risk of endophthalmitis following intravitreal triamcinolone injection in the DRCRnet and SCORE clinical trials. American Journal of Ophthalmology 2007;**144**(3):454-56

24. Blodi BA, Domalpally A, Scott IU, et al. Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study system for evaluation of

- stereoscopic color fundus photographs and fluorescein angiograms:
SCORE Study Report 9. Archives of ophthalmology 2010;**128**(9):1140-45
25. Chan CK, Ip MS, VanVeldhuisen PC, et al. SCORE Study report #11: incidences
of neovascular events in eyes with retinal vein occlusion. Ophthalmology
2011;**118**(7):1364-72
26. Ip M, Oden N, VanVeldhuisen P, Scott I, Blodi B. The Standard Care vs.
Corticosteroid for Retinal Vein Occlusion Study: Design and Baseline
Characteristics. American Academy of Ophthalmology 2008:260
27. Ip MS, Oden NL, Scott IU, et al. SCORE Study report 3: study design and
baseline characteristics. Ophthalmology 2009;**116**(9):1770-77
28. Ip MS, Scott IU, VanVeldhuisen PC, et al. A randomized trial comparing the
efficacy and safety of intravitreal triamcinolone with observation to treat
vision loss associated with macular edema secondary to central retinal
vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein
Occlusion (SCORE) study report 5. Archives of ophthalmology
2009;**127**(9):1101-14
29. Myers D, Blodi B, Ip M, Scott I, Warren K. Reading Center Evaluation of OCT
Images From Patients Enrolled in the Standard Care vs. Corticosteroid for
Retinal Vein Occlusion (SCORE) Study. Iovs. 2006;**47**:ARVO
30. Oden NL, Veldhuisen PC, Scott IU, Ip MS, Blodi BA. Temporal Variability of
OCT in Retinal Vein Occlusion Participants in the SCORE Study. Iovs.
2007;**48**:ARVO
31. Scott IU, Blodi BA, Ip MS, et al. SCORE Study Report 2: Interobserver
agreement between investigator and reading center classification of
retinal vein occlusion type. Ophthalmology 2009;**116**(4):756-61

32. Scott IU, Oden NL, VanVeldhuisen PC, et al. SCORE Study Report 7: incidence of intravitreal silicone oil droplets associated with staked-on vs luer cone syringe design. American Journal of Ophthalmology 2009;**148**(5):725-32
33. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline predictors of visual acuity and retinal thickness outcomes in patients with retinal vein occlusion: Standard Care Versus Corticosteroid for Retinal Vein Occlusion Study report 10. Ophthalmology 2011;**118**(2):345-52
34. Scott IU, VanVeldhuisen PC, Oden NL, et al. SCORE Study report 1: baseline associations between central retinal thickness and visual acuity in patients with retinal vein occlusion. Ophthalmology 2009;**116**(3):504-12
35. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline Characteristics and Response to Treatment of Participants With Hemiretinal Compared With Branch Retinal or Central Retinal Vein Occlusion in the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study: SCORE Study Report 14. Archives of ophthalmology 2012;**130**(12):1517-24
36. Warren K, Blodi BA, Oden N, Veldhuisen P, Scott IU, Ip M. Reading Center Evaluation of Baseline Retinal Images in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. Iovs. 2008:ARVO
37. Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. BMJ (Clinical research ed.) 2005;**331**(7521):897-900 doi: 10.1136/bmj.331.7521.897[published Online First: Epub Date]].
38. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. Statistics in medicine 2004;**23**(20):3105-24 doi: 10.1002/sim.1875[published Online First: Epub Date]].

39. Spiegelhalter D, Thomas A, Best N, Lunn D. *WinBUGS User Manual: Version 1.4*. Cambridge, Mass: MRC Biostatistics Unit, 2003.
40. Brooks SP, Gelman A. General Methods for Monitoring Convergence of Iterative Simulations. *Journal of Computational and Graphical Statistics* 1998;**7**(4):434-55 doi: 10.1080/10618600.1998.10474787[published Online First: Epub Date]].
41. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;**64**(2):163-71 doi: 10.1016/j.jclinepi.2010.03.016[published Online First: Epub Date]].
42. Epstein D, Algvere P, Von WG, Seregard S, Kvanta A. Long-term benefit from bevacizumab for macular edema in central retinal vein occlusion: 12-month results of a prospective study. *Acta Ophthalmologica* 2012;**90**:48 doi: 10.1111/j.1755-3768.2012.02549.x[published Online First: Epub Date]].
43. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Benefit from bevacizumab for macular edema in central retinal vein occlusion: twelve-month results of a prospective, randomized study. *Ophthalmology* 2012;**119**(12):2587-91
44. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Bevacizumab for macular edema in central retinal vein occlusion: a prospective, randomized, double-masked clinical study. *Ophthalmology* 2012;**119**(6):1184-89

45. Ford JA, Clar C, Lois N, et al. Treatments for macular oedema following central retinal vein occlusion: systematic review. BMJ open 2014;**4**(2):e004120 doi: 10.1136/bmjopen-2013-004120[published Online First: Epub Date]].

46. FDA. Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators. Secondary Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators 2011. <http://www.fda.gov/RegulatoryInformation/Guidances/ucm126501.htm>.

47. Martin DF, Maguire MG, Fine SL, et al. Ranibizumab and bevacizumab for treatment of neovascular age-related macular degeneration: two-year results. Ophthalmology 2012;**119**(7):1388-98 doi: 10.1016/j.ophtha.2012.03.053[published Online First: Epub Date]].

48. Chakravarthy U, Harding SP, Rogers CA, et al. Ranibizumab versus bevacizumab to treat neovascular age-related macular degeneration: one-year findings from the IVAN randomized trial. Ophthalmology 2012;**119**(7):1399-411 doi: S0161-6420(12)00358-2 [pii];10.1016/j.ophtha.2012.04.015 [doi][published Online First: Epub Date]].

49. Ford JA, Elders A, Shyangdan D, Royle P, Waugh N. The relative clinical effectiveness of ranibizumab and bevacizumab in diabetic macular oedema: an indirect comparison in a systematic review. BMJ (Clinical research ed.) 2012;**345**:e5182

50. Chakravarthy U, Harding SP, Rogers CA, et al. Alternative treatments to inhibit VEGF in age-related choroidal neovascularisation: 2-year findings of the IVAN randomised controlled trial. *Lancet* 2013 doi: 10.1016/s0140-6736(13)61501-9[published Online First: Epub Date]].
51. Heier JS, Brown DM, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology* 2012;**119**(12):2537-48 doi: 10.1016/j.ophtha.2012.09.006[published Online First: Epub Date]].
52. Papadopoulos N, Martin J, Ruan Q, et al. Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* 2012;**15**(2):171-85 doi: 10.1007/s10456-011-9249-6[published Online First: Epub Date]].
53. FDA. VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee. Secondary VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee 2011. <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/DermatologicandOphthalmicDrugsAdvisoryCommittee/UCM259143.pdf>.
54. Stewart MW, Rosenfeld PJ. Predicted biological activity of intravitreal VEGF Trap. *The British journal of ophthalmology* 2008;**92**(5):667-8 doi: 10.1136/bjo.2007.134874[published Online First: Epub Date]].
55. Bakall B, Folk JC, Boldt HC, et al. Aflibercept therapy for exudative age-related macular degeneration resistant to bevacizumab and ranibizumab. *Am J Ophthalmol* 2013;**156**(1):15-22.e1 doi: 10.1016/j.ajo.2013.02.017[published Online First: Epub Date]].

56. Cho H, Shah CP, Weber M, Heier JS. Aflibercept for exudative AMD with persistent fluid on ranibizumab and/or bevacizumab. The British journal of ophthalmology 2013;**97**(8):1032-5 doi: 10.1136/bjophthalmol-2013-303344[published Online First: Epub Date]].

57. Raftery J, Clegg A, Jones J, Tan SC, Lotery A. Ranibizumab (Lucentis) versus bevacizumab (Avastin): modelling cost effectiveness. The British journal of ophthalmology 2007;**91**(9):1244-6 doi: 10.1136/bjo.2007.116616[published Online First: Epub Date]].

Confidential peer review only



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	3-4
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5-6
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5-6
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	1
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	7
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	41
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	7-8
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7-8
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	7-8
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	7-8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	8-9
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2 for each meta-analysis).	8-9



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	8-9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	8-9
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11+27
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	20-22
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	25-26
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	23-24
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	29-31
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	11-13
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	NA
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	14-18
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	14-18
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	14-18
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	1

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

John A Ford, Deepson Shyangdan, Olalekan A. Uthman, Noemi Lois, Norman Waugh

John A Ford, Academic Clinical Fellow in Public Health, Department of Population Health and Primary Care, University of East Anglia
Deepson Shyangdan, Research Fellow, Warwick Evidence, University of Warwick
Olalekan Uthman, Assistant Professor in Applied Research, University of Warwick
Noemi Lois, Professor of Ophthalmology, Centre for Vision and Vascular Science, Queen's University
Norman Waugh, Professor of Health Technology Assessment, University of Warwick

Corresponding author
Dr. John Ford
Norwich Medical School
University of East Anglia
Chancellor Drive
Norwich
NR4 7TJ
John.ford@uea.ac.uk
Tel: 01603 591269

Protocol – no published protocol exists for this study.

Ethics approval – not required

Funding – no external funding required

Data sharing: no additional data available

Word count: 3,331 words

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

What is already known on this subject

Anti-VEGF drugs (ranibizumab, bevacizumab and aflibercept) and corticosteroids (dexamethasone and triamcinolone), given intravitreally, have all been shown to be effective compared to placebo for the treatment of macular oedema secondary to central retinal vein occlusion.

There are no head-to-head trials.

What this study adds

There was no evidence of a difference in the effectiveness of aflibercept, ranibizumab, bevacizumab and triamcinolone for improving vision.

Clinicians may prefer aflibercept because steroids are associated with cataract formation and ranibizumab might require more frequent injections.

For peer review only

Abstract

Objective: To indirectly compare aflibercept, bevacizumab, dexamethasone, ranibizumab and triamcinolone for treatment of macular oedema secondary to central retinal vein occlusion using a network meta-analysis.

Design: Network meta-analysis

Data sources: The following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE; CDSR, DARE, HTA, NHSEED, CENTRAL; Science Citation Index and Conference Proceedings Citation Index-Science

Eligibility criteria for selecting studies: Only randomized controlled trials assessing patients with macular oedema secondary to central retinal vein occlusion were included. Studies had to report either proportions of patients gaining more than or equal to 3 lines, losing more than or equal to 3 lines, or mean change in best corrected visual acuity. Two authors screened titles and abstracts, extracted data and undertook risk of bias assessment. Bayesian network meta-analysis was used to compare the different interventions.

Results: Seven studies, assessing five drugs, were judged to be sufficiently comparable for inclusion in the NMA. For the proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2mg had a higher probability of being more effective than sham and dexamethasone. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision compared to those treated with sham. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1.25mg and aflibercept 2mg had a higher probability of improvement in mean best correct visual acuity compared to those treated with sham injections.

Conclusions: We found no evidence of differences between ranibizumab, aflibercept, bevacizumab and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections

Systematic review registration – Not registered

Strengths and limitations of this study

- Important topic area, with significant policy implications
- Robust method used to identify studies
- Network meta-analysis are based on a number of assumptions
- Network meta-analysis is the best method to compare interventions in the absence of head to head trials

Introduction

Central retinal vein occlusion (CRVO) dramatically reduces an individual's functioning and quality of life.[1] It is estimated that the 15 year cumulative incidence of central retinal vein occlusion is 0.5%.[2] Visual loss is caused by thrombosis of the central retinal vein which leads to a rise in venous pressure and an increase in vascular endothelial growth factor (VEGF), consequently causing an increase in vascular permeability. Macular oedema subsequently ensues with varying degrees of ischaemia and neo-vascularisation. Although CRVO is generally classified as ischaemic or non-ischaemic, ischaemia should be regarded as a spectrum.[3] Cases with ischaemia carry a considerably worse prognosis as in around a third of them, neovascular glaucoma may develop; the most devastating complication of CRVO.[4]

CRVO is more common in older people with risk factors such as diabetes, hypertension or hyperlipidaemia, but can occur in young people with inflammatory disorders. Hayreh and colleagues in a 27-year cohort study found that only 13% of people with CRVO were under 45 years of age.[3] In 95% of cases CRVO affects only one eye.[3] However visual loss in this already co-morbid patient group significantly compounds their already impaired functioning and quality of life. Patients can lose confidence, struggle with daily activities and become increasingly dependent on friends and family.[1]

For many years, laser photocoagulation was the only effective therapeutic strategy that could be used in the management of patients with CRVO. It was only useful for reducing the risk of neovascular glaucoma, but not effective for the treatment of macular oedema in CRVO.[5] Over the past decade a number of drugs to treat macular oedema have been introduced, including the steroids, triamcinolone and dexamethasone, and the anti-VEGFs, ranibizumab, bevacizumab, pegaptanib and aflibercept. Dexamethasone,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ranibizumab and aflibercept have been assessed in large commercially funded trials.[6-13] Bevacizumab was originally developed as an anti-cancer drug and has been found to be effective in treating macular oedema secondary to age-related macular degeneration,[14] diabetic macular oedema, [15] branch retinal vein occlusion[16] and central retinal vein occlusion.[17] Like triamcinolone, bevacizumab is used off licence in the eye. Ranibizumab is a derived from the same parent molecule of the bevacizumab monoclonal antibody and was developed and commercially marketed specifically for use in the eye.

In the United Kingdom, the National Institute of Health and Care Excellence (NICE) has recommended the use of dexamethasone, ~~and~~ ranibizumab and aflibercept for the treatment of macular oedema secondary to CRVO in separate appraisals[18-20] ~~and it is currently evaluating aflibercept. If aflibercept is also endorsed and with no head-to-head trials comparing these drugs, Therefore~~ clinicians ~~will be in the position of~~ having three NICE-recommended treatments for CRVO without head-to-head trials or clear guidance on which one may be best for their patients. On this basis, the aim of this study was to indirectly compare in a network meta-analysis the clinical effectiveness of aflibercept, ranibizumab, bevacizumab, dexamethasone and triamcinolone for the treatment of macular oedema secondary to CRVO.

Methods

Information sources and search strategy

To identify suitable studies, initially for a systematic review of treatment of macular oedema after CRVO (submitted for publication) the following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE (all via OVID); CDSR, DARE, HTA, NHSEED, CENTRAL (all via The Cochrane Library); Science Citation Index and Conference Proceedings Citation Index-Science (via Web of Knowledge). The MEDLINE search strategy is shown in appendix 1. This search strategy was modified for other databases. In addition to the bibliographic database searching, supplementary searches were undertaken to look for recent and unpublished studies in the WHO International Clinical Trials Registry Platform and ophthalmology conference websites (American Academy of Ophthalmology, Association for Research in Vision and Ophthalmology from 2010 to 2012).

Study selection

Only randomised controlled trials which included patients with macular oedema secondary to central retinal vein occlusion were included. It was acceptable for a study to include both branch retinal vein occlusion and central retinal vein occlusion provided that the central retinal vein occlusion group was reported separately. The following drugs were included: dexamethasone, triamcinolone, ranibizumab, bevacizumab and aflibercept. Pegaptanib was not included because it is not used routinely in clinical practice. Only doses which are used in clinical practice were included. Studies had to report at least one of the following outcomes: proportions of patients gaining more than or equal to 3 lines from baseline to six months, proportions of patients losing more than or equal to 3 lines from baseline to six months and mean change in best corrected visual acuity (BCVA) from baseline to six months

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Risk of bias assessment

The Cochrane Collaboration’s tool for assessing risk of bias was used.[21] The trials were graded (unclear, high or low risk of bias) based on: (i) sequence generation, (ii) allocation concealment, (iii) blinding of outcome assessor, (iv) incomplete outcome data, and (v) selective outcome reporting.

Study selection and data abstraction

Two authors independently assessed the eligibility and methodological quality of the studies identified during the literature search. Two authors extracted and compared the data. For each study identified that met the selection criteria, details on study design, study population characteristics, intervention, outcome measures, and study quality were extracted. Discrepancies were resolved by consensus through discussion. Studies were assessed for comparability based on the populations included, trial arms, outcome measures and duration of follow-up. Common comparators were identified from the trials and a network diagram was created.

Summary measures

The primary measures of treatment effects were relative risk (RR) for the proportions of patients gaining more than or equal to 3 lines of vision, proportions of patients losing more than or equal to 3 lines of vision and weighted mean difference (WMD) for mean change BCVA. We used the following methods to calculate standard deviations, when incompletely reported: (1) contact with the corresponding author; or (2) estimation of the standard deviation on the basis of the sample size, median, and range as suggested by Hozo and colleagues[22] or on the basis of the sample size and P value.

In one trial (SCORE),[23-36] six month data was not available because patients were followed up every four months. For the dichotomous outcomes i.e. proportions of patients gaining and losing ≥ 3 lines, we averaged four and eight month data to get the six months follow-up data. For the third outcome i.e. mean change BCVA, again data from two time-points were used. Weighted mean and SDs for each treatment arm was calculated using mean and SDs of two time-points.

Data synthesis and model implementation

Bayesian network meta-analysis [37 38] (NMA) was used to compare the different interventions. Network meta-analysis is a generalization of meta-analysis methods because they allow comparisons of agents not addressed within individual primary trials. Bayesian statistical inference provides probability distributions for treatment effect parameters (RR and WMD), with 95% credible intervals (95% CrI), rather than 95% confidence intervals (95% CI). A 95% credible interval can be interpreted as there being a 95% probability that the parameter takes a value in the specified range.[37 38]

All analyses were conducted using a Bayesian Markov Chain Monte Carlo (MCMC) method and fitted in the freely available Bayesian software, WinBUGS 1.4.3.[39] Two Markov chains were run simultaneously using different initial values. Convergence to a stable solution was checked by viewing plots of the sampled simulations and using the Brooks-Gelman-Rubin diagnostic tool.[40] Convergence was found to be adequate after running 20 000 samples for both chains. These samples were then discarded and a further 70 000 sampled simulation was then run, on which the results were based. [We also calculated the probability of treatment being the most effective \(first best\), the second best, the third best, and so on, and presented the results graphically with rankograms. \(Salanti\)](#)[41]:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Like standard meta-analysis comparison, a NMA can be either a fixed- or a random-effect models. We used the Bayesian Deviation Information Criterion (DIC) to compare fixed and random effect models. The most appropriate NMA model can be identified as the one with the lowest DIC. The DIC measures the fit of the model while penalizing it for the number of effective parameters. The fixed - effect model was chosen because of the small number of trials available for each comparison and difficulty in estimating between studies variance if random-effect model was implemented and the difference in DIC is less than 5.

For peer review only

Results

Study selection and characteristics

The literature search identified 945 articles, as shown in Figure 1. Seven studies were judged to be sufficiently comparable to be included in the network meta-analysis. Tables 1 and 2 present the characteristics and results of the included trials. Two studies [11-13] compared aflibercept 2 mg against sham; two identical studies [6-8] compared dexamethasone 0.7 mg (Ozurdex) against sham; one study [9 10] compared ranibizumab 0.5 mg against sham; one study [42-44] compared bevacizumab 1.25 mg against sham, and finally one study [23-36] compared triamcinolone 4 mg against observation. Sham or observation were used as the common comparator. The number of included participants varied from 60 [42-44] to 437 [6-8]. Most studies required patients to be treatment naive and have macular oedema with retinal thickness measuring at least 250 or 300 μm on optical coherence tomography. Sham injection was undertaken by placing a needleless syringe onto the eye. All studies, except for Epstein and colleagues 2012[42-44], were multi-centre, international studies. Most studies had an extension phase after the primary outcome, but this was not included in the network meta-analysis.

The sufficiently comparable studies were combined into a network analysis based on a common comparator. The network for the proportions of patients gaining more than or equal to 3 lines is shown in Figure 2. This network is the same for the other two outcomes, but without dexamethasone because the trial did not report these outcomes.

Risk of bias of included trials

Risk of bias is shown in Table 3. Included studies were generally of high quality, with all studies being judged to be of low or unclear bias for all criteria. The non-commercially

funded bevacizumab trial had fewer patients and inevitably results had wider confidence intervals.[42-44] In no study does it appear that patients were asked at the end of the trial what arm they thought they had been assigned. It is unclear how many could distinguish injections (intervention arm) from punctureless pressure (sham arm).

Effects of interventions on proportions of patients gaining ≥ 3 lines

Figure 3 displays a forest plot of the risk ratio and 95% credible interval in proportions of patients gaining more than or equal to 3 lines for all the possible pairwise comparisons. In terms of proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of being more effective than a sham and dexamethasone (eFigure 41). There was no difference in the proportions of patients gaining more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg.

Effects of interventions on proportions of patients losing ≥ 3 lines

Figure 54 displays forest plot of the risk ratio and 95% credible interval of proportions of patients losing more than or equal to 3 lines for all the possible pairwise comparisons. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision than those treated with sham. There was no difference in the proportions of patients losing more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25 mg and aflibercept 2mg. eFigure 62 shows ranking for efficacy in terms of proportions of patients losing ≥ 3 lines.

Effects of interventions on mean change in BCVA

Figure 75 displays a forest plot of the mean changes and 95% credible intervals of

improvement in BCVA for all the possible pairwise comparisons. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of improvement in BCVA compared to those treated with sham injections. Patients treated with aflibercept 2mg had a higher probability of improvement in BCVA compared with those treated with triamcinolone 4mg (eFigure 83). There was no difference in mean change in BCVA from baseline between patients treated with ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2 mg.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

Statement of principal findings

Our results show no evidence of a difference in effectiveness between aflibercept, ranibizumab and triamcinolone. Bevacizumab was similar to these drugs in terms of letters gained and mean change in BCVA. Dexamethasone was less effective compared to these drugs.

Strengths and limitations

This is the first study providing an indirect comparison of drugs to treat macular oedema secondary to CRVO. A robust search strategy, screening process and data extraction was used, and this analysis drew on a systematic review. The studies included had, in general, a low risk of bias. Safety was not considered in this study but is described in detail elsewhere.[45] Five different drugs were suitable for network meta-analysis. Unpublished data was obtained from one author. [42-44] Bayesian methods were used for the NMA. There was good model fit and convergence within the analysis.

However pre-specified outcomes were not reported in all studies and the sample size varied considerably. For example Epstein 2012, assessing bevacizumab, only included 30 participants in each arm.[42-44] This resulted in wide credible intervals from the network meta-analysis which may lead to a type 1 error especially with regards to the proportions of patients losing more than or equal to 3 lines. The SCORE study compared triamcinolone to observation.[23-36] The NMA assumes a [11] similar effect of sham and observation and this may result in a small degree of bias. Only six months of data was included, and the long term effects are not known. Using a six-month follow-up period may disadvantage dexamethasone because peak effect in the GENEVA trials was seen at 90 days, and by six months, benefits had been largely lost.[6-8]

As with most network meta-analyses, methodological heterogeneity was present. There were some differences amongst the trials. For example CRUISE[9 10], assessing ranibizumab, did not include as many patients with ischaemic CRVO as the aflibercept trials.[12 13] There were also some small differences in the chronicity of macular oedema and the mean BCVA at baseline.

Meaning of the study: possible explanations and implications for clinicians and policymakers

No head-to-head trials comparing aflibercept, bevacizumab, ranibizumab, triamcinolone or dexamethasone have been published in central retinal vein occlusion. Part of the reason for this is that the Food and Drug Administration require proof of the safety and effectiveness of a drug.[46] The easiest and quickest method for pharmaceutical companies to produce this is through placebo controlled trials. Trials comparing new medications to current best treatment would be considerably more useful to clinicians and patients.

Head-to-head trials comparing some of these drugs are available in other conditions. For example a comparison of ranibizumab and bevacizumab was undertaken in age related macular degeneration in the Comparison of Age-related macular degeneration Treatment Trials (CATT)[47] and alternative treatments to Inhibit VEGF in patients with Age-related choroidal Neovascularisation (IVAN)[48] trials. Both of these trials found no difference in effectiveness between ranibizumab and bevacizumab. Furthermore an indirect comparison of ranibizumab and bevacizumab found no evidence of a difference between these drugs.[49] Thus, it is highly probable that this may also apply in CRVO. The difference seen in our results regarding bevacizumab may be due to the low number of patients included in Epstein 2012.[42-44] In the CATT trial, more patients were

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

hospitalized in the bevacizumab arm, but the authors did not believe that this was explained by a direct effect of bevacizumab.[47] The 2-year results from the IVAN showed little difference in cardiovascular events, with the number being insignificantly lower with bevacizumab.[50] Ranibizumab and aflibercept were directly compared in two similarly designed trials, VEGF Trap-eye: investigation of Efficacy and safety in Wet age-related macular degeneration (VIEW 1 and 2).[51] Similar efficacy and safety was found in both drugs.

From the included trials it is clear that intraocular steroids are associated with complications, including increased intra-ocular pressure and cataract formation.[6-8 23-36] These are substantial drawbacks for using steroids to treat macular oedema in CRVO. However, many affected patients may be already pseudophakic and, on these, the use of intraocular steroids may be reasonable. Steroids may have a place in the treatment pathway of patients who have failed on anti-VEGF therapy, but this has yet to be tested. The anti-VEFG drugs have a good safety profile and do not cause cataract formation.[9-13 42-44] For this reason are likely to be more favoured by clinicians than steroids.

Aflibercept, compared with ranibizumab and bevacizumab, targets a wider range of cytokines and may have a stronger binding affinity.[52] Initial results suggested that aflibercept would require fewer injections than ranibizumab.[51] Heier and colleagues compared aflibercept and ranibizumab in two similarly designed randomised controlled trials in age related macular degeneration. They found that 2 mg aflibercept administered every eight weeks produced similar effects at 96 weeks to 0.5 mg ranibizumab every four weeks.[51] This was reflected in the FDA Dermatologic and Ophthalmic Drugs Advisory Committee recommendation that aflibercept should be given every two months following three initial monthly doses in age related macular

oedema.[53] This may be because aflibercept also appears to last longer in the eye than ranibizumab.[54] Age related macular degeneration is a more aggressive condition than central retinal vein occlusion and so it is unlikely that more frequent dosing would be needed. Therefore aflibercept may be preferred because it would reduce pressure on out-patient clinics. Furthermore there is some evidence from patients with age-related macular degeneration that aflibercept may be effective in patients who have not responded to ranibizumab.[55 56] This may be due to the higher affinity and wider number of cytokines that are targeted. There is no reason to suspect that these effects be any different for the macular oedema caused by central retinal vein occlusion. However we have as yet no evidence as to whether ranibizumab would be effective after aflibercept has failed.

The National Institute of Health and Care Excellence has recommended dexamethasone and ranibizumab,[18 19] and is currently appraising aflibercept. Until these technologies are reviewed together and compared with each other, clinicians may be left with three recommended drugs for macular oedema secondary to central retinal vein occlusion. It should be noted that during the appraisal of ranibizumab the evidence review group found that in the cost-effectiveness analysis dexamethasone was extendedly dominated by ranibizumab (an intervention is judged not be cost-effective because it has an ICER that is greater than that of a more effective intervention). The committee appraising ranibizumab did not re-consider the previous appraisal decision on dexamethasone.

Our results show that dexamethasone was not as effective as ranibizumab or aflibercept, at six months follow-up and with the dosing regimens in the trials. However these results do not assess quality of life or cost effectiveness. Bevacizumab is likely to prove more cost effective than both aflibercept and ranibizumab because it costs substantially

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

less.[57] However the National Institute for Health and Care Excellence has not issued guidance on bevacizumab because it does not have a license for use in the eye.

Unanswered questions and future research

Not all patients benefit from the use of anti-VEGF drugs; only about 60% gain 15 or more letters. It is not clear why some patients benefit more than others. Future research should focus on identifying subgroups of patients who are likely to benefit. Only a few of these trials included ischaemic patients, and in these trials only a few patients with ischaemia were included.[11-13] More research assessing the effectiveness of these drugs in severely ischaemic patients is needed.

Head-to-head trials comparing ranibizumab, aflibercept, bevacizumab and triamcinolone are needed. These should include assessment of cost effectiveness. To assist this, a better measure of quality of life is needed for patients with eye conditions. The widely-used EQ5D may not be sensitive enough to measure changes which are important to patients, such as the ability to drive.

In conclusion, we have found no evidence of difference between ranibizumab, bevacizumab, aflibercept and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections.

BMJ Open: first published as 10.1136/bmjopen-2014-005292 on 23 July 2014. Downloaded from <http://bmjopen.bmj.com/> on June 9, 2025 at Agence Bibliographique de l'Enseignement Supérieur (ABES).
Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Acknowledgements

We thank Christine Clar, Sian Thomas and Rachel Court for assisting with searches, screening and data extraction for the systematic review which precede this study. We thank the authors of the Epstein 2012 trial for providing additional data.

Copyright statement

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above."

Declaration of competing interests

"All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work."

Contribution statement

NW conceived the idea. All authors contributed to the design of the study. DS and OU undertook the statistical analysis. JF, DS and OU wrote the first draft of the manuscript. All authors redrafted and agreed the final article. JF is the guarantor.

Transparency statement

JF affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

Table 1: Baseline characteristics and results of all included studies

Study	Participants	Intervention / Outcomes
DEXAMETHASONE		
GENEVA 2010[6-8] International	N: CRVO – 437 eyes of 437 patients randomised; 94% follow-up at 6 months	1. Dexamethasone 0.7 mg (n=136) Single dose
Setting: multicentre (167 centres in 24 countries, so a mean of 2.6 patients per centre)	Participants: adults with visual acuity reduced because of macular oedema due to CRVO or BRVO	2. Dexamethasone 0.35 mg (n=154) Single dose
Design: 2 identical double-blind, sham-controlled RCTs, phase 3		3. Sham (n=147) Single dose - a needleless applicator was placed against the conjunctiva to simulate the placement of study medication.
Follow-up: primary endpoint for the masked trial: 6 months; primary endpoint for the open-label extension: 12 months		Primary end point: gain of ≥15 ETDRS letters; for the open-label extension: safety

TRIAMCINOLONE		
SCORE 2009[23-36] USA Setting: multicentre Design: RCT Follow-up: primary end point 12 months, FU planned up to 36 months	N: 271 eyes of 271 patients randomised; 83% (observation) and 90% (triamcinolone) completed 12 months Participants: centre-involved macular oedema secondary to CRVO	1. Triamcinolone 1 mg (n=92) Every 4 months depending on retreatment regimen (ave 2.2 injections at 12 months) 2. Triamcinolone 4 mg (n=91) Every 4 months depending on retreatment regimen (ave 2.0 injections at 12 months) (The form of triamcinolone used was Trivaris, no longer available. It was made by the manufacturer of Ozurdex (Allergan)) 3. Observation (n=88) Primary end point: gain of ≥ 15 ETDRS letters
AFLIBERCEPT		
COPERNICUS 2012[12 13] International Setting: multicentre, 70 sites in North and South America, India and Israel. Mean 2.7 patients per centre. Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU 2 years	N: 189 eyes of 189 patients randomised; 95.7% (aflibercept) and 81.1% (sham) completed 24 weeks; 93% (aflibercept) and 77% (sham) completed 52 weeks Participants: adult patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=114) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=73) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥ 15 ETDRS letters

GALILEO 2012[11] International Setting: multicentre, 10 countries in Europe and Asia; 63 centres in total Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU up to 12 months, planned up to 76 weeks	N: 177 eyes of 177 patients randomised; 90.6% (aflibercept) and 78.9% (sham) completed 24 weeks Participants: treatment-naïve patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=103) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=71) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥15 ETDRS letters
--	--	--

RANIBIZUMAB		
CRUISE 2010[9 10] USA Setting: multicentre Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 6 months, FU up to 12 months	N: 392 eyes of 392 patients randomised; 97.7% (ranibizumab 0.3 mg), 91.5% (ranibizumab 0.5 mg), and 88.5% (sham) completed 6 months Participants: patients with foveal centre-involved macular oedema secondary to CRVO diagnosed within 12 months	1. Ranibizumab 0.3 mg (n=132) Every 4 weeks for 6 months (ave number not available) 2. Ranibizumab 0.5 mg (n=130) Every 4 weeks for 6 months (ave number not available) 3. Sham (n=130) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to the injection site) Primary end point: mean change from baseline BCVA
BEVACIZUMAB		
Epstein 2012[42-44] Sweden Setting: Single centre; St. Eriks Eye Hospital Stockholm Design: sham-injection controlled, double masked RCT Follow-up: primary end-point 6 months; open label extension up to 12 months	N: 60 eyes of 60 patients randomised; 93% completed open label extension Participants: patients with CRVO of ≤ 6 months	1. Bevacizumab 1.25 mg (n=30) Every 6 weeks for 6 months (ave number not available) 2. Sham (n=30) Every 6 weeks for 6 months (ave number not available) (syringe without needle pressed to the globe) Primary end point: gain of ≥ 15 ETDRS letters

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

FU= follow-up, RCT = randomised controlled trial, N = number, CRVO = central retinal vein occlusion, ETDRS = Early Treatment Diabetic Retinopathy Study, BRVO = branch retinal vein occlusion

For peer review only
Confidential

Table 2: Baseline characteristics and results of included trials

	COPERNICUS[12 13]	GALILEO[11]	CRUISE[9 10]	GENEVA[6-8]	Epstein et al (2012)[42-44]	SCORE[23-36]
BASELINE SIMILARITIES						
Number (%) of patients						
	Aflib 2 mg: 114 Sham: 73	Aflib 2 mg: 103 Sham: 68	Rani 0.5 mg: 130 Sham: 130	Dexa 0.7 mg: 136 Sham: 147	Beva 1.25 mg: 30 Sham: 30	Triam 4 mg: 91 Obser: 88
Age (years)						
	Aflib 2 mg: 65.5 SD13.6 Sham: 67.5 SD14.3	Aflib 2 mg: 59.9 SD12.4 Sham: 63.8 SD13.3	Rani 0.5 mg: 67.6 SD12.4 Sham: 65.4 SD13.1	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 70.6 SD 12.6 Sham: 70.4 SD 10.4	Triam 4 mg: 67.5 SD 12.0 Obser: 69.2 SD 12.8
BCVA at baseline (SD)						
	Aflib 2 mg: 50.7 SD13.90 Sham: 48.9 SD14.42	Aflib 2 mg: 53.6 SD15.8 Sham: 50.9 SD15.4	Rani 0.5 mg: 48.1 SD14.6 Sham: 49.2 SD14.7	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 44.4 SD 15.3 Sham: 43.6 SD 16.0	Triam 4 mg: 51.0 SD 14.4 Obser: 52.1 SD 13.1
Duration of MO from diagnosis to screening						
	Aflib 2 mg: 2.73 SD3.09(in months) Sham: 1.88 SD2.19 (in months)	Aflib 2 mg: 50.9 SD15.4)(in days) Sham: 87.6 SD79.1 (in days)	Rani 0.5 mg: - Sham: -	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: NR Sham: NR	Triam 4 mg: 4.2 SD 3.6 (in months) Obser: 4.2 SD 3.1 (in months)
RESULTS						
Number (%) of patients gaining ≥ 15 letters improvement from baseline to 6 months						
	Aflib 2 mg: 64 (56.1)	Aflib 2 mg: 62 (60.2)	Rani 0.5 mg: 62 (47.7)	Dexa 0.7 mg: 25 (18)	Beva 1.25 mg: 18 (60%)	Triam 4 mg: 18 (19.5%) (avg of 4 and 8 mths)

	Sham: 9 (12.3)	Sham: 15 (22.1)	Sham: 22 (16.9)	Sham: 18 (12)	Sham: 6 (20%)	Obser: 3 (4%) (avg of 4 and 8 mths)
Number (%) of patients losing ≥15 letters of BCVA from baseline to 6 months						
	Aflib 2 mg: 2 (1.8)	Aflib 2 mg: 8 (7.8)	Rani 0.5 mg: 2 (1.5)	Dexa 0.7 mg: NR	Beva 1.25 mg: 2 (6.7%)	Triam 4 mg: 19 (20.5%) (avg of 4 and 8 mths)
	Sham: 20 (27.4)	Sham: 15 (22.1)	Sham: 20 (15.4)	Sham: NR	Sham: 7 (23.3%)	Obser: 31 (35.5%) (avg of 4 and 8 mths)
Mean change (SD) from baseline in BCVA						
	Aflib 2 mg: 17.3 (12.8)	Aflib 2 mg: 18.0 (12.2)	Rani 0.5 mg: 14.9 (13.2)	Dexa 0.7 mg: 0.1 (NR)	Beva 1.25 mg: 14.1 SD 18.7	Triam 4 mg: -0.15 SD20.67 (n=85) (weight mean and SD of 4 and 8 months)
	Sham: -4 (18)	Sham: 3.3 (14.1)	Sham: 0.8 (16.2)	Sham: -1.8 (NR)	Sham: -2.0 SD 20.5	Obser: -9.66 SD18.04 (n=75) (weighted mean and SD of 4 and 8 months)

NR = not reported, Aflib = aflibercept, Rani = ranibizumab, Dexa = dexamethasone, Triam = triamcinolone, Obser = observation, SD = standard deviation, avg = average

Table 3: Risk of bias

Study (author and year)	Adequate sequence generation	Allocation concealment	Masking	Incomplete outcome data addressed	Free of selective reporting	Free of other bias (e.g. similarity at baseline, power assessment)	Funder
GENEVA 2010[6-8]	Low	Low	Partial: patients and assessors of efficacy variables	Low: ITT analysis, 94% FU at 6 months	Low	<i>Power</i> : 81% power to detect difference in primary outcome with n=495 for each trial <i>Similarity at baseline</i> : yes	Allergan Inc.
SCORE 2009[23-36]	Low	Unclear	Partial (physicians and patients masked to dose but not triamcinolone versus observation)	Low: ITT analysis, 83 to 90% FU at 12 months	Low	<i>Power</i> : 80% power to detect difference in primary outcome with n=486 (but only 271 randomised) <i>Similarity at baseline</i> : yes	National Eye Institute grants, Allergan
COPERNICUS 2012[12 13]	Low	Unclear	Low: double-blind	Low: ITT analysis, 89.9% assessed at primary end point	Low	<i>Power</i> : 90% power to detect difference in primary outcome with n=165 <i>Similarity at baseline</i> : yes	Bayer HealthCare, Regeneron Pharmaceuticals
GALILEO 2012[11]	Unclear	Unclear	Low: double-blind	Low: ITT analysis, 86% assessed at primary end point	Low	<i>Power</i> : 90% power to detect difference in primary outcome with n=150 <i>Similarity at baseline</i> : yes	Bayer HealthCare, Regeneron Pharmaceuticals

CRUISE 2010[9 10]	Low	Unclear	Low: patients and evaluating examiners, injecting physicians masked to dose	Low: ITT analysis, 88.5 to 97.7% completed 6 months	Low	<i>Power: not reported</i> <i>Similarity at baseline: yes</i>	Genentech Inc.
Epstein 2012[42-44]	Unclear	Low	Low: patients, outcome assessors	Low: ITT analysis; missing data for 2 patients (primary endpoint)	Low	<i>Power: 80% power to detect difference in primary outcome with n=24 per group</i> <i>Similarity at baseline: yes</i>	Unclear; authors are consultants for Allergan, Novartis, Alcon, Bayer

ITT= intention to treat, FU = follow-up

Figure 1: study selection flow diagram





Figure 2: Network of randomized controlled trials comparing different treatments for proportions of gaining 3 or more lines of vision



Copyright only

Figure 3: Proportions of patients gaining 3 lines or more from baseline to six months




Figure 4: Rankogram for gaining ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions




Figure 5: Proportions of patients losing 3 lines or more from baseline to six months





Figure 6: Rankogram for losing ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions



BMJ Open: first published as 10.1136/bmjopen-2014-005292 on 23 July 2014. Downloaded from <http://bmjopen.bmj.com/> on June 9, 2025 at Agence Bibliographique de l'Enseignement Supérieur (ABES). Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Figure 7: Mean BCVA change from baseline to 6 months.





Figure 8: Rankogram for mean change in BCVA - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions



Appendix: MEDLINE search strategy

Ovid MEDLINE(R) 1946 to March Week 1 2013, searched on 20 March 2013

- 1 CRVO.mp.
- 2 Retinal Vein Occlusion/
- 3 retinal vein occlusion.mp.
- 4 retinal vein obstruction.mp.
- 5 retinal venous occlusion.mp.
- 6 retinal venous obstruction.mp.
- 7 retina*.mp.
- 8 ("central vein occlusion" or "central vein obstruction" or "central venous occlusion" or "central venous obstruction").mp.
- 9 7 and 8
- 10 1 or 2 or 3 or 4 or 5 or 6 or 9
- 11 randomized controlled trial.pt.
- 12 (random* or "controlled trial*" or "clinical trial*" or rct).tw.
- 13 11 or 12
- 14 (metaanalys* or "meta analys*" or "meta-analys*").tw.
- 15 "systematic review*".tw.
- 16 meta analysis.pt.
- 17 14 or 15 or 16
- 18 10 and 13
- 19 10 and 17
- 20 18 or 19
- 21 limit 20 to yr="2005 -Current"

References

1. Deramo VA, Cox TA, Syed AB, Lee PP, Fekrat S. Vision-related quality of life in people with central retinal vein occlusion using the 25-item National Eye Institute Visual Function Questionnaire. Arch.Ophthalmol. 2003;**121**(9):1297-302 doi: 10.1001/archopht.121.9.1297 [doi];121/9/1297 [pii][published Online First: Epub Date]].

2. Klein R, Moss SE, Meuer SM, Klein BE. The 15-year cumulative incidence of retinal vein occlusion: the Beaver Dam Eye Study. Archives of ophthalmology 2008;**126**(4):513-8 doi: 10.1001/archopht.126.4.513[published Online First: Epub Date]].

3. Hayreh SS, Podhajsky PA, Zimmerman MB. Natural history of visual outcome in central retinal vein occlusion. Ophthalmology 2011;**118**(1):119-33 doi: S0161-6420(10)00447-1 [pii];10.1016/j.ophtha.2010.04.019 [doi][published Online First: Epub Date]].

4. McIntosh RL, Rogers SL, Lim L, et al. Natural history of central retinal vein occlusion: an evidence-based systematic review. Ophthalmology 2010;**117**(6):1113-23 doi: S0161-6420(10)00133-8 [pii];10.1016/j.ophtha.2010.01.060 [doi][published Online First: Epub Date]].

5. The Royal College of Ophthalmologists. Interim Guidelines for Management of Retinal Vein Occlusion. Secondary Interim Guidelines for Management of Retinal Vein Occlusion 2010. http://www.rcophth.ac.uk/core/core_picker/download.asp?id=728&file_title=Interim+Guidelines+for+Management+of+Retinal+Vein+Occlusion+2010.

6. Haller JA, Bandello F, Belfort R, Jr., et al. Randomized, sham-controlled trial of dexamethasone intravitreal implant in patients with macular edema due to retinal vein occlusion. *Ophthalmology* 2010;**117**(6):1134-46
7. Haller JA, Bandello F, Belfort R, Jr., et al. Dexamethasone intravitreal implant in patients with macular edema related to branch or central retinal vein occlusion twelve-month study results. *Ophthalmology* 2011;**118**(12):2453-60
8. Yeh WS, Haller JA, Lanzetta P, et al. Effect of the duration of macular edema on clinical outcomes in retinal vein occlusion treated with dexamethasone intravitreal implant. *Ophthalmology* 2012;**119**(6):1190-98
9. Brown DM, Campochiaro PA, Singh RP, et al. Ranibizumab for macular edema following central retinal vein occlusion: six-month primary end point results of a phase III study. *Ophthalmology* 2010;**117**(6):1124-33
10. Campochiaro PA, Brown DM, Awh CC, et al. Sustained benefits from ranibizumab for macular edema following central retinal vein occlusion: twelve-month outcomes of a phase III study. *Ophthalmology* 2011;**118**(10):2041-49
11. Holz FG, Roeder J, Ogura Y, et al. VEGF Trap-Eye for macular oedema secondary to central retinal vein occlusion: 6-month results of the phase III GALILEO study. *British Journal of Ophthalmology* 2013;**97**(3):278-84
12. Boyer D, Heier J, Brown DM, et al. Vascular endothelial growth factor Trap-Eye for macular edema secondary to central retinal vein occlusion: six-month results of the phase 3 COPENICUS study. *Ophthalmology* 2012;**119**(5):1024-32

13. Brown DM, Heier JS, Clark WL, et al. Intravitreal Aflibercept Injection for Macular Edema Secondary to Central Retinal Vein Occlusion: 1-Year Results From the Phase 3 COPENICUS Study. *American Journal of Ophthalmology* 2013;**155**(3):429-37

14. Michels S, Rosenfeld PJ, Puliafito CA, Marcus EN, Venkatraman AS. Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration twelve-week results of an uncontrolled open-label clinical study. *Ophthalmology* 2005;**112**(6):1035-47 doi: 10.1016/j.ophtha.2005.02.007[published Online First: Epub Date]].

15. Arevalo JF, Fromow-Guerra J, Quiroz-Mercado H, et al. Primary intravitreal bevacizumab (Avastin) for diabetic macular edema: results from the Pan-American Collaborative Retina Study Group at 6-month follow-up. *Ophthalmology* 2007;**114**(4):743-50 doi: 10.1016/j.ophtha.2006.12.028[published Online First: Epub Date]].

16. Rabena MD, Pieramici DJ, Castellarin AA, Nasir MA, Avery RL. Intravitreal bevacizumab (Avastin) in the treatment of macular edema secondary to branch retinal vein occlusion. *Retina (Philadelphia, Pa.)* 2007;**27**(4):419-25 doi: 10.1097/IAE.0b013e318030e77e[published Online First: Epub Date]].

17. Algvere PV, Epstein D, von Wendt G, Seregard S, Kvanta A. Intravitreal bevacizumab in central retinal vein occlusion: 18-month results of a prospective clinical trial. *European journal of ophthalmology* 2011;**21**(6):789-95 doi: 10.5301/ejo.2011.6522[published Online First: Epub Date]].

18. National Institute for Health and Care Excellence. Ranibizumab for treating visual impairment caused by macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 283. 2013.
<http://guidance.nice.org.uk/TA283/Guidance/pdf/English>.
19. National Institute for Health and Clinical Excellence. Dexamethasone intravitreal implant for the treatment of macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 229. 2011.
<http://guidance.nice.org.uk/TA229/Guidance/pdf/English>.
20. National Institute for Health and Care Excellence. Aflibercept for treating visual impairment caused by macular oedema secondary to central retinal vein occlusion. NICE technology appraisal guidance 305. 2014.
<http://guidance.nice.org.uk/TA305>.
21. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. BMJ (Clinical research ed.) 2011;**343**:d5928 doi: 10.1136/bmj.d5928[published Online First: Epub Date]].
22. Hozo S, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. BMC Medical Research Methodology 2005;**5**(1):13
23. Bhavsar AR, Ip MS, Glassman AR, Groups DatSS. The risk of endophthalmitis following intravitreal triamcinolone injection in the DRCRnet and SCORE clinical trials. American Journal of Ophthalmology 2007;**144**(3):454-56
24. Blodi BA, Domalpally A, Scott IU, et al. Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study system for evaluation of

stereoscopic color fundus photographs and fluorescein angiograms:
SCORE Study Report 9. Archives of ophthalmology 2010;**128**(9):1140-45

25. Chan CK, Ip MS, VanVeldhuisen PC, et al. SCORE Study report #11: incidences
of neovascular events in eyes with retinal vein occlusion. Ophthalmology
2011;**118**(7):1364-72

26. Ip M, Oden N, VanVeldhuisen P, Scott I, Blodi B. The Standard Care vs.
Corticosteroid for Retinal Vein Occlusion Study: Design and Baseline
Characteristics. American Academy of Ophthalmology 2008:260

27. Ip MS, Oden NL, Scott IU, et al. SCORE Study report 3: study design and
baseline characteristics. Ophthalmology 2009;**116**(9):1770-77

28. Ip MS, Scott IU, VanVeldhuisen PC, et al. A randomized trial comparing the
efficacy and safety of intravitreal triamcinolone with observation to treat
vision loss associated with macular edema secondary to central retinal
vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein
Occlusion (SCORE) study report 5. Archives of ophthalmology
2009;**127**(9):1101-14

29. Myers D, Blodi B, Ip M, Scott I, Warren K. Reading Center Evaluation of OCT
Images From Patients Enrolled in the Standard Care vs. Corticosteroid for
Retinal Vein Occlusion (SCORE) Study. Iovs. 2006;**47**:ARVO

30. Oden NL, Veldhuisen PC, Scott IU, Ip MS, Blodi BA. Temporal Variability of
OCT in Retinal Vein Occlusion Participants in the SCORE Study. Iovs.
2007;**48**:ARVO

31. Scott IU, Blodi BA, Ip MS, et al. SCORE Study Report 2: Interobserver
agreement between investigator and reading center classification of
retinal vein occlusion type. Ophthalmology 2009;**116**(4):756-61

32. Scott IU, Oden NL, VanVeldhuisen PC, et al. SCORE Study Report 7: incidence of intravitreal silicone oil droplets associated with staked-on vs luer cone syringe design. American Journal of Ophthalmology 2009;**148**(5):725-32
33. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline predictors of visual acuity and retinal thickness outcomes in patients with retinal vein occlusion: Standard Care Versus Corticosteroid for Retinal Vein Occlusion Study report 10. Ophthalmology 2011;**118**(2):345-52
34. Scott IU, VanVeldhuisen PC, Oden NL, et al. SCORE Study report 1: baseline associations between central retinal thickness and visual acuity in patients with retinal vein occlusion. Ophthalmology 2009;**116**(3):504-12
35. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline Characteristics and Response to Treatment of Participants With Hemiretinal Compared With Branch Retinal or Central Retinal Vein Occlusion in the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study: SCORE Study Report 14. Archives of ophthalmology 2012;**130**(12):1517-24
36. Warren K, Blodi BA, Oden N, Veldhuisen P, Scott IU, Ip M. Reading Center Evaluation of Baseline Retinal Images in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. Iovs. 2008:ARVO
37. Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. BMJ (Clinical research ed.) 2005;**331**(7521):897-900 doi: 10.1136/bmj.331.7521.897[published Online First: Epub Date]].
38. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. Statistics in medicine 2004;**23**(20):3105-24 doi: 10.1002/sim.1875[published Online First: Epub Date]].

39. Spiegelhalter D, Thomas A, Best N, Lunn D. *WinBUGS User Manual: Version 1.4*. Cambridge, Mass: MRC Biostatistics Unit, 2003.
40. Brooks SP, Gelman A. General Methods for Monitoring Convergence of Iterative Simulations. *Journal of Computational and Graphical Statistics* 1998;**7**(4):434-55 doi: 10.1080/10618600.1998.10474787[published Online First: Epub Date]].
41. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;**64**(2):163-71 doi: 10.1016/j.jclinepi.2010.03.016[published Online First: Epub Date]].
42. Epstein D, Algvere P, Von WG, Seregard S, Kvanta A. Long-term benefit from bevacizumab for macular edema in central retinal vein occlusion: 12-month results of a prospective study. *Acta Ophthalmologica* 2012;**90**:48 doi: 10.1111/j.1755-3768.2012.02549.x[published Online First: Epub Date]].
43. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Benefit from bevacizumab for macular edema in central retinal vein occlusion: twelve-month results of a prospective, randomized study. *Ophthalmology* 2012;**119**(12):2587-91
44. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Bevacizumab for macular edema in central retinal vein occlusion: a prospective, randomized, double-masked clinical study. *Ophthalmology* 2012;**119**(6):1184-89

45. Ford JA, Clar C, Lois N, et al. Treatments for macular oedema following central retinal vein occlusion: systematic review. BMJ open 2014;**4**(2):e004120 doi: 10.1136/bmjopen-2013-004120[published Online First: Epub Date]].
46. FDA. Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators. Secondary Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators 2011.
<http://www.fda.gov/RegulatoryInformation/Guidances/ucm126501.htm>
47. Martin DF, Maguire MG, Fine SL, et al. Ranibizumab and bevacizumab for treatment of neovascular age-related macular degeneration: two-year results. Ophthalmology 2012;**119**(7):1388-98 doi: 10.1016/j.ophtha.2012.03.053[published Online First: Epub Date]].
48. Chakravarthy U, Harding SP, Rogers CA, et al. Ranibizumab versus bevacizumab to treat neovascular age-related macular degeneration: one-year findings from the IVAN randomized trial. Ophthalmology 2012;**119**(7):1399-411 doi: S0161-6420(12)00358-2 [pii];10.1016/j.ophtha.2012.04.015 [doi][published Online First: Epub Date]].
49. Ford JA, Elders A, Shyangdan D, Royle P, Waugh N. The relative clinical effectiveness of ranibizumab and bevacizumab in diabetic macular oedema: an indirect comparison in a systematic review. BMJ (Clinical research ed.) 2012;**345**:e5182

50. Chakravarthy U, Harding SP, Rogers CA, et al. Alternative treatments to inhibit VEGF in age-related choroidal neovascularisation: 2-year findings of the IVAN randomised controlled trial. *Lancet* 2013 doi: 10.1016/s0140-6736(13)61501-9[published Online First: Epub Date]].
51. Heier JS, Brown DM, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology* 2012;**119**(12):2537-48 doi: 10.1016/j.ophtha.2012.09.006[published Online First: Epub Date]].
52. Papadopoulos N, Martin J, Ruan Q, et al. Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* 2012;**15**(2):171-85 doi: 10.1007/s10456-011-9249-6[published Online First: Epub Date]].
53. FDA. VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee. Secondary VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee 2011. <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/DermatologicandOphthalmicDrugsAdvisoryCommittee/UCM259143.pdf>.
54. Stewart MW, Rosenfeld PJ. Predicted biological activity of intravitreal VEGF Trap. *The British journal of ophthalmology* 2008;**92**(5):667-8 doi: 10.1136/bjo.2007.134874[published Online First: Epub Date]].
55. Bakall B, Folk JC, Boldt HC, et al. Aflibercept therapy for exudative age-related macular degeneration resistant to bevacizumab and ranibizumab. *Am J Ophthalmol* 2013;**156**(1):15-22.e1 doi: 10.1016/j.ajo.2013.02.017[published Online First: Epub Date]].

56. Cho H, Shah CP, Weber M, Heier JS. Aflibercept for exudative AMD with persistent fluid on ranibizumab and/or bevacizumab. The British journal of ophthalmology 2013;**97**(8):1032-5 doi: 10.1136/bjophthalmol-2013-303344[published Online First: Epub Date]].
57. Raftery J, Clegg A, Jones J, Tan SC, Lotery A. Ranibizumab (Lucentis) versus bevacizumab (Avastin): modelling cost effectiveness. The British journal of ophthalmology 2007;**91**(9):1244-6 doi: 10.1136/bjo.2007.116616[published Online First: Epub Date]].

Confidential peer review only

BMJ Open

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-005292.R2
Article Type:	Research
Date Submitted by the Author:	27-Jun-2014
Complete List of Authors:	Ford, John; University of East Anglia, Public Health Shyangdan, Deepson; University of Warwick, Warwick Evidence, Warwick Medical School Uthman, Olalekan; Warwick Medical School, Lois, Noemi; Queen's University, Centre for Vision and Vascular Science, Waugh, Norman; University of Warwick, Warwick Evidence
Primary Subject Heading:	Ophthalmology
Secondary Subject Heading:	Health policy
Keywords:	Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Medical retina < OPHTHALMOLOGY, Medical ophthalmology < OPHTHALMOLOGY

SCHOLARONE™
Manuscripts

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

John A Ford, Deepson Shyangdan, Olalekan A. Uthman, Noemi Lois, Norman Waugh

John A Ford, Academic Clinical Fellow in Public Health, Department of Population Health and Primary Care, University of East Anglia
Deepson Shyangdan, Research Fellow, Warwick Evidence, University of Warwick
Olalekan Uthman, Assistant Professor in Applied Research, University of Warwick
Noemi Lois, Professor of Ophthalmology, Centre for Vision and Vascular Science, Queen's University
Norman Waugh, Professor of Health Technology Assessment, University of Warwick

Corresponding author
Dr. John Ford
Norwich Medical School
University of East Anglia
Chancellor Drive
Norwich
NR4 7TJ
John.ford@uea.ac.uk
Tel: 01603 591269

Protocol – no published protocol exists for this study.

Word count: 3,331 words

What is already known on this subject

Anti-VEGF drugs (ranibizumab, bevacizumab and aflibercept) and corticosteroids (dexamethasone and triamcinolone), given intravitreally, have all been shown to be effective compared to placebo for the treatment of macular oedema secondary to central retinal vein occlusion.

There are no head-to-head trials.

What this study adds

There was no evidence of a difference in the effectiveness of aflibercept, ranibizumab, bevacizumab and triamcinolone for improving vision.

Clinicians may prefer aflibercept because steroids are associated with cataract formation and ranibizumab might require more frequent injections.

Abstract

Objective: To indirectly compare aflibercept, bevacizumab, dexamethasone, ranibizumab and triamcinolone for treatment of macular oedema secondary to central retinal vein occlusion using a network meta-analysis.

Design: Network meta-analysis

Data sources: The following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE; CDSR, DARE, HTA, NHSEED, CENTRAL; Science Citation Index and Conference Proceedings Citation Index-Science

Eligibility criteria for selecting studies: Only randomized controlled trials assessing patients with macular oedema secondary to central retinal vein occlusion were included. Studies had to report either proportions of patients gaining more than or equal to 3 lines, losing more than or equal to 3 lines, or mean change in best corrected visual acuity. Two authors screened titles and abstracts, extracted data and undertook risk of bias assessment. Bayesian network meta-analysis was used to compare the different interventions.

Results: Seven studies, assessing five drugs, were judged to be sufficiently comparable for inclusion in the NMA. For the proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2mg had a higher probability of being more effective than sham and dexamethasone. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision compared to those treated with sham. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab

1.25mg and aflibercept 2mg had a higher probability of improvement in mean best correct visual acuity compared to those treated with sham injections.

Conclusions: We found no evidence of differences between ranibizumab, aflibercept, bevacizumab and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections

Systematic review registration – Not registered

Strengths and limitations of this study

- Important topic area, with significant policy implications
- Robust method used to identify studies
- Network meta-analysis are based on a number of assumptions
- Network meta-analysis is the best method to compare interventions in the absence of head to head trials

Introduction

Central retinal vein occlusion (CRVO) dramatically reduces an individual's functioning and quality of life.[1] It is estimated that the 15 year cumulative incidence of central retinal vein occlusion is 0.5%.[2] Visual loss is caused by thrombosis of the central retinal vein which leads to a rise in venous pressure and an increase in vascular endothelial growth factor (VEGF), consequently causing an increase in vascular permeability. Macular oedema subsequently ensues with varying degrees of ischaemia and neo-vascularisation. Although CRVO is generally classified as ischaemic or non-ischaemic, ischaemia should be regarded as a spectrum.[3] Cases with ischaemia carry a considerably worse prognosis as in around a third of them, neovascular glaucoma may develop; the most devastating complication of CRVO.[4]

CRVO is more common in older people with risk factors such as diabetes, hypertension or hyperlipidaemia, but can occur in young people with inflammatory disorders. Hayreh and colleagues in a 27-year cohort study found that only 13% of people with CRVO were under 45 years of age.[3] In 95% of cases CRVO affects only one eye.[3] However visual loss in this already co-morbid patient group significantly compounds their already impaired functioning and quality of life. Patients can lose confidence, struggle with daily activities and become increasingly dependent on friends and family.[1]

For many years, laser photocoagulation was the only effective therapeutic strategy that could be used in the management of patients with CRVO. It was only useful for reducing the risk of neovascular glaucoma, but not effective for the treatment of macular oedema in CRVO.[5] Over the past decade a number of drugs to treat macular oedema have been introduced, including the steroids, triamcinolone and dexamethasone, and the anti-VEGFs, ranibizumab, bevacizumab, pegaptanib and aflibercept. Dexamethasone,

ranibizumab and aflibercept have been assessed in large commercially funded trials.[6-13] Bevacizumab was originally developed as an anti-cancer drug and has been found to be effective in treating macular oedema secondary to age-related macular degeneration,[14] diabetic macular oedema, [15] branch retinal vein occlusion[16] and central retinal vein occlusion.[17] Like triamcinolone, bevacizumab is used off licence in the eye. Ranibizumab is a derived from the same parent molecule of the bevacizumab monoclonal antibody and was developed and commercially marketed specifically for use in the eye.

In the United Kingdom, the National Institute of Health and Care Excellence (NICE) has recommended the use of dexamethasone, ranibizumab and aflibercept for the treatment of macular oedema secondary to CRVO in separate appraisals[18-20] Therefore clinicians have three NICE-recommended treatments for CRVO without head-to-head trials or clear guidance on which one may be best for their patients. On this basis, the aim of this study was to indirectly compare in a network meta-analysis the clinical effectiveness of aflibercept, ranibizumab, bevacizumab, dexamethasone and triamcinolone for the treatment of macular oedema secondary to CRVO.

Methods

Information sources and search strategy

To identify suitable studies, initially for a systematic review of treatment of macular oedema after CRVO (submitted for publication) the following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE (all via OVID); CDSR, DARE, HTA, NHSEED, CENTRAL (all via The Cochrane Library); Science Citation Index and Conference Proceedings Citation Index-Science (via Web of Knowledge). The MEDLINE search strategy is shown in appendix 1. This search strategy was modified for other databases. In addition to the bibliographic database searching, supplementary searches were undertaken to look for recent and unpublished studies in the WHO International Clinical Trials Registry Platform and ophthalmology conference websites (American Academy of Ophthalmology, Association for Research in Vision and Ophthalmology from 2010 to 2012).

Study selection

Only randomised controlled trials which included patients with macular oedema secondary to central retinal vein occlusion were included. It was acceptable for a study to include both branch retinal vein occlusion and central retinal vein occlusion provided that the central retinal vein occlusion group was reported separately. The following drugs were included: dexamethasone, triamcinolone, ranibizumab, bevacizumab and aflibercept. Pegaptanib was not included because it is not used routinely in clinical practice. Only doses which are used in clinical practice were included. Studies had to report at least one of the following outcomes: proportions of patients gaining more than or equal to 3 lines from baseline to six months, proportions of patients losing more than or equal to 3 lines from baseline to six months and mean change in best corrected visual acuity (BCVA) from baseline to six months

Risk of bias assessment

The Cochrane Collaboration's tool for assessing risk of bias was used.[21] The trials were graded (unclear, high or low risk of bias) based on: (i) sequence generation, (ii) allocation concealment, (iii) blinding of outcome assessor, (iv) incomplete outcome data, and (v) selective outcome reporting.

Study selection and data abstraction

Two authors independently assessed the eligibility and methodological quality of the studies identified during the literature search. Two authors extracted and compared the data. For each study identified that met the selection criteria, details on study design, study population characteristics, intervention, outcome measures, and study quality were extracted. Discrepancies were resolved by consensus through discussion. Studies were assessed for comparability based on the populations included, trial arms, outcome measures and duration of follow-up. Common comparators were identified from the trials and a network diagram was created.

Summary measures

The primary measures of treatment effects were relative risk (RR) for the proportions of patients gaining more than or equal to 3 lines of vision, proportions of patients losing more than or equal to 3 lines of vision and weighted mean difference (WMD) for mean change BCVA. We used the following methods to calculate standard deviations, when incompletely reported: (1) contact with the corresponding author; or (2) estimation of the standard deviation on the basis of the sample size, median, and range as suggested by Hozo and colleagues[22] or on the basis of the sample size and P value.

In one trial (SCORE),[23-36] six month data was not available because patients were followed up every four months. For the dichotomous outcomes i.e. proportions of patients gaining and losing ≥ 3 lines, we averaged four and eight month data to get the six months follow-up data. For the third outcome i.e. mean change BCVA, again data from two time-points were used. Weighted mean and SDs for each treatment arm was calculated using mean and SDs of two time-points.

Data synthesis and model implementation

Bayesian network meta-analysis [37 38] (NMA) was used to compare the different interventions. Network meta-analysis is a generalization of meta-analysis methods because they allow comparisons of agents not addressed within individual primary trials. Bayesian statistical inference provides probability distributions for treatment effect parameters (RR and WMD), with 95% credible intervals (95% CrI), rather than 95% confidence intervals (95% CI). A 95% credible interval can be interpreted as there being a 95% probability that the parameter takes a value in the specified range.[37 38]

All analyses were conducted using a Bayesian Markov Chain Monte Carlo (MCMC) method and fitted in the freely available Bayesian software, WinBUGS 1.4.3.[39] Two Markov chains were run simultaneously using different initial values. Convergence to a stable solution was checked by viewing plots of the sampled simulations and using the Brooks-Gelman-Rubin diagnostic tool.[40] Convergence was found to be adequate after running 20 000 samples for both chains. These samples were then discarded and a further 70 000 sampled simulation was then run, on which the results were based. We also calculated the probability of treatment being the most effective (first best), the second best, the third best, and so on, and presented the results graphically with rankograms.[41]

Like standard meta-analysis comparison, a NMA can be either a fixed- or a random-effect models. We used the Bayesian Deviation Information Criterion (DIC) to compare fixed and random effect models. The most appropriate NMA model can be identified as the one with the lowest DIC. The DIC measures the fit of the model while penalizing it for the number of effective parameters. The fixed - effect model was chosen because of the small number of trials available for each comparison and difficulty in estimating between studies variance if random-effect model was implemented and the difference in DIC is less than 5.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Study selection and characteristics

The literature search identified 945 articles, as shown in Figure 1. Seven studies were judged to be sufficiently comparable to be included in the network meta-analysis. Tables 1 and 2 present the characteristics and results of the included trials. Two studies [11-13] compared aflibercept 2 mg against sham; two identical studies [6-8] compared dexamethasone 0.7 mg (Ozurdex) against sham; one study [9 10] compared ranibizumab 0.5 mg against sham; one study [42-44] compared bevacizumab 1.25 mg against sham, and finally one study [23-36] compared triamcinolone 4 mg against observation. Sham or observation were used as the common comparator. The number of included participants varied from 60 [42-44] to 437 [6-8]. Most studies required patients to be treatment naive and have macular oedema with retinal thickness measuring at least 250 or 300 µm on optical coherence tomography. Sham injection was undertaken by placing a needleless syringe onto the eye. All studies, except for Epstein and colleagues 2012[42-44], were multi-centre, international studies. Most studies had an extension phase after the primary outcome, but this was not included in the network meta-analysis.

The sufficiently comparable studies were combined into a network analysis based on a common comparator. The network for the proportions of patients gaining more than or equal to 3 lines is shown in Figure 2. This network is the same for the other two outcomes, but without dexamethasone because the trial did not report these outcomes.

Risk of bias of included trials

Risk of bias is shown in Table 3. Included studies were generally of high quality, with all studies being judged to be of low or unclear bias for all criteria. The non-commercially

funded bevacizumab trial had fewer patients and inevitably results had wider confidence intervals.[42-44] In no study does it appear that patients were asked at the end of the trial what arm they thought they had been assigned. It is unclear how many could distinguish injections (intervention arm) from punctureless pressure (sham arm).

Effects of interventions on proportions of patients gaining ≥ 3 lines

Figure 3 displays a forest plot of the risk ratio and 95% credible interval in proportions of patients gaining more than or equal to 3 lines for all the possible pairwise comparisons. In terms of proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of being more effective than a sham and dexamethasone (Figure 4). There was no difference in the proportions of patients gaining more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg.

Effects of interventions on proportions of patients losing ≥ 3 lines

Figure 5 displays forest plot of the risk ratio and 95% credible interval of proportions of patients losing more than or equal to 3 lines for all the possible pairwise comparisons. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision than those treated with sham. There was no difference in the proportions of patients losing more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25 mg and aflibercept 2mg. Figure 6 shows ranking for efficacy in terms of proportions of patients losing ≥ 3 lines.

Effects of interventions on mean change in BCVA

Figure 7 displays a forest plot of the mean changes and 95% credible intervals of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

improvement in BCVA for all the possible pairwise comparisons. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of improvement in BCVA compared to those treated with sham injections. Patients treated with aflibercept 2mg had a higher probability of improvement in BCVA compared with those treated with triamcinolone 4mg (Figure 8). There was no difference in mean change in BCVA from baseline between patients treated with ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2 mg.

For peer review only

Discussion

Statement of principal findings

Our results show no evidence of a difference in effectiveness between aflibercept, ranibizumab and triamcinolone. Bevacizumab was similar to these drugs in terms of letters gained and mean change in BCVA. Dexamethasone was less effective compared to these drugs.

Strengths and limitations

This is the first study providing an indirect comparison of drugs to treat macular oedema secondary to CRVO. A robust search strategy, screening process and data extraction was used, and this analysis drew on a systematic review. The studies included had, in general, a low risk of bias. Safety was not considered in this study but is described in detail elsewhere.[45] Five different drugs were suitable for network meta-analysis. Unpublished data was obtained from one author. [42-44] Bayesian methods were used for the NMA. There was good model fit and convergence within the analysis.

However pre-specified outcomes were not reported in all studies and the sample size varied considerably. For example Epstein 2012, assessing bevacizumab, only included 30 participants in each arm.[42-44] This resulted in wide credible intervals from the network meta-analysis which may lead to a type 1 error especially with regards to the proportions of patients losing more than or equal to 3 lines. The SCORE study compared triamcinolone to observation.[23-36] The NMA assumes a [11] similar effect of sham and observation and this may result in a small degree of bias. Only six months of data was included, and the long term effects are not known. Using a six-month follow-up period may disadvantage dexamethasone because peak effect in the GENEVA trials was seen at 90 days, and by six months, benefits had been largely lost.[6-8]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

As with most network meta-analyses, methodological heterogeneity was present. There were some differences amongst the trials. For example CRUISE[9 10], assessing ranibizumab, did not include as many patients with ischaemic CRVO as the aflibercept trials.[12 13] There were also some small differences in the chronicity of macular oedema and the mean BCVA at baseline.

Meaning of the study: possible explanations and implications for clinicians and policymakers

No head-to-head trials comparing aflibercept, bevacizumab, ranibizumab, triamcinolone or dexamethasone have been published in central retinal vein occlusion. Part of the reason for this is that the Food and Drug Administration require proof of the safety and effectiveness of a drug.[46] The easiest and quickest method for pharmaceutical companies to produce this is through placebo controlled trials. Trials comparing new medications to current best treatment would be considerably more useful to clinicians and patients.

Head-to-head trials comparing some of these drugs are available in other conditions. For example a comparison of ranibizumab and bevacizumab was undertaken in age related macular degeneration in the Comparison of Age-related macular degeneration Treatment Trials (CATT)[47] and alternative treatments to Inhibit VEGF in patients with Age-related choroidal Neovascularisation (IVAN)[48] trials. Both of these trials found no difference in effectiveness between ranibizumab and bevacizumab. Furthermore an indirect comparison of ranibizumab and bevacizumab found no evidence of a difference between these drugs.[49] Thus, it is highly probable that this may also apply in CRVO. The difference seen in our results regarding bevacizumab may be due to the low number of patients included in Epstein 2012.[42-44] In the CATT trial, more patients were

hospitalized in the bevacizumab arm, but the authors did not believe that this was explained by a direct effect of bevacizumab.[47] The 2-year results from the IVAN showed little difference in cardiovascular events, with the number being insignificantly lower with bevacizumab.[50] Ranibizumab and aflibercept were directly compared in two similarly designed trials, VEGF Trap-eye: investigation of Efficacy and safety in Wet age-related macular degeneration (VIEW 1 and 2).[51] Similar efficacy and safety was found in both drugs.

From the included trials it is clear that intraocular steroids are associated with complications, including increased intra-ocular pressure and cataract formation.[6-8 23-36] These are substantial drawbacks for using steroids to treat macular oedema in CRVO. However, many affected patients may be already pseudophakic and, on these, the use of intraocular steroids may be reasonable. Steroids may have a place in the treatment pathway of patients who have failed on anti-VEGF therapy, but this has yet to be tested. The anti-VEGF drugs have a good safety profile and do not cause cataract formation.[9-13 42-44] For this reason are likely to be more favoured by clinicians than steroids.

Aflibercept, compared with ranibizumab and bevacizumab, targets a wider range of cytokines and may have a stronger binding affinity.[52] Initial results suggested that aflibercept would require fewer injections than ranibizumab.[51] Heier and colleagues compared aflibercept and ranibizumab in two similarly designed randomised controlled trials in age related macular degeneration. They found that 2 mg aflibercept administered every eight weeks produced similar effects at 96 weeks to 0.5 mg ranibizumab every four weeks.[51] This was reflected in the FDA Dermatologic and Ophthalmic Drugs Advisory Committee recommendation that aflibercept should be given every two months following three initial monthly doses in age related macular

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

oedema.[53] This may be because aflibercept also appears to last longer in the eye than ranibizumab.[54] Age related macular degeneration is a more aggressive condition than central retinal vein occlusion and so it is unlikely that more frequent dosing would be needed. Therefore aflibercept may be preferred because it would reduce pressure on out-patient clinics. Furthermore there is some evidence from patients with age-related macular degeneration that aflibercept may be effective in patients who have not responded to ranibizumab.[55 56] This may be due to the higher affinity and wider number of cytokines that are targeted. There is no reason to suspect that these effects be any different for the macular oedema caused by central retinal vein occlusion. However we have as yet no evidence as to whether ranibizumab would be effective after aflibercept has failed.

The National Institute of Health and Care Excellence has recommended dexamethasone, ranibizumab and aflibercept as options in the treatment of macular oedema secondary to CRVO[18-20]. Until these technologies are reviewed together and compared with each other, clinicians are left with three recommended drugs. It should be noted that during the appraisal of ranibizumab the evidence review group found that in the cost-effectiveness analysis dexamethasone was extendedly dominated by ranibizumab (an intervention is judged not be cost-effective because it has an ICER that is greater than that that of a more effective intervention). The committee appraising ranibizumab did not re-consider the previous appraisal decision on dexamethasone.

Our results show that dexamethasone was not as effective as ranibizumab or aflibercept, at six months follow-up and with the dosing regimens in the trials. However these results do not assess quality of life or cost effectiveness. Bevacizumab is likely to prove more cost effective than both aflibercept and ranibizumab because it costs substantially less.[57] However the National Institute for Health and Care Excellence has not issued

guidance on bevacizumab because it does not have a license for use in the eye.

Unanswered questions and future research

Not all patients benefit from the use of anti-VEGF drugs; only about 60% gain 15 or more letters. It is not clear why some patients benefit more than others. Future research should focus on identifying subgroups of patients who are likely to benefit. Only a few of these trials included ischaemic patients, and in these trials only a few patients with ischaemia were included.[11-13] More research assessing the effectiveness of these drugs in severely ischaemic patients is needed.

Head-to-head trials comparing ranibizumab, aflibercept, bevacizumab and triamcinolone are needed. These should include assessment of cost effectiveness. To assist this, a better measure of quality of life is needed for patients with eye conditions. The widely-used EQ5D may not be sensitive enough to measure changes which are important to patients, such as the ability to drive.

In conclusion, we have found no evidence of difference between ranibizumab, bevacizumab, aflibercept and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

We thank Christine Clar, Sian Thomas and Rachel Court for assisting with searches, screening and data extraction for the systematic review which precede this study. We thank the authors of the Epstein 2012 trial for providing addition data.

Copyright statement

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above."

Declaration of competing interests

"All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work."

Contribution statement

NW conceived the idea. All authors contributed to the design of the study. DS and OU undertook the statistical analysis. JF, DS and OU wrote the first draft of the manuscript. All authors redrafted and agreed the final article. JF is the guarantor.

Transparency statement

JF affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

Data Sharing Statement

No additional data available

Ethics approval – not required

Funding – no external funding required

References

1. Deramo VA, Cox TA, Syed AB, et al. Vision-related quality of life in people with central retinal vein occlusion using the 25-item National Eye Institute Visual Function Questionnaire. Arch.Ophthalmol. 2003;**121**(9):1297-302 doi: 10.1001/archopht.121.9.1297 [doi];121/9/1297 [pii][published Online First: Epub Date]].
2. Klein R, Moss SE, Meuer SM, et al. The 15-year cumulative incidence of retinal vein occlusion: the Beaver Dam Eye Study. Archives of ophthalmology 2008;**126**(4):513-8 doi: 10.1001/archopht.126.4.513[published Online First: Epub Date]].
3. Hayreh SS, Podhajsky PA, Zimmerman MB. Natural history of visual outcome in central retinal vein occlusion. Ophthalmology 2011;**118**(1):119-33 doi: S0161-6420(10)00447-1 [pii];10.1016/j.ophtha.2010.04.019 [doi][published Online First: Epub Date]].
4. McIntosh RL, Rogers SL, Lim L, et al. Natural history of central retinal vein occlusion: an evidence-based systematic review. Ophthalmology 2010;**117**(6):1113-23 doi: S0161-6420(10)00133-8 [pii];10.1016/j.ophtha.2010.01.060 [doi][published Online First: Epub Date]].
5. The Royal College of Ophthalmologists. Interim Guidelines for Management of Retinal Vein Occlusion. Secondary Interim Guidelines for Management of Retinal Vein Occlusion 2010.
http://www.rcophth.ac.uk/core/core_picker/download.asp?id=728&file_title=Interim+Guidelines+for+Management+of+Retinal+Vein+Occlusion+2010.

6. Haller JA, Bandello F, Belfort R, Jr., et al. Randomized, sham-controlled trial of dexamethasone intravitreal implant in patients with macular edema due to retinal vein occlusion. *Ophthalmology* 2010;**117**(6):1134-46
7. Haller JA, Bandello F, Belfort R, Jr., et al. Dexamethasone intravitreal implant in patients with macular edema related to branch or central retinal vein occlusion twelve-month study results. *Ophthalmology* 2011;**118**(12):2453-60
8. Yeh WS, Haller JA, Lanzetta P, et al. Effect of the duration of macular edema on clinical outcomes in retinal vein occlusion treated with dexamethasone intravitreal implant. *Ophthalmology* 2012;**119**(6):1190-98
9. Brown DM, Campochiaro PA, Singh RP, et al. Ranibizumab for macular edema following central retinal vein occlusion: six-month primary end point results of a phase III study. *Ophthalmology* 2010;**117**(6):1124-33
10. Campochiaro PA, Brown DM, Awh CC, et al. Sustained benefits from ranibizumab for macular edema following central retinal vein occlusion: twelve-month outcomes of a phase III study. *Ophthalmology* 2011;**118**(10):2041-49
11. Holz FG, Roeder J, Ogura Y, et al. VEGF Trap-Eye for macular oedema secondary to central retinal vein occlusion: 6-month results of the phase III GALILEO study. *British Journal of Ophthalmology* 2013;**97**(3):278-84
12. Boyer D, Heier J, Brown DM, et al. Vascular endothelial growth factor Trap-Eye for macular edema secondary to central retinal vein occlusion: six-month results of the phase 3 COPENICUS study. *Ophthalmology* 2012;**119**(5):1024-32

13. Brown DM, Heier JS, Clark WL, et al. Intravitreal Aflibercept Injection for Macular Edema Secondary to Central Retinal Vein Occlusion: 1-Year Results From the Phase 3 COPENICUS Study. *American Journal of Ophthalmology* 2013;**155**(3):429-37
14. Michels S, Rosenfeld PJ, Puliafito CA, et al. Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration twelve-week results of an uncontrolled open-label clinical study. *Ophthalmology* 2005;**112**(6):1035-47 doi: 10.1016/j.ophtha.2005.02.007[published Online First: Epub Date]].
15. Arevalo JF, Fromow-Guerra J, Quiroz-Mercado H, et al. Primary intravitreal bevacizumab (Avastin) for diabetic macular edema: results from the Pan-American Collaborative Retina Study Group at 6-month follow-up. *Ophthalmology* 2007;**114**(4):743-50 doi: 10.1016/j.ophtha.2006.12.028[published Online First: Epub Date]].
16. Rabena MD, Pieramici DJ, Castellarin AA, et al. Intravitreal bevacizumab (Avastin) in the treatment of macular edema secondary to branch retinal vein occlusion. *Retina (Philadelphia, Pa.)* 2007;**27**(4):419-25 doi: 10.1097/IAE.0b013e318030e77e[published Online First: Epub Date]].
17. Algvere PV, Epstein D, von Wendt G, et al. Intravitreal bevacizumab in central retinal vein occlusion: 18-month results of a prospective clinical trial. *European journal of ophthalmology* 2011;**21**(6):789-95 doi: 10.5301/ejo.2011.6522[published Online First: Epub Date]].
18. National Institute for Health and Care Excellence. Ranibizumab for treating visual impairment caused by macular oedema secondary to retinal vein

occlusion: NICE technology appraisal guidance 283. 2013.
<http://guidance.nice.org.uk/TA283/Guidance/pdf/English>.

19. National Institute for Health and Clinical Excellence. Dexamethasone intravitreal implant for the treatment of macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 229. 2011.
<http://guidance.nice.org.uk/TA229/Guidance/pdf/English>.

20. National Institute for Health and Care Excellence. Aflibercept for treating visual impairment caused by macular oedema secondary to central retinal vein occlusion. NICE technology appraisal guidance 305. 2014.
<http://guidance.nice.org.uk/TA305>.

21. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. BMJ (Clinical research ed.) 2011;**343**:d5928 doi: 10.1136/bmj.d5928[published Online First: Epub Date]].

22. Hozo S, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. BMC Medical Research Methodology 2005;**5**(1):13

23. Bhavsar AR, Ip MS, Glassman AR, Groups DatSS. The risk of endophthalmitis following intravitreal triamcinolone injection in the DRCRnet and SCORE clinical trials. American Journal of Ophthalmology 2007;**144**(3):454-56

24. Blodi BA, Domalpally A, Scott IU, et al. Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study system for evaluation of stereoscopic color fundus photographs and fluorescein angiograms: SCORE Study Report 9. Archives of ophthalmology 2010;**128**(9):1140-45

25. Chan CK, Ip MS, VanVeldhuisen PC, et al. SCORE Study report #11: incidences of neovascular events in eyes with retinal vein occlusion. *Ophthalmology* 2011;**118**(7):1364-72
26. Ip M, Oden N, VanVeldhuisen P, Scott I, et al. The Standard Care vs. Corticosteroid for Retinal Vein Occlusion Study: Design and Baseline Characteristics. *American Academy of Ophthalmology* 2008:260
27. Ip MS, Oden NL, Scott IU, et al. SCORE Study report 3: study design and baseline characteristics. *Ophthalmology* 2009;**116**(9):1770-77
28. Ip MS, Scott IU, VanVeldhuisen PC, et al. A randomized trial comparing the efficacy and safety of intravitreal triamcinolone with observation to treat vision loss associated with macular edema secondary to central retinal vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) study report 5. *Archives of ophthalmology* 2009;**127**(9):1101-14
29. Myers D, Blodi B, Ip M, et al. Reading Center Evaluation of OCT Images From Patients Enrolled in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. *Iovs.* 2006;**47**:ARVO
30. Oden NL, Veldhuisen PC, Scott IU, et al. Temporal Variability of OCT in Retinal Vein Occlusion Participants in the SCORE Study. *Iovs.* 2007;**48**:ARVO
31. Scott IU, Blodi BA, Ip MS, et al. SCORE Study Report 2: Interobserver agreement between investigator and reading center classification of retinal vein occlusion type. *Ophthalmology* 2009;**116**(4):756-61
32. Scott IU, Oden NL, VanVeldhuisen PC, et al. SCORE Study Report 7: incidence of intravitreal silicone oil droplets associated with staked-on vs luer cone syringe design. *American Journal of Ophthalmology* 2009;**148**(5):725-32

33. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline predictors of visual acuity and retinal thickness outcomes in patients with retinal vein occlusion: Standard Care Versus COrticosteroid for REtinal Vein Occlusion Study report 10. *Ophthalmology* 2011;**118**(2):345-52
34. Scott IU, VanVeldhuisen PC, Oden NL, et al. SCORE Study report 1: baseline associations between central retinal thickness and visual acuity in patients with retinal vein occlusion. *Ophthalmology* 2009;**116**(3):504-12
35. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline Characteristics and Response to Treatment of Participants With Hemiretinal Compared With Branch Retinal or Central Retinal Vein Occlusion in the Standard Care vs COrticosteroid for REtinal Vein Occlusion (SCORE) Study: SCORE Study Report 14. *Archives of ophthalmology* 2012;**130**(12):1517-24
36. Warren K, Blodi BA, Oden N, et al. Reading Center Evaluation of Baseline Retinal Images in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. *Iovs.* 2008:ARVO
37. Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *BMJ (Clinical research ed.)* 2005;**331**(7521):897-900 doi: 10.1136/bmj.331.7521.897[published Online First: Epub Date]].
38. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. *Statistics in medicine* 2004;**23**(20):3105-24 doi: 10.1002/sim.1875[published Online First: Epub Date]].
39. Spiegelhalter D, Thomas A, Best N, et al. *WinBUGS User Manual: Version 1.4*. Cambridge, Mass: MRC Biostatistics Unit, 2003.

40. Brooks SP, Gelman A. General Methods for Monitoring Convergence of Iterative Simulations. *Journal of Computational and Graphical Statistics* 1998;**7**(4):434-55 doi: 10.1080/10618600.1998.10474787[published Online First: Epub Date]].
41. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;**64**(2):163-71 doi: 10.1016/j.jclinepi.2010.03.016[published Online First: Epub Date]].
42. Epstein D, Algvere P, Von WG, et al. Long-term benefit from bevacizumab for macular edema in central retinal vein occlusion: 12-month results of a prospective study. *Acta Ophthalmologica* 2012;**90**:48 doi: 10.1111/j.1755-3768.2012.02549.x[published Online First: Epub Date]].
43. Epstein DL, Algvere PV, Von WG, et al. Benefit from bevacizumab for macular edema in central retinal vein occlusion: twelve-month results of a prospective, randomized study. *Ophthalmology* 2012;**119**(12):2587-91
44. Epstein DL, Algvere PV, Von WG, S et al. Bevacizumab for macular edema in central retinal vein occlusion: a prospective, randomized, double-masked clinical study. *Ophthalmology* 2012;**119**(6):1184-89
45. Ford JA, Clar C, Lois N, et al. Treatments for macular oedema following central retinal vein occlusion: systematic review. *BMJ open* 2014;**4**(2):e004120 doi: 10.1136/bmjopen-2013-004120[published Online First: Epub Date]].
46. FDA. Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators. Secondary Drug Study Designs -

Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators 2011.

<http://www.fda.gov/RegulatoryInformation/Guidances/ucm126501.htm>

47. Martin DF, Maguire MG, Fine SL, et al. Ranibizumab and bevacizumab for treatment of neovascular age-related macular degeneration: two-year results. *Ophthalmology* 2012;**119**(7):1388-98 doi: 10.1016/j.opthta.2012.03.053[published Online First: Epub Date]].
48. Chakravarthy U, Harding SP, Rogers CA, et al. Ranibizumab versus bevacizumab to treat neovascular age-related macular degeneration: one-year findings from the IVAN randomized trial. *Ophthalmology* 2012;**119**(7):1399-411 doi: S0161-6420(12)00358-2 [pii];10.1016/j.opthta.2012.04.015 [doi][published Online First: Epub Date]].
49. Ford JA, Elders A, Shyangdan D, et al. The relative clinical effectiveness of ranibizumab and bevacizumab in diabetic macular oedema: an indirect comparison in a systematic review. *BMJ (Clinical research ed.)* 2012;**345**:e5182
50. Chakravarthy U, Harding SP, Rogers CA, et al. Alternative treatments to inhibit VEGF in age-related choroidal neovascularisation: 2-year findings of the IVAN randomised controlled trial. *Lancet* 2013 doi: 10.1016/s0140-6736(13)61501-9[published Online First: Epub Date]].
51. Heier JS, Brown DM, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology*

- 2012;**119**(12):2537-48 doi: 10.1016/j.opthta.2012.09.006[published Online First: Epub Date]].
52. Papadopoulos N, Martin J, Ruan Q, et al. Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* 2012;**15**(2):171-85 doi: 10.1007/s10456-011-9249-6[published Online First: Epub Date]].
53. FDA. VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee. Secondary VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee 2011. <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/DermatologicandOphthalmicDrugsAdvisoryCommittee/UCM259143.pdf>.
54. Stewart MW, Rosenfeld PJ. Predicted biological activity of intravitreal VEGF Trap. *The British journal of ophthalmology* 2008;**92**(5):667-8 doi: 10.1136/bjo.2007.134874[published Online First: Epub Date]].
55. Bakall B, Folk JC, Boldt HC, et al. Aflibercept therapy for exudative age-related macular degeneration resistant to bevacizumab and ranibizumab. *Am J Ophthalmol* 2013;**156**(1):15-22.e1 doi: 10.1016/j.ajo.2013.02.017[published Online First: Epub Date]].
56. Cho H, Shah CP, Weber M, et al. Aflibercept for exudative AMD with persistent fluid on ranibizumab and/or bevacizumab. *The British journal of ophthalmology* 2013;**97**(8):1032-5 doi: 10.1136/bjophthalmol-2013-303344[published Online First: Epub Date]].
57. Raftery J, Clegg A, Jones J, et al. Ranibizumab (Lucentis) versus bevacizumab (Avastin): modelling cost effectiveness. *The British journal of*

ophthalmology 2007;**91**(9):1244-6 doi:
10.1136/bjo.2007.116616[published Online First: Epub Date]].

For peer review only

Figure Legend

Figure 1: study selection flow diagram

Figure 2: Network of randomized controlled trials comparing different treatments for proportions of gaining 3 or more lines of vision

Figure 3: Proportions of patients gaining 3 lines or more from baseline to six months

Figure 4: Rankogram for gaining ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions

Figure 5: Proportions of patients losing 3 lines or more from baseline to six months

Figure 6: Rankogram for losing ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions

Figure 7: Mean BCVA change from baseline to 6 months.

Figure 8: Rankogram for mean change in BCVA - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions

Table 1: Baseline characteristics and results of all included studies

Study	Participants	Intervention / Outcomes
DEXAMETHASONE		
GENEVA 2010[6-8] International	N: CRVO – 437 eyes of 437 patients randomised; 94% follow-up at 6 months	1. Dexamethasone 0.7 mg (n=136) Single dose
Setting: multicentre (167 centres in 24 countries, so a mean of 2.6 patients per centre)	Participants: adults with visual acuity reduced because of macular oedema due to CRVO or BRVO	2. Dexamethasone 0.35 mg (n=154) Single dose
Design: 2 identical double-blind, sham-controlled RCTs, phase 3		3. Sham (n=147) Single dose - a needleless applicator was placed against the conjunctiva to simulate the placement of study medication.
Follow-up: primary endpoint for the masked trial: 6 months; primary endpoint for the open-label extension: 12 months		Primary end point: gain of ≥15 ETDRS letters; for the open-label extension: safety

TRIAMCINOLONE		
SCORE 2009[23-36] USA Setting: multicentre Design: RCT Follow-up: primary end point 12 months, FU planned up to 36 months	N: 271 eyes of 271 patients randomised; 83% (observation) and 90% (triamcinolone) completed 12 months Participants: centre-involved macular oedema secondary to CRVO	1. Triamcinolone 1 mg (n=92) Every 4 months depending on retreatment regimen (ave 2.2 injections at 12 months) 2. Triamcinolone 4 mg (n=91) Every 4 months depending on retreatment regimen (ave 2.0 injections at 12 months) (The form of triamcinolone used was Trivaris, no longer available. It was made by the manufacturer of Ozurdex (Allergan)) 3. Observation (n=88) Primary end point: gain of ≥ 15 ETDRS letters
AFLIBERCEPT		
COPERNICUS 2012[12 13] International Setting: multicentre, 70 sites in North and South America, India and Israel. Mean 2.7 patients per centre. Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU 2 years	N: 189 eyes of 189 patients randomised; 95.7% (aflibercept) and 81.1% (sham) completed 24 weeks; 93% (aflibercept) and 77% (sham) completed 52 weeks Participants: adult patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=114) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=73) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥ 15 ETDRS letters

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

GALILEO 2012[11] International Setting: multicentre, 10 countries in Europe and Asia; 63 centres in total Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU up to 12 months, planned up to 76 weeks	N: 177 eyes of 177 patients randomised; 90.6% (aflibercept) and 78.9% (sham) completed 24 weeks Participants: treatment-naïve patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=103) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=71) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥15 ETDRS letters
--	--	--

RANIBIZUMAB		
CRUISE 2010[9 10] USA Setting: multicentre Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 6 months, FU up to 12 months	N: 392 eyes of 392 patients randomised; 97.7% (ranibizumab 0.3 mg), 91.5% (ranibizumab 0.5 mg), and 88.5% (sham) completed 6 months Participants: patients with foveal centre-involved macular oedema secondary to CRVO diagnosed within 12 months	1. Ranibizumab 0.3 mg (n=132) Every 4 weeks for 6 months (ave number not available) 2. Ranibizumab 0.5 mg (n=130) Every 4 weeks for 6 months (ave number not available) 3. Sham (n=130) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to the injection site) Primary end point: mean change from baseline BCVA
BEVACIZUMAB		
Epstein 2012[42-44] Sweden Setting: Single centre; St. Eriks Eye Hospital Stockholm Design: sham-injection controlled, double masked RCT Follow-up: primary end-point 6 months; open label extension up to 12 months	N: 60 eyes of 60 patients randomised; 93% completed open label extension Participants: patients with CRVO of ≤ 6 months	1. Bevacizumab 1.25 mg (n=30) Every 6 weeks for 6 months (ave number not available) 2. Sham (n=30) Every 6 weeks for 6 months (ave number not available) (syringe without needle pressed to the globe) Primary end point: gain of ≥ 15 ETDRS letters

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

FU= follow-up, RCT = randomised controlled trial, N = number, CRVO = central retinal vein occlusion, ETDRS = Early Treatment Diabetic Retinopathy Study, BRVO = branch retinal vein occlusion

For peer review only
Confidential

Table 2: Baseline characteristics and results of included trials

	COPERNICUS[12 13]	GALILEO[11]	CRUISE[9 10]	GENEVA[6-8]	Epstein et al (2012)[42-44]	SCORE[23-36]
BASELINE SIMILARITIES						
Number (%) of patients						
	Aflib 2 mg: 114 Sham: 73	Aflib 2 mg: 103 Sham: 68	Rani 0.5 mg: 130 Sham: 130	Dexa 0.7 mg: 136 Sham: 147	Beva 1.25 mg: 30 Sham: 30	Triam 4 mg: 91 Obser: 88
Age (years)						
	Aflib 2 mg: 65.5 SD13.6 Sham: 67.5 SD14.3	Aflib 2 mg: 59.9 SD12.4 Sham: 63.8 SD13.3	Rani 0.5 mg: 67.6 SD12.4 Sham: 65.4 SD13.1	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 70.6 SD 12.6 Sham: 70.4 SD 10.4	Triam 4 mg: 67.5 SD 12.0 Obser: 69.2 SD 12.8
BCVA at baseline (SD)						
	Aflib 2 mg: 50.7 SD13.90 Sham: 48.9 SD14.42	Aflib 2 mg: 53.6 SD15.8 Sham: 50.9 SD15.4	Rani 0.5 mg: 48.1 SD14.6 Sham: 49.2 SD14.7	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 44.4 SD 15.3 Sham: 43.6 SD 16.0	Triam 4 mg: 51.0 SD 14.4 Obser: 52.1 SD 13.1
Duration of MO from diagnosis to screening						
	Aflib 2 mg: 2.73 SD3.09(in months) Sham: 1.88 SD2.19 (in months)	Aflib 2 mg: 50.9 SD15.4)(in days) Sham: 87.6 SD79.1 (in days)	Rani 0.5 mg: - Sham: -	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: NR Sham: NR	Triam 4 mg: 4.2 SD 3.6 (in months) Obser: 4.2 SD 3.1 (in months)
RESULTS						
Number (%) of patients gaining ≥ 15 letters improvement from baseline to 6 months						
	Aflib 2 mg: 64 (56.1)	Aflib 2 mg: 62 (60.2)	Rani 0.5 mg: 62 (47.7)	Dexa 0.7 mg: 25 (18)	Beva 1.25 mg: 18 (60%)	Triam 4 mg: 18 (19.5%) (avg of 4 and 8 mths)

	Sham: 9 (12.3)	Sham: 15 (22.1)	Sham: 22 (16.9)	Sham: 18 (12)	Sham: 6 (20%)	Obser: 3 (4%) (avg of 4 and 8 mths)
Number (%) of patients losing ≥15 letters of BCVA from baseline to 6 months						
	Aflib 2 mg: 2 (1.8)	Aflib 2 mg: 8 (7.8)	Rani 0.5 mg: 2 (1.5)	Dexa 0.7 mg: NR	Beva 1.25 mg: 2 (6.7%)	Triam 4 mg: 19 (20.5%) (avg of 4 and 8 mths)
	Sham: 20 (27.4)	Sham: 15 (22.1)	Sham: 20 (15.4)	Sham: NR	Sham: 7 (23.3%)	Obser: 31 (35.5%) (avg of 4 and 8 mths)
Mean change (SD) from baseline in BCVA						
	Aflib 2 mg: 17.3 (12.8)	Aflib 2 mg: 18.0 (12.2)	Rani 0.5 mg: 14.9 (13.2)	Dexa 0.7 mg: 0.1 (NR)	Beva 1.25 mg: 14.1 SD 18.7	Triam 4 mg: -0.15 SD20.67 (n=85) (weight mean and SD of 4 and 8 months)
	Sham: -4 (18)	Sham: 3.3 (14.1)	Sham: 0.8 (16.2)	Sham: -1.8 (NR)	Sham: -2.0 SD 20.5	Obser: -9.66 SD18.04 (n=75) (weighted mean and SD of 4 and 8 months)

NR = not reported, Aflib = aflibercept, Rani = ranibizumab, Dexa = dexamethasone, Triam = triamcinolone, Obser = observation, SD = standard deviation, avg = average

Table 3: Risk of bias

Study (author and year)	Adequate sequence generation	Allocation concealment	Masking	Incomplete outcome data addressed	Free of selective reporting	Free of other bias (e.g. similarity at baseline, power assessment)	Funder
GENEVA 2010[6-8]	Low	Low	Partial: patients and assessors of efficacy variables	Low: ITT analysis, 94% FU at 6 months	Low	<i>Power</i> : 81% power to detect difference in primary outcome with n=495 for each trial <i>Similarity at baseline</i> : yes	Allergan Inc.
SCORE 2009[23-36]	Low	Unclear	Partial (physicians and patients masked to dose but not triamcinolone versus observation)	Low: ITT analysis, 83 to 90% FU at 12 months	Low	<i>Power</i> : 80% power to detect difference in primary outcome with n=486 (but only 271 randomised) <i>Similarity at baseline</i> : yes	National Eye Institute grants, Allergan
COPERNICUS 2012[12 13]	Low	Unclear	Low: double-blind	Low: ITT analysis, 89.9% assessed at primary end point	Low	<i>Power</i> : 90% power to detect difference in primary outcome with n=165 <i>Similarity at baseline</i> : yes	Bayer HealthCare, Regeneron Pharmaceuticals
GALILEO 2012[11]	Unclear	Unclear	Low: double-blind	Low: ITT analysis, 86% assessed at primary end point	Low	<i>Power</i> : 90% power to detect difference in primary outcome with n=150 <i>Similarity at baseline</i> : yes	Bayer HealthCare, Regeneron Pharmaceuticals

CRUISE 2010[9 10]	Low	Unclear	Low: patients and evaluating examiners, injecting physicians masked to dose	Low: ITT analysis, 88.5 to 97.7% completed 6 months	Low	Power: not reported Similarity at baseline: yes	Genentech Inc.
Epstein 2012[42-44]	Unclear	Low	Low: patients, outcome assessors	Low: ITT analysis; missing data for 2 patients (primary endpoint)	Low	Power: 80% power to detect difference in primary outcome with n=24 per group Similarity at baseline: yes	Unclear; authors are consultants for Allergan, Novartis, Alcon, Bayer

ITT= intention to treat, FU = follow-

For peer review only

Confidential

Drug treatment of macular oedema secondary to central retinal vein occlusion: a network meta-analysis

John A Ford, Deepson Shyangdan, Olalekan A. Uthman, Noemi Lois, Norman Waugh

John A Ford, Academic Clinical Fellow in Public Health, Department of Population Health and Primary Care, University of East Anglia
Deepson Shyangdan, Research Fellow, Warwick Evidence, University of Warwick
Olalekan Uthman, Assistant Professor in Applied Research, University of Warwick
Noemi Lois, Professor of Ophthalmology, Centre for Vision and Vascular Science, Queen’s University
Norman Waugh, Professor of Health Technology Assessment, University of Warwick

Corresponding author
Dr. John Ford
Norwich Medical School
University of East Anglia
Chancellor Drive
Norwich
NR4 7TJ
john.ford@uea.ac.uk
Tel: 01603 591269

Field Code Changed

Protocol – no published protocol exists for this study.

Ethics approval – not required

Funding – no external funding required

Data sharing: no additional data available

Word count: 3,331 words

What is already known on this subject

Anti-VEGF drugs (ranibizumab, bevacizumab and aflibercept) and corticosteroids (dexamethasone and triamcinolone), given intravitreally, have all been shown to be effective compared to placebo for the treatment of macular oedema secondary to central retinal vein occlusion.

There are no head-to-head trials.

What this study adds

There was no evidence of a difference in the effectiveness of aflibercept, ranibizumab, bevacizumab and triamcinolone for improving vision.

Clinicians may prefer aflibercept because steroids are associated with cataract formation and ranibizumab might require more frequent injections.

Abstract

Objective: To indirectly compare aflibercept, bevacizumab, dexamethasone, ranibizumab and triamcinolone for treatment of macular oedema secondary to central retinal vein occlusion using a network meta-analysis.

Design: Network meta-analysis

Data sources: The following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE; CDSR, DARE, HTA, NHSEED, CENTRAL; Science Citation Index and Conference Proceedings Citation Index-Science

Eligibility criteria for selecting studies: Only randomized controlled trials assessing patients with macular oedema secondary to central retinal vein occlusion were included. Studies had to report either proportions of patients gaining more than or equal to 3 lines, losing more than or equal to 3 lines, or mean change in best corrected visual acuity. Two authors screened titles and abstracts, extracted data and undertook risk of bias assessment. Bayesian network meta-analysis was used to compare the different interventions.

Results: Seven studies, assessing five drugs, were judged to be sufficiently comparable for inclusion in the NMA. For the proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2mg had a higher probability of being more effective than sham and dexamethasone. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision compared to those treated with sham. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab

1.25mg and aflibercept 2mg had a higher probability of improvement in mean best correct visual acuity compared to those treated with sham injections.

Conclusions: We found no evidence of differences between ranibizumab, aflibercept, bevacizumab and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections

Systematic review registration – Not registered

Strengths and limitations of this study

- Important topic area, with significant policy implications
- Robust method used to identify studies
- Network meta-analysis are based on a number of assumptions
- Network meta-analysis is the best method to compare interventions in the absence of head to head trials

Introduction

Central retinal vein occlusion (CRVO) dramatically reduces an individual's functioning and quality of life.[1] It is estimated that the 15 year cumulative incidence of central retinal vein occlusion is 0.5%.[2] Visual loss is caused by thrombosis of the central retinal vein which leads to a rise in venous pressure and an increase in vascular endothelial growth factor (VEGF), consequently causing an increase in vascular permeability. Macular oedema subsequently ensues with varying degrees of ischaemia and neo-vascularisation. Although CRVO is generally classified as ischaemic or non-ischaemic, ischaemia should be regarded as a spectrum.[3] Cases with ischaemia carry a considerably worse prognosis as in around a third of them, neovascular glaucoma may develop; the most devastating complication of CRVO.[4]

CRVO is more common in older people with risk factors such as diabetes, hypertension or hyperlipidaemia, but can occur in young people with inflammatory disorders. Hayreh and colleagues in a 27-year cohort study found that only 13% of people with CRVO were under 45 years of age.[3] In 95% of cases CRVO affects only one eye.[3] However visual loss in this already co-morbid patient group significantly compounds their already impaired functioning and quality of life. Patients can lose confidence, struggle with daily activities and become increasingly dependent on friends and family.[1]

For many years, laser photocoagulation was the only effective therapeutic strategy that could be used in the management of patients with CRVO. It was only useful for reducing the risk of neovascular glaucoma, but not effective for the treatment of macular oedema in CRVO.[5] Over the past decade a number of drugs to treat macular oedema have been introduced, including the steroids, triamcinolone and dexamethasone, and the anti-VEGFs, ranibizumab, bevacizumab, pegaptanib and aflibercept. Dexamethasone,

ranibizumab and aflibercept have been assessed in large commercially funded trials.[6-13] Bevacizumab was originally developed as an anti-cancer drug and has been found to be effective in treating macular oedema secondary to age-related macular degeneration,[14] diabetic macular oedema, [15] branch retinal vein occlusion[16] and central retinal vein occlusion.[17] Like triamcinolone, bevacizumab is used off licence in the eye. Ranibizumab is a derived from the same parent molecule of the bevacizumab monoclonal antibody and was developed and commercially marketed specifically for use in the eye.

In the United Kingdom, the National Institute of Health and Care Excellence (NICE) has recommended the use of dexamethasone, ranibizumab and aflibercept for the treatment of macular oedema secondary to CRVO in separate appraisals[18-20] Therefore clinicians have three NICE-recommended treatments for CRVO without head-to-head trials or clear guidance on which one may be best for their patients. On this basis, the aim of this study was to indirectly compare in a network meta-analysis the clinical effectiveness of aflibercept, ranibizumab, bevacizumab, dexamethasone and triamcinolone for the treatment of macular oedema secondary to CRVO.

Methods

Information sources and search strategy

To identify suitable studies, initially for a systematic review of treatment of macular oedema after CRVO (submitted for publication) the following databases were searched from January 2005 to March 2013: MEDLINE, MEDLINE In-process, EMBASE (all via OVID); CDSR, DARE, HTA, NHSEED, CENTRAL (all via The Cochrane Library); Science Citation Index and Conference Proceedings Citation Index-Science (via Web of Knowledge). The MEDLINE search strategy is shown in appendix 1. This search strategy was modified for other databases. In addition to the bibliographic database searching, supplementary searches were undertaken to look for recent and unpublished studies in the WHO International Clinical Trials Registry Platform and ophthalmology conference websites (American Academy of Ophthalmology, Association for Research in Vision and Ophthalmology from 2010 to 2012).

Study selection

Only randomised controlled trials which included patients with macular oedema secondary to central retinal vein occlusion were included. It was acceptable for a study to include both branch retinal vein occlusion and central retinal vein occlusion provided that the central retinal vein occlusion group was reported separately. The following drugs were included: dexamethasone, triamcinolone, ranibizumab, bevacizumab and aflibercept. Pegaptanib was not included because it is not used routinely in clinical practice. Only doses which are used in clinical practice were included. Studies had to report at least one of the following outcomes: proportions of patients gaining more than or equal to 3 lines from baseline to six months, proportions of patients losing more than or equal to 3 lines from baseline to six months and mean change in best corrected visual acuity (BCVA) from baseline to six months

Risk of bias assessment

The Cochrane Collaboration's tool for assessing risk of bias was used.[21] The trials were graded (unclear, high or low risk of bias) based on: (i) sequence generation, (ii) allocation concealment, (iii) blinding of outcome assessor, (iv) incomplete outcome data, and (v) selective outcome reporting.

Study selection and data abstraction

Two authors independently assessed the eligibility and methodological quality of the studies identified during the literature search. Two authors extracted and compared the data. For each study identified that met the selection criteria, details on study design, study population characteristics, intervention, outcome measures, and study quality were extracted. Discrepancies were resolved by consensus through discussion. Studies were assessed for comparability based on the populations included, trial arms, outcome measures and duration of follow-up. Common comparators were identified from the trials and a network diagram was created.

Summary measures

The primary measures of treatment effects were relative risk (RR) for the proportions of patients gaining more than or equal to 3 lines of vision, proportions of patients losing more than or equal to 3 lines of vision and weighted mean difference (WMD) for mean change BCVA. We used the following methods to calculate standard deviations, when incompletely reported: (1) contact with the corresponding author; or (2) estimation of the standard deviation on the basis of the sample size, median, and range as suggested by Hozo and colleagues[22] or on the basis of the sample size and P value.

In one trial (SCORE),[23-36] six month data was not available because patients were followed up every four months. For the dichotomous outcomes i.e. proportions of patients gaining and losing ≥ 3 lines, we averaged four and eight month data to get the six months follow-up data. For the third outcome i.e. mean change BCVA, again data from two time-points were used. Weighted mean and SDs for each treatment arm was calculated using mean and SDs of two time-points.

Data synthesis and model implementation

Bayesian network meta-analysis [37 38] (NMA) was used to compare the different interventions. Network meta-analysis is a generalization of meta-analysis methods because they allow comparisons of agents not addressed within individual primary trials. Bayesian statistical inference provides probability distributions for treatment effect parameters (RR and WMD), with 95% credible intervals (95% CrI), rather than 95% confidence intervals (95% CI). A 95% credible interval can be interpreted as there being a 95% probability that the parameter takes a value in the specified range.[37 38]

All analyses were conducted using a Bayesian Markov Chain Monte Carlo (MCMC) method and fitted in the freely available Bayesian software, WinBUGS 1.4.3.[39] Two Markov chains were run simultaneously using different initial values. Convergence to a stable solution was checked by viewing plots of the sampled simulations and using the Brooks-Gelman-Rubin diagnostic tool.[40] Convergence was found to be adequate after running 20 000 samples for both chains. These samples were then discarded and a further 70 000 sampled simulation was then run, on which the results were based. We also calculated the probability of treatment being the most effective (first best), the second best, the third best, and so on, and presented the results graphically with rankograms.[41]

Like standard meta-analysis comparison, a NMA can be either a fixed- or a random-effect models. We used the Bayesian Deviation Information Criterion (DIC) to compare fixed and random effect models. The most appropriate NMA model can be identified as the one with the lowest DIC. The DIC measures the fit of the model while penalizing it for the number of effective parameters. The fixed - effect model was chosen because of the small number of trials available for each comparison and difficulty in estimating between studies variance if random-effect model was implemented and the difference in DIC is less than 5.

1

2

3

4

5

6 **Results**

7

8

9

10 **Study selection and characteristics**

11 The literature search identified 945 articles, as shown in Figure 1. Seven studies were

12 judged to be sufficiently comparable to be included in the network meta-analysis. Tables

13 1 and 2 present the characteristics and results of the included trials. Two studies [11-

14 13] compared aflibercept 2 mg against sham; two identical studies [6-8] compared

15 dexamethasone 0.7 mg (Ozurdex) against sham; one study [9 10] compared

16 ranibizumab 0.5 mg against sham; one study [42-44] compared bevacizumab 1.25 mg

17 against sham, and finally one study [23-36] compared triamcinolone 4 mg against

18 observation. Sham or observation were used as the common comparator. The number of

19 included participants varied from 60 [42-44] to 437 [6-8]. Most studies required

20 patients to be treatment naive and have macular oedema with retinal thickness

21 measuring at least 250 or 300 µm on optical coherence tomography. Sham injection was

22 undertaken by placing a needleless syringe onto the eye. All studies, except for Epstein

23 and colleagues 2012[42-44], were multi-centre, international studies. Most studies had

24 an extension phase after the primary outcome, but this was not included in the network

25 meta-analysis.

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41 The sufficiently comparable studies were combined into a network analysis based on a

42 common comparator. The network for the proportions of patients gaining more than or

43 equal to 3 lines is shown in Figure 2. This network is the same for the other two

44 outcomes, but without dexamethasone because the trial did not report these outcomes.

45

46

47

48

49

50 **Risk of bias of included trials**

51 Risk of bias is shown in Table 3. Included studies were generally of high quality, with all

52 studies being judged to be of low or unclear bias for all criteria. The non-commercially

funded bevacizumab trial had fewer patients and inevitably results had wider confidence intervals.[42-44] In no study does it appear that patients were asked at the end of the trial what arm they thought they had been assigned. It is unclear how many could distinguish injections (intervention arm) from punctureless pressure (sham arm).

Effects of interventions on proportions of patients gaining ≥ 3 lines

Figure 3 displays a forest plot of the risk ratio and 95% credible interval in proportions of patients gaining more than or equal to 3 lines for all the possible pairwise comparisons. In terms of proportions of patients gaining more than or equal to 3 lines, triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of being more effective than a sham and dexamethasone (Figure 4). There was no difference in the proportions of patients gaining more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg.

Effects of interventions on proportions of patients losing ≥ 3 lines

Figure 5 displays forest plot of the risk ratio and 95% credible interval of proportions of patients losing more than or equal to 3 lines for all the possible pairwise comparisons. A smaller proportions of patients treated with triamcinolone 4mg, ranibizumab 0.5mg or aflibercept 2mg lost more than or equal to 3 lines of vision than those treated with sham. There was no difference in the proportions of patients losing more than or equal to 3 lines between triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25 mg and aflibercept 2mg. Figure 6 shows ranking for efficacy in terms of proportions of patients losing ≥ 3 lines.

Effects of interventions on mean change in BCVA

Figure 7 displays a forest plot of the mean changes and 95% credible intervals of

improvement in BCVA for all the possible pairwise comparisons. Patients treated with triamcinolone 4mg, ranibizumab 0.5mg, bevacizumab 1.25mg, aflibercept 2mg had a higher probability of improvement in BCVA compared to those treated with sham injections. Patients treated with aflibercept 2mg had a higher probability of improvement in BCVA compared with those treated with triamcinolone 4mg (Figure 8). There was no difference in mean change in BCVA from baseline between patients treated with ranibizumab 0.5mg, bevacizumab 1.25mg and aflibercept 2 mg.

Discussion

Statement of principal findings

Our results show no evidence of a difference in effectiveness between aflibercept, ranibizumab and triamcinolone. Bevacizumab was similar to these drugs in terms of letters gained and mean change in BCVA. Dexamethasone was less effective compared to these drugs.

Strengths and limitations

This is the first study providing an indirect comparison of drugs to treat macular oedema secondary to CRVO. A robust search strategy, screening process and data extraction was used, and this analysis drew on a systematic review. The studies included had, in general, a low risk of bias. Safety was not considered in this study but is described in detail elsewhere.[45] Five different drugs were suitable for network meta-analysis. Unpublished data was obtained from one author. [42-44] Bayesian methods were used for the NMA. There was good model fit and convergence within the analysis.

However pre-specified outcomes were not reported in all studies and the sample size varied considerably. For example Epstein 2012, assessing bevacizumab, only included 30 participants in each arm.[42-44] This resulted in wide credible intervals from the network meta-analysis which may lead to a type 1 error especially with regards to the proportions of patients losing more than or equal to 3 lines. The SCORE study compared triamcinolone to observation.[23-36] The NMA assumes a [11] similar effect of sham and observation and this may result in a small degree of bias. Only six months of data was included, and the long term effects are not known. Using a six-month follow-up period may disadvantage dexamethasone because peak effect in the GENEVA trials was seen at 90 days, and by six months, benefits had been largely lost.[6-8]

As with most network meta-analyses, methodological heterogeneity was present. There were some differences amongst the trials. For example CRUISE[9 10], assessing ranibizumab, did not include as many patients with ischaemic CRVO as the aflibercept trials.[12 13] There were also some small differences in the chronicity of macular oedema and the mean BCVA at baseline.

Meaning of the study: possible explanations and implications for clinicians and policymakers

No head-to-head trials comparing aflibercept, bevacizumab, ranibizumab, triamcinolone or dexamethasone have been published in central retinal vein occlusion. Part of the reason for this is that the Food and Drug Administration require proof of the safety and effectiveness of a drug.[46] The easiest and quickest method for pharmaceutical companies to produce this is through placebo controlled trials. Trials comparing new medications to current best treatment would be considerably more useful to clinicians and patients.

Head-to-head trials comparing some of these drugs are available in other conditions. For example a comparison of ranibizumab and bevacizumab was undertaken in age related macular degeneration in the Comparison of Age-related macular degeneration Treatment Trials (CATT)[47] and alternative treatments to Inhibit VEGF in patients with Age-related choroidal Neovascularisation (IVAN)[48] trials. Both of these trials found no difference in effectiveness between ranibizumab and bevacizumab. Furthermore an indirect comparison of ranibizumab and bevacizumab found no evidence of a difference between these drugs.[49] Thus, it is highly probable that this may also apply in CRVO. The difference seen in our results regarding bevacizumab may be due to the low number of patients included in Epstein 2012.[42-44] In the CATT trial, more patients were

hospitalized in the bevacizumab arm, but the authors did not believe that this was explained by a direct effect of bevacizumab.[47] The 2-year results from the IVAN showed little difference in cardiovascular events, with the number being insignificantly lower with bevacizumab.[50] Ranibizumab and aflibercept were directly compared in two similarly designed trials, VEGF Trap-eye: investigation of Efficacy and safety in Wet age-related macular degeneration (VIEW 1 and 2).[51] Similar efficacy and safety was found in both drugs.

From the included trials it is clear that intraocular steroids are associated with complications, including increased intra-ocular pressure and cataract formation.[6-8 23-36] These are substantial drawbacks for using steroids to treat macular oedema in CRVO. However, many affected patients may be already pseudophakic and, on these, the use of intraocular steroids may be reasonable. Steroids may have a place in the treatment pathway of patients who have failed on anti-VEGF therapy, but this has yet to be tested. The anti-VEGF drugs have a good safety profile and do not cause cataract formation.[9-13 42-44] For this reason are likely to be more favoured by clinicians than steroids.

Aflibercept, compared with ranibizumab and bevacizumab, targets a wider range of cytokines and may have a stronger binding affinity.[52] Initial results suggested that aflibercept would require fewer injections than ranibizumab.[51] Heier and colleagues compared aflibercept and ranibizumab in two similarly designed randomised controlled trials in age related macular degeneration. They found that 2 mg aflibercept administered every eight weeks produced similar effects at 96 weeks to 0.5 mg ranibizumab every four weeks.[51] This was reflected in the FDA Dermatologic and Ophthalmic Drugs Advisory Committee recommendation that aflibercept should be given every two months following three initial monthly doses in age related macular

oedema.[53] This may be because aflibercept also appears to last longer in the eye than ranibizumab.[54] Age related macular degeneration is a more aggressive condition than central retinal vein occlusion and so it is unlikely that more frequent dosing would be needed. Therefore aflibercept may be preferred because it would reduce pressure on out-patient clinics. Furthermore there is some evidence from patients with age-related macular degeneration that aflibercept may be effective in patients who have not responded to ranibizumab.[55 56] This may be due to the higher affinity and wider number of cytokines that are targeted. There is no reason to suspect that these effects be any different for the macular oedema caused by central retinal vein occlusion. However we have as yet no evidence as to whether ranibizumab would be effective after aflibercept has failed.

The National Institute of Health and Care Excellence has recommended dexamethasone, ~~and ranibizumab~~ and aflibercept as options in the treatment of macular oedema secondary to CRVO. [18-20-19] ~~and is currently appraising aflibercept~~. Until these technologies are reviewed together and compared with each other, clinicians ~~may be~~ left with three recommended drugs ~~for macular oedema secondary to central retinal vein occlusion~~. It should be noted that during the appraisal of ranibizumab the evidence review group found that in the cost-effectiveness analysis dexamethasone was extendedly dominated by ranibizumab (an intervention is judged not be cost-effective because it has an ICER that is greater than that that of a more effective intervention). The committee appraising ranibizumab did not re-consider the previous appraisal decision on dexamethasone.

Our results show that dexamethasone was not as effective as ranibizumab or aflibercept, at six months follow-up and with the dosing regimens in the trials. However these results do not assess quality of life or cost effectiveness. Bevacizumab is likely to prove

more cost effective than both aflibercept and ranibizumab because it costs substantially less.[57] However the National Institute for Health and Care Excellence has not issued guidance on bevacizumab because it does not have a license for use in the eye.

Unanswered questions and future research

Not all patients benefit from the use of anti-VEGF drugs; only about 60% gain 15 or more letters. It is not clear why some patients benefit more than others. Future research should focus on identifying subgroups of patients who are likely to benefit. Only a few of these trials included ischaemic patients, and in these trials only a few patients with ischaemia were included.[11-13] More research assessing the effectiveness of these drugs in severely ischaemic patients is needed.

Head-to-head trials comparing ranibizumab, aflibercept, bevacizumab and triamcinolone are needed. These should include assessment of cost effectiveness. To assist this, a better measure of quality of life is needed for patients with eye conditions. The widely-used EQ5D may not be sensitive enough to measure changes which are important to patients, such as the ability to drive.

In conclusion, we have found no evidence of difference between ranibizumab, bevacizumab, aflibercept and triamcinolone for improving vision. The anti-VEGFs are likely to be favoured because they are not associated with steroid-induced cataract formation. Aflibercept may be preferred by clinicians because it might require fewer injections.

1
2
3
4
5
6 **Acknowledgements**
7

8 *We thank Christine Clar, Sian Thomas and Rachel Court for assisting with searches,*
9 *screening and data extraction for the systematic review which precede this study. We thank*
10 *the authors of the Epstein 2012 trial for providing addition data.*

11
12 **Copyright statement**

13 *"The Corresponding Author has the right to grant on behalf of all authors and does grant*
14 *on behalf of all authors, a worldwide licence to the Publishers and its licensees in*
15 *perpetuity, in all forms, formats and media (whether known now or created in the future),*
16 *to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the*
17 *Contribution into other languages, create adaptations, reprints, include within collections*
18 *and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other*
19 *derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the*
20 *Contribution, v) the inclusion of electronic links from the Contribution to third party*
21 *material where-ever it may be located; and, vi) licence any third party to do any or all of*
22 *the above."*

23
24 **Declaration of competing interests**

25 "All authors have completed the ICMJE uniform disclosure form at
26 www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the
27 submitted work; no financial relationships with any organisations that might have an
28 interest in the submitted work in the previous three years; no other relationships or
29 activities that could appear to have influenced the submitted work."

30
31 **Contribution statement**

32
33 NW conceived the idea. All authors contributed to the design of the study. DS and OU
34 undertook the statistical analysis. JF, DS and OU wrote the first draft of the manuscript.
35 All authors redrafted and agreed the final article. JF is the guarantor.

36
37 **Transparency statement**

38 JF affirms that the manuscript is an honest, accurate, and transparent account of the
39 study being reported; that no important aspects of the study have been omitted; and
40 that any discrepancies from the study as planned (and, if relevant, registered) have been
41 explained.

42
43 All authors had full access to all of the data in the study and can take responsibility for
44 the integrity of the data and the accuracy of the data analysis

Table 1: Baseline characteristics and results of all included studies

Study	Participants	Intervention / Outcomes
DEXAMETHASONE		
GENEVA 2010[6-8] International	N: CRVO – 437 eyes of 437 patients randomised; 94% follow-up at 6 months Participants: adults with visual acuity reduced because of macular oedema due to CRVO or BRVO	1. Dexamethasone 0.7 mg (n=136) Single dose 2. Dexamethasone 0.35 mg (n=154) Single dose 3. Sham (n=147) Single dose - a needleless applicator was placed against the conjunctiva to simulate the placement of study medication. Primary end point: gain of ≥ 15 ETDRS letters; for the open-label extension: safety
Setting: multicentre (167 centres in 24 countries, so a mean of 2.6 patients per centre)		
Design: 2 identical double-blind, sham-controlled RCTs, phase 3		
Follow-up: primary endpoint for the masked trial: 6 months; primary endpoint for the open-label extension: 12 months		

TRIAMCINOLONE		
SCORE 2009[23-36] USA Setting: multicentre Design: RCT Follow-up: primary end point 12 months, FU planned up to 36 months	N: 271 eyes of 271 patients randomised; 83% (observation) and 90% (triamcinolone) completed 12 months Participants: centre-involved macular oedema secondary to CRVO	1. Triamcinolone 1 mg (n=92) Every 4 months depending on retreatment regimen (ave 2.2 injections at 12 months) 2. Triamcinolone 4 mg (n=91) Every 4 months depending on retreatment regimen (ave 2.0 injections at 12 months) (The form of triamcinolone used was Trivaris, no longer available. It was made by the manufacturer of Ozurdex (Allergan)) 3. Observation (n=88) Primary end point: gain of ≥15 ETDRS letters
AFLIBERCEPT		
COPERNICUS 2012[12 13] International Setting: multicentre, 70 sites in North and South America, India and Israel. Mean 2.7 patients per centre. Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU 2 years	N: 189 eyes of 189 patients randomised; 95.7% (aflibercept) and 81.1% (sham) completed 24 weeks; 93% (aflibercept) and 77% (sham) completed 52 weeks Participants: adult patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=114) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=73) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥15 ETDRS letters

Field Code Changed

GALILEO 2012[11] International Setting: multicentre, 10 countries in Europe and Asia; 63 centres in total Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 24 weeks, FU up to 12 months, planned up to 76 weeks	N: 177 eyes of 177 patients randomised; 90.6% (aflibercept) and 78.9% (sham) completed 24 weeks Participants: treatment-naïve patients with centre-involved CRVO for a maximum of 9 months	1. Aflibercept 2mg (n=103) Every 4 weeks for 6 months (ave number not available) 2. Sham (n=71) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to conjunctival surface) Primary end point: gain of ≥ 15 ETDRS letters
--	--	--

RANIBIZUMAB		
CRUISE 2010 [9 10] USA Setting: multicentre Design: double-blind, sham-controlled RCT, phase 3 Follow-up: primary end point 6 months, FU up to 12 months	N: 392 eyes of 392 patients randomised; 97.7% (ranibizumab 0.3 mg), 91.5% (ranibizumab 0.5 mg), and 88.5% (sham) completed 6 months Participants: patients with foveal centre-involved macular oedema secondary to CRVO diagnosed within 12 months	1. Ranibizumab 0.3 mg (n=132) Every 4 weeks for 6 months (ave number not available) 2. Ranibizumab 0.5 mg (n=130) Every 4 weeks for 6 months (ave number not available) 3. Sham (n=130) Every 4 weeks for 6 months (ave number not available) (empty syringe without needle pressed to the injection site) Primary end point: mean change from baseline BCVA
BEVACIZUMAB		
Epstein 2012 [42-44] Sweden Setting: Single centre; St. Eriks Eye Hospital Stockholm Design: sham-injection controlled, double masked RCT Follow-up: primary end-point 6 months; open label extension up to 12 months	N: 60 eyes of 60 patients randomised; 93% completed open label extension Participants: patients with CRVO of ≤6 months	1. Bevacizumab 1.25 mg (n=30) Every 6 weeks for 6 months (ave number not available) 2. Sham (n=30) Every 6 weeks for 6 months (ave number not available) (syringe without needle pressed to the globe) Primary end point: gain of ≥15 ETDRS letters

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

FU= follow-up, RCT = randomised controlled trial, N = number, CRVO = central retinal vein occlusion, ETDRS = Early Treatment Diabetic Retinopathy Study, BRVO = branch retinal vein occlusion

Table 2: Baseline characteristics and results of included trials

	COPERNICUS[12 13]	GALILEO[11]	CRUISE[9 10]	GENEVA[6- 8]	Epstein et al (2012)[42- 44]	SCORE[23-36]
BASELINE SIMILARITIES						
Number (%) of patients						
	Aflib 2 mg: 114 Sham: 73	Aflib 2 mg: 103 Sham: 68	Rani 0.5 mg: 130 Sham: 130	Dexa0.7 mg: 136 Sham: 147	Beva 1.25 mg: 30 Sham: 30	Triam 4 mg: 91 Obser: 88
Age (years)						
	Aflib 2 mg: 65.5 SD13.6 Sham: 67.5 SD14.3	Aflib 2 mg: 59.9 SD12.4 Sham: 63.8 SD13.3	Rani 0.5 mg: 67.6 SD12.4 Sham: 65.4 SD13.1	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 70.6 SD 12.6 Sham: 70.4 SD 10.4	Triam 4 mg: 67.5 SD 12.0 Obser: 69.2 SD 12.8
BCVA at baseline (SD)						
	Aflib 2 mg: 50.7 SD13.90 Sham: 48.9 SD14.42	Aflib 2 mg: 53.6 SD15.8 Sham: 50.9 SD15.4	Rani 0.5 mg: 48.1 SD14.6 Sham: 49.2 SD14.7	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: 44.4 SD 15.3 Sham: 43.6 SD 16.0	Triam 4 mg: 51.0 SD 14.4 Obser: 52.1 SD 13.1
Duration of MO from diagnosis to screening						
	Aflib 2 mg: 2.73 SD3.09(in months) Sham: 1.88 SD2.19 (in months)	Aflib 2 mg: 50.9 SD15.4)(in days) Sham: 87.6 SD79.1 (in days)	Rani 0.5 mg: - Sham: -	Dexa 0.7 mg: NR Sham: NR	Beva 1.25 mg: NR Sham: NR	Triam 4 mg: 4.2 SD 3.6 (in months) Obser: 4.2 SD 3.1 (in months)
RESULTS						
Number (%) of patients gaining ≥15 letters improvement from baseline to 6 months						
	Aflib 2 mg: 64 (56.1)	Aflib 2 mg: 62 (60.2)	Rani 0.5 mg: 62 (47.7)	Dexa 0.7 mg: 25 (18)	Beva 1.25 mg: 18 (60%)	Triam 4 mg: 18 (19.5%) (avg of 4 and 8 mths)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

	Sham: 9 (12.3)	Sham: 15 (22.1)	Sham: 22 (16.9)	Sham: 18 (12)	Sham: 6 (20%)	Obser: 3 (4%) (avg of 4 and 8 mths)
	Number (%) of patients losing ≥ 15 letters of BCVA from baseline to 6 months					
	Aflib 2 mg: 2 (1.8)	Aflib 2 mg: 8 (7.8)	Rani 0.5 mg: 2 (1.5)	Dexa 0.7 mg: NR	Beva 1.25 mg: 2 (6.7%)	Triam 4 mg: 19 (20.5%) (avg of 4 and 8 mths)
	Sham: 20 (27.4)	Sham: 15 (22.1)	Sham: 20 (15.4)	Sham: NR	Sham: 7 (23.3%)	Obser: 31 (35.5%) (avg of 4 and 8 mths)
	Mean change (SD) from baseline in BCVA					
	Aflib 2 mg: 17.3 (12.8)	Aflib 2 mg: 18.0 (12.2)	Rani 0.5 mg: 14.9 (13.2)	Dexa 0.7 mg: 0.1 (NR)	Beva 1.25 mg: 14.1 SD 18.7	Triam 4 mg: -0.15 SD 20.67 (n=85) (weight mean and SD of 4 and 8 months)
	Sham: -4 (18)	Sham: 3.3 (14.1)	Sham: 0.8 (16.2)	Sham: -1.8 (NR)	Sham: -2.0 SD 20.5	Obser: -9.66 SD 18.04 (n=75) (weighted mean and SD of 4 and 8 months)

NR = not reported, Aflib = aflibercept, Rani = ranibizumab, Dexa = dexamethasone, Triam = triamcinolone, Obser = observation, SD = standard deviation, avg = average

Table 3: Risk of bias

Study (author and year)	Adequate sequence generation	Allocation concealment	Masking	Incomplete outcome data addressed	Free of selective reporting	Free of other bias (e.g. similarity at baseline, power assessment)	Funder
GENEVA 2010[6-8]	Low	Low	Partial: patients and assessors of efficacy variables	Low: ITT analysis, 94% FU at 6 months	Low	<i>Power:</i> 81% power to detect difference in primary outcome with n=495 for each trial <i>Similarity at baseline:</i> yes	Allergan Inc.
SCORE 2009[23-36]	Low	Unclear	Partial (physicians and patients masked to dose but not triamcinolone versus observation)	Low: ITT analysis, 83 to 90% FU at 12 months	Low	<i>Power:</i> 80% power to detect difference in primary outcome with n=486 (but only 271 randomised) <i>Similarity at baseline:</i> yes	National Eye Institute grants, Allergan
COPERNICUS 2012[12 13]	Low	Unclear	Low: double-blind	Low: ITT analysis, 89.9% assessed at primary end point	Low	<i>Power:</i> 90% power to detect difference in primary outcome with n=165 <i>Similarity at baseline:</i> yes	Bayer HealthCare, Regeneron Pharmaceuticals
GALILEO 2012[11]	Unclear	Unclear	Low: double-blind	Low: ITT analysis, 86% assessed at primary end point	Low	<i>Power:</i> 90% power to detect difference in primary outcome with n=150 <i>Similarity at baseline:</i> yes	Bayer HealthCare, Regeneron Pharmaceuticals

CRUISE 2010[9 10]	Low	Unclear	Low: patients and evaluating examiners, injecting physicians masked to dose	Low: ITT analysis, 88.5 to 97.7% completed 6 months	Low	<i>Power</i> : not reported <i>Similarity at baseline</i> : yes	Genentech Inc.
Epstein 2012[42-44]	Unclear	Low	Low: patients, outcome assessors	Low: ITT analysis; missing data for 2 patients (primary endpoint)	Low	<i>Power</i> : 80% power to detect difference in primary outcome with n=24 per group <i>Similarity at baseline</i> : yes	Unclear; authors are consultants for Allergan, Novartis, Alcon, Bayer

ITT= intention to treat, FU = follow-up

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legend

- Figure 1: study selection flow diagram
- Figure 2: Network of randomized controlled trials comparing different treatments for proportions of gaining 3 or more lines of vision
- Figure 3: Proportions of patients gaining 3 lines or more from baseline to six months
- Figure 4: Rankogram for gaining ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions
- Figure 5: Proportions of patients losing 3 lines or more from baseline to six months
- Figure 6: Rankogram for losing ≥ 3 lines - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions
- Figure 7: Mean BCVA change from baseline to 6 months.
- Figure 8: Rankogram for mean change in BCVA - distribution of the probabilities of every treatment being ranked at each of the possible 6 positions

Appendix: MEDLINE search strategy

Ovid MEDLINE(R) 1946 to March Week 1 2013, searched on 20 March 2013

- 1 CRVO.mp.
- 2 Retinal Vein Occlusion/
- 3 retinal vein occlusion.mp.
- 4 retinal vein obstruction.mp.
- 5 retinal venous occlusion.mp.
- 6 retinal venous obstruction.mp.
- 7 retina*.mp.
- 8 ("central vein occlusion" or "central vein obstruction" or "central venous occlusion" or "central venous obstruction").mp.
- 9 7 and 8
- 10 1 or 2 or 3 or 4 or 5 or 6 or 9
- 11 randomized controlled trial.pt.
- 12 (random* or "controlled trial*" or "clinical trial*" or rct).tw.
- 13 11 or 12
- 14 (metaanalys* or "meta analys*" or "meta-analys*").tw.
- 15 "systematic review*".tw.
- 16 meta analysis.pt.
- 17 14 or 15 or 16
- 18 10 and 13
- 19 10 and 17
- 20 18 or 19
- 21 limit 20 to yr="2005 -Current"

References

1. Deramo VA, Cox TA, Syed AB, Lee PP, Fekrat S. Vision-related quality of life in people with central retinal vein occlusion using the 25-item National Eye Institute Visual Function Questionnaire. Arch.Ophthalmol. 2003;**121**(9):1297-302 doi: 10.1001/archophth.121.9.1297 [doi];121/9/1297 [pii][published Online First: Epub Date]].

2. Klein R, Moss SE, Meuer SM, Klein BE. The 15-year cumulative incidence of retinal vein occlusion: the Beaver Dam Eye Study. Archives of ophthalmology 2008;**126**(4):513-8 doi: 10.1001/archophth.126.4.513[published Online First: Epub Date]].

3. Hayreh SS, Podhajsky PA, Zimmerman MB. Natural history of visual outcome in central retinal vein occlusion. Ophthalmology 2011;**118**(1):119-33 doi: S0161-6420(10)00447-1 [pii];10.1016/j.ophtha.2010.04.019 [doi][published Online First: Epub Date]].

4. McIntosh RL, Rogers SL, Lim L, et al. Natural history of central retinal vein occlusion: an evidence-based systematic review. Ophthalmology 2010;**117**(6):1113-23 doi: S0161-6420(10)00133-8 [pii];10.1016/j.ophtha.2010.01.060 [doi][published Online First: Epub Date]].

5. The Royal College of Ophthalmologists. Interim Guidelines for Management of Retinal Vein Occlusion. Secondary Interim Guidelines for Management of Retinal Vein Occlusion 2010. http://www.rcophth.ac.uk/core/core_picker/download.asp?id=728&filetitle=Interim+Guidelines+for+Management+of+Retinal+Vein+Occlusion+2010.

6. Haller JA, Bandello F, Belfort R, Jr., et al. Randomized, sham-controlled trial of dexamethasone intravitreal implant in patients with macular edema due to retinal vein occlusion. *Ophthalmology* 2010;**117**(6):1134-46
7. Haller JA, Bandello F, Belfort R, Jr., et al. Dexamethasone intravitreal implant in patients with macular edema related to branch or central retinal vein occlusion twelve-month study results. *Ophthalmology* 2011;**118**(12):2453-60
8. Yeh WS, Haller JA, Lanzetta P, et al. Effect of the duration of macular edema on clinical outcomes in retinal vein occlusion treated with dexamethasone intravitreal implant. *Ophthalmology* 2012;**119**(6):1190-98
9. Brown DM, Campochiaro PA, Singh RP, et al. Ranibizumab for macular edema following central retinal vein occlusion: six-month primary end point results of a phase III study. *Ophthalmology* 2010;**117**(6):1124-33
10. Campochiaro PA, Brown DM, Awh CC, et al. Sustained benefits from ranibizumab for macular edema following central retinal vein occlusion: twelve-month outcomes of a phase III study. *Ophthalmology* 2011;**118**(10):2041-49
11. Holz FG, Roeder J, Ogura Y, et al. VEGF Trap-Eye for macular oedema secondary to central retinal vein occlusion: 6-month results of the phase III GALILEO study. *British Journal of Ophthalmology* 2013;**97**(3):278-84
12. Boyer D, Heier J, Brown DM, et al. Vascular endothelial growth factor Trap-Eye for macular edema secondary to central retinal vein occlusion: six-month results of the phase 3 COPERNICUS study. *Ophthalmology* 2012;**119**(5):1024-32

13. Brown DM, Heier JS, Clark WL, et al. Intravitreal Aflibercept Injection for Macular Edema Secondary to Central Retinal Vein Occlusion: 1-Year Results From the Phase 3 COPENICUS Study. *American Journal of Ophthalmology* 2013;**155**(3):429-37

14. Michels S, Rosenfeld PJ, Puliafito CA, Marcus EN, Venkatraman AS. Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration twelve-week results of an uncontrolled open-label clinical study. *Ophthalmology* 2005;**112**(6):1035-47 doi: 10.1016/j.ophtha.2005.02.007[published Online First: Epub Date]].

15. Arevalo JF, Fromow-Guerra J, Quiroz-Mercado H, et al. Primary intravitreal bevacizumab (Avastin) for diabetic macular edema: results from the Pan-American Collaborative Retina Study Group at 6-month follow-up. *Ophthalmology* 2007;**114**(4):743-50 doi: 10.1016/j.ophtha.2006.12.028[published Online First: Epub Date]].

16. Rabena MD, Pieramici DJ, Castellarin AA, Nasir MA, Avery RL. Intravitreal bevacizumab (Avastin) in the treatment of macular edema secondary to branch retinal vein occlusion. *Retina (Philadelphia, Pa.)* 2007;**27**(4):419-25 doi: 10.1097/IAE.0b013e318030e77e[published Online First: Epub Date]].

17. Algvere PV, Epstein D, von Wendt G, Seregard S, Kvanta A. Intravitreal bevacizumab in central retinal vein occlusion: 18-month results of a prospective clinical trial. *European journal of ophthalmology* 2011;**21**(6):789-95 doi: 10.5301/ejo.2011.6522[published Online First: Epub Date]].

18. National Institute for Health and Care Excellence. Ranibizumab for treating visual impairment caused by macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 283. 2013.
<http://guidance.nice.org.uk/TA283/Guidance/pdf/English>.
19. National Institute for Health and Clinical Excellence. Dexamethasone intravitreal implant for the treatment of macular oedema secondary to retinal vein occlusion: NICE technology appraisal guidance 229. 2011.
<http://guidance.nice.org.uk/TA229/Guidance/pdf/English>.
20. National Institute for Health and Care Excellence. Aflibercept for treating visual impairment caused by macular oedema secondary to central retinal vein occlusion. NICE technology appraisal guidance 305. 2014.
<http://guidance.nice.org.uk/TA305>.
21. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ (Clinical research ed.)* 2011;**343**:d5928 doi: 10.1136/bmj.d5928[published Online First: Epub Date]].
22. Hozo S, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Medical Research Methodology* 2005;**5**(1):13
23. Bhavsar AR, Ip MS, Glassman AR, Groups DatSS. The risk of endophthalmitis following intravitreal triamcinolone injection in the DRCRnet and SCORE clinical trials. *American Journal of Ophthalmology* 2007;**144**(3):454-56
24. Blodi BA, Domalpally A, Scott IU, et al. Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study system for evaluation of

stereoscopic color fundus photographs and fluorescein angiograms:
SCORE Study Report 9. Archives of ophthalmology 2010;**128**(9):1140-45

25. Chan CK, Ip MS, VanVeldhuisen PC, et al. SCORE Study report #11: incidences of neovascular events in eyes with retinal vein occlusion. Ophthalmology 2011;**118**(7):1364-72

26. Ip M, Oden N, VanVeldhuisen P, Scott I, Blodi B. The Standard Care vs. Corticosteroid for Retinal Vein Occlusion Study: Design and Baseline Characteristics. American Academy of Ophthalmology 2008:260

27. Ip MS, Oden NL, Scott IU, et al. SCORE Study report 3: study design and baseline characteristics. Ophthalmology 2009;**116**(9):1770-77

28. Ip MS, Scott IU, VanVeldhuisen PC, et al. A randomized trial comparing the efficacy and safety of intravitreal triamcinolone with observation to treat vision loss associated with macular edema secondary to central retinal vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) study report 5. Archives of ophthalmology 2009;**127**(9):1101-14

29. Myers D, Blodi B, Ip M, Scott I, Warren K. Reading Center Evaluation of OCT Images From Patients Enrolled in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. Iovs. 2006;**47**:ARVO

30. Oden NL, Veldhuisen PC, Scott IU, Ip MS, Blodi BA. Temporal Variability of OCT in Retinal Vein Occlusion Participants in the SCORE Study. Iovs. 2007;**48**:ARVO

31. Scott IU, Blodi BA, Ip MS, et al. SCORE Study Report 2: Interobserver agreement between investigator and reading center classification of retinal vein occlusion type. Ophthalmology 2009;**116**(4):756-61

32. Scott IU, Oden NL, VanVeldhuisen PC, et al. SCORE Study Report 7: incidence of intravitreal silicone oil droplets associated with staked-on vs luer cone syringe design. *American Journal of Ophthalmology* 2009;**148**(5):725-32
33. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline predictors of visual acuity and retinal thickness outcomes in patients with retinal vein occlusion: Standard Care Versus Corticosteroid for Retinal Vein Occlusion Study report 10. *Ophthalmology* 2011;**118**(2):345-52
34. Scott IU, VanVeldhuisen PC, Oden NL, et al. SCORE Study report 1: baseline associations between central retinal thickness and visual acuity in patients with retinal vein occlusion. *Ophthalmology* 2009;**116**(3):504-12
35. Scott IU, VanVeldhuisen PC, Oden NL, et al. Baseline Characteristics and Response to Treatment of Participants With Hemiretinal Compared With Branch Retinal or Central Retinal Vein Occlusion in the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) Study: SCORE Study Report 14. *Archives of ophthalmology* 2012;**130**(12):1517-24
36. Warren K, Blodi BA, Oden N, Veldhuisen P, Scott IU, Ip M. Reading Center Evaluation of Baseline Retinal Images in the Standard Care vs. Corticosteroid for Retinal Vein Occlusion (SCORE) Study. *Iovs*. 2008:ARVO
37. Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *BMJ (Clinical research ed.)* 2005;**331**(7521):897-900 doi: 10.1136/bmj.331.7521.897[published Online First: Epub Date]].
38. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. *Statistics in medicine* 2004;**23**(20):3105-24 doi: 10.1002/sim.1875[published Online First: Epub Date]].

39. Spiegelhalter D, Thomas A, Best N, Lunn D. *WinBUGS User Manual: Version 1.4*. Cambridge, Mass: MRC Biostatistics Unit, 2003.

40. Brooks SP, Gelman A. General Methods for Monitoring Convergence of Iterative Simulations. *Journal of Computational and Graphical Statistics* 1998;**7**(4):434-55 doi: 10.1080/10618600.1998.10474787[published Online First: Epub Date]].

41. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology* 2011;**64**(2):163-71 doi: 10.1016/j.jclinepi.2010.03.016[published Online First: Epub Date]].

42. Epstein D, Algvere P, Von WG, Seregard S, Kvanta A. Long-term benefit from bevacizumab for macular edema in central retinal vein occlusion: 12-month results of a prospective study. *Acta Ophthalmologica* 2012;**90**:48 doi: 10.1111/j.1755-3768.2012.02549.x[published Online First: Epub Date]].

43. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Benefit from bevacizumab for macular edema in central retinal vein occlusion: twelve-month results of a prospective, randomized study. *Ophthalmology* 2012;**119**(12):2587-91

44. Epstein DL, Algvere PV, Von WG, Seregard S, Kvanta A. Bevacizumab for macular edema in central retinal vein occlusion: a prospective, randomized, double-masked clinical study. *Ophthalmology* 2012;**119**(6):1184-89

45. Ford JA, Clar C, Lois N, et al. Treatments for macular oedema following central retinal vein occlusion: systematic review. BMJ open 2014;**4**(2):e004120 doi: 10.1136/bmjopen-2013-004120[published Online First: Epub Date]].
46. FDA. Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators. Secondary Drug Study Designs - Information Sheet. Guidance for Institutional Review Boards and Clinical Investigators 2011.
<http://www.fda.gov/RegulatoryInformation/Guidances/ucm126501.htm>.
47. Martin DF, Maguire MG, Fine SL, et al. Ranibizumab and bevacizumab for treatment of neovascular age-related macular degeneration: two-year results. Ophthalmology 2012;**119**(7):1388-98 doi: 10.1016/j.ophtla.2012.03.053[published Online First: Epub Date]].
48. Chakravarthy U, Harding SP, Rogers CA, et al. Ranibizumab versus bevacizumab to treat neovascular age-related macular degeneration: one-year findings from the IVAN randomized trial. Ophthalmology 2012;**119**(7):1399-411 doi: S0161-6420(12)00358-2 [pii];10.1016/j.ophtla.2012.04.015 [doi][published Online First: Epub Date]].
49. Ford JA, Elders A, Shyangdan D, Royle P, Waugh N. The relative clinical effectiveness of ranibizumab and bevacizumab in diabetic macular oedema: an indirect comparison in a systematic review. BMJ (Clinical research ed.) 2012;**345**:e5182

50. Chakravarthy U, Harding SP, Rogers CA, et al. Alternative treatments to inhibit VEGF in age-related choroidal neovascularisation: 2-year findings of the IVAN randomised controlled trial. *Lancet* 2013 doi: 10.1016/s0140-6736(13)61501-9[published Online First: Epub Date]].

51. Heier JS, Brown DM, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology* 2012;**119**(12):2537-48 doi: 10.1016/j.optha.2012.09.006[published Online First: Epub Date]].


52. Papadopoulos N, Martin J, Ruan Q, et al. Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* 2012;**15**(2):171-85 doi: 10.1007/s10456-011-9249-6[published Online First: Epub Date]].

53. FDA. VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee. Secondary VEGF TRAP-EYE (aflibercept ophthalmic solution). Ophthalmologic drugs advisory committee 2011. <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/DermatologicandOphthalmicDrugsAdvisoryCommittee/UCM259143.pdf>.


54. Stewart MW, Rosenfeld PJ. Predicted biological activity of intravitreal VEGF Trap. *The British journal of ophthalmology* 2008;**92**(5):667-8 doi: 10.1136/bjo.2007.134874[published Online First: Epub Date]].

55. Bakall B, Folk JC, Boldt HC, et al. Aflibercept therapy for exudative age-related macular degeneration resistant to bevacizumab and ranibizumab. *Am J Ophthalmol* 2013;**156**(1):15-22.e1 doi: 10.1016/j.ajo.2013.02.017[published Online First: Epub Date]].


56. Cho H, Shah CP, Weber M, Heier JS. Aflibercept for exudative AMD with persistent fluid on ranibizumab and/or bevacizumab. *The British journal of ophthalmology* 2013;**97**(8):1032-5 doi: 10.1136/bjophthalmol-2013-303344[published Online First: Epub Date]].
57. Raftery J, Clegg A, Jones J, Tan SC, Lotery A. Ranibizumab (Lucentis) versus bevacizumab (Avastin): modelling cost effectiveness. *The British journal of ophthalmology* 2007;**91**(9):1244-6 doi: 10.1136/bjo.2007.116616[published Online First: Epub Date]].




170x172mm (300 x 300 DPI)




125x59mm (300 x 300 DPI)




170x152mm (300 x 300 DPI)




170x88mm (300 x 300 DPI)




170x90mm (300 x 300 DPI)



170x90mm (300 x 300 DPI)



170x155mm (300 x 300 DPI)



170x90mm (300 x 300 DPI)

Appendix: MEDLINE search strategy

Ovid MEDLINE(R) 1946 to March Week 1 2013, searched on 20 March 2013

- 1 CRVO.mp.
- 2 Retinal Vein Occlusion/
- 3 retinal vein occlusion.mp.
- 4 retinal vein obstruction.mp.
- 5 retinal venous occlusion.mp.
- 6 retinal venous obstruction.mp.
- 7 retina*.mp.
- 8 ("central vein occlusion" or "central vein obstruction" or "central venous occlusion" or "central venous obstruction").mp.
- 9 7 and 8
- 10 1 or 2 or 3 or 4 or 5 or 6 or 9
- 11 randomized controlled trial.pt.
- 12 (random* or "controlled trial*" or "clinical trial*" or rct).tw.
- 13 11 or 12
- 14 (metaanalys* or "meta analys*" or "meta-analys*").tw.
- 15 "systematic review*".tw.
- 16 meta analysis.pt.
- 17 14 or 15 or 16
- 18 10 and 13
- 19 10 and 17
- 20 18 or 19
- 21 limit 20 to yr="2005 -Current"



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	3-4
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5-6
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5-6
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	1
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	7
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	41
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	7-8
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7-8
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	7-8
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	7-8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	8-9
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2 for each meta-analysis).	8-9



PRISMA 2009 Checklist

Page 1 of 2

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	8-9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	8-9
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	11+27
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	20-22
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	25-26
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	23-24
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	29-31
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	11-13
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	NA
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	14-18
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	14-18
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	14-18
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	1

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Page 2 of 2

Protected by copyright, including for uses related to text and data mining, AI training and similar technologies.