

BMJ Open

UK newspaper and online news representations of women's and men's 'binge' drinking: a challenge for communicating evidence-based messages about single episodic drinking?

Journal:	BMJ Open
Manuscript ID	bmjopen-2016-013124
Article Type:	Research
Date Submitted by the Author:	21-Jun-2016
Complete List of Authors:	Patterson, Chris; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit Emslie, Carol; Glasgow Caledonian University Institute for Applied Health Research Mason, Olivia; Durham University, Department of Geography Fergie, Gillian; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit Hilton, Shona; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit
Primary Subject Heading:	Communication
Secondary Subject Heading:	Sociology, Public health
Keywords:	PUBLIC HEALTH, QUALITATIVE RESEARCH, SOCIAL MEDICINE

SCHOLARONE™
Manuscripts

UK newspaper and online news representations of women’s and men’s ‘binge’ drinking: a challenge for communicating evidence-based messages about single episodic drinking?

Patterson^{1*}, Emslie², Mason³, Fergie¹, Hilton¹

¹MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB UK

²Institute for Applied Health Research/School of Health and Life Sciences, Glasgow Caledonian University, Cowcaddens Road, Glasgow G4 0BA UK

²Department of Geography, Durham University, Lower Mountjoy South Road, Durham DH1 3LE UK

Corresponding author:

Chris Patterson

MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB UK

chris.patterson@glasgow.ac.uk

0141 353 7631

Keywords: gender; media; alcohol; binge drinking

Running head: Media representations of ‘binge’ drinking

3,674 words

ABSTRACT

Objectives: In the UK, Men's alcohol-related morbidity and mortality still greatly exceed women's, despite an increase in women's alcohol consumption in recent decades. Newly proposed UK alcohol guidelines introduce gender-neutral weekly alcohol consumption limits. This study explores how UK newspaper and online news represent women's and men's 'binge' drinking to identify opportunities to better align reporting of harmful drinking with evidence.

Design: Quantitative and qualitative content analysis of 308 articles published in seven UK national newspapers and the BBC News website between 1 January 2012 and 31 December 2013.

Results: Articles associated women with 'binge' drinking more frequently than men and problematised women's drinking more. Men were more frequently characterised as violent or disorderly, while women were characterised as out of control, putting themselves in danger, harming their physical appearance and burdening men. Judgemental descriptions of female 'binge' drinkers' clothing and appearance were frequent.

Conclusions: The UK news media's disproportionate focus on women's 'binge' drinking is at odds with epidemiological evidence, may reproduce harmful gender stereotypes and may obstruct public understandings of the gender-neutral weekly consumption limits in newly-proposed alcohol guidelines. In order to better align reporting of harmful drinking with current evidence, public health advocates may engage with the media with a view to shifting media framing of 'binge' drinking away from specific groups (young people; females) and contexts (public drinking) and towards the health risks of specific drinking behaviours, which affect all groups regardless of context.

Strengths and limitations of this study

- Quantitative and qualitative content analysis of a large, comprehensive, two-year sample of national newspaper and online news about ‘binge’ drinking.
- The findings illustrate how media portrayals of ‘binge’ drinking are harmful and identify opportunities for these portrayals to be better aligned with evidence.
- Content analysis facilitates understandings of the messages being presented to the public, but cannot determine the extent to which audiences’ understandings of specific issues correlate with media representations.
- Some findings may have differed if the data analysed had gone beyond text articles, for example by incorporating images, video and social media content.

INTRODUCTION

Despite the increase in women's consumption of alcohol over the last 30 years in the UK,[1] men still drink more than women, experience more drink-related health and social problems and have double women's rates of alcohol-related deaths (15.9 to 7.8 deaths per 100,000 population respectively in 2012[2]). New UK drinking guidelines propose equal weekly alcohol consumption limits of 14 units for both men and women spread evenly over three days or more.[3] These proposed gender-neutral limits differ from previous guidelines, which, in 1995, suggested a weekly maximum of 21 units for men (with no more than 4 units in any day) and 14 units for women (with no more than 3 units in any day).[4]

'Binge' drinking, also called single episodic drinking, is associated with increased accidental injury, poor health outcomes and wider social harms.[3] Herring and Berridge[5] describe binge drinking as a 'confused concept', the definitions of which has changed over time and continues to be interpreted inconsistently. They suggest that 'binge' drinking is currently portrayed as a youth issue, despite evidence that single episodic drinking is performed by various age groups.[2] Qualitative evidence indicates that young women in Scotland define binge drinking in terms of types of behaviour rather than the quantity of alcohol consumed.[6] As with most health issues, popular perceptions of binge drinking are likely influenced by mass media representations,[7, 8] and differences in representations of women's and men's drinking is heavily influenced by how societies regulate gender roles.[9] Exploring media representations of men's and women's binge drinking allows us to examine how shared cultural values around alcohol are articulated and constructed[10] to inform efforts to improve media representations, and therefore public understandings, of harmful drinking behaviours.[11] In light of the gender-neutral consumption limits in the newly-proposed guidelines,[3] it is useful to better understand how women's and men's binge drinking are represented by the media.

Research illustrates a clear gender divide in media portrayals of drinking behaviours, with men's drinking normalised and women's problematised. Day and colleagues[12] found that UK newspapers frequently characterised women as departing from idealised notions of femininity in terms of appearance (weight gain, deeper voices, loss of good looks) and motherhood (reduced fertility and unborn children). They found men were framed as violent, but women, framed as sexual predators invading traditionally male-dominated drinking environments, were partially responsible for men's violence. Conversely, Wood and colleagues[13] found that UK newspaper coverage of proposed minimum unit pricing (MUP) policies presented women as being at 'risk of harm' from male aggression. Lyons, Dalton & Hoy[14] found that both young women's and young men's magazines framed binge drinking as normative, adult and professional. Young men's magazines associated men's drinking with traditional masculine images but derided young women's drinks and drinking behaviours. Nicholls[15] found that UK television and newspaper news associated men's drinking with violence (both as perpetrators or as victims) and women's drunkenness with unfeminine and undignified behaviours, such as loss of consciousness or partial nudity. Similarly, Atkinson and colleagues'[16] found that UK magazines targeted at teenagers depicted women's drinking as more problematic than men's, and that women were both portrayed as 'behaving like men' ('ladettes' appropriating male space) but also sexualised and highly emotional.

This study comprises a comprehensive, mixed-methods content analysis of two years of UK media coverage of binge drinking, designed to contribute new insights to a growing body of literature about gendered media representations of alcohol, with a particular focus on binge drinking. Our aim is to improve understandings of UK newspaper and online news representations of women's and men's 'binge' drinking in the context of changing alcohol guidelines. These improved understandings may inform efforts to better align media

representations of harmful drinking with current evidence, which could improve public understandings of the risks of single episodic drinking.

METHODS

We selected seven highly-circulated[17, 18] UK national newspapers, including their Sunday counterparts and, and three UK online news websites with large readerships[19] (Table 1).

Publications represented three genres: quality, middle-market tabloid and tabloid. This typology helps to ensure the sample represents diverse readerships in terms of age, social class and political alignment.[20]

We selected a search period of 1 January 2012 to 31 December 2013, during which alcohol was subject to heightened media interest due to debates in the UK and Scottish parliaments around MUP.[21] Relevant newspaper articles, including articles from newspapers' online editions, were identified using the databases Nexis and Newsbank. BBC News articles were identified using the search function on the BBC website. The search term used identified articles that contain both three or more mentions of "binge" and one or more mentions of "drink OR drinker OR drinkers OR drinking". Duplicate articles were removed, and articles were manually excluded if they did were not predominantly focused on binge drinking in the UK, or if they were not in the news feature or editorial format. Initial searches identified 537 articles, of which 308 met the inclusion criteria and were eligible for detailed coding and analysis.

Table 1. Summary of publications and articles in the sample

Genre / medium	Publication	All articles		Article format					
				Standard		Feature		Editorial	
		n	%	n	%	n	%	n	%
Quality (n=86)	Guardian / Observer	17	5.5	7	3.0	9	17.3	1	5.3
	Independent / Independent on Sunday	17	5.5	11	4.6	5	9.6	1	5.3
	Daily Telegraph / Sunday Telegraph	52	16.9	44	18.6	5	9.6	3	15.8
Middle-market tabloids (n=39)	Daily Mail / Mail on Sunday	26	8.4	12	5.1	11	21.2	3	15.8
	Express / Sunday Express	13	4.2	8	3.4	3	5.8	2	10.5
Tabloids (n=75)	Daily Mirror / Sunday Mirror	13	4.2	10	4.2	2	3.8	1	5.3
	The Sun / News of the World	62	20.1	46	19.4	9	17.3	7	36.8
Online (n=108)	BBC News website	39	12.7	37	15.6	1	1.9	1	5.3
	Guardian.co.uk	41	13.3	41	17.3	0	0.0	0	0.0
	MailOnline	28	9.1	21	8.9	7	13.5	0	0.0
Total:		308	100	237	100	52	100	19	100

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

SH and CE read each article to identify emergent themes, integrating these within a quantitative coding frame comprising three broad thematic categories: how is binge drinking described and which sections of the population are associated with this behaviour?; how are the drivers of the binge drinking described and who is to blame?; and what are the consequences of binge drinking? Using the coding frame, OM coded each article quantitatively. To ensure consistent coding GF coded a random subsample of 39 articles (12.7%). Linearly-weighted kappa tests of inter-rater agreement were performed on each variable, and those that returned a coefficient below 0.8 were discarded to ensure only variables with strong agreement were analysed.

Statistical procedures comprised: a simple linear regression examining the relationship between publication year and the count of articles published per year; chi-square tests of whether thematic variables varied significantly by gender-focus; and paired t-tests comparing the means of thematic variables. Following quantitative analysis, a subsample of 46 articles that mentioned 'harms to appearance' were analysed thematically[22] by GF and CP using NVivo 10.

RESULTS

Quantitative findings

Between 1 January 2012 and 31 December 2013, 308 articles about binge drinking were published in the seven print publications and three websites included in the sample. Of these articles, 86 (27.9%) were published in 'quality' newspapers, 39 (12.7%) in middle-market tabloids, 75 (24.4%) in tabloids and 108 (35.1%) on websites (Table 1). The majority of articles were standard news format (n=237, 76.9%), while 52 (16.9%) were feature articles and 19 (6.2%) editorials. The yearly count of articles decreased significantly (coefficient - 0.702, p=0.027) across the sample period (Figure 1). There was a peak of 56 articles in March

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2012, 66.1% of which were related to MUP, in particular the UK Prime Minister’s announcement of plans to introduce MUP. There was elevated reporting from November 2012 to January 2013 (n=62), during which 53.2% of articles mentioned MUP, largely related to opposition to the measure within the UK Cabinet. In total, 133 (43.2%) articles mentioned MUP.

To examine associations with gender, we coded articles according to whether they mentioned binge drinking in females exclusively (n=68, 22.1%), males exclusively (n=30, 9.7%), both males and females (n=43, 14.0%), or neither males nor females explicitly (n=167, 54.2%). There was no significant association between gender category and publication genre (p=0.382) or between between gender category and article format (p=0.303).

Articles were coded to identify the characteristics of binge drinkers and the specific drinks associated with binge drinking. Thirty-two (10.4%) of all 308 articles associated mothers (including pregnant women) with binge drinking, while only 11 (3.6%) mentioned fathers (Table 2). There were no statistically significant relationships between focusing on a specific gender and mentioning younger or older people. Articles focused exclusively on women were significantly (p=0.047) more likely to refer to home drinking than those that focused exclusively on men (Table 2). No other location was significantly associated with gender category. Wine (n=58, 18.8%) was the only type of drinks significantly associated more with one gender than the other, being more likely to be mentioned in articles focusing exclusively on females (p=0.004).

Table 2. Social groups, locations and harms associated with binge drinking

	All articles (n=308)		Only females (n=68)		Only males (n=30)		Both (n=43)		Neither (n=167)		χ ² P-value (all)	χ ² P-value (Female vs. Male)
	n	%	n	%	n	%	n	%	n	%		
Who is binge drinking?												
Mothers (inc. pregnant women)	32	10.4	23	7.5	0	0.0	9.0	2.9	0.0	0.0	0.000	0.000
Fathers	11	3.6	0	0.0	0	0.0	11.0	3.6	0.0	0.0	0.000	-
Young people (inc. children and students)	119	38.6	24.0	35.3	8.0	26.7	18.0	41.9	69.0	41.3	0.419	0.401
Older people	17	5.5	5	1.6	0	0.0	3.0	1.0	9.0	2.9	0.499	0.127
Which locations of binge drinking are mentioned?												
At home	46	14.9	14	4.5	1	0.3	10.0	3.2	21.0	6.8	0.047	0.029
On city streets	60	19.5	12	3.9	5	1.6	10.0	3.2	33.0	10.7	0.087	0.906
Pubs, bars and clubs	56	18.2	12	3.9	5	1.6	6.0	1.9	33.0	10.7	0.836	0.906
What types of drink are associated with binge drinking?												
Wine	58	18.8	23	7.5	2	0.6	11.0	3.6	22.0	7.1	0.001	0.004
Spirits	58	18.8	22	7.1	12	3.9	4.0	1.3	20.0	6.5	0.000	0.464
Beer	51	16.6	10	3.2	8	2.6	10.0	3.2	23.0	7.5	0.193	0.159
Cider	26	8.4	4	1.3	5	1.6	4.0	1.3	13.0	4.2	0.343	0.880
Alcopops	9	2.9	5	1.6	0	0.0	1.0	0.3	3.0	1.0	0.093	0.127
Shots	5	1.6	3	1.0	1	0.3	0.0	0.0	1.0	0.3	0.129	0.804
Cocktails	4	1.3	2	0.6	1	0.3	0.0	0.0	1.0	0.3	0.308	0.917
Fortified wine	4	1.3	1	0.3	0	0.0	2.0	0.6	1.0	0.3	0.186	0.504
Sparkling wine	3	1.0	1	0.3	0	0.0	1.0	0.3	1.0	0.3	0.676	0.504
Liqueurs	2	0.6	1	0.3	0	0.0	0.0	0.0	1.0	0.3	0.754	0.504
Lambrini	2	0.6	2	0.6	0	0.0	0.0	0.0	0.0	0.0	0.069	0.343
What harms of binge drinking are mentioned?												
Any personal harms	38	12.3	11	3.6	6	1.9	8	2.6	13	4.2	0.062	0.645
Harms to relationships	21	6.8	5	1.6	6	1.9	2	0.6	8	2.6	0.022	0.068
Harms to parenting	24	7.8	7	2.3	0	0.0	7	2.3	10	3.2	0.042	0.068
Any economic harms	107	34.7	16	5.2	8	2.6	17	5.5	66	21.4	0.080	0.739
Harms to the NHS	103	33.4	16	5.2	8	2.6	16	5.2	63	20.5	0.153	0.739
Harms to economic productivity	11	3.6	2	0.6	0	0.0	1	0.3	8	2.6	0.551	0.343
Any social harms	138	44.8	25	8.1	15	4.9	17	5.5	81	26.3	0.319	0.219
Social disorder and violence	128	41.6	19	6.2	14	4.5	16	5.2	79	25.6	0.044	0.071
Sexual assault and rape	19	6.2	9	2.9	4	1.3	2	0.6	4	1.3	0.005	0.989
Fear	15	4.9	0	0.0	0	0.0	2	0.6	13	4.2	0.044	-
Any health harms	192	62.3	40	13.0	21	6.8	26	8.4	105	34.1	0.755	0.293
Death	77	25.0	8	2.6	16	5.2	10	3.2	43	14.0	0.000	0.000
Physical health	138	44.8	32	10.4	11	3.6	14	4.5	81	26.3	0.215	0.339
Harms to appearance	46	14.9	19	6.2	0	0.0	8	2.6	19	6.2	0.001	0.001

Almost two-thirds of articles (n=192, 62.3%) described health harms related to binge drinking, while one-quarter of articles (n=77) mentioned risk of death. Mentions of specific categories of harm were recorded (Table 2), and relationships between gender and specific harms were identified. Mentioning of harms to appearance was significantly related to gender category (p=0.001), with 19 of the 46 articles mentioning harms to appearance mentioned females exclusively, while none of those 46 articles mentioned males exclusively. Similarly, harms to parenting were mentioned in seven articles (10.3%) exclusively mentioning females, but none exclusively mentioning males. Sexual assault and rape were mentioned in nine (2.9%) articles exclusively mentioning females, and four (13.3%) of the articles exclusively mentioning males. Significantly (p=0.008) more articles directly associated the risk of death from drinking with males (n=24, 7.8%) than with females (n=10, 3.3%). Conversely, non-fatal health harms of drinking were associated with females (n=69, 22.4%) significantly (p<0.000) more frequently than males (n=25, 8.1%).

Qualitative findings

Stark differences were evident in how men and women were described in the 46 articles that reported on ‘harms to appearance’. This broad category incorporated aspects of bodily appearance, clothing, self-presentation and behaviours. Two key themes emerged from thematic analysis of these 46 articles: physical appearance and loss of self-control.

Physical Appearance

Various harms to complexion were reported in relation to women, including the use of make-up to hide grey skin. Damage to hair, lips, noses and teeth were also discussed specifically in relation to women, as well as the use of less-specific terms such as ‘ravaged’ (The Sun, 13

Feb 2012). An article headlined '*Lambrini ruined my looks in 5 months*' (The Sun, 27 Sept 2013) associated an '*attractive*' woman becoming '*a swollen, haggard wreck*' with a brand of perry overtly marketed to young women. Weight gain caused by alcohol consumption was cited as both a product of alcohol use and a cause of 'drunkorexia', the practice of eating less to offset calories gained from alcohol drinks.[23] Drunkorexia was typically associated with women; one article stated that it '*affects mainly young women*' (Daily Mail, 19 Jul 2012) while another defined it as a condition in which '*calorie-conscious women skip meals in order to binge drink*' (Mirror, 20 Dec 2012).

For men, partial nudity was frequently related to drunken behaviours, such as lost and ripped clothes, and intentionally revealed genitals. Typifying this, the Daily Mail describes a male student '*flashing his penis in the street and laughing at his own loutish behaviour*' (19 June 2012). By comparison, partial nudity in women was more frequently related to clothing choices likely made while sober, with frequent mention of revealing clothing, particularly miniskirts, and impractical shoes. One article described a fancy dress event in which '*undergraduate girls take to the streets dressed in little more than their underwear*' (Daily Mail, 8 Oct 2012). Women were variously described as '*scantily-clad*' (Daily Mail, 22 Aug 2012), '*halfnaked*' (The Sun, 3 Mar 2013), and '*nearly bearing all*' (Daily Mail, 8 Oct 2012). Aside from choices made in sobriety, harms to appearance that could befall women during drinking episodes included smeared makeup, vomit in hair, stained clothes and gravel in knees. Articles also made frequent mentions of unintentional exposure of underwear or body parts by women, in contrast to their male counterparts, who typically exposed their genitals deliberately.

Loss of self-control

Descriptions of women’s and men’s loss of physical control differed. Some articles depicted drunken women burdening male partners. For example, two articles described women vomiting on their boyfriends, while one article described female students needing ‘*to be carried back to their rooms by boyfriends*’ (Daily Mail, 19 Jun 2012). No articles characterised men as relying on or burdening their partners while binge drinking.

Women’s aggressive behaviours were limited to verbal conflict, with drunken women characterised as noisy, argumentative, emotional and hard to control. An article describing Cardiff nightlife reported that drunken women ‘*scream at their boyfriends in shop doorways*’ and ‘*sob in public*’ (Daily Mail, 21 Aug 2012). Men were more typically associated with aggressive behaviours, including violent language and actions, indecent exposure, damaging property and ‘loutish’ (Daily Mail, 19 Jun 2012) behaviour. While male aggression was typically presented as endangering others, women were characterised as putting themselves at risk, sometimes of undesirable sexual situations and assault. One article reproduced Office of National Statistics data about fines for drunk and disorderly behaviour, presenting men as much more likely than women to receive such fines, but suggesting that the number of women receiving such fines is increasing more rapidly than men (Telegraph, 14 Jun 2009).

DISCUSSION

We examined UK newspaper and online news representations of ‘binge’ drinking, examining differences and similarities in representations of women’s and men’s drinking behaviours. Our quantitative analysis identified that women were associated with binge drinking more frequently than men, and that women’s drinking was problematised more than men’s. Both of these representations conflict with epidemiological evidence.[2] We found gender differences in the locations that binge drinking is described as taking place in, the health risks faced by binge drinkers, and mentions of parents’ binge drinking. Female binge drinking was

disproportionately associated with drinking wine, drinking in the home, and harms to appearance. Motherhood was mentioned in relation to binge drinking more frequently than fatherhood. There were significant gender differences in the attribution of health harms, with male binge drinking frequently associated with mortality and female binge drinking frequently associated with morbidity. Notably, harms to appearance were associated with women's binge drinking much more frequently than men's.

Our qualitative analysis focused on depictions of binge drinking as negatively affecting physical appearance (comprising bodily and other aspects of appearance) and self-control. Women engaged in binge drinking were presented as helpless, physically incapacitated, transgressive and burdening male partners, who were sometimes cast as carers for drunken women. Binge drinking men were associated with physical and verbal aggression, but not behaviours that marked them as weak or vulnerable; while women were characterised as endangering themselves, men were more likely to endanger others. The empowering, yet derogatory, 'ladette' narrative was notably absent, with articles instead depicting women as less able than men to maintain socially acceptable behaviour during episodic drinking. Binge drinking was characterised as impacting women's physical appearance more than men's. Articles associated binge drinking with both immediate and long-term damage to physical characteristics and self-presentation. Descriptions of binge-drinking women's dress were often judgemental in tone, typically characterising them as partially nude. Our analysis suggests that media representations of women's binge drinking do not focus solely on the effects of binge drinking on women, but also reflect broader social expectations about women's public behaviours.

As Measham & Ostergaard [24] argue, *'The public face of binge drinking, [...] a staple of early 21st century tabloid newspapers, became the young woman emulating male consumption patterns [...] with clothes askew stumbling around the city centre streets at*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

night' (p417). There may be unintended consequences of this disproportionate focus on women's – as opposed to men's - single episodic drinking in media reports. For example, it may reinforce harmful gender stereotypes by suggesting that drinking is more problematic for women than for men, and may encourage victim blaming in relation to sexual assaults after drinking.[25]

Our findings build on the growing body of literature about gendered media representations of alcohol, by specifically focusing on portrayals of binge drinking. Depictions of women as unable to control themselves physically and emotionally echoed the findings of Atkinson and colleagues,[16] while the burden this weakness was depicted as placing on men could be seen as corroborating Day and colleagues'[12] finding that traditionally male-dominated drinking environments and activities are perceived to be threatened by women's increasing involvement. Day and colleagues'[12] finding that women were presented as making themselves vulnerable to male aggression was replicated in our finding that articles presented women as in danger and men as dangerous. The depictions of women's dress and behaviours found in our analysis echoed Nicholls'[15] findings. We also identified that, when male whole or partial nudity was mentioned, it was typical presented as frivolous or ridiculous, while women's partial nudity was presented with an underlying morally-loaded, judgemental tone, perhaps resulting from women being viewed as publically breaking social conventions. Our analysis supports Herring and Berridge's[5] observation that binge drinking is popularly conceived in terms of specific social contexts, rather than being defined purely in terms of exceeding a certain quantity of alcohol in a limited time. Media framing of binge drinking as an activity of young men and women, often in public spaces, disproportionately emphasises the context of single episodic drinking and contributes to constructing binge drinking as something many people (including those who are older, or drinking in private settings) might not associate with. The disproportionate association of binge drinking with young peoples'

public, often anti-social (albeit, performed in social groups), drinking could have a damaging influence if it leads those who engage in single episodic drinking outside of these contexts not to identify their behaviours as harmful. The dominant narrative of binge drinking may have become unhelpful as a health concept. As an alternative, it may be useful to draw a clear distinction in health communications between 'binge drinking', as an inconsistently-defined, context-specific and value-laden term, and single episodic drinking, as a specific and widespread practice with a range of health consequences documented by research evidence.

Our findings have relevance to the development of alcohol guidelines. New guidelines propose reducing men's weekly maximum alcohol consumption to match women's, but media representations of binge drinking reinforce the idea that women and men differ in their binge drinking behaviours and the types of risks that they face. Portrayals of women as the more problematic binge drinkers, and men's binge drinking as aligned with male gender roles and therefore less problematic, may contradict the recommendation that men should drink no more than women, therefore representing a barrier to public understanding and acceptance of the proposed guidelines.

Our conclusions are subject to limitations. While mass media's influence on public perceptions is well-evidenced and well-understood[7], content analysis alone cannot determine the extent to which audiences' understandings correlate with media representations; audiences are subject to many influences beyond print and online news media, and do not consume media in a passive, non-critical way. Studying other forms of media, such as television or social media, could help create a more complete understanding of media representations of binge drinking. Furthermore, a larger sample size, achieved by choosing a longer timeframe or including more publications, could have increased the validity of our conclusions. A potential limitation of the data used is that the Nexis and Newsbank databases store articles in text-only format, without the images used in the original

versions. Coders noted that text-only articles that appear factual and impartial could be perceived as more morally loaded with the original photographs included. Similarly, some articles may only reveal implicit gender bias when viewed with the original photographs included; this is illustrated in an article about binge drinking on The Sun website that contains seven images of women drinking and none of men, while the text makes no mention of gender.[26] With these limitations taken into account, this comprehensive content analysis of a two-year period of coverage usefully contributes to understandings of gendered media representations of drinking, with a specific focus on binge drinking.

In conclusion, our analysis suggests that popular representations of binge drinking may be harmful in three ways. Firstly, associating binge drinking with certain demographics and situations could lead the public to underestimate the health risks of single episodic drinking among those not included in the ‘binge drinking’ demographics, particularly older people. The morally-loaded representations of women’s binge drinking may reproduce harmful stereotypes and stigma about the vulnerability of drunken women and the social unacceptability of female drunkenness. Finally, media content reinforcing a skewed representation of binge drinking may present a challenge to public health stakeholders seeking to promote evidence-based information, recommendations and policies with the goal of reducing alcohol-related health harms.

The misrepresentations identified by this research highlight a need for health advocates to engage with mass media to promote clear, evidence-informed messages about single episodic drinking to help better align reporting on harmful drinking with evidence, with the ultimate goal of improving public understandings of harmful drinking behaviours. Media communications strategies may seek to avoid stereotypes of ‘binge drinking’ that implicitly define binge drinking in terms of social contexts and behaviours not directly related to alcohol consumption, instead offering clear, value-free definitions of single episodic drinking

based on specific quantities of alcohol and specific, time-bound episodes. Identifying the need to improve communication of single episodic drinking is particularly relevant given the current public debate around new alcohol guidelines,[3] which may or may not specify recommended limits to single-episode alcohol consumption, depending on the outcome of the current consultation.

Contributorship statement

CE and SH conceived of the study and initiated the study design. OM and GF performed quantitative coding. CP and GF performed qualitative coding. OM prepared the initial draft of the paper. All authors contributed substantially to redrafting. The final manuscript was prepared by CP and approved by CE, GF, OM and SH.

Competing interests

The authors declare no competing interests.

Funding

This work was supported by the UK Medical Research Council as part of the Understandings and Uses of Public Health Research programme [MC_U130085862, MC_UU_12017/6] and the Informing Healthy Public Policy programme [MC_UU_12017/15] and by the Chief Scientist Office of the Scottish Government Health Directorates [SPHSU15] at the MRC/CSO Social and Public Health Sciences Unit, University of Glasgow. The funding body had no role in the design, collection, analysis or interpretation of this study.

Data sharing statement

No additional unpublished data from the study have been made available.

REFERENCES

1. Smith L, Foxcroft DR. Drinking in the UK: An exploration of trends: Joseph Rowntree Foundation, 2009.
2. Office for National Statistics. Adult Drinking Habits in Great Britain, 2013: Office for National Statistics, 2015.
3. Department of Health. Alcohol Guidelines Review: report from the guidelines development group to the UK Chief Medical Officers: Department of Health, 2015.
4. Department of Health. Sensible Drinking: The Report of an Inter-Departmental Working Group, 1995.
5. Herring R, Berridge V, Thom B. Binge drinking: an exploration of a confused concept. *J. Epidemiol. Community Health* 2008;**62**(6):476-79
6. Guise JMF, Gill JS. 'Binge drinking? It's good, it's harmless fun': a discourse analysis of accounts of female undergraduate drinking in Scotland. *Health Educ. Res.* 2007;**22**(6):895-906 doi: 10.1093/her/cym034published Online First: Epub Date]].
7. McCombs ME, Shaw DL. The agenda-setting function of mass media. *Public Opin. Q.* 1972;**36**(2):176-87
8. Scheufele DA, Tewksbury D. Framing, agenda setting, and priming: The evolution of three media effects models. *J. Commun.* 2007;**57**(1):9-20
9. Wilsnack RW, Wilsnack SC. *Gender and alcohol: Individual and social perspectives*: Rutgers Center of Alcohol Studies, 1997.
10. Casswell S. Public Discourse on Alcohol. *Health Prom. Int.* 1997;**12**(3):251-57
11. Dorfman L. Studying the news on public health: How content analysis supports media advocacy. *American Journal of Health Behavior* 2003;**27**(Supplement 3):S217-S26
12. Day K, Gough B, McFadden M. "Warning! alcohol can seriously damage your feminine health" a discourse analysis of recent British newspaper coverage of women and drinking. *Feminist media studies* 2004;**4**(2):165-83
13. Wood K, Patterson C, Katikireddi SV, Hilton S. Harms to 'others' from alcohol consumption in the minimum unit pricing policy debate: a qualitative content analysis of UK newspapers (2005–12). *Addiction* 2014;**109**(4):578-84 doi: 10.1111/add.12427published Online First: Epub Date]].
14. Lyons AC, Dalton SI, Hoy A. 'Hardcore Drinking': Portrayals of Alcohol Consumption in Young Women's and Men's Magazines. *Journal of health psychology* 2006;**11**(2):223-32
15. Nicholls J. UK news reporting of alcohol: An analysis of television and newspaper coverage. *Drugs (Abingdon Engl)* 2011;**18**(3):200-06
16. Atkinson A, Kirton A, Sumnall H. The gendering of alcohol in consumer magazines: An analysis of male and female targeted publications. *Journal of gender studies* 2012;**21**(4):365-86
17. National Readership Survey. National Readership Survey. Secondary National Readership Survey 2012. <http://www.nrs.co.uk/>.
18. Newsworks. Newsworks. Secondary Newsworks 2012. www.nmauk.co.uk.
19. Ofcom. Communications Market Report 2015. London: Ofcom, 2015.
20. Hilton S, Patterson C, Teyhan A. Escalating coverage of obesity in UK newspapers: The evolution and framing of the 'obesity epidemic' from 1996 to 2010. *Obesity (Silver Spring)* 2012;**20**(8):1688–95 doi: doi:10.1038/oby.2012.27published Online First: Epub Date]].
21. Patterson C, Katikireddi SV, Wood K, Hilton S. Representations of minimum unit pricing for alcohol in UK newspapers: a case study of a public health policy debate. *Journal*

- of *Public Health* 2015;**37**(1):40-49 doi: 10.1093/pubmed/fdu078published Online First: Epub Date]].
22. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative research in psychology* 2006;**3**(2):77-101
23. Barry AE, Piazza-Gardner AK. Drunkorexia: Understanding the Co-occurrence of Alcohol Consumption and Eating/Exercise Weight Management Behaviors. *J. Am. Coll. Health* 2012;**60**(3):236-43 doi: 10.1080/07448481.2011.587487published Online First: Epub Date]].
24. Measham F, Østergaard J. The public face of binge drinking: British and Danish young women, recent trends in alcohol consumption and the European binge drinking debate. *Probation Journal* 2009;**56**(4):415-34
25. Fitzgerald N, Angus K, Emslie C, Shipton D, Bauld L. Gender differences in the impact of population-level alcohol policy interventions: evidence synthesis of systematic reviews. *Addiction* in press
26. The Sun. Your friends, bad habits or anxiety... take our quiz to find YOUR booze binge trigger. Secondary Your friends, bad habits or anxiety... take our quiz to find YOUR booze binge trigger 2016.
<http://www.thesun.co.uk/sol/homepage/fabulous/7112837/Your-friends-bad-habits-or-anxiety-take-our-quiz-to-find-YOUR-booze-binge-trigger.html>.




Figure 1. Frequency of publication of articles about 'binge' drinking by quarter
(Figure 1)
272x193mm (300 x 300 DPI)

BMJ Open

A content analysis of UK newspaper and online news representations of women's and men's 'binge' drinking: a challenge for communicating evidence-based messages about single-episodic drinking?

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-013124.R1
Article Type:	Research
Date Submitted by the Author:	04-Oct-2016
Complete List of Authors:	Patterson, Chris; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit Emslie , Carol; Glasgow Caledonian University Institute for Applied Health Research Mason, Olivia; Durham University, Department of Geography Fergie, Gillian; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit Hilton, Shona; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit
Primary Subject Heading:	Communication
Secondary Subject Heading:	Sociology, Public health
Keywords:	PUBLIC HEALTH, QUALITATIVE RESEARCH, SOCIAL MEDICINE

SCHOLARONE™
Manuscripts

A content analysis of UK newspaper and online news representations of women’s and men’s ‘binge’ drinking: a challenge for communicating evidence-based messages about single-episodic drinking?

Patterson^{1*}, Emslie², Mason³, Fergie¹, Hilton¹

¹MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB UK

²Institute for Applied Health Research/School of Health and Life Sciences, Glasgow Caledonian University, Cowcaddens Road, Glasgow G4 0BA UK

²Department of Geography, Durham University, Lower Mountjoy South Road, Durham DH1 3LE UK

Corresponding author:

Chris Patterson

MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB UK

chris.patterson@glasgow.ac.uk

0141 353 7631

Keywords: gender; media; alcohol; binge drinking

Running head: Media representations of ‘binge’ drinking

ABSTRACT

Objectives: In the UK, men's alcohol-related morbidity and mortality still greatly exceeds women's, despite an increase in women's alcohol consumption in recent decades. Recently-proposed UK alcohol guidelines introduce gender-neutral weekly alcohol consumption limits. This study explores how UK newspaper and online news represent women's and men's 'binge' drinking to identify opportunities to better align reporting of harmful drinking with evidence.

Design: Quantitative and qualitative content analysis of 308 articles published in seven UK national newspapers and the BBC News website between 1 January 2012 and 31 December 2013.

Results: Articles associated women with 'binge' drinking more frequently than men and presented women's drinking as more problematic. Men were more frequently characterised as violent or disorderly, while women were characterised as out of control, putting themselves in danger, harming their physical appearance and burdening men. Moralistic descriptions of female 'binge' drinkers' clothing and appearance were typical.

Conclusions: The UK news media's disproportionate focus on women's 'binge' drinking is at odds with epidemiological evidence, may reproduce harmful gender stereotypes and may obstruct public understandings of the gender-neutral weekly consumption limits in newly-proposed alcohol guidelines. In order to better align reporting of harmful drinking with current evidence, public health advocates may engage with the media with a view to shifting media framing of 'binge' drinking away from specific groups (young people; females) and contexts (public drinking) and towards the health risks of specific drinking behaviours, which affect all groups regardless of context.

Strengths and limitations of this study

- Quantitative and qualitative content analysis of a large, comprehensive, two-year sample of UK national newspaper and online news about ‘binge’ drinking.
- The findings illustrate how media portrayals of ‘binge’ drinking could be harmful, and identify opportunities for these portrayals to be better aligned with evidence.
- Content analysis facilitates understandings of the messages being presented to the public, but cannot determine the extent to which audiences’ understandings are influenced by media representations of specific issues.
- Some findings could have differed if the data analysed had gone beyond text articles, for example by incorporating images, video and social media content.

INTRODUCTION

While the gap between women's and men's excessive and harmful consumption of alcohol has narrowed in the UK in recent years,[1] men still drink more than women[2], experience more drink-related health and social problems and face twice women's alcohol-related mortality (15.9 to 7.8 deaths per 100,000 population respectively in 2012[2]). New UK drinking guidelines propose equal weekly alcohol consumption limits of 14 units for both men and women spread evenly over three days or more.[3] These proposed gender-neutral limits differ from previous guidelines, which, in 1995, suggested a weekly maximum of 21 units for men (with no more than 4 units in any day) and 14 units for women (with no more than 3 units in any day).[4] Different limits for men and women are the norm in alcohol guidelines internationally, but there is precedent for gender-neutral limits in Australian and Portuguese recommendations.[5] The gender-neutral limits proposed by the UK Chief Medical Officers are rationalised by the observation that, while women can experience more long-term health harms from alcohol, men are at more risk of immediate health harms.[3] In light of the shift towards gender-neutrality in alcohol consumption recommendations, it is timely to consider public perceptions of gender differences in drinking behaviours.

As with most health issues, popular perceptions of drinking are likely to be influenced by mass media representations,[6, 7] and, in turn, differences in representations of women's and men's drinking are heavily influenced by how societies regulate gender roles.[8] Media content analysis cannot tell us if, and how, specific content influences audiences' understandings and behaviours, and content analysis findings must be considered with that inherent limitation in mind. Nonetheless, exploring media representations of men's and women's drinking behaviours allows us to examine how shared cultural values around alcohol are articulated and constructed[9], which might inform efforts to improve media representations, and therefore public understandings, of harmful drinking behaviours.[10]

Understanding gender differences in media representations of drinking behaviours is of heightened research interest in light of the gender-neutral consumption limits in the newly-proposed guidelines.[3]

Research illustrates a clear gender divide in media portrayals of drinking behaviours, with men's drinking normalised and women's problematised. Day and colleagues[11] found that UK newspapers frequently characterised women as departing from idealised notions of femininity in terms of appearance (e.g. weight gain, deeper voices, loss of good looks) and motherhood (e.g. reduced fertility and unborn children). They found that men were framed as violent, but partial responsibility for men's violence was attributed to women, who were framed as sexual predators invading traditionally male-dominated drinking environments. Conversely, Wood and colleagues[12] found that UK newspaper coverage of proposed minimum unit pricing (MUP) policies presented women as being at 'risk of harm' from male aggression. Lyons, Dalton & Hoy[13] found that both young women's and young men's magazines framed binge drinking as normative, adult and professional, with young men's magazines associating men's drinking with traditional masculine images but deriding young women's drinking behaviours. Nicholls[14] found that UK television and newspaper news associated men's drinking with violence (both as perpetrators and victims) and women's drunkenness with unfeminine and undignified behaviours, such as loss of consciousness or partial nudity. Similarly, Atkinson and colleagues[15] found that UK magazines targeted at teenagers depicted women's drinking as more problematic than men's, and portrayed women as 'behaving like men' in male spaces, sexualised and highly emotional.

As Nicholls[14] found, 'binge drinking', also called single-episodic drinking, is a key focus of media representations of harmful drinking; *"problem drinking is less commonly associated with dependence and more commonly associated with binge, harmful and hazardous drinking"*. Herring and Berridge[16] describe binge drinking as a 'confused concept' that has

historically been, and continues to be, defined inconsistently. They suggest that ‘binge’ drinking is currently portrayed as a youth issue, despite evidence that single-episodic drinking is performed by various age groups[2], highlighting a potential area of media misrepresentation of population drinking behaviours. Qualitative evidence indicates that young women in Scotland define ‘binge’ drinking in terms of types of behaviour rather than the quantity of alcohol consumed,[17] which may present an obstacle to clearly and objectively defining ‘binge’ drinking. Mixed-method evidence suggests that both male and female students in England perceive binge drinking and public drunkenness as masculine behaviours[18]. Two studies based on qualitative data from New Zealand have highlighted the role of alcohol-related behaviours in young people’s gender identities. Firstly, Willott and Lyons[19] found that young men and women perceived consuming large quantities of alcohol in a single episode (among other behaviours) as a key performance of masculinity, and highlighted the identity negotiations undertaken by men who do not engage in these normative, masculine drinking behaviours. Secondly, Hutton and colleagues[20] examined the challenges that young women experience in curating social media personae that balance engagement in “the culture of intoxication” (p.88) with maintaining respectability. Given the primacy of binge drinking in UK media coverage of harmful alcohol use,[14] binge drinking is an important lens through which to examine gendered representations of drinking behaviours.

This study comprises a comprehensive, mixed-methods content analysis of two years of UK media coverage of binge drinking, designed to contribute new insights to a growing body of literature about gendered media representations of alcohol, with a particular focus on binge drinking. Our aim is to improve understandings of UK newspaper and online news representations of women’s and men’s ‘binge’ drinking, focusing not on how ‘binge’ drinking is defined, but rather on how different types of drinking behaviour are gendered in

media content wherein those behaviours are labelled as ‘binge’ drinking. While the limitations of media content analysis must be taken into account, the improved understandings produced by this research may help to inform efforts to better align media representations of harmful drinking with current evidence, which could in turn improve public understandings of the risks of single-episodic drinking.

METHODS

We selected seven highly-circulated[21, 22] UK national newspapers, including their Sunday counterparts, and the most-read exclusively-online news website[23] (Table 1). The chosen newspaper publications represented three genres: quality, middle-market tabloid and tabloid. This typology helps to ensure a sample that represents diverse readerships in terms of age, social class and political alignment.[24] Quality newspapers are those that were traditionally printed in broadsheet format, have predominantly middle-class audiences, are politically diverse and are serious in tone. Middle-market tabloids are printed in tabloid format and are less serious in tone than quality-genre newspapers, and have a predominantly older, middle-class, right-wing audience. Tabloids are less serious than middle-market tabloids, typically more sensationalist, politically diverse and have predominantly working-class audiences.

We selected a search period of 1 January 2012 to 31 December 2013, during which alcohol was subject to heightened media interest due to debates in the UK and Scottish parliaments around MUP.[25] Much of the media coverage of MUP was related to ‘binge’ drinking due to the UK Government’s Alcohol Strategy[26] overtly positioning ‘binge’ drinking as the primary target of MUP. Relevant newspaper articles, were identified using the Nexis database including articles from online editions of the two of the selected publications (The Guardian and The Daily Mail) that are archived in Nexis. BBC News articles were identified using the search function on the BBC website. The search string identified articles that contain both

three or more mentions of “binge” and one or more mentions of “drink OR drinker OR drinkers OR drinking”. Duplicate articles were removed, and articles were manually excluded if they did not predominantly focus on binge drinking in the UK, or if they were not in the news, feature or editorial formats. Initial searches identified 537 articles, of which 308 met the inclusion criteria and were eligible for detailed coding and analysis.

For peer review only

Table 1. Summary of publications and articles in the sample

Genre / medium	Publication	All articles		Article format					
				Standard		Feature		Editorial	
		n	%	n	%	n	%	n	%
Quality (n=86)	Guardian / Observer	58	18.8	48	20.3	9	17.3	1	5.3
	Independent / Independent on Sunday	17	5.5	11	4.6	5	9.6	1	5.3
	Daily Telegraph / Sunday Telegraph	52	16.9	44	18.6	5	9.6	3	15.8
Middle-market tabloids (n=39)	Daily Mail / Mail on Sunday	54	17.5	33	13.9	18	34.6	3	15.8
	Express / Sunday Express	13	4.2	8	3.4	3	5.8	2	10.5
Tabloids (n=75)	Daily Mirror / Sunday Mirror	13	4.2	10	4.2	2	3.8	1	5.3
	The Sun / News of the World	62	20.1	46	19.4	9	17.3	7	36.8
Online (n=39)	BBC News website	39	12.7	37	15.6	1	1.9	1	5.3
Total:		308	100	237	100	52	100	19	100

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

To systematically and comprehensively code and analyse the media content, we used a mixed-methods content analysis, first using quantitative content analysis to measure the frequency of content within the articles across the whole sample, and secondly using qualitative content analysis for more in-depth, reflexive analysis of the meanings of the content of a subsample of articles.[27] From this perspective, quantitative content analysis is concerned with measuring and analysing manifest content (the surface-level content of the articles, the coding of which does not require interpretation on the part of the coder, such that it can be recorded relatively objectively), while qualitative content analysis is concerned with latent content (the underlying meanings of the text, as interpreted by coders in an inherently subjective process).[28] The mixed-methods approach in this research comprised the following steps: constructing a coding frame; coding manifest content using the coding frame; establishing the reliability of the data collected and excluding unreliable data; analysing the quantitative the data; identifying aspects of manifest content to examine further using qualitative analysis; and finally performing thematic analysis of the latent content of articles containing the manifest content of interest.

To construct a coding frame with which to code manifest content, SH and CE read randomly-selected articles from the sample in batches of 20, recording potential thematic categories relevant to the research topic as they emerged. At the point where a batch of 20 articles was read without any new categories emerging, the list of categories was deemed to have reached saturation. The collected thematic categories were grouped within three broad thematic categories: how is binge drinking described and which sections of the population are associated with this behaviour?; how are the drivers of the binge drinking described and who is to blame?; and what are the consequences of binge drinking? These categories were chosen by a combination of *a priori* knowledge of the research topic and understandings of the content of the sample that emerged from close reading of the articles.[25] The final coding

frame comprised the three broad thematic categories and their sub-categories, as well as codes for more routine aspects of articles, such as publication, article format and word count. In addition, coders recorded whether each article mentioned males and females, allowing articles to be divided into four gender categories: those that mention females exclusively, those that mention males exclusively, those that mention both males and females and those that mention neither males nor females explicitly. These codes enabled analysis of whether themes varied significantly by the gender focus of articles.

To collect quantitative data, OM read each article in turn, using the coding frame to record whether each theme was present within its manifest content. To ensure consistency of coding, GF coded a random subsample of 39 articles (12.7%). Linearly-weighted kappa tests of inter-rater agreement between OM and GF were then performed on each variable across those 39 double-coded articles, and variables that returned a coefficient below 0.8 were discarded from the study to ensure that only variables with strong agreement were retained for analysis. The coding frame data were then entered into SPSS for analysis. Statistical procedures comprised: a simple linear regression examining the relationship between publication quarter and the count of articles published; chi-square tests of whether thematic variables varied significantly by gender-focus, publication genre or article format; and paired t-tests comparing the means of thematic variables. Where statistical significance is presented in the results, p-values are reported in terms of being significant ($p<0.05$) or highly significant ($p<0.01$).

Following quantitative analysis, the thematic category of ‘harms to appearance’ was deemed noteworthy and suitable for qualitative coding and analysis. The decision to focus on that theme was informed by stark gender differences in content that emerged from the quantitative analysis, and an expectation that valuable understandings could be gained from deeper analysis. To analyse the latent content of the 46 articles that had been coded as mentioning ‘harms to appearance’, GF and CP employed a thematic analysis approach[29] using NVivo

10, closely reading each article to generate initial codes, which were then collated into potential themes. GF and CP collaborated closely to assure the themes were defined clearly and that they worked across the 46 articles. The findings from the thematic analysis are presented in terms of two broad themes, using typical quotations to illustrate article content.

RESULTS

Quantitative findings

Between 1 January 2012 and 31 December 2013, 308 articles about binge drinking were published in the seven print publications and one website included in the sample. Of these articles, 86 (27.9%) were published in 'quality' newspapers, 39 (12.7%) in middle-market tabloids, 75 (24.4%) in tabloids and 108 (35.1%) on BBC News (Table 1). The majority of articles were standard news format (n=237, 76.9%), while 52 (16.9%) were feature articles and 19 (6.2%) editorials. The frequency of articles published per quarter decreased across the two-year period (Figure 1), but publication quarter was not a statistically significant predictor of article frequency (coefficient -6.714, p=0.088). There was a peak of 56 articles in March 2012, 66.1% of which were related to MUP, in particular the UK Prime Minister's announcement of plans to introduce MUP. There was elevated reporting from November 2012 to January 2013 (n=62), during which 53.2% of articles mentioned MUP, largely related to opposition to MUP within the UK Cabinet. In total, 133 (43.2%) articles mentioned MUP.

[Figure 1]

Articles were coded according to whether they mentioned binge drinking in females exclusively (n=68, 22.1%), males exclusively (n=30, 9.7%), both males and females (n=43,

14.0%), or neither males nor females explicitly (n=167, 54.2%). To be coded as mentioning both males and females, an article had to include specific discussion of each gender individually. There was no significant association between gender category and publication genre (p=0.382) or between gender category and article format (p=0.303). Furthermore, neither publication genre nor article format were significantly associated with differences in reporting on any of the thematic categories listed in Table 2.

Articles were coded to identify the characteristics of binge drinkers and the specific drinks associated with binge drinking. Thirty-two (10.4%) of all 308 articles associated mothers (including pregnant women) with binge drinking, highly significantly (p<0.000) more than mentioned fathers (n=11, 3.6%) (Table 2). There were no statistically significant relationships between gender focus and mentioning either younger or older people. Articles focused exclusively on women were significantly (p=0.029) more likely to refer to 'binge' drinking in the home than those that focused exclusively on men (Table 2). No other location was significantly associated with gender category. Wine (n=58, 18.8%) was the only type of drink significantly associated more with one gender than the other, being predominantly mentioned in articles focusing exclusively on females (p=0.004).

Table 2. Social groups, locations and harms associated with binge drinking

	All articles (n=308)		Only females (n=68)		Only males (n=30)		Both (n=43)		Neither (n=167)		χ ² P-value (all)	χ ² P-value (Female vs. Male)
	n	%	n	%	n	%	n	%	n	%		
Who is binge drinking?												
Mothers (inc. pregnant women)	32	10.4	23	7.5	0	0.0	9.0	2.9	0.0	0.0	0.000**	0.000**
Fathers	11	3.6	0	0.0	0	0.0	11.0	3.6	0.0	0.0	0.000**	-
Young people (inc. children and students)	119	38.6	24.0	35.3	8.0	26.7	18.0	41.9	69.0	41.3	0.419	0.401
Older people	17	5.5	5	1.6	0	0.0	3.0	1.0	9.0	2.9	0.499	0.127
Which locations of binge drinking are mentioned?												
At home	46	14.9	14	4.5	1	0.3	10.0	3.2	21.0	6.8	0.047*	0.029*
On city streets	60	19.5	12	3.9	5	1.6	10.0	3.2	33.0	10.7	0.087	0.906
Pubs, bars and clubs	56	18.2	12	3.9	5	1.6	6.0	1.9	33.0	10.7	0.836	0.906
What types of drink are associated with binge drinking?												
Wine	58	18.8	23	7.5	2	0.6	11.0	3.6	22.0	7.1	0.001**	0.004*
Spirits	58	18.8	22	7.1	12	3.9	4.0	1.3	20.0	6.5	0.000**	0.464
Beer	51	16.6	10	3.2	8	2.6	10.0	3.2	23.0	7.5	0.193	0.159
Cider	26	8.4	4	1.3	5	1.6	4.0	1.3	13.0	4.2	0.343	0.880
Alcopops	9	2.9	5	1.6	0	0.0	1.0	0.3	3.0	1.0	0.093	0.127
Shots	5	1.6	3	1.0	1	0.3	0.0	0.0	1.0	0.3	0.129	0.804
Cocktails	4	1.3	2	0.6	1	0.3	0.0	0.0	1.0	0.3	0.308	0.917
Fortified wine	4	1.3	1	0.3	0	0.0	2.0	0.6	1.0	0.3	0.186	0.504
Sparkling wine	3	1.0	1	0.3	0	0.0	1.0	0.3	1.0	0.3	0.676	0.504
Liqueurs	2	0.6	1	0.3	0	0.0	0.0	0.0	1.0	0.3	0.754	0.504
Lambrini	2	0.6	2	0.6	0	0.0	0.0	0.0	0.0	0.0	0.069	0.343
What harms of binge drinking are mentioned?												
Any personal harms	38	12.3	11	3.6	6	1.9	8	2.6	13	4.2	0.062	0.645
Harms to relationships	21	6.8	5	1.6	6	1.9	2	0.6	8	2.6	0.022*	0.068
Harms to parenting	24	7.8	7	2.3	0	0.0	7	2.3	10	3.2	0.042*	0.068
Any economic harms	107	34.7	16	5.2	8	2.6	17	5.5	66	21.4	0.080	0.739
Harms to the NHS	103	33.4	16	5.2	8	2.6	16	5.2	63	20.5	0.153	0.739
Harms to economic productivity	11	3.6	2	0.6	0	0.0	1	0.3	8	2.6	0.551	0.343
Any social harms	138	44.8	25	8.1	15	4.9	17	5.5	81	26.3	0.319	0.219
Social disorder and violence	128	41.6	19	6.2	14	4.5	16	5.2	79	25.6	0.044*	0.071
Sexual assault and rape	19	6.2	9	2.9	4	1.3	2	0.6	4	1.3	0.005**	0.989
Fear	15	4.9	0	0.0	0	0.0	2	0.6	13	4.2	0.044*	-
Any health harms	192	62.3	40	13.0	21	6.8	26	8.4	105	34.1	0.755	0.293
Death	77	25.0	8	2.6	16	5.2	10	3.2	43	14.0	0.000**	0.000**
Physical health	138	44.8	32	10.4	11	3.6	14	4.5	81	26.3	0.215	0.339
Harms to appearance	46	14.9	19	6.2	0	0.0	8	2.6	19	6.2	0.001**	0.001**

*Statistically significant at the 0.05 level

**Statistically significant at the 0.01 level

Almost two-thirds of articles (n=192, 62.3%) described health harms related to binge drinking, while one-quarter of articles (n=77) mentioned risk of death. Mentions of specific categories of harm were recorded (Table 2), and relationships between gender and specific harms were identified. Mentioning harms to appearance was highly significantly related to gender category (p=0.001); 19 of the 46 articles that mentioned harms to appearance mentioned females exclusively, while none of those 46 articles mentioned males exclusively. Harms to parenting were mentioned more frequently in articles (10.3%) that exclusively mentioned females, and none that exclusively mentioned males (0=0.042), but the difference was not significant (p=0.068). Sexual assault and rape were mentioned more frequently in articles exclusively mentioning females (n=9, 2.9%) than those exclusively mentioning males (n=4, 13.3%), but not significantly (p=0.989). More articles directly associated the risk of death from drinking with males (n=16, 20.8%) than with females (n=8, 10.4%), and this difference was highly significant (p<0.000). Conversely, non-fatal physical health harms were associated with females (n=32, 23.2%) more frequently than males (n=11, 8.0%), albeit not significantly so (p=0.339).

Qualitative findings

As identified in the quantitative findings, just eight of the 46 articles that reported on ‘harms to appearance’ mentioned men, and none did so exclusively. These 46 articles were subjected to thematic analysis. A broad conceptualisation of ‘harms to appearance’ was used that included behavioural as well as aesthetic aspects of physical appearance that may be influenced (or perceived to be influenced) by single-episodic drinking. Using this broad definition of ‘harms to appearance’, two key themes emerged from thematic analysis of the subsample: physical appearance and loss of self-control.

Physical Appearance

Various harms to complexion were reported in relation to women, including the use of make-up to hide grey skin. Articles described specific damage to women's features, including hair, lips, noses and teeth, as well as using less-specific descriptions of physical damage such as 'ravaged' (The Sun, 13 Feb 2012). An article headlined '*Lambrini ruined my looks in 5 months*' (The Sun, 27 Sept 2013) associated an 'attractive' woman becoming '*a swollen, haggard wreck*' with a brand of perry overtly marketed to young women. Weight gain caused by alcohol consumption was cited as both a product of alcohol use and a cause of 'drunkorexia', the practice of eating less to offset calories gained from alcoholic drinks.[30] Drunkorexia was typically associated with women; one article stated that it '*affects mainly young women*' (Daily Mail, 19 Jul 2012) while another defined it as a condition in which '*calorie-conscious women skip meals in order to binge drink*' (Mirror, 20 Dec 2012).

Within the few articles in the subsample that mentioned men, male partial nudity was related to drunken behaviours, such as lost and ripped clothes, and intentional indecent exposure. Typifying this, the Daily Mail described a male student '*flashing his penis in the street and laughing at his own loutish behaviour*' (19 June 2012). By comparison, partial nudity in women was more typically related to clothing choices likely made while sober, with descriptions of revealing clothing, particularly miniskirts, and impractical shoes. One article described a fancy dress event in which '*undergraduate girls take to the streets dressed in little more than their underwear*' (Daily Mail, 8 Oct 2012). Women were variously described as '*scantily-clad*' (Daily Mail, 22 Aug 2012), '*half-naked*' (The Sun, 3 Mar 2013), and '*nearly bearing all*' (Daily Mail, 8 Oct 2012). Aside from choices of attire made in sobriety, harms to appearance that could befall women during drinking episodes included smeared makeup, vomit in hair, stained clothes and gravel embedded in knees. Articles also made

1
2
3 mention of unintentional exposure of underwear or body parts by women, which was in
4
5 contrast to males, who were typically described as exposed their genitalia deliberately.
6
7

8
9 Loss of self-control

10
11 Descriptions of women's and men's loss of physical control differed. Some articles depicted
12
13 drunken women as burdening male partners. For example, two articles described women
14
15 vomiting on their boyfriends, while one described female students needing '*to be carried*
16
17 *back to their rooms by boyfriends*' (Daily Mail, 19 Jun 2012). No articles characterised
18
19 'binge' drinking men as relying on, or burdening, their partners.
20
21

22
23 Women's aggressive behaviours were limited to verbal conflict, with drunken women
24
25 characterised as noisy, argumentative, emotional and hard to control. An article describing
26
27 Cardiff nightlife reported that drunken women '*scream at their boyfriends in shop doorways*'
28
29 and '*sob in public*' (Daily Mail, 21 Aug 2012). Men were more typically associated with
30
31 aggressive behaviours, including violent language and actions, damaging property and non-
32
33 specific 'loutish' (Daily Mail, 19 Jun 2012) behaviour. While male aggression was presented
34
35 as endangering others, women were characterised as putting themselves at risk, sometimes of
36
37 undesirable sexual situations and assault. One article reproduced Office of National Statistics
38
39 data about fines for drunk and disorderly behaviour, presenting men as much more likely than
40
41 women to receive such fines, but suggesting that the number of women receiving the fines is
42
43 increasing more rapidly than men (Telegraph, 14 Jun 2009).
44
45
46
47

48
49 **DISCUSSION**

50
51 We examined UK newspaper and online news representations of 'binge' drinking, exploring
52
53 differences and similarities in representations of women's and men's drinking behaviours.
54
55 Our quantitative analysis identified that women were associated with binge drinking more
56
57
58
59
60

frequently than men, which conflicts with epidemiological evidence[2] and young people's perceptions of 'binge' drinking as a masculine activity.[18] We found gender differences in the locations in which that 'binge' drinking is described as taking place, the health risks faced by binge drinkers, and discussion of parents' binge drinking. Female 'binge' drinking was typically associated with drinking wine, drinking in the home, and harms to appearance. Motherhood was mentioned in relation to 'binge' drinking more frequently than fatherhood. There were significant gender differences in the attribution of health harms, with male 'binge' drinking frequently associated with mortality and female 'binge' drinking frequently associated with morbidity. Women's drinking was typically presented as more problematic than men's, and, notably, harms to appearance were associated with women's 'binge' drinking much more frequently than men's. The articles in our sample associated wine with women's 'binge' drinking significantly more than men's, but our data did not identify significant gender associations with other specific types of drink.

Our qualitative analysis focused on depictions of the relationship between 'binge' drinking and appearance, comprising aspects of both physical appearance and behaviour. Women engaged in 'binge' drinking were presented as helpless, physically incapacitated and transgressive, and as burdens to male partners, who were sometimes cast as carers for drunk women. 'Binge' drinking men were associated with physical and verbal aggression, but not behaviours that marked them as weak or vulnerable; while women were characterised as endangering themselves, men were more likely to endanger others. Articles typically depicted women as less able than men to maintain socially acceptable behaviour during single-episodic drinking. 'Binge' drinking was characterised as impacting women's physical appearance more than men's. Articles associated binge drinking with both immediate and long-term damage to physical characteristics and self-presentation. Descriptions of binge-drinking women's dress had a moralistic tone that was absent in representations of men, with

articles typically focusing on degrees of nudity and implicitly questioning the propriety of women's chosen attire. Our analysis suggests that media representations of women's 'binge' drinking do not focus solely on the effects of binge drinking on women, but also reflect broader social expectations about women's public behaviours.

Measham & Ostergaard [31] suggest that newspapers have constructed 'binge' drinking as a problem of young women adopting male behaviours:

The public face of binge drinking, [...] a staple of early 21st century tabloid newspapers, became the young woman emulating male consumption patterns [...] with clothes askew stumbling around the city centre streets at night' (p417).

There may be unintended consequences of this disproportionate focus on women's – as opposed to men's – single-episodic drinking in media reports. For example, it may reinforce harmful gender stereotypes by suggesting that drinking is more of a problem for women than for men, and may encourage victim blaming in relation to sexual assaults after drinking.[32] It is notable that the media's disproportionately frequent association of women with single-episodic drinking in our sample is at odds with young adults' perceptions of these types of drinking behaviours as being masculine.[18, 19] One potential explanation for this is that news producers regard women's 'binge' drinking as being of greater interest than men's because it is a deviation from gender norms. Whether disproportionate media focus on women's single-episodic drinking might influence a shift in public perceptions of how such behaviours correspond to gender identities may be a question for further research.

Our findings build on the growing body of literature about gendered media representations of alcohol by focusing specifically on portrayals of binge drinking. Depictions of women as unable to control themselves physically and emotionally echoed the findings of Atkinson and colleagues,[15] while the burden this weakness was depicted as placing on men could be seen

as corroborating Day and colleagues'[11] finding that traditionally male-dominated drinking environments and activities are perceived to be threatened by women's increasing involvement. Day and colleagues'[11] finding that women were presented as making themselves vulnerable to male aggression was replicated in our finding that articles presented women as in danger, and men as dangerous. The depictions of women's dress and behaviours found in our analysis echoed Nicholls'[14] findings. We found that whole or partial nudity in males was typically presented as frivolous or ridiculous, while women's partial nudity was presented with an underlying morally-loaded tone, perhaps resulting from women being perceived as publically breaking social conventions.

Our analysis supports Herring and Berridge's[16] observation that 'binge' drinking is popularly conceived in terms of specific social contexts, rather than being defined purely in terms of exceeding a certain quantity of alcohol in a limited time. Media framing of 'binge' drinking as an activity of young men and women, often in public spaces, disproportionately emphasises the context of single-episodic drinking and contributes to constructing binge drinking as something many people (including older people or those who drink in private settings) might not associate with. The disproportionate association of 'binge' drinking with young peoples' public, often anti-social (although performed within in social groups), drinking could have a damaging influence if it leads those who engage in single-episodic drinking outside of these contexts not to identify their behaviours as harmful. The dominant narrative of 'binge' drinking may have become unhelpful as a health concept. As an alternative, it may be useful to draw a clear distinction in health communications between 'binge drinking', as an inconsistently-defined, context-specific and value-laden term, and single-episodic drinking, as a specific, widespread and context-independent practice with a range of health consequences documented by research evidence.

Our findings have relevance to the development of alcohol guidelines. While the new guidelines propose gender-neutral consumption limits, the Department of Health report acknowledges that men and women vary in their drinking behaviours and the long- and short-term health risks they face[3]. Media associations of men with alcohol-related mortality and women with alcohol-related morbidity is in line with epidemiological evidence[3]. However, differences in how media coverage problematises women’s and men’s ‘binge’ drinking could promote perceptions of men’s ‘binge’ drinking as less harmful than women’s, potentially exacerbating men’s harmful drinking behaviours and hindering public understandings and acceptance of the proposed guidelines.

Our conclusions are subject to limitations. While mass media’s influence on public perceptions is well-evidenced and well-understood[6], content analysis alone cannot determine the extent to which audiences’ understandings correlate with media representations; audiences are subject to many influences beyond print and online news media, and do not consume media content in a passive, non-critical way. Studying other forms of media, such as television or social media, could help create a more complete understanding of media representations of ‘binge’ drinking. Furthermore, a larger sample size, achieved by choosing a longer timeframe or including more publications, could have increased the validity of our conclusions. A potential limitation of the data used is that the Nexis database stores articles in text-only format, without the images used in the original articles. Coders noted that text-only articles that appear factual and impartial could be perceived as more morally-loaded with the original photographs included. Similarly, some articles may only reveal implicit gender bias when viewed with the original photographs included; this is illustrated in an article about ‘binge’ drinking on The Sun’s website that contains seven images of women drinking and none of men, while the text makes no mention of gender.[33] Finally, as the article search period concluded two years prior to the proposal

of new alcohol guidelines, our analysis cannot tell us how representations changed immediately preceding, during or following the proposal. Further research might analyse how media coverage of the guidelines represented 'binge' drinking and gender roles.

Alternatively, comparative research could investigate whether media representations differed before and after the proposal. With these limitations taken into account, this comprehensive content analysis of a two-year period of coverage contributes to understandings of gendered media representations of drinking, with a specific focus on binge drinking.

In conclusion, our analysis suggests that popular representations of binge drinking may be harmful in three ways. Firstly, if public audiences' understandings are influenced by media coverage of 'binge' drinking, associations of 'binge' drinking with certain demographics and situations could lead the public to underestimate the health risks of single-episodic drinking among those not typically depicted in 'binge' drinking narratives, particularly older people. Morally-loaded representations of women's binge drinking may reproduce harmful stereotypes and stigma about the vulnerability of drunken women and the social unacceptability of female drunkenness. Finally, media content reinforcing a skewed representation of binge drinking may present a challenge to public health stakeholders seeking to promote evidence-based information, recommendations and policies with the goal of reducing alcohol-related health harms.

The misrepresentations identified by this research suggest that there may be a need for health advocates to engage with mass media to promote clear, evidence-informed messages about single-episodic drinking to help better align reporting on harmful drinking with evidence, with the ultimate goal of improving public understandings of harmful drinking behaviours.

Media communications strategies may seek to avoid stereotypes of 'binge drinking' that implicitly define binge drinking in terms of social contexts and behaviours not directly related to alcohol consumption, instead offering clear, value-free definitions of single-

episodic drinking based on specific quantities of alcohol and specific, time-bound episodes. Identifying the need to improve communication of single-episodic drinking is particularly relevant given the current public debate around new alcohol guidelines,[3] which may or may not specify recommended limits to single-episode alcohol consumption, depending on the outcome of the current consultation.

Contributorship statement

CE and SH conceived of the study and initiated the study design. OM and GF performed quantitative coding. CP and GF performed qualitative coding. OM prepared the initial draft of the paper. All authors contributed substantially to redrafting. The final manuscript was prepared by CP and approved by CE, GF, OM and SH.

Competing interests

The authors declare no competing interests.

Funding

This work was supported by the UK Medical Research Council as part of the Understandings and Uses of Public Health Research programme [MC_U130085862, MC_UU_12017/6] and the Informing Healthy Public Policy programme [MC_UU_12017/15] and by the Chief Scientist Office of the Scottish Government Health Directorates [SPHSU15] at the MRC/CSO Social and Public Health Sciences Unit, University of Glasgow. The funding body had no role in the design, collection, analysis or interpretation of this study.

Data sharing statement

No additional unpublished data from the study have been made available.

REFERENCES

1. Smith L, Foxcroft DR. Drinking in the UK: An exploration of trends: Joseph Rowntree Foundation, 2009.
2. Office for National Statistics. Adult drinking habits in Great Britain, 2013: Office for National Statistics, 2015.
3. Department of Health. Alcohol Guidelines Review: report from the guidelines development group to the UK Chief Medical Officers: Department of Health, 2015.
4. Department of Health. Sensible Drinking: The Report of an Inter-Departmental Working Group, 1995.
5. Furtwaengler NA, De Visser RO. Lack of international consensus in low-risk drinking guidelines. *Drug Alcohol Rev* 2013;**32**(1):11-18
6. McCombs ME, Shaw DL. The agenda-setting function of mass media. *Public Opin Q* 1972;**36**(2):176-87
7. Scheufele DA, Tewksbury D. Framing, agenda setting, and priming: The evolution of three media effects models. *J Commun* 2007;**57**(1):9-20
8. Wilsnack RW, Wilsnack SC. *Gender and alcohol: Individual and social perspectives*: Rutgers Center of Alcohol Studies, 1997.
9. Casswell S. Public Discourse on Alcohol. *Health Promot Int* 1997;**12**(3):251-57
10. Dorfman L. Studying the news on public health: How content analysis supports media advocacy. *Am J Health Behav* 2003;**27**(Supplement 3):S217-S26
11. Day K, Gough B, McFadden M. "Warning! alcohol can seriously damage your feminine health" a discourse analysis of recent British newspaper coverage of women and drinking. *Fem Media Stud* 2004;**4**(2):165-83
12. Wood K, Patterson C, Katikireddi SV, Hilton S. Harms to 'others' from alcohol consumption in the minimum unit pricing policy debate: a qualitative content analysis of UK newspapers (2005–12). *Addict* 2014;**109**(4):578-84
13. Lyons AC, Dalton SI, Hoy A. 'Hardcore Drinking': Portrayals of Alcohol Consumption in Young Women's and Men's Magazines. *J Health Psychol* 2006;**11**(2):223-32
14. Nicholls J. UK news reporting of alcohol: An analysis of television and newspaper coverage. *Drugs (Abingdon Engl)* 2011;**18**(3):200-06
15. Atkinson A, Kirton A, Sumnall H. The gendering of alcohol in consumer magazines: An analysis of male and female targeted publications. *J Gend Stud* 2012;**21**(4):365-86
16. Herring R, Berridge V, Thom B. Binge drinking: an exploration of a confused concept. *J Epidemiol Community Health* 2008;**62**(6):476-79
17. Guise JMF, Gill JS. 'Binge drinking? It's good, it's harmless fun': a discourse analysis of accounts of female undergraduate drinking in Scotland. *Health Educ Res* 2007;**22**(6):895-906
18. De Visser RO, McDonnell EJ. 'That's OK. He's a guy': A mixed-methods study of gender double-standards for alcohol use. *Psychol Health* 2012;**27**(5):618-39
19. Willott S, Lyons AC. Consuming male identities: Masculinities, gender relations and alcohol consumption in Aotearoa New Zealand. *J Community Appl Soc Psychol* 2012;**22**(4):330-45
20. Hutton F, Griffin C, Lyons A, Niland P, McCreanor T. "Tragic girls" and "crack whores": Alcohol, femininity and Facebook. *Fem Psychol* 2016;**26**(1):73-93
21. National Readership Survey. National Readership Survey. Secondary National Readership Survey 2012. <http://www.nrs.co.uk/> (accessed 19th September 2012)
22. Newsworks. Newsworks. Secondary Newsworks 2012. www.nmauk.co.uk (accessed 19th September 2012)

23. Ofcom. Communications Market Report 2015. London: Ofcom, 2015.

24. Hilton S, Patterson C, Teyhan A. Escalating coverage of obesity in UK newspapers: the evolution and framing of the “obesity epidemic” from 1996 to 2010. *Obes* 2012;**20**(8):1688-95

25. Patterson C, Katikireddi SV, Wood K, Hilton S. Representations of minimum unit pricing for alcohol in UK newspapers: a case study of a public health policy debate. *J Public Health* 2015;**37**(1):40-49

26. HM Government Home Department. The Government's Alcohol Strategy, 2012.

27. Altheide DL, Schneider CJ. *Qualitative media analysis*: Sage, 2012.

28. Neuendorf KA, Kumar A. Content analysis. *The International Encyclopedia of Political Communication* 2006

29. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;**3**(2):77-101

30. Barry AE, Piazza-Gardner AK. Drunkorexia: Understanding the Co-occurrence of Alcohol Consumption and Eating/Exercise Weight Management Behaviors. *J Am Coll Health* 2012;**60**(3):236-43 doi: 10.1080/07448481.2011.587487published Online First: Epub Date]].

31. Measham F, Østergaard J. The public face of binge drinking: British and Danish young women, recent trends in alcohol consumption and the European binge drinking debate. *Probat J* 2009;**56**(4):415-34


32. Fitzgerald N, Angus K, Emslie C, Shipton D, Bauld L. Gender differences in the impact of population-level alcohol policy interventions: evidence synthesis of systematic reviews. *Addict* in press

33. The Sun. Your friends, bad habits or anxiety... take our quiz to find YOUR booze binge trigger. Secondary Your friends, bad habits or anxiety... take our quiz to find YOUR booze binge trigger 2016.
<http://www.thesun.co.uk/sol/homepage/fabulous/7112837/Your-friends-bad-habits-or-anxiety-take-our-quiz-to-find-YOUR-booze-binge-trigger.html>.(accessed 13 May 2016)

Figure legends

Figure 1: Frequency of publication of articles about ‘binge’ drinking by quarter

For peer review only



Frequency of publication of articles about 'binge' drinking by quarter
[Figure 1]
146x100mm (300 x 300 DPI)

BMJ Open

A content analysis of UK newspaper and online news representations of women's and men's 'binge' drinking: a challenge for communicating evidence-based messages about single-episodic drinking?

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-013124.R2
Article Type:	Research
Date Submitted by the Author:	09-Nov-2016
Complete List of Authors:	Patterson, Chris; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit Emslie, Carol; Glasgow Caledonian University Institute for Applied Health Research Mason, Olivia; Durham University, Department of Geography Fergie, Gillian; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit Hilton, Shona; University of Glasgow, MRC/CSO Social & Public Health Sciences Unit
Primary Subject Heading:	Communication
Secondary Subject Heading:	Sociology, Public health
Keywords:	PUBLIC HEALTH, QUALITATIVE RESEARCH, SOCIAL MEDICINE

SCHOLARONE™
Manuscripts

A content analysis of UK newspaper and online news representations of women’s and men’s ‘binge’ drinking: a challenge for communicating evidence-based messages about single-episodic drinking?

Patterson^{1*}, Emslie², Mason³, Fergie¹, Hilton¹

¹MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB UK

²Institute for Applied Health Research/School of Health and Life Sciences, Glasgow Caledonian University, Cowcaddens Road, Glasgow G4 0BA UK

²Department of Geography, Durham University, Lower Mountjoy South Road, Durham DH1 3LE UK

Corresponding author:

Chris Patterson

MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB UK

chris.patterson@glasgow.ac.uk

0141 353 7631

Keywords: gender; media; alcohol; binge drinking

Running head: Media representations of ‘binge’ drinking

ABSTRACT

Objectives: In the UK, men's alcohol-related morbidity and mortality still greatly exceeds women's, despite an increase in women's alcohol consumption in recent decades. New UK alcohol guidelines introduce gender-neutral low-risk alcohol consumption guidance. This study explores how UK newspaper and online news represent women's and men's 'binge' drinking to identify opportunities to better align reporting of harmful drinking with evidence.

Design: Quantitative and qualitative content analysis of 308 articles published in seven UK national newspapers and the BBC News website between 1 January 2012 and 31 December 2013.

Results: Articles associated women with 'binge' drinking more frequently than men, and presented women's drinking as more problematic. Men were more frequently characterised as violent or disorderly, while women were characterised as out of control, putting themselves in danger, harming their physical appearance and burdening men. Descriptions of female 'binge' drinkers' clothing and appearance were typically moralistic.

Conclusions: The UK news media's disproportionate focus on women's 'binge' drinking is at odds with epidemiological evidence, may reproduce harmful gender stereotypes and may obstruct public understandings of the gender-neutral weekly consumption limits in newly-proposed alcohol guidelines. In order to better align reporting of harmful drinking with current evidence, public health advocates may engage with the media with a view to shifting media framing of 'binge' drinking away from specific groups (young people; females) and contexts (public drinking) and towards the health risks of specific drinking behaviours, which affect all groups regardless of context.

Strengths and limitations of this study

- Quantitative and qualitative content analysis of a large, comprehensive, two-year sample of UK national newspaper and online news about ‘binge’ drinking.
- The findings illustrate how media portrayals of ‘binge’ drinking could be harmful, and identify opportunities for these portrayals to be better aligned with evidence.
- Content analysis facilitates understandings of the messages being presented to the public, but cannot determine the extent to which audiences’ understandings are influenced by media representations of specific issues.
- Some findings may have differed if the data analysed had gone beyond text articles, for example by incorporating images, video and social media content.

INTRODUCTION

While the gap between women's and men's excessive and harmful consumption of alcohol in the UK has narrowed in recent years,[1] men still drink more than women[2], experience more drink-related health and social problems and face twice women's alcohol-related mortality (15.9 to 7.8 deaths per 100,000 population respectively in 2012[2]). National guidelines have typically issued different alcohol consumption guidance for men and women, with women being advised to drink less than men, but the UK have recently joined Australia and Portugal[3, 4] in issuing the same low-risk consumption guidance for both men and women, drawing on evidence that the health risks posed to each gender are similar at low-risk levels of consumption.[5] In light of this shift towards gender-neutrality in alcohol consumption guidelines, it is timely to consider how gender differences in drinking behaviours are represented and perceived.

As with most health issues, popular perceptions of drinking are likely to be influenced by mass media representations[6-8], and, in turn, differences in representations of women's and men's drinking are heavily influenced by how societies regulate gender roles.[9] Media content analysis cannot tell us if, and how, specific content influences audiences' understandings and behaviours, and content analysis findings must be considered with that inherent limitation in mind. Nonetheless, exploring media representations of men's and women's drinking behaviours allows us to examine how shared cultural values around alcohol are articulated and constructed[10], which might inform efforts to improve media representations, and therefore public understandings, of harmful drinking behaviours.[11]

Research illustrates a clear gender divide in media portrayals of drinking behaviours, with men's drinking normalised and women's problematised. Day and colleagues[12] found that UK newspapers frequently characterised women as departing from idealised notions of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

femininity in terms of appearance (e.g. weight gain, deeper voices, loss of good looks) and motherhood (e.g. reduced fertility and unborn children). They found that men were framed as violent, but partial responsibility for men’s violence was attributed to women, who were framed as sexual predators invading traditionally male-dominated drinking environments. Conversely, Wood and colleagues[13] found that UK newspaper coverage of proposed minimum unit pricing (MUP) policies presented women as being at ‘risk of harm’ from male aggression. Lyons, Dalton & Hoy[14] found that both young women’s and young men’s magazines framed binge drinking as normative, adult and professional, with young men’s magazines associating men’s drinking with traditional masculine images and deriding young women’s drinking behaviours. Nicholls[15] found that UK television and newspaper news associated men’s drinking with violence (both as perpetrators and victims) and women’s drunkenness with unfeminine and undignified behaviours, such as loss of consciousness or partial nudity. Similarly, Atkinson and colleagues[16] found that UK magazines targeted at teenagers depicted women’s drinking as more problematic than men’s, and portrayed women as ‘behaving like men’ in male spaces, sexualised and highly emotional.

As Nicholls[15] found, ‘binge drinking’, also called single-episodic drinking, is a key focus of media representations of harmful drinking; *“problem drinking is less commonly associated with dependence and more commonly associated with binge, harmful and hazardous drinking”*. Herring and Berridge[17] describe binge drinking as a ‘confused concept’ that has historically been, and continues to be, defined inconsistently. They suggest that ‘binge’ drinking is currently portrayed as a youth issue, despite evidence that single-episodic drinking is performed by various age groups[2], highlighting a potential area of media misrepresentation of population drinking behaviours. Qualitative evidence indicates that young women in Scotland define ‘binge’ drinking in terms of types of behaviour rather than the quantity of alcohol consumed,[18] which may present an obstacle to clearly and

objectively defining 'binge' drinking. Mixed-method evidence suggests that both male and female students in England perceive binge drinking and public drunkenness as masculine behaviours[19]. Two studies based on qualitative data from New Zealand have highlighted the role of alcohol-related behaviours in young people's gender identities. Firstly, Willott and Lyons[20] found that young men and women perceived consuming large quantities of alcohol in a single episode (among other behaviours) as a key performance of masculinity, and highlighted the identity negotiations undertaken by men who do not engage in these normative, masculine drinking behaviours. Secondly, Hutton and colleagues[21] examined the challenges that young women experience in curating social media personae that balance engagement in "the culture of intoxication" (p.88) with maintaining respectability. Given the primacy of binge drinking in UK media coverage of harmful alcohol use,[15] binge drinking is an important lens through which to examine gendered representations of drinking behaviours.

This study comprises a comprehensive, mixed-methods content analysis of two years of UK media coverage of binge drinking, designed to contribute new insights to a growing body of literature about gendered media representations of alcohol, with a particular focus on binge drinking. Our aim is to improve understandings of UK newspaper and online news representations of women's and men's 'binge' drinking, focusing not on how 'binge' drinking is defined, but rather on how different types of drinking behaviour are gendered in media content wherein those behaviours are labelled as 'binge' drinking. While the limitations of media content analysis must be taken into account, the improved understandings produced by this research may help to inform efforts to better align media representations of harmful drinking with current evidence, which could in turn improve public understandings of the risks of single-episodic drinking.

METHODS

We selected seven highly-circulated[22, 23] UK national newspapers, including their Sunday counterparts, and the most-read exclusively-online news website[24] (Table 1). The chosen newspaper publications represented three genres: quality, middle-market tabloid and tabloid. This typology helps to ensure a sample that represents diverse readerships in terms of age, social class and political alignment.[25] Quality newspapers are those that were traditionally printed in broadsheet format, have predominantly middle-class audiences, are politically diverse and are serious in tone. Middle-market tabloids are printed in tabloid format and are less serious in tone than quality-genre newspapers, and have a predominantly older, middle-class, right-wing audience. Tabloids are less serious and typically more sensationalist than middle-market tabloids, are politically diverse and have predominantly working-class audiences.

We selected a search period of 1 January 2012 to 31 December 2013, during which alcohol was subject to heightened media interest due to debates in the UK and Scottish parliaments around MUP.[26] Much of the media coverage of MUP was related to ‘binge’ drinking due to the UK Government’s Alcohol Strategy[27] overtly positioning ‘binge’ drinking as the primary target of MUP. Relevant newspaper articles, were identified using the Nexis database including articles from online editions of the two of the selected publications (The Guardian and The Daily Mail) that are archived in Nexis. BBC News articles were identified using the search function on the BBC website. The search string identified articles that contain both three or more mentions of “binge” and one or more mentions of “drink OR drinker OR drinkers OR drinking”. Duplicate articles were removed, and articles were manually excluded if they did not predominantly focus on binge drinking in the UK, or if they were not in the news, feature or editorial formats. Initial searches identified 537 articles, of which 308 met the inclusion criteria and were eligible for detailed coding and analysis.

Table 1. Summary of publications and articles in the sample

Genre / medium	Publication	All articles		Article format					
		n	%	Standard		Feature		Editorial	
		n	%	n	%	n	%	n	%
Quality (n=86)	Guardian / Observer	58	18.8	48	20.3	9	17.3	1	5.3
	Independent / Independent on Sunday	17	5.5	11	4.6	5	9.6	1	5.3
	Daily Telegraph / Sunday Telegraph	52	16.9	44	18.6	5	9.6	3	15.8
Middle-market tabloids (n=39)	Daily Mail / Mail on Sunday	54	17.5	33	13.9	18	34.6	3	15.8
	Express / Sunday Express	13	4.2	8	3.4	3	5.8	2	10.5
Tabloids (n=75)	Daily Mirror / Sunday Mirror	13	4.2	10	4.2	2	3.8	1	5.3
	The Sun / News of the World	62	20.1	46	19.4	9	17.3	7	36.8
Online (n=39)	BBC News website	39	12.7	37	15.6	1	1.9	1	5.3
Total:		308	100	237	100	52	100	19	100

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

To systematically and comprehensively code and analyse the media content, we used a mixed-methods content analysis. Firstly, we used quantitative content analysis to measure the frequency of content within the articles across the whole sample, and, secondly we performed qualitative content analysis of the content of a subsample of articles for more in-depth, reflexive analysis.[28] From this perspective, quantitative content analysis is concerned with measuring and analysing manifest content (the surface-level content of the articles, the coding of which does not require interpretation on the part of the coder, such that it can be recorded relatively objectively), while qualitative content analysis is concerned with latent content (the underlying meanings of the text, as interpreted by coders in an inherently subjective process).[29] The mixed-methods approach in this research comprised the following steps: constructing a coding frame; coding manifest content using the coding frame; establishing the reliability of the data collected and excluding unreliable data; analysing the quantitative the data; identifying aspects of manifest content to examine further using qualitative analysis; and finally performing thematic analysis of the latent content of articles containing the manifest content of interest.

To construct a coding frame with which to code manifest content, SH and CE read randomly-selected articles from the sample in batches of 20, recording potential thematic categories relevant to the research topic as they emerged. At the point where a batch of articles was read without any new categories emerging, the list of categories was deemed to have reached saturation. The collected thematic categories were grouped into three broad thematic categories: how is binge drinking described and which sections of the population are associated with this behaviour?; how are the drivers of the binge drinking described and who is to blame?; and what are the consequences of binge drinking? These categories were chosen by a combination of *a priori* knowledge of the research topic and understandings of the content of the sample that emerged from close reading of the articles.[26] The final coding

frame comprised the three broad thematic categories and their sub-categories, as well as fields to record more routine details of articles, such as publication, article format and word count. In addition, coders recorded whether each article mentioned males and females, allowing articles to be divided into four gender categories: those that mention females exclusively, those that mention males exclusively, those that mention both males and females and those that mention neither males nor females explicitly. These codes enabled analysis of whether themes varied by the gender focus of articles.

To collect quantitative data, OM read each article in turn, using the coding frame to record whether each theme was present within its manifest content. To ensure consistency of coding, GF coded a random subsample of 39 articles (12.7%). Linearly-weighted kappa tests of inter-rater agreement between OM and GF were then performed on each variable across those 39 double-coded articles, and variables that returned a coefficient below 0.8 were discarded from the study to ensure that only variables with strong agreement were retained for analysis. The coding frame data were then entered into SPSS for analysis. Statistical procedures comprised: a simple linear regression examining the relationship between publication quarter and the count of articles published; chi-square tests of whether thematic variables varied by gender-focus, publication genre or article format; and paired t-tests comparing the means of thematic variables. The threshold of statistical significance is set at 0.01 throughout to mitigate the risk of type-1 errors.

Following quantitative analysis, the thematic category of 'harms to appearance' was deemed noteworthy and suitable for qualitative coding and analysis. The decision to focus on that theme was informed by stark gender differences in content that emerged from the quantitative analysis, and an expectation that valuable understandings could be gained from deeper analysis. To analyse the latent content of the 46 articles that had been coded as mentioning 'harms to appearance', GF and CP employed a thematic analysis approach[30] using NVivo

10, closely reading each article to generate initial codes, which were then collated into potential themes. GF and CP collaborated closely to assure the themes were defined clearly and that they worked across the 46 articles. The findings from the thematic analysis are presented in terms of two broad themes, using typical quotations to illustrate article content.

RESULTS

Quantitative findings

Between 1 January 2012 and 31 December 2013, 308 articles about binge drinking were published in the seven print publications and one website included in the sample. Of these articles, 86 (27.9%) were published in ‘quality’ newspapers, 39 (12.7%) in middle-market tabloids, 75 (24.4%) in tabloids and 108 (35.1%) on BBC News (Table 1). The majority of articles were standard news format (n=237, 76.9%), while 52 (16.9%) were feature articles and 19 (6.2%) editorials. The frequency of articles published per quarter decreased across the two-year period (Figure 1), but publication quarter was not a statistically significant predictor of article frequency (coefficient -6.714, p=0.088). There was a peak of 56 articles in March 2012, 66.1% of which were related to MUP, in particular the UK Prime Minister’s announcement of plans to introduce MUP. There was elevated reporting from November 2012 to January 2013 (n=62), during which 53.2% of articles mentioned MUP, largely related to opposition to MUP within the UK Cabinet. In total, 133 (43.2%) articles mentioned MUP.

[Figure 1]

Articles were coded according to whether they mentioned binge drinking in females exclusively (n=68, 22.1%), males exclusively (n=30, 9.7%), both males and females (n=43,

14.0%), or neither males nor females explicitly (n=167, 54.2%). To be coded as mentioning both males and females, an article had to include specific discussion of each gender individually. There were not significant associations between gender category and either genre (p=0.382) or article format (p=0.303). Furthermore, neither publication genre nor article format were significantly associated with differences in reporting on any of the thematic categories listed in Table 2.

Articles were coded to identify the characteristics of binge drinkers and the specific drinks associated with binge drinking. Thirty-two (10.4%) of all 308 articles associated mothers (including pregnant women) with binge drinking, which differed significantly (p<0.000) from the 11 (3.6%) that mentioned fathers (Table 2). There were no statistically significant relationships between gender focus and mentioning either younger or older people. While more articles that mentioned binge drinking in the home were focused exclusively on women, no drinking location was significantly associated with gender category (Table 2). Wine (n=58, 18.8%) was the only type of drink significantly associated more with one gender than the other, being predominantly mentioned in articles focusing exclusively on females (p=0.004).

Table 2. Social groups, locations and harms associated with binge drinking

	All articles (n=308)		Only females (n=68)		Only males (n=30)		Both (n=43)		Neither (n=167)		χ ² P-value (all)	χ ² P-value (Female vs. Male)
	n	%	n	%	n	%	n	%	n	%		
Who is binge drinking?												
Mothers (inc. pregnant women)	32	10.4	23	7.5	0	0.0	9.0	2.9	0.0	0.0	0.000*	0.000*
Fathers	11	3.6	0	0.0	0	0.0	11.0	3.6	0.0	0.0	0.000*	-
Young people (inc. children and students)	119	38.6	24.0	35.3	8.0	26.7	18.0	41.9	69.0	41.3	0.419	0.401
Older people	17	5.5	5	1.6	0	0.0	3.0	1.0	9.0	2.9	0.499	0.127
Which locations of binge drinking are mentioned?												
At home	46	14.9	14	4.5	1	0.3	10.0	3.2	21.0	6.8	0.047	0.029
On city streets	60	19.5	12	3.9	5	1.6	10.0	3.2	33.0	10.7	0.087	0.906
Pubs, bars and clubs	56	18.2	12	3.9	5	1.6	6.0	1.9	33.0	10.7	0.836	0.906
What types of drink are associated with binge drinking?												
Wine	58	18.8	23	7.5	2	0.6	11.0	3.6	22.0	7.1	0.001*	0.004*
Spirits	58	18.8	22	7.1	12	3.9	4.0	1.3	20.0	6.5	0.000*	0.464
Beer	51	16.6	10	3.2	8	2.6	10.0	3.2	23.0	7.5	0.193	0.159
Cider	26	8.4	4	1.3	5	1.6	4.0	1.3	13.0	4.2	0.343	0.880
Alcopops	9	2.9	5	1.6	0	0.0	1.0	0.3	3.0	1.0	0.093	0.127
Shots	5	1.6	3	1.0	1	0.3	0.0	0.0	1.0	0.3	0.129	0.804
Cocktails	4	1.3	2	0.6	1	0.3	0.0	0.0	1.0	0.3	0.308	0.917
Fortified wine	4	1.3	1	0.3	0	0.0	2.0	0.6	1.0	0.3	0.186	0.504
Sparkling wine	3	1.0	1	0.3	0	0.0	1.0	0.3	1.0	0.3	0.676	0.504
Liqueurs	2	0.6	1	0.3	0	0.0	0.0	0.0	1.0	0.3	0.754	0.504
Lambrini	2	0.6	2	0.6	0	0.0	0.0	0.0	0.0	0.0	0.069	0.343
What harms of binge drinking are mentioned?												
Any personal harms	38	12.3	11	3.6	6	1.9	8	2.6	13	4.2	0.062	0.645
Harms to relationships	21	6.8	5	1.6	6	1.9	2	0.6	8	2.6	0.022	0.068
Harms to parenting	24	7.8	7	2.3	0	0.0	7	2.3	10	3.2	0.042	0.068
Any economic harms	107	34.7	16	5.2	8	2.6	17	5.5	66	21.4	0.080	0.739
Harms to the NHS	103	33.4	16	5.2	8	2.6	16	5.2	63	20.5	0.153	0.739
Harms to economic productivity	11	3.6	2	0.6	0	0.0	1	0.3	8	2.6	0.551	0.343
Any social harms	138	44.8	25	8.1	15	4.9	17	5.5	81	26.3	0.319	0.219
Social disorder and violence	128	41.6	19	6.2	14	4.5	16	5.2	79	25.6	0.044	0.071
Sexual assault and rape	19	6.2	9	2.9	4	1.3	2	0.6	4	1.3	0.005*	0.989
Fear	15	4.9	0	0.0	0	0.0	2	0.6	13	4.2	0.044	-
Any health harms	192	62.3	40	13.0	21	6.8	26	8.4	105	34.1	0.755	0.293
Death	77	25.0	8	2.6	16	5.2	10	3.2	43	14.0	0.000*	0.000*
Physical health	138	44.8	32	10.4	11	3.6	14	4.5	81	26.3	0.215	0.339
Harms to appearance	46	14.9	19	6.2	0	0.0	8	2.6	19	6.2	0.001*	0.001*

*Statistically significant at the 0.01 level

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Almost two-thirds of articles (n=192, 62.3%) described health harms related to binge drinking, while one-quarter (n=77) mentioned risk of death. Mentions of specific categories of harm were recorded (Table 2), and relationships between gender and specific harms were identified. Mentioning harms to appearance was significantly related to gender category (p=0.001); 19 of the 46 articles that mentioned harms to appearance mentioned females exclusively, while none of those 46 articles mentioned males exclusively. Harms to parenting were mentioned in seven (10.3%) of the 68 articles that exclusively mentioned females, and none that exclusively mentioned males, but the difference was not significant (p=0.068). Sexual assault and rape were mentioned more frequently in articles exclusively mentioning females (n=9, 2.9%) than those exclusively mentioning males (n=4, 13.3%), but not significantly so (p=0.989). More articles directly associated the risk of death from drinking with males (n=16, 20.8%) than with females (n=8, 10.4%), and this difference was significant (p<0.000). Conversely, non-fatal physical health harms were associated with females (n=32, 23.2%) more frequently than males (n=11, 8.0%), albeit not significantly so (p=0.339).

Qualitative findings

As identified in the quantitative findings, just eight of the 46 articles that reported on 'harms to appearance' mentioned men, and none did so exclusively. These 46 articles were subjected to thematic analysis. A broad conceptualisation of 'harms to appearance' was used that included behavioural as well as aesthetic aspects of physical appearance that may be influenced (or perceived to be influenced) by single-episodic drinking. Using this broad definition, two key themes emerged from thematic analysis of the subsample: physical appearance and loss of self-control.

Physical Appearance

Various harms to complexion were reported in relation to women, including the use of make-up to hide grey skin. Articles described specific damage to women's features, including hair, lips, noses and teeth, as well as using less-specific descriptions of physical damage such as 'ravaged' (The Sun, 13 Feb 2012). An article headlined *'Lambrini ruined my looks in 5 months'* (The Sun, 27 Sept 2013) associated an 'attractive' woman becoming 'a swollen, haggard wreck' with the consumption of a perry overtly marketed to young women. Weight gain caused by alcohol consumption was cited as both a product of alcohol use and a cause of 'drunkorexia', the practice of eating less to offset calories gained from alcoholic drinks.[31] Drunkorexia was typically associated with women; one article stated that it 'affects mainly young women' (Daily Mail, 19 Jul 2012) while another defined it as a condition in which 'calorie-conscious women skip meals in order to binge drink' (Mirror, 20 Dec 2012).

Within the few articles in the subsample that mentioned men, male partial nudity was related to drunken behaviours, such as lost and ripped clothes, and intentional indecent exposure. Typifying this, the Daily Mail described a male student *'flashing his penis in the street and laughing at his own loutish behaviour'* (19 June 2012). Conversely, partial nudity in women was more typically related to clothing choices presumably made while sober, with descriptions of revealing clothing, particularly miniskirts, and impractical shoes. One article described a fancy dress event in which *'undergraduate girls take to the streets dressed in little more than their underwear'* (Daily Mail, 8 Oct 2012). Women were variously described as 'scantily-clad' (Daily Mail, 22 Aug 2012), 'half-naked' (The Sun, 3 Mar 2013), and 'nearly bearing all' (Daily Mail, 8 Oct 2012). Aside from choices of attire made in sobriety, harms to appearance that could befall women during drinking episodes included smeared makeup, vomit in hair, stained clothes and gravel embedded in knees. Articles also made mention of unintentional exposure of underwear or body parts by women, which was in contrast to males, who were typically described as exposed their genitalia deliberately.

1
2
3 Loss of self-control
4

5
6 Descriptions of women's and men's loss of physical control differed. Some articles depicted
7
8 drunken women as burdening male partners. For example, two articles described women
9
10 vomiting on their boyfriends, while one described female students needing *'to be carried*
11
12 *back to their rooms by boyfriends'* (Daily Mail, 19 Jun 2012). No articles characterised
13
14 'binge' drinking men as relying on, or burdening, their partners.
15
16

17
18 Women's aggressive behaviours were limited to verbal conflict, with drunken women
19
20 characterised as noisy, argumentative, emotional and hard to control. An article describing
21
22 Cardiff nightlife reported that drunken women *'scream at their boyfriends in shop doorways'*
23
24 and *'sob in public'* (Daily Mail, 21 Aug 2012). Men were more typically associated with
25
26 aggressive behaviours, including violent language and actions, damaging property and non-
27
28 specific 'loutish' (Daily Mail, 19 Jun 2012) behaviour. While male aggression was presented
29
30 as endangering others, women were characterised as putting themselves at risk, sometimes of
31
32 undesirable sexual situations and assault. One article reproduced Office of National Statistics
33
34 data about fines for drunk and disorderly behaviour, presenting men as much more likely than
35
36 women to receive such fines, but suggesting that the number of women receiving the fines is
37
38 increasing more rapidly than men (Telegraph, 14 Jun 2009).
39
40
41
42

43 DISCUSSION

44
45
46 We examined UK newspaper and online news representations of 'binge' drinking, exploring
47
48 differences and similarities in representations of women's and men's drinking behaviours.

49
50 Our quantitative analysis identified that women were associated with binge drinking more
51
52 frequently than men, which conflicts with epidemiological evidence[2] and young people's
53
54 perceptions of 'binge' drinking as a masculine activity.[19] We found various differences in
55
56 how binge drinking was represented in association with different genders. Motherhood was
57
58
59
60

mentioned in relation to ‘binge’ drinking more frequently than fatherhood. There were significant gender differences in the attribution of health harms, with male ‘binge’ drinking frequently associated with mortality and female ‘binge’ drinking more frequently associated with morbidity. Women’s drinking was typically presented as more problematic than men’s, and, notably, harms to appearance were associated with women’s ‘binge’ drinking much more frequently than men’s. The articles in our sample associated wine with women’s ‘binge’ drinking significantly more than men’s, but our data did not identify significant gender associations with other specific types of drink.

Our qualitative analysis focused on depictions of the relationship between ‘binge’ drinking and appearance, comprising aspects of both physical appearance and behaviour. Women engaged in ‘binge’ drinking were presented as helpless, physically incapacitated and transgressive, and as burdens to male partners, who were sometimes cast as carers for drunk women. ‘Binge’ drinking men were associated with physical and verbal aggression, but not behaviours that marked them as weak or vulnerable; while women were characterised as endangering themselves, men were more likely to endanger others. Articles typically depicted women as less able than men to maintain socially acceptable behaviour during single-episodic drinking. ‘Binge’ drinking was characterised as impacting women’s physical appearance more than men’s. Articles associated binge drinking with both immediate and long-term damage to physical characteristics and self-presentation. Descriptions of binge-drinking women’s dress had a moralistic tone that was absent in representations of men, with articles typically focusing on degrees of nudity and implicitly questioning the propriety of women’s chosen attire. Our analysis suggests that media representations of women’s ‘binge’ drinking do not focus solely on the effects of binge drinking on women, but also reflect broader social expectations about women’s public behaviours.

Measham & Ostergaard [32] suggest that newspapers have constructed ‘binge’ drinking as a problem of young women adopting male behaviours:

The public face of binge drinking, [...] a staple of early 21st century tabloid newspapers, became the young woman emulating male consumption patterns [...] with clothes askew stumbling around the city centre streets at night’ (p417).

There may be unintended consequences of this disproportionate focus on women’s – as opposed to men’s – single-episodic drinking in media reports. For example, it may reinforce harmful gender stereotypes by suggesting that drinking is more problematic for women than men, and may encourage victim blaming in relation to sexual assaults after drinking.[33] It is notable that the media’s disproportionately frequent association of women with single-episodic drinking in our sample is at odds with young adults’ perceptions of these types of drinking behaviours as being masculine.[19, 20] One potential explanation for this is that news producers regard women’s ‘binge’ drinking as being of greater interest than men’s because it is a deviation from gender norms. Whether disproportionate media focus on women’s single-episodic drinking might influence a shift in public perceptions of how such behaviours correspond to gender identities may be a question for further research.

Our findings build on the growing body of literature about gendered media representations of alcohol by focusing specifically on portrayals of binge drinking. Depictions of women as unable to control themselves physically and emotionally echoed the findings of Atkinson and colleagues,[16] while the burden this weakness was depicted as placing on men could be seen as corroborating Day and colleagues’[12] finding that traditionally male-dominated drinking environments and activities are perceived to be threatened by women’s increasing involvement. Day and colleagues’[12] finding that women were presented as making themselves vulnerable to male aggression was replicated in our finding that articles presented

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

women as in danger, and men as dangerous. The depictions of women’s dress and behaviours found in our analysis echoed Nicholls’[15] findings. We found that whole or partial nudity in males was typically presented as frivolous or ridiculous, while women’s partial nudity was presented with an underlying morally-loaded tone, perhaps resulting from women being perceived as publically breaking social conventions.

Our analysis supports Herring and Berridge’s[17] observation that ‘binge’ drinking is popularly conceived in terms of specific social contexts, rather than being defined purely in terms of exceeding a certain quantity of alcohol in a limited time. Media framing of ‘binge’ drinking as an activity of young men and women, often in public spaces, disproportionately emphasises the context of single-episodic drinking and contributes to constructing binge drinking as something many (including older people or those who drink in private settings) might not associate themselves with. The disproportionate association of ‘binge’ drinking with young peoples’ public, often anti-social (although performed within in social groups), drinking could have a damaging influence if it leads those who engage in single-episodic drinking outside of these contexts to misidentify their behaviours as harmless. The dominant narrative of ‘binge’ drinking may have become unhelpful as a health concept. As an alternative, it may be useful to draw a clear distinction in health communications between ‘binge drinking’, as an inconsistently-defined, context-specific and value-laden term, and single-episodic drinking, as a specific, widespread and context-independent practice with a range of health consequences documented by research evidence.

Our findings have relevance to the development of alcohol guidelines. While the new UK guidelines propose gender-neutral low-risk consumption guidelines, the Department of Health acknowledges that men and women vary in their drinking behaviours and the long- and short-term health risks they face[5]. Media associations of men with alcohol-related mortality and women with alcohol-related morbidity are in line with epidemiological

evidence[5]. However, differences in how media coverage problematises women's and men's 'binge' drinking could promote perceptions of men's 'binge' drinking as less harmful than women's, potentially exacerbating men's harmful drinking behaviours and hindering public understandings and acceptance of the proposed guidelines.

Our conclusions are subject to limitations. While mass media's influence on public perceptions of health issues is extensively researched and well established[6], content analysis alone cannot determine the extent to which audiences' understandings correlate with media representations; audiences are subject to many influences beyond print and online news media, and do not consume media content in a passive, non-critical way. Studying other forms of media, such as television or social media, could help create a more complete understanding of media representations of 'binge' drinking. Furthermore, a larger sample size, achieved by choosing a longer timeframe or including more publications, could have increased the validity of our conclusions. A potential limitation of the data used is that the Nexis database stores articles in text-only format, omitting any images included in the original articles. Coders noted that text-only articles that appear factual and impartial could be perceived as more morally-loaded with the original photographs included. Similarly, some articles may only reveal implicit gender bias when viewed with the original photographs included; this is illustrated in an article about 'binge' drinking on The Sun's website that contains seven images of women drinking and none of men, while the text makes no mention of gender.[34] Finally, as the article search period concluded two years prior to the proposal of new alcohol guidelines, our analysis cannot tell us how representations changed immediately preceding, during or following the proposal. Further research might analyse how media coverage of alcohol guidelines represents 'binge' drinking and gender roles.

Alternatively, comparative research could investigate whether media representations differed before and after the introduction of new guidelines. With these limitations taken into account,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

this comprehensive content analysis of a two-year period of coverage contributes to understandings of gendered media representations of drinking, with a specific focus on ‘binge’ drinking.

In conclusion, our analysis suggests that popular representations of binge drinking may be harmful in three ways. Firstly, if public audiences’ understandings are influenced by media coverage of ‘binge’ drinking, associations of ‘binge’ drinking with certain demographics and situations could lead the public to underestimate the health risks of single-episodic drinking among those not typically depicted in ‘binge’ drinking narratives, particularly older people. Morally-loaded representations of women’s binge drinking may reproduce harmful stereotypes and stigma about the vulnerability of drunken women and the social unacceptability of female drunkenness. Finally, media content reinforcing a skewed representation of binge drinking may present a challenge to public health stakeholders seeking to promote evidence-based information, recommendations and policies with the goal of reducing alcohol-related health harms.

The misrepresentations identified by this research suggest that there may be a need for health advocates to engage with mass media to promote clear, evidence-informed messages about single-episodic drinking to help better align reporting on harmful drinking with evidence, with the ultimate goal of improving public understandings of harmful drinking behaviours. Media communications strategies may seek to avoid stereotypes of ‘binge drinking’ that implicitly define binge drinking in terms of social contexts and behaviours not directly related to alcohol consumption, instead offering clear, value-free definitions of single-episodic drinking based on specific quantities of alcohol and specific, time-bound episodes. Identifying the need to improve communication of single-episodic drinking is particularly relevant given the recent public consultation about new alcohol guidelines,[5] and the UK

Government's decision not include specific guidance on single-occasion alcohol consumption within the final guidelines.

Contributorship statement

CE and SH conceived of the study and initiated the study design. OM and GF performed quantitative coding. CP and GF performed qualitative coding. OM prepared the initial draft of the paper. All authors contributed substantially to redrafting. The final manuscript was prepared by CP and approved by CE, GF, OM and SH.

Competing interests

The authors declare no competing interests.

Funding

This work was supported by the UK Medical Research Council as part of the Understandings and Uses of Public Health Research programme [MC_U130085862, MC_UU_12017/6] and the Informing Healthy Public Policy programme [MC_UU_12017/15] and by the Chief Scientist Office of the Scottish Government Health Directorates [SPHSU15] at the MRC/CSO Social and Public Health Sciences Unit, University of Glasgow. The funding body had no role in the design, collection, analysis or interpretation of this study.

Data sharing statement

No additional unpublished data from the study have been made available.

REFERENCES

1. Smith L, Foxcroft DR. Drinking in the UK: An exploration of trends: Joseph Rowntree Foundation, 2009.
2. Office for National Statistics. Adult drinking habits in Great Britain, 2013: Office for National Statistics, 2015.
3. Furtwaengler NA, De Visser RO. Lack of international consensus in low-risk drinking guidelines. *Drug Alcohol Rev* 2013;**32**(1):11-18
4. Department of Health. Sensible Drinking: The Report of an Inter-Departmental Working Group, 1995.
5. Department of Health. Alcohol Guidelines Review: report from the guidelines development group to the UK Chief Medical Officers: Department of Health, 2015.
6. McCombs ME, Shaw DL. The agenda-setting function of mass media. *Public Opin Q* 1972;**36**(2):176-87
7. Scheufele DA, Tewksbury D. Framing, agenda setting, and priming: The evolution of three media effects models. *J Commun* 2007;**57**(1):9-20
8. Seale C. *Media and health*: SAGE Publications Limited, 2002.
9. Wilsnack RW, Wilsnack SC. *Gender and alcohol: Individual and social perspectives*: Rutgers Center of Alcohol Studies, 1997.
10. Casswell S. Public Discourse on Alcohol. *Health Promot Int* 1997;**12**(3):251-57
11. Dorfman L. Studying the news on public health: How content analysis supports media advocacy. *Am J Health Behav* 2003;**27**(Supplement 3):S217-S26
12. Day K, Gough B, McFadden M. "Warning! alcohol can seriously damage your feminine health" a discourse analysis of recent British newspaper coverage of women and drinking. *Fem Media Stud* 2004;**4**(2):165-83
13. Wood K, Patterson C, Katikireddi SV, Hilton S. Harms to 'others' from alcohol consumption in the minimum unit pricing policy debate: a qualitative content analysis of UK newspapers (2005–12). *Addict* 2014;**109**(4):578-84
14. Lyons AC, Dalton SI, Hoy A. 'Hardcore Drinking': Portrayals of Alcohol Consumption in Young Women's and Men's Magazines. *J Health Psychol* 2006;**11**(2):223-32
15. Nicholls J. UK news reporting of alcohol: An analysis of television and newspaper coverage. *Drugs (Abingdon Engl)* 2011;**18**(3):200-06
16. Atkinson A, Kirton A, Sumnall H. The gendering of alcohol in consumer magazines: An analysis of male and female targeted publications. *J Gend Stud* 2012;**21**(4):365-86
17. Herring R, Berridge V, Thom B. Binge drinking: an exploration of a confused concept. *J Epidemiol Community Health* 2008;**62**(6):476-79
18. Guise JMF, Gill JS. 'Binge drinking? It's good, it's harmless fun': a discourse analysis of accounts of female undergraduate drinking in Scotland. *Health Educ Res* 2007;**22**(6):895-906
19. De Visser RO, McDonnell EJ. 'That's OK. He's a guy': A mixed-methods study of gender double-standards for alcohol use. *Psychol Health* 2012;**27**(5):618-39
20. Willott S, Lyons AC. Consuming male identities: Masculinities, gender relations and alcohol consumption in Aotearoa New Zealand. *J Community Appl Soc Psychol* 2012;**22**(4):330-45
21. Hutton F, Griffin C, Lyons A, Niland P, McCreanor T. "Tragic girls" and "crack whores": Alcohol, femininity and Facebook. *Fem Psychol* 2016;**26**(1):73-93
22. National Readership Survey. National Readership Survey. Secondary National Readership Survey 2012. <http://www.nrs.co.uk/>. (accessed 19th September 2012)


23. Newsworks. Newsworks. Secondary Newsworks 2012. www.nmauk.co.uk (accessed 19th September 2012)
24. Ofcom. Communications Market Report 2015. London: Ofcom, 2015.
25. Hilton S, Patterson C, Teyhan A. Escalating coverage of obesity in UK newspapers: the evolution and framing of the “obesity epidemic” from 1996 to 2010. *Obes* 2012;**20**(8):1688-95
26. Patterson C, Katikireddi SV, Wood K, Hilton S. Representations of minimum unit pricing for alcohol in UK newspapers: a case study of a public health policy debate. *J Public Health* 2015;**37**(1):40-49
27. HM Government Home Department. The Government's Alcohol Strategy, 2012.
28. Altheide DL, Schneider CJ. *Qualitative media analysis*: Sage, 2012.
29. Neuendorf KA, Kumar A. Content analysis. *The International Encyclopedia of Political Communication* 2006
30. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;**3**(2):77-101
31. Barry AE, Piazza-Gardner AK. Drunkorexia: Understanding the Co-occurrence of Alcohol Consumption and Eating/Exercise Weight Management Behaviors. *J Am Coll Health* 2012;**60**(3):236-43 doi: 10.1080/07448481.2011.587487published Online First: Epub Date].
32. Measham F, Østergaard J. The public face of binge drinking: British and Danish young women, recent trends in alcohol consumption and the European binge drinking debate. *Probat J* 2009;**56**(4):415-34
33. Fitzgerald N, Angus K, Emslie C, Shipton D, Bauld L. Gender differences in the impact of population-level alcohol policy interventions: evidence synthesis of systematic reviews. *Addict* in press
34. The Sun. Your friends, bad habits or anxiety... take our quiz to find YOUR booze binge trigger. Secondary Your friends, bad habits or anxiety... take our quiz to find YOUR booze binge trigger 2016. <http://www.thesun.co.uk/sol/homepage/fabulous/7112837/Your-friends-bad-habits-or-anxiety-take-our-quiz-to-find-YOUR-booze-binge-trigger.html> (accessed 13 May 2016)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legends

Figure 1: Frequency of publication of articles about ‘binge’ drinking by quarter

For peer review only



Frequency of publication of articles about 'binge' drinking by quarter
[Figure 1]
146x100mm (300 x 300 DPI)

Correction

Patterson C, Emslie C, Mason O, *et al.* Content analysis of UK newspaper and online news representations of women's and men's 'binge' drinking: a challenge for communicating evidence-based messages about single-episodic drinking? *BMJ Open* 2016;**6**: e013124. doi: 10.1136/bmjopen-2016-013124

In [Table 1](#), the figures in parentheses in the first column are incorrect. The corrected [Table 1](#) is shown below.

Table 1 Summary of publications and articles in the sample

Genre / medium	Publication	All articles		Article format					
				Standard		Feature		Editorial	
		n	%	n	%	n	%	n	%
Quality (n=127)	Guardian / Observer	58	18.8	48	20.3	9	17.3	1	5.3
	Independent / Independent on Sunday	17	5.5	11	4.6	5	9.6	1	5.3
	Daily Telegraph / Sunday Telegraph	52	16.9	44	18.6	5	9.6	3	15.8
Middle-market tabloids (n=67)	Daily Mail / Mail on Sunday	54	17.5	33	13.9	18	34.6	3	15.8
	Express / Sunday Express	13	4.2	8	3.4	3	5.8	2	10.5
Tabloids (n=75)	Daily Mirror / Sunday Mirror	13	4.2	10	4.2	2	3.8	1	5.3
	The Sun / News of the World	62	20.1	46	19.4	9	17.3	7	36.8
Online (n=39)	BBC News website	39	12.7	37	15.6	1	1.9	1	5.3
Total:		308	100	237	100	52	100	19	100

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

BMJ Open 2017;**7**:e013124corr1. doi:10.1136/bmjopen-2016-013124corr1



CrossMark