

BMJ Open

Time trends in educational inequalities in cancer mortality in Colombia, 1998-2012: is increasing healthcare insurance coverage in a middle-income country reflected in reducing inequalities in premature cancer mortality?

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2015-008985
Article Type:	Research
Date Submitted by the Author:	03-Jun-2015
Complete List of Authors:	de Vries, Esther; National Cancer Institute, Cancer Surveillance and Epidemiology Arroyave, Ivan; National School of Public Health, Department of Specific Sciences in Public Health; Erasmus MC University Medical Center, Public Health Pardo, Constanza; National Cancer Institute, Cancer surveillance and epidemiology
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Global health, Health policy, Oncology, Sociology
Keywords:	Epidemiology < ONCOLOGY, PUBLIC HEALTH, EPIDEMIOLOGY, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, SOCIAL MEDICINE

SCHOLARONE™
Manuscripts

Time trends in educational inequalities in cancer mortality in Colombia, 1998-2012: is increasing healthcare insurance coverage in a middle-income country reflected in reducing inequalities in premature cancer mortality?

Authors:

ESTHER DE VRIES^{1,2,#,*}, IVAN ARROYAVE^{2,3,#}, CONSTANZA PARDO⁴

: Shared first authorship, both authors contributed equally;

* : Corresponding author

Affiliations:

1. Directorate for Research, Surveillance and Cancer Control, National Cancer Institute, Calle 1 No.9-85, Bogota, Colombia
2. Department of Public Health, Erasmus University Medical Center. Dr. Molewaterplein 50, 3015 GE, Rotterdam, The Netherlands
3. Department of Specific Sciences in Public Health, National School of Public Health, University of Antioquia, Calle 62 N° 52-59, Medellin, Colombia
4. Cancer Surveillance and Epidemiology Group, National Cancer Institute, Calle 1 No.9-85, Bogota, Colombia

Corresponding Author:

Esther de Vries
Director for Research, Surveillance and Cancer Control
National Cancer Institute, Bogotá, Colombia

Data sharing statement: No additional data are available

Funding statement: This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. The salaries of the investigators were provided directly by their employers.

Competing interests statement: The authors have no competing interests to report

STRUCTURED ABSTRACT

Background: Premature cancer mortality in Colombia was documented to decline for all educational levels, although strongest in the higher educated levels.

Methods: In this descriptive study we linked Colombian national mortality data of age groups 20-64 years to census data to obtain age-standardized cancer mortality rates by educational level. Using Poisson regression we modelled premature mortality by educational level estimating rate ratios (RR) and relative index of inequality (RII) and its absolute counterpart, the Slope Index of Inequality (SII).

Results: RR show increased risks of dying among the lower educated compared to the highest educated, this tendency was stronger in women ($RR_{\text{primary}} 1.35$; $RR_{\text{secondary}} 1.11$, both $p < 0.0001$) than in men ($RR_{\text{primary}} 1.49$; $RR_{\text{secondary}} 1.22$, both $p < 0.0001$). In absolute terms (SII) cancer caused a difference of 20.5 per 100.000 male deaths, and 28.5 per 100.000 female deaths between the highest and lowest educated. RII was significantly higher for women than for men and in the younger age categories and initially decreased from 1998-2002 to 2003-2007, but then in the period 2008-2012 they increased significantly back to its previous levels. Among women, no significant increases or declines in cancer mortality over time were observed in recent periods in the lowest educated group, whereas strong recent declines were observed in those with secondary education or higher.

Conclusion: Documented educational inequalities in cancer mortality in Colombia are not only persistent, but increasing, fastest among women, and concentrated in young age categories. This trend was not curbed by the increase in healthcare insurance coverage. These results should urge policy makers to improve equal access to prevention, early detection, diagnostic and treatment facilities.

Strengths and limitations

- Population-based mortality databases with information on educational level provide a unique data source to evaluate educational differences
 - Definition of the variable for educational level does not guarantee having terminated the indicated level, there may be variance within the educational groups
 - Time period included covers a period of important changes in Colombian society, expected to be reflected in cancer mortality rates
 - Multiple imputation improved statistical power and precision of analyses
 - The underlying causes of these trends are unknown
-

Contriburship statement

- Obtaining data: Iván Arroyave, Constanza Pardo
- Study Design: Iván Arroyave, Esther de Vries
- Programming and analysis: Iván Arroyave, Esther de Vries
- Manuscript preparation and revision: Esther de Vries, Iván Arroyave, Constanza Pardo

For peer review only

Introduction

Colombia is a country with very large socioeconomic inequalities, causing, among others, large differences in health-related topics such as life expectancy and incidence, prognosis and mortality of disease [1-3]. In 1993, a major health care reform in Colombia introduced mandatory health insurance coverage (HIC) [4]. Citizens are assigned to two major schemes based on income: (i) the contributory scheme, which covers workers and their families with an income above the cut-off and is financed through payroll and employer's contributions; and (ii) the subsidized scheme, mainly funded via taxes, which subsidizes the poor as identified through a proxy means test [5]. HIC was 23.7% in 1993 just before the reform [6] and rose to 37.7% in 1994, just afterwards [4]. Initially, the coverage slowly improved [4] and a subsequent reform led in 2002 to a noticeable increase of healthcare coverage (reaching around 96% by 2008) by improving efficiency in the use of resources and by reducing its reliance on national budgets [7] (Figure 1). In the poorest income quartile, HIC increased from 6% to over 70% between 1993 and 2009 [5], attributable to the subsidized scheme [8], which shows that the program reached the poorer income groups more aggressively. This increased HIC is expected to contribute to a reduction in health inequalities [4, 5], mainly by improving the situation for the poor: ensuring timely care and bringing patients in closer contact with the health care system [9, 10]. It has been argued, however, that the reform increased the complexity of the system which could potentially lead to delays in certain types of care [11] and reduced spending in prevention and public health [12].

In Colombia, three different periods in the implementation of universal HIC can be discerned (Figure 1): During the period 1998-2002, HIC was relatively low and stable (around 70% during all the period); the second period (2003-2007) covered the years of rapid increase in coverage among the poor (6.9% average annual growth in HIC, 12.2% in subsidized scheme); while during the third period (2008-2012) universal coverage remained stable according to official figures at on average 96.1%. The discrepancy in HIC trends between these three periods offers a natural experiment to examine the impact of HIC on socioeconomic disparities in mortality.

Few studies have examined the impact of different health insurance status on inequalities in mortality and even less is known about inequalities in mortality in middle-income countries, many of whom introduced major health care system reforms during the last decades. As in Colombia the increase of HIC towards universal coverage was deliberately addressed to the poorest population [4, 5], and educational level is a good proxy variable for Socioeconomic Status (SAS) [13], we hypothesize that increased HIC contributed to reducing inequalities in cancer mortality. Cancer mortality is a particularly interesting indicator of the effect of insurance coverage due to the expected increased access to

prevention measures, early detection, timely treatment therapies, and high-cost interventions [14-16].

Indeed, previous analyses of time trends of differences in cancer mortality by attained educational level in Colombia (1998-2007) were promising, particularly for gastric and cervical cancer where faster declines in mortality occurred in the lowest educated groups and therefore socioeconomic differences were expected to decrease in the near future [3]. In this paper we evaluate if the seemingly positive trend in cancer mortality by educational level continued, by using a slightly different definition of educational level which allowed us to extend the previous analyses (up to 2007) by 5 extra years, covering 1998-2012.

Methods

Data

Education level criteria

Educational level was categorized in three groups based on the highest educational level accessed (not necessarily graduated) by the deceased: (a) Primary school or less, (b) Secondary school, and (c) Tertiary (post-secondary education). This definition differs from the previously used attained (graduated) educational level variable [1, 3], which was discontinued in the national databases after 2007.

Deaths

National mortality data for cancer deaths (International Classification of Diseases (ICD-10) codes C00-C96, see Appendix-Table 1) for the period 1998-2012 were obtained from the National Administrative Department of Statistics (DANE), which routinely registers information on sex, date and cause of death as well as educational level from death certificates. Although the exact wording for the variable "accessed education" changed in the DANE database since 2008, the new categories were consistent with those of the databases of previous years and could be used. Trends of counts were found to be continuous and regular.

We focused on adult premature mortality (mortality below age 65) because it is known that information on educational level from mortality statistics is unreliable at ages 65 and beyond [17]. Additionally, premature mortality is an indicator of population health believed to be strongly influenced by social, economic and environmental factors [18], and is a common indicator of health system performance [19].

Data on age and sex were available for more than 99% of all deaths, while data on educational level was missing for approximately 16.5% of 190,993 cancer deaths. We used multiple imputation methods [20] implemented in SAS through the IMPUTE procedure to impute educational level for these cases in order to avoid bias due to the potentially higher rates of missing education for lower educated individuals, and to minimize the potential for numerator/denominator bias [17]. In short, this procedure fits a sequence of regression models and draws values from the corresponding predictive distributions. The sequential regression procedure was applied based on a model that included sex, region, urban/rural residence and marital status as covariates. Details of this method are

explained elsewhere [20]. The imputation procedure was successful in 90.8% of cases with missing information, resulting in a total of 188,091 (98.5%) cancer deaths for our analyses.

Population

Data on mid-year population counts by age, sex and educational level were obtained from the Colombian Demography Health Surveys (DHS) [21] which contain periodical information on the distribution of education by age, sex and calendar year (1995, 2000, 2005 and 2010). Age 20 was chosen as the lower age limit of this study as almost 100% of individuals accessed their highest educational level by this age.

The resulting proportions of individuals in each educational level were multiplied with the total population numbers per year, age and sex which were obtained from census combined with statistical projections from DANE [22]. These two information sources were combined to estimate the annual population size in each educational group (Appendix-Figure 1). We performed demographic projections to obtain population counts for years in-between every lustrum using the demographic Software PASEX [23]. Additional details on the procedure are available elsewhere [23].

Analysis

All analyses were conducted in each of the five multiple databases generated by the multiple imputation process. Since results were nearly identical for all imputations, we used standard techniques as implemented in the PROC MIANALYZE procedure in SAS to combine estimates from all databases and adjust standard errors to account for uncertainty in the imputation [24]. This procedure reads the parameter estimates and associated covariance matrix for each imputed data set, and then derives valid multivariate inferences for these parameters. This allows for valid statistical inference that appropriately reflects uncertainty due to missing values [24].

All analyses were conducted in SAS® version 9.2.

Age Standardised Mortality Rates (ASMR)

Data on population counts were combined with data on deaths to obtain a complete database of death counts by educational level, sex and five-year age group. We first calculated age-standardized mortality rates by educational level, sex and year using the World Health population of 1997 [25] as standard population, resulting in Age

Standardised Mortality Rates (ASMR) expressed per 100,000 person-years (Figure 2). We also calculated ASMR by sex and year (Figure 2) and by educational level and sex (Table 1).

Annual trends in ASMR by sex and educational level were quantified by calculating the estimated Annual Percentage Change (EAPC) which measures the average rate of change in the mortality rate per year (negative EAPC: decreasing trend, positive EAPC : increasing trend). To test whether an apparent change in mortality trends was statistically significant, we used joinpoint regression, which fits a series of joined straight lines to age-adjusted rates and uses a Monte Carlo Permutation method to identify the best-fitting point (called joinpoint, year in which a significant change in the mortality trend occurred), where the rate of increase or decrease changes significantly [26]. EAPC and joinpoints were determined based on the log-transformed ASMRs and their standard errors. We specified a maximum of 2 joinpoints with at least 4 observation points to either extreme of the data using joinpoint modelling based on the Joinpoint program [27] (Figure 2).

Regression models

We implemented separate Poisson regression models separately by sex with number of deaths as dependent variable and the natural log of person-years as offset variable, incorporating age and educational level as independent variables. We first estimated Rate Ratios (RR) of mortality by educational level, which compared the mortality of all educational groups to the mortality in the tertiary education group. Changes in the rate ratio over time result from changes in both risks and the distribution of educational level [28]. To assess changes in disparities 'controlling' for changes in the educational distribution, we estimated the Relative Index of Inequality (RII), regressing mortality on the mid-point of the cumulative distribution of education, thereby taking into account the size of each educational group [28, 29]. Values higher than 1 indicate educational inequalities favouring the higher educated.

We evaluated if RRs significantly changed over time and within periods on continuous scales, which was not the case (results not shown). We calculated RII for three 5-year periods (1998-2002, 2003-2007, and 2008-2012) and for three age groups (20-34, 35-49, and 50-64) in order to identify differences in inequalities along age (Figure 2).

To compare the contribution of each cause of death to mortality disparities for non-communicable diseases (classified by their ICD-10 codes, see Appendix-table 1) we calculated the Slope Index of Inequality (SII) (Figure 4). The SII is a measure of absolute disparities that represents the difference in mortality between the population at the top and the bottom of the educational distribution [28]. Further details on the RII and SII are available elsewhere [28].

Results

Table 1 shows counts of cancer deaths and population at ages 20-64 years between 1998 and 2012 in Colombia. After excluding non-imputed cases (1.5%), 106,199 cancer deaths occurred from 1998 to 2012 among women while 81,892 occurred among men, with age-standardized cancer mortality rates (ASMR, per 100,000 person-years) of 58 among women and 49 among men. Men and women with lower levels of education had higher cancer mortality rates than their higher-educated counterparts. In addition, ASMR were larger among women than among men in each educational level.

The risk of dying was significantly and consistently higher among the lower educated. The RRs show clearly and statistically significantly increased risks of dying among the lower educated compared to the highest educated, this tendency was stronger in women ($RR_{primary}=1.35$; $RR_{secondary}=1.11$ both $p<0.0001$) than in men ($RR_{primary}=1.49$; $RR_{secondary}=1.22$, both $p<0.0001$).

During the 15 years of observation, the general tendency was for age-standardized cancer mortality rates to decline, with recent stabilizations, reaching an average ASMR in 2012 of 47.3 per 100,000 males and 55.2 per 100,000 females. Average EAPC for males in the periods defined by joinpoint were: 1998-2000: 2.4% (95% CI -2.0; 7.0), 2000-2012: -0.92% (95% CI -1.2; -0.7); for females 1998-2003: 0.7 (95% CI -0.7; 2.1), 2003-2006: -2.8% (95% CI -8.2; 2.9), 2006-2012 -0.6 (95% CI -1.5; 0.3). The general tendency for a declining trend in mortality was not reflected equally by educational level and sex (figure 2). Women had higher mortality rates than men. No significant increases or declines over time were observed in the lowest educated groups, whereas in the middle and high educated groups there was at least one period with significantly declining cancer mortality rates. In these groups, tendencies were for mortality to decrease substantially in the period after 2002/2003, with the exception of the highest educated males who showed recent increases. The strongest declines in cancer mortality over time were observed in the group of women with secondary or tertiary education. The most consistent declines in cancer mortality rates for all educational levels in both sexes were observed in the period 2003-2007, the period with rapid increases in HIC, particularly among the poor.

RII initially decreased from 1998-2002 to 2003-2007 (reaching significance among women), but then in the period 2008-2012 they increased back to its previous levels (Figure 3, left panel). Trends in the joinpoint for RII are in line with this finding: no significant change was found for educational inequalities among men, whereas there was an initial significant decline (-7.6%) among women up to 2004 followed by a significant increase (+3.8%) in the period 2004-2012 (Appendix-Figure 2). RII also showed that inequalities were significantly higher for the younger age categories compared to older ones and tended to be higher for women (Figure 3, right panel).

Figure 4 shows the contribution to absolute differences in ASMR by education as measured by the SII for non-communicable diseases (appendix-table 1). Cancer was the second largest cause, after cardiovascular disease: explaining 20% of male differences (SII=20.5 deaths per 100,000 population) and 24% of female differences (SII=28.5). Absolute differences in cancer mortality were 39% larger for women than for men (28.5 vs. 20.5), while this difference was only 14% for all non-communicable diseases (SII= 103.2 and 120.3 respectively).

For peer review only

Discussion

We observed strong and persistent educational differences in cancer mortality rates in Colombia. These differences appeared to be diminishing in the period 2002-2007 [3] when healthcare insurance coverage largely grew, especially for the most disadvantaged part of the population. But in recent years, despite the achievement of almost universal HIC, inequalities were widening, particularly among women, with rates in the lowest educational groups being stable or increasing and those in the higher educated groups gradually declining. This means that the lowest educated part of the population is increasingly lagging behind in the progress against cancer, despite the increase in insurance coverage.

Explanation of results

Educational level was found to be previously as a good proxy variable for Socioeconomic Status (SES) [13]. Lower SES is associated with a higher prevalence of cancer risk factors (such as smoking, alcohol, obesity, occupational risk factors, housing circumstances), and less healthcare utilization (because of lower resources but also lower awareness and poorer access due to longer distances or high impact of family income when taking time off to visit a doctor) [30-32].

Not only absolute cancer mortality rates, but also inequalities in mortality were largely higher for women than for men. In a previous paper we showed this to be almost entirely attributable to cervical cancer mortality, which accounted for 51% of inequalities in total female cancer mortality in 1998-2007 at 25-64 [3]. Several Colombian studies have identified that both characteristics of the health system such as access and delays in diagnosis and treatment, and characteristics of the populations of lower socioeconomic classes contribute to the higher burden of cervical cancer mortality [33, 34]. A previous study showed that lower access to mammography screening for early detection of breast cancer was associated with lower education, affiliation to the subsidized regime or being unaffiliated [35]. In general, financial access to healthcare has been more limited to women [36, 37] which potentially limits access to expensive cancer therapies, particularly for most deprived women.

We also observed clearly increased educational differences in cancer mortality for the youngest age groups, in which mortality from cancer is a relatively rare event. Among males, leukaemias, stomach cancers and Non-Hodgkin lymphomas are important cancer causes of deaths in the age groups 15-39 years (representing 44% of all cancer deaths in this age group); for females these include cervical cancer and leukaemias as well as non-Hodgkin lymphomas (representing 38%) [38]. Previous studies have shown that mortality

or survival of these cancer types is very much dependent on socioeconomic status [3, 14, 39]. As these tumours explain an important proportion of cancer deaths in these ages, this probably explains the high RII in the younger age groups as compared to the higher age categories.

Our results clearly illustrate that an almost complete coverage does not necessarily reduce inequalities in health [40, 41]. Particularly, having health insurance may be universal, but depending on income the type of health insurance is different (subsidized, contributory and special or exceptional regimes), with the most wealthy population often buying additional private health assurance to ensure rapid and more broad access [42]. Complete coverage does not automatically guarantee quality of the insurance (translating into early and timely diagnosis and adequate treatment), as seems to be the case: To warrant rights to get access to expensive treatments and medication, as usual in cancer, exceptional legal mechanisms are frequently launched in Colombia: Technical-scientific committees of health assurers, and an action for protection so-called “tutela” [42, 43]. A study shows that those affiliated to the contributory regime are more likely to warrant additional rights more efficiently [43], potentially increasing inequalities between regimes.

In parallel with the increases in HIC, other changes in the Colombian health care system occurred, including increases in investments in care [5, 44]. Of course, our results are consequence of the impact of the entire reform rather than only the increased coverage in health insurance. However, despite the large reforms, inequalities in cancer mortality in Colombia are increasing. This is not unique to Colombia; in several countries, as diverse as Taiwan, Thailand and European countries, inequalities in health increased upon reaching almost universal coverage [45-48].

Strengths and weaknesses

This is one of few studies investigating educational differences in mortality using population-based mortality and population-based educational level. Despite several strengths, some limitations should be considered in our study: We did not correct for misclassification within the cancer groups, but since we studied the group of cancer deaths overall, regardless of the subtype of cancer, most of these errors are cancelled out. In general “cancer” as cause of death is correctly coded [49], and most likely particularly in setting of ages under 65.

Data on mortality were obtained from mortality registries, while data on the population distribution by education were obtained from censuses and demographic projections. This may have led to the so-called numerator/denominator bias, which generally results in an overestimation of disparities [17]. Furthermore, for some years, data on population size

were obtained from demographic projections combined with distributions of education from surveys. To assess the impact of this potential bias, we experimented with different education distributions from multiple data sources [50-52], showing that the overall trends in our study were robust to different assumptions on the distribution of education [3].

Potentially, there has been substantial under-registration of deaths in some regions. Previous studies comparing national mortality rates with indirect estimates from census [53, 54] suggest that underregistration is particularly important in the poorest regions, implying that estimates of disparities in mortality by educational level are likely to be an underestimation, because those with lower education are more likely to live in areas with higher underregistration rates. This may also have led to underestimation of the extent to which inequalities have increased, because underregistration decreased over the study period [53, 54]. Our results, therefore, are indicative of potentially larger inequalities in mortality by education, which continue to increase.

Information on education was missing for 16.5% of cancer deaths, potentially leading to an underestimation of disparities, as missing values are usually more common in the least educated [55]. However, we imputed values for educational level for individuals with missing educational information based on a information on age, sex marital status, region and urban/rural residence, thereby limiting the potential impact of this source of bias.

Unanswered questions and future research

Socioeconomic inequalities in mortality of non-communicable diseases have been associated with the unequal distribution of behavioural risk factors, such as smoking, alcohol consumption, an unhealthy diet, sedentary lifestyle higher risk of injuries [2, 56]. This implies underlying inequalities in exposure to risk factors and incidence of these diseases. However, data on the incidence of disease and prevalence distribution of all those behavioural and poverty risk factors by socioeconomic status are largely lacking in Colombia and most developing countries. Studies on the mechanisms underlying the way these risk factors contribute to inequalities in mortality are scarce, even in developed countries. We previously described not only large inequalities in mortality of cancers with a primary infectious aetiology such as gastric cancer [3], but also documented inequalities in survival of gastric cancer measured by both type of health insurance and socioeconomic stratum [14]. These findings indicate that both access and quality of care (health insurance) as other factors related to socioeconomic status independently affect survival for the lower socioeconomic classes negatively. Detailed studies are needed on both types of factors: which barriers exist for timely diagnosis and adequate treatment in situations

with universal health coverage and how can they be tackled? And on which other factors related to socioeconomic stratum can interventions be designed? The answers to these questions will be various and, depending on the specific disease, will be quite different in character. It is of high importance that data on socioeconomic indicators are collected in a standard way, so that for example cancer registries can calculate incidence rates by socioeconomic stratum, which, at this moment, is not possible in Colombia [57].

Socioeconomic differences were substantially larger for women than for men, in our previous study this difference was almost entirely attributable to inequalities in cervical cancer mortality [3]. In this more recent study we see that the gender inequalities grew despite the virtually universal HIC since 2008. Gender inequalities were larger for cancer than for other non-communicable diseases according SII; more detailed analyses are needed to assess to which extent the growing inequalities are due to an increase in cervical cancer mortality amongst less favoured.

Conclusions

The recent negative curbs in mortality among the lowest educated groups, resulting in increasingly differences between educational groups, are a call for attention by the Colombian health authorities to take measures in both primary prevention, access to early detection and timely and adequate treatment. Underlying causes of the observed differences are multiple, from differences in housing, lifestyle, health-related beliefs, to differences in access to diagnostic facilities and high-quality treatment, which all should be of concern to those in charge [58]. Insurance coverage is clearly not enough to counterbalance the deleterious effects of decreases in spending on primary prevention and other social and economic conditions which determine the growing levels of educational inequalities in cancer.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

We are grateful to Mauricio Avendaño and Alex Burdorf for their contributions at initial stages of the analyses and for their continued support in this project.

For peer review only

References

1. Arroyave I, Cardona D, Burdorf A, Avendaño M: **The Impact of Increasing Health Insurance Coverage on Disparities in Mortality: Health Care Reform in Colombia, 1998–2007.** *American Journal of Public Health* 2013:e1-e7.
2. Arroyave I, Cardona D, Burdorf A, Avendaño M: **Socioeconomic inequalities in premature mortality in Colombia, 1998-2007: The double burden of non-communicable diseases and injuries.** *Preventive Medicine* 2014(64C):41-47.
3. de Vries E, Arroyave I, Pardo EC, Wiesner C, Murillo R, Forman D, Burdorf L, Avendaño M: **Trends in inequalities in premature cancer mortality by educational level in Colombia, 1998-2007.** *Journal of Epidemiology & Community Health* 2015, **69**(5):408-415.
4. Agudelo-Calderón CA, Cardona-Botero J, Ortega-Bolaños J, Robledo-Martínez R: **Sistema de salud en Colombia: 20 años de logros y problemas.** *Ciência & Saúde Coletiva* 2011, **16**:2817-2828.
5. Giedion U, Uribe MV: **Colombia's Universal Health Insurance System.** *Health Affairs* 2009, **28**(3):853-863.
6. Céspedes-Londoño JE, Jaramillo-Pérez I, Castaño-Yepes RA: **Impacto de la reforma del sistema de seguridad social sobre la equidad en los servicios de salud en Colombia.** *Cadernos de Saúde Pública* 2002, **18**(4):1003-1024.
7. Nieto-Parra S, Olivera M: **Making Reform Happen in Colombia: The Process of Regional Transfer Reform.** In: *OECD Development Centre Working Papers*. Organisation for Economic Co-operation and Development (OECD); 2012: 39.
8. Ministerio de la Protección Social: **Informe de Actividades 2009-2010 - Informe Cuatrienio 2006 - 2010, al Honorable Congreso de la República.** In: Edited by Dirección General de Planeación y Análisis de Política, Grupo de Estudios y Planeación Sectorial. Bogotá D.C.; 2010.
9. Ruiz F, Amaya L, Venegas S: **Progressive segmented health insurance: Colombian health reform and access to health services.** *Health Economics* 2007, **16**(1):3-18.
10. Miller G, Pinto DM, Vera-Hernández M: **Risk Protection, Service Use, and Health Outcomes Under Colombia's Health Insurance Program for the Poor.** *American Economic Journal: Applied Economics* 2013, **5**(4):61-91.
11. Abadia CE, Oviedo DG: **Bureaucratic Itineraries in Colombia. A theoretical and methodological tool to assess managed-care health care systems.** *Social Science & Medicine* 2009, **68**(6):1153-1160.
12. De Groote T, De Paepe P, Unger J-P: **Colombia: in vivo test of health sector privatization in the developing world.** *International Journal of Health Services* 2005, **35**(1):125 - 141.
13. Oakes JM, Rossi PH: **The measurement of SES in health research: current practice and steps toward a new approach.** *Social Science & Medicine* 2003, **56**(4):769-784.
14. de Vries E, Uribe C, Pardo C, Lemmens V, Van de Poel E, Forman D: **Gastric cancer survival and affiliation to health insurance in a middle-income setting.** *Cancer Epidemiology* 2014.
15. Smith JK, Ng SC, Zhou Z, Carroll JE, McDade TP, Shah SA, Tseng JF: **Does increasing insurance improve outcomes for US cancer patients?** *Journal of Surgical Research* 2013, **185**(1):15-20.
16. Kim S, Kwon S: **Has the National Health Insurance improved the inequality in the use of tertiary-care hospitals in Korea?** *Health Policy* 2014, **118**(3):377-385.
17. Kunst AE, Groenhouf F, Borgan J, Costa G, Desplanques G, Faggiano F, Hemström O, Martikainen P, Vågerö D, Valkonen T *et al*: **Socio-economic inequalities in mortality.**

Methodological problems illustrated with three examples from Europe. *Revue d'epidemiologie et de sante publique* 1998, **46**(6):467-479.

18. World Health Organization: **Closing the gap in a generation: Health equity through action on the social determinants of health**. In. Geneva, Switzerland: World Health Organization; 2008.

19. Smith PC: **Performance measurement for health system improvement: experiences, challenges and prospects**: Cambridge University Press; 2010.

20. Raghunathan T, Lepkowski J, Van Hoewyk J, Solenberger P: **A Multivariate Technique for Multiply Imputing Missing Values Using a Sequence of Regression Models**. *Survey Methodology* 2001, **27**(1):85-95.

21. Profamilia: **Encuesta Nacional de Demografía y Salud**. In. Bogotá D.C.; 1990, 1995, 2000, 2005, 2010.

22. Departamento Administrativo Nacional de Estadística (DANE): **Colombia: Tablas de mortalidad nacionales y departamentales 1985-2020**. In., Sep. 2007 edn. Bogotá D.C.; 2007.

23. United States Census Bureau: **Population Analysis System (PASEX)**. In: *Population Analysis Spreadsheets*. 2.04g edn. Washington DC: International Programs Center (IPC) - United States Census Bureau; 2011.

24. SAS Institute I: **The MIANALYZE Procedure**. In: *User's Guide SAS/STAT® 92*. Edited by Inc. SI, Version 8 edn. Cary, NC: SAS Institute Inc.; 2008: 201-233.

25. Ahmad OB, Boschi-Pinto C, Lopez AD, Murray CJ, Lozano R, Inoue M: **Age standardization of rates: A new WHO standard**. In: *GPE Discussion Paper Series*. Geneva: World Health Organization; 2001.

26. National Cancer Institute, Statistical Research and Applications Branch: **Joinpoint Regression Program**. In., 4.0.4 edn; 2013.

27. Kim HJ, Fay MP, Feuer EJ, Midthune DN: **Permutation tests for joinpoint regression with applications to cancer rates**. *Statistics in Medicine* 2000, **19**(3):335-351.

28. Mackenbach JP, Kunst AE: **Measuring the magnitude of socio-economic inequalities in health: An overview of available measures illustrated with two examples from Europe**. *Social Science & Medicine* 1997, **44**(6):757-771.

29. Harper S, Lynch J, Meersman SC, Breen N, Davis WW, Reichman ME: **An overview of methods for monitoring social disparities in cancer with an example using trends in lung cancer incidence by area-socioeconomic position and race-ethnicity, 1992-2004**. *Am J Epidemiol* 2008, **167**(8):889-899.

30. Arias SA: **Inequidad y cáncer: una revisión conceptual**. *Revista de la Facultad Nacional de Salud Pública* 2009, **27**(3):341-348.

31. Woods L, Rachet B, Coleman M: **Origins of socio-economic inequalities in cancer survival: a review**. *Annals of Oncology* 2006, **17**(1):5-19.

32. Auvinen A, Karjalainen S: **Possible explanations for social class differences in cancer patient survival**. In: *Social Inequalities and Cancer*. edn. Edited by Kogevinas M, Pearce N, Susser M, Boffetta P. Lyon; 1997: 377-397.

33. Piñeros M, Cendales R, Murillo R, Wiesner C, Tovar S: **Cobertura de la Citología de Cuello Uterino y Factores Relacionados en Colombia, 2005**. *Revista de Salud Pública* 2007, **9**(3):327-341.

34. Wiesner-Ceballos C, Cendales-Duarte R, Murillo R, Piñeros M, Tovar-Murillo SL: **Seguimiento de mujeres con anormalidad citológica de cuello uterino, en Colombia**. *Revista de Salud Pública* 2010, **12**(1):1-13.

35. Charry LCd, Carrasquilla G, Roca S: **Equidad en la Detección del Cáncer de Seno en Colombia.** *Revista de Salud Pública* 2008, **10**(4):571-582.
36. Ewig C, Bello AH: **Gender equity and health sector reform in Colombia: Mixed state-market model yields mixed results.** *Social Science & Medicine* 2009, **68**(6):1145-1152.
37. Guarnizo-Herreño CC, Agudelo C: **Equidad de Género en el Acceso a los Servicios de Salud en Colombia.** *Revista de Salud Pública* 2008, **10**(1):44-57.
38. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F: **GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet].** In. Lyon, France: International Agency for Research on Cancer; 2013.
39. Petridou ET, Sergentanis T, Perlepe C, Papathoma P, Tsilimidos G, Kontogeorgi E, Kourti M, Baka M, Moschovi M, Polychronopoulou S: **Socio-economic disparities in survival from childhood leukemia in the USA and globally: a meta-analysis.** *Annals of Oncology* 2014, **26**(3):589-597.
40. Franks P, Clancy CM, Gold MR: **Health insurance and mortality: Evidence from a national cohort.** *JAMA: The Journal of the American Medical Association* 1993, **270**(6):737-741.
41. Katz SJ, Hofer TP: **Socioeconomic disparities in preventive care persist despite universal coverage. Breast and cervical cancer screening in Ontario and the United States.** *JAMA* 1994, **272**(7):530-534.
42. Guerrero R, Gallego AI, Becerril-Montekio V, Vásquez J: **Sistema de salud de Colombia.** *Salud Pública de México* 2011, **53**:s144-s155.
43. Uprimny R, Durán J: **Equidad y protección judicial del derecho a la salud en Colombia.** In: *Políticas Sociales.* Santiago de Chile: Comisión Económica para América Latina y el Caribe (CEPAL); 2014: 70.
44. Restrepo Zea JH: **El seguro de salud en Colombia: ¿cobertura universal?** *Revista Gerencia y Políticas de Salud* 2002(Nº 2):25-40.
45. Shou-Hsia C, Tung-Liang C: **The Effect of Universal Health Insurance on Health Care Utilization in Taiwan.** *JAMA: The Journal of the American Medical Association* 1997, **278**(2):89-93.
46. Suraratdecha C, Saithanu S, Tangcharoensathien V: **Is universal coverage a solution for disparities in health care?: Findings from three low-income provinces of Thailand.** *Health Policy* 2005, **73**(3):272-284.
47. Mackenbach JP, Stirbu I, Roskam A-JR, Schaap MM, Menvielle G, Leinsalu M, Kunst AE: **Socioeconomic Inequalities in Health in 22 European Countries.** *New England Journal of Medicine* 2008, **358**(23):2468-2481.
48. Avendaño M, Kunst AE, Huisman M, van Lenthe F, Bopp M, Regidor E, Glickman M, Costa G, Spadea T, Deboosere P *et al*: **Socioeconomic status and ischaemic heart disease mortality in 10 western European populations during the 1990s.** *Heart* 2006, **92**(4):461-467.
49. Harteloh P, de Bruin K, Kardaun J: **The reliability of cause-of-death coding in The Netherlands.** *European journal of epidemiology* 2010, **25**(8):531-538.
50. International Institute of Applied System Analysis (IIASA), Vienna Institute of Demography of the Austrian Academy of Sciences (VID): **IVEP-1.0 (IIASA/VID education database).** In. Viena; 2010.
51. **World Development Indicators (WDI)** [<http://data.worldbank.org/data-catalog/world-development-indicators>]

52. Barro RJ, Lee J-W: **A new data set of educational attainment in the world, 1950–2010**. In: *NBER Working Paper Series*. vol. Working Paper 15902. Cambridge, MA: National Bureau of Economic Research; 2010.

53. Patiño-Villada FA, Arango-Vélez EF, Quintero-Velásquez MA, Arenas-Sosa MM: **Factores de riesgo cardiovascular en una población urbana de Colombia**. *Revista de Salud Pública* 2011, **13**:433-445.

54. Storr CL, Cheng H, Posada-Villa J, Aguilar-Gaxiola S, Anthony JC: **Adult smokers in Colombia: who isn't giving it up?** *Addict Behav* 2008, **33**(3):412-421.

55. Rodríguez-García J: **Desigualdades socioeconómicas entre departamentos y su asociación con indicadores de mortalidad en Colombia en 2000**. *Revista Panamericana de Salud Pública* 2007, **21**:111-124.


56. Adler NE, Newman K: **Socioeconomic Disparities In Health: Pathways And Policies**. *Health Affairs* 2002, **21**(2):60-76.

57. De Vries E, Murillo R: **Regarding articles about Cali Cancer Registry 2**. *Salud Pública de México* 2015, **57**(3).

58. Ottersen OP, Dasgupta J, Blouin C, Buss P, Chongsuvivatwong V, Frenk J, Fukuda-Parr S, Gawanas BP, Giacaman R, Gyapong J: **The political origins of health inequity: prospects for change**. *The Lancet* 2014, **383**(9917):630-667.

FIGURES AND TABLES

Figure 1: Trends in national healthcare insurance coverage (HIC), Colombia, 1998-2012



Other schemes include primarily members of the military and teacher and oil workers syndicate members. Based on Annual reports of the Ministry of Health and Social Protection

Table 1: Descriptives of the study: deaths, population and rates separately by sex and educational level, Colombia, 1998-2012


Sex	Educational level ^a	Cancer deaths ^b	Population size	ASMR ^c (95% Confidence Intervals)	
Men	Primary	50,126	67,815,336	54.0	(53.5, 54.5)
	Secondary	22,273	67,742,806	45.1	(44.4, 45.8)
	Tertiary	9,493	30,715,631	41.4	(40.5, 42.3)
	Total	81,892	166,273,773	48.9	(48.8, 49.1)
Women	Primary	65,336	68,637,666	64.7	(64.2, 65.2)
	Secondary	30,414	72,387,214	54.2	(53.5, 54.9)
	Tertiary	10,449	35,149,442	47.2	(46.2, 48.3)
	Total	106,199	176,174,322	57.8	(57.6, 58.0)

^a: Educational attainment values registered after final imputation. Primary = up to elementary or primary school; secondary = any level of high-school; tertiary = any level of postsecondary education after high school including college and university.

^b: Cancer deaths after imputation.


^c: ASMR: Age standardized mortality rates per 100,000 population; Estimates (WHO standard population 1997) for educational level combine results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values.

Figure 2: Age-standardized cancer mortality trends, including APC based on joinpoint models, by sex and educational level




Markers: Observed age-standardized cancer mortality rates. Lines: modelled age-standardized cancer mortality trends. Estimates are results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values. The points represent ASMR, lines represent the trendlines between joinpoints. Blue, diamonds: maximum primary education; Red, triangles: secondary education; Green, circles: tertiary education. Numbers adjacent to the lines represent estimates annual percent change (EAPC) during the corresponding periods, based on joinpoint modelling; a star indicates statistical significance at $\alpha 0.05$.

Figure 3: Sex-specific Relative Index of Inequality by period and age group



Estimates from combined results from 5 databases resulting from multiple imputations, appropriately reflecting uncertainty attributable to missing values.


Figure 4. Slope Index of Inequality (SII) for non-communicable diseases



See Appendix-Table 1 for International Classification of Diseases (ICD-10) codes for non-communicable diseases

APPENDIX

Appendix-Figure 1. Educational level trends over time, Colombia, 1998-2012




The thicker markers are those from the year with real data available (source: Colombian Demography Health Surveys in 1995, 2000, 2005 and 2010). The points in between were interpolated (see methods)

Appendix-Table 1. Classification of Causes of death, International Classification of Diseases (ICD-10) codes for non-communicable diseases

Non-Communicable Diseases		ICD-10 codes
Cardiovascular diseases		I00-I99
Malignant neoplasm		C00-D48
Diabetes mellitus		E10-E14
Chronic lower respiratory diseases (CLRD)		J40-J47
Other Non-Communicable Diseases	Mental and behavioural disorders	F00-F99
	Diseases of the nervous system	G00-G99
	Diseases of the digestive system	K00-K93
	Diseases of the genitourinary system	N00-N99
	Congenital malformations, deformations and chromosomal abnormalities	Q00-Q99

Appendix-Figure 2: Sex-specific Relative Index of Inequality (RII) trends including APC based on joinpoint models



Markers: Annual observed cancer mortality RII. Lines: modelled RII trends. Estimates are results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values.

BMJ Open

Time trends in educational inequalities in cancer mortality in Colombia, 1998-2012

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2015-008985.R1
Article Type:	Research
Date Submitted by the Author:	06-Aug-2015
Complete List of Authors:	de Vries, Esther; National Cancer Institute, Cancer Surveillance and Epidemiology Arroyave, Ivan; National School of Public Health, Department of Specific Sciences in Public Health; Erasmus MC University Medical Center, Public Health Pardo, Constanza; National Cancer Institute, Cancer surveillance and epidemiology
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Global health, Health policy, Oncology, Sociology
Keywords:	Epidemiology < ONCOLOGY, PUBLIC HEALTH, EPIDEMIOLOGY, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, SOCIAL MEDICINE

SCHOLARONE™
Manuscripts

Time trends in educational inequalities in cancer mortality in Colombia, 1998-2012

Authors:

ESTHER DE VRIES ^{ERROR! REFERENCE SOURCE NOT FOUND.,1,#,*}, IVAN ARROYAVE ^{1,3,#}, CONSTANZA PARDO ¹

: Shared first authorship, both authors contributed equally;

* : Corresponding author

Affiliations:

1. Cancer Surveillance and Epidemiology Group, National Cancer Institute, Calle 1 No.9-85, Bogota, Colombia
2. Department of Public Health, Erasmus University Medical Center. Dr. Molewaterplein 50, 3015 GE, Rotterdam, The Netherlands
3. Department of Specific Sciences in Public Health, National School of Public Health, University of Antioquia, Calle 62 N° 52-59, Medellin, Colombia

Corresponding Author:

Esther de Vries

Director for Research, Surveillance and Cancer Control
National Cancer Institute, Bogotá, Colombia

Data sharing statement: No additional data are available

Funding statement: This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. The salaries of the investigators were provided directly by their employers.

Competing interests statement: The authors have no competing interests to report

Ethics committee approval: This article is based on secondary analysis of data on deaths and population counts in aggregate form made publically available by the National Statistics Office in Colombia. Ethical approval for this study was not required.

STRUCTURED ABSTRACT

Objectives: To evaluate trends in premature cancer mortality in Colombia by educational level in three periods; 1998-2002 with low healthcare insurance coverage, 2003-2007 with rapidly increasing coverage, and finally 2008-2012 with almost universal coverage (2008-2012).

Setting: Colombian population-based, national secondary mortality data..

Participants: We included all (n=188,091 cancer deaths occurring in age groups 20-64 years between 1998 and 2012, excluding only cases with low levels of quality of registration (n=2,902, 1.5%).

Primary and secondary outcome measures: In this descriptive study we linked mortality data of ages 20-64 years to census data to obtain age-standardized cancer mortality rates by educational level. Using Poisson regression we modelled premature mortality by educational level estimating rate ratios (RR), relative index of inequality (RII) and the Slope Index of Inequality (SII).

Results: Relative measures showed increased risks of dying among the lower educated compared to the highest educated, this tendency was stronger in women ($RR_{\text{primary}} 1.49$; $RR_{\text{secondary}} 1.22$, both $p < 0.0001$) than in men ($RR_{\text{primary}} 1.35$; $RR_{\text{secondary}} 1.11$, both $p < 0.0001$). In absolute terms (SII) cancer caused a difference per 100,000 deaths between the highest and lowest educated of 20.5 in males and 28.5 in females. RII was significantly higher among women and the younger age categories. RII decreased between the first and second period, afterwards (2008-2012) increased significantly back to their previous levels. Among women, no significant increases or declines in cancer mortality over time were observed in recent periods in the lowest educated group, whereas strong recent declines were observed in those with secondary education or higher.

Conclusions: Educational inequalities in cancer mortality in Colombia are increasing, in absolute and relative terms concentrated in young age categories. This trend was not curbed by increases in healthcare insurance coverage. Policy makers should focus on improve equal access to prevention, early detection, diagnostic and treatment facilities.

Strengths and limitations

- Population-based mortality databases with information on educational level provide a unique data source to evaluate educational differences
- Definition of the variable for educational level does not guarantee having terminated the indicated level, there may be variance within the educational groups
- Time period included covers a period of important changes in Colombian society, expected to be reflected in cancer mortality rates
- Multiple imputation improved statistical power and precision of analyses
- The underlying causes of these trends are unknown

Contributorship statement

Esther de Vries performed the joinpoint analyses, adapted tables and figures and was the main writer of the text. Ivan Arroyave performed the main part of the analyses apart from the joinpoints, discussed with Esther de Vries all initial tables and figures, and commented on the text. Constanza Pardo contributed to content and improvement of the text and figures.

Introduction

Colombia is a country with very large socioeconomic inequalities, causing, among others, large differences in health-related topics such as life expectancy and incidence, prognosis and mortality of disease ¹⁻³. In 1993, a major health care reform in Colombia introduced mandatory health insurance coverage (HIC) ⁴. Citizens are assigned to two major schemes based on income: (i) the contributory scheme, which covers workers and their families with an income above the cut-off and is financed through payroll and employer's contributions; and (ii) the subsidized scheme, mainly funded via taxes, which subsidizes the poor as identified through a proxy means test ⁵. HIC was 23.7% in 1993 just before the reform ⁶ and rose to 37.7% in 1994, just afterwards ⁴. Initially, the coverage slowly improved ⁴ and a subsequent reform led in 2002 to a noticeable increase of healthcare coverage (reaching around 96% by 2008) by improving efficiency in the use of resources and by reducing its reliance on national budgets ⁷ (Figure 1). In the poorest income quartile, HIC increased from 6% to over 70% between 1993 and 2009 ⁵, attributable to the subsidized scheme ⁸, which shows that the program reached the poorer income groups more aggressively. This increased HIC is expected to contribute to a reduction in health inequalities ^{4 5}, mainly by improving the situation for the poor: ensuring timely care and bringing patients in closer contact with the health care system ^{9 10}. It has been argued, however, that the reform increased the complexity of the system which could potentially lead to delays in certain types of care ¹¹ and reduced spending in prevention and public health ¹².

In Colombia, three different periods in the implementation of universal HIC can be discerned (Figure 1): During the period 1998-2002, HIC was relatively low and stable (around 70% during all the period); the second period (2003-2007) covered the years of rapid increase in coverage among the poor (6.9% average annual growth in HIC, 12.2% in subsidized scheme); while during the third period (2008-2012) universal coverage remained stable according to official figures at on average 96.1%. The discrepancy in HIC trends between these three periods offers a natural experiment to examine the impact of HIC on socioeconomic disparities in mortality.

Few studies have examined the impact of different health insurance status on inequalities in mortality and even less is known about inequalities in mortality in middle-income countries, many of whom introduced major health care system reforms during the last decades. As in Colombia the increase of HIC towards universal coverage was deliberately addressed to the poorest population ^{4 5}, and educational level is a good proxy variable for Socioeconomic Status (SAS) ¹³, we hypothesize that increased HIC contributed to reducing inequalities in cancer mortality. Cancer mortality is a particularly interesting indicator of

the effect of insurance coverage due to the expected increased access to prevention measures, early detection, timely treatment therapies, and high-cost interventions¹⁴⁻¹⁶.

Indeed, previous analyses of time trends of differences in cancer mortality by attained educational level in Colombia (1998-2007) were promising, particularly for gastric and cervical cancer where faster declines in mortality occurred in the lowest educated groups and therefore socioeconomic differences were expected to decrease in the near future³. In this paper we evaluate if the seemingly positive trend in cancer mortality by educational level continued, by using a slightly different definition of educational level which allowed us to extend the previous analyses (up to 2007) by 5 extra years, covering 1998-2012.

Methods

Data

Education level criteria

Educational level was defined as the highest level in which the individual has been enrolled during his life (i.e. the person accessed but not necessarily graduated this level), and was categorized in three groups based on the highest educational level accessed by the deceased: (a) Primary school or less, (b) Secondary school, and (c) Tertiary (post-secondary education). In previous papers educational level has been used, based on the highest educational level attained (i.e. completed) by the deceased¹³ but this category is restricted to the period 1998-2007. In our calculations we found both approaches to be similar in terms of the results yielded.

Deaths

National mortality data for cancer deaths (International Classification of Diseases (ICD-10) codes C00-C96, see Appendix-Table 1) for the period 1998-2012 were obtained from the National Administrative Department of Statistics (DANE), which routinely registers information on sex, date and cause of death as well as educational level from death certificates. Although the exact wording for the variable "accessed education" changed in the DANE database since 2008, the new categories were consistent with those of the databases of previous years and could be used. Trends of counts were found to be continuous and regular.

We focused on adult premature mortality (mortality below age 65) because it is known that information on educational level from mortality statistics is unreliable at ages 65 and beyond¹⁷. Additionally, premature mortality is an indicator of population health believed to be strongly influenced by social, economic and environmental factors¹⁸, and is a common indicator of health system performance¹⁹.

Data on age and sex were available for more than 99% of all deaths, while data on educational level was missing for approximately 16.5% of 190,993 cancer deaths. We used multiple imputation methods²⁰ implemented in SAS through the IMPUTE procedure to impute educational level for these cases in order to avoid bias due to the potentially higher rates of missing education for lower educated individuals, and to minimize the potential for numerator/denominator bias¹⁷. In short, this procedure fits a sequence of regression models and draws values from the corresponding predictive distributions. The

sequential regression procedure was applied based on a model that included sex, region, urban/rural residence and marital status as covariates. Details of this method are explained elsewhere²⁰. The imputation procedure was successful in 90.8% of cases with missing information, resulting in a total of 188,091 (98.5%) cancer deaths for our analyses.

Population

Data on mid-year population counts by age, sex and educational level were obtained from the Colombian Demography Health Surveys (DHS)²¹, which contain periodical information on the distribution of education by age, sex and calendar year (1995, 2000, 2005 and 2010). Age 20 was chosen as the lower age limit of this study as almost 100% of individuals accessed their highest educational level by this age.

The resulting proportions of individuals in each educational level were multiplied with the total population numbers per year, age and sex which were obtained from census combined with statistical projections from DANE²². These two information sources were combined to estimate the annual population size in each educational group (Appendix-Figure 1). We performed demographic projections to obtain population counts for years in-between every lustrum using the demographic Software PASEX²³. Additional details on the procedure are available elsewhere²³.

Analysis

All analyses were conducted in each of the five multiple databases generated by the multiple imputation process. Since results were nearly identical for all imputations, we used standard techniques as implemented in the PROC MIANALYZE procedure in SAS to combine estimates from all databases and adjust standard errors to account for uncertainty in the imputation²⁴. This procedure reads the parameter estimates and associated covariance matrix for each imputed data set, and then derives valid multivariate inferences for these parameters. This allows for valid statistical inference that appropriately reflects uncertainty due to missing values²⁴.

All analyses were conducted in SAS® version 9.2.

Age Standardised Mortality Rates (ASMR)

Data on population counts were combined with data on deaths to obtain a complete database of death counts by educational level, sex and five-year age group. We first calculated age-standardized mortality rates by educational level, sex and year using the

World Health population of 1997²⁵ as standard population, resulting in Age Standardised Mortality Rates (ASMR) expressed per 100,000 person-years (Figure 2). We also calculated ASMR by sex and year (Figure 2) and by educational level and sex (Table 1).

Annual trends in ASMR by sex and educational level were quantified by calculating the estimated Annual Percentage Change (EAPC) which measures the average rate of change in the mortality rate per year (negative EAPC: decreasing trend, positive EAPC : increasing trend). To test whether an apparent change in mortality trends was statistically significant, we used joinpoint regression, which fits a series of joined straight lines to age-adjusted rates and uses a Monte Carlo Permutation method to identify the best-fitting point (called joinpoint, year in which a significant change in the mortality trend occurred), where the rate of increase or decrease changes significantly²⁶. EAPC and joinpoints were determined based on the log-transformed ASMRs and their standard errors. We specified a maximum of 2 joinpoints with at least 4 observation points to either extreme of the data using joinpoint modelling based on the Joinpoint program²⁷ (Figure 2).

Regression models

We implemented separate Poisson regression models separately by sex with number of deaths as dependent variable and the natural log of person-years as offset variable, incorporating age and educational level as independent variables. We first estimated Rate Ratios (RR) of mortality by educational level, which compared the mortality of all educational groups to the mortality in the tertiary education group. Changes in the rate ratio over time result from changes in both risks and the distribution of educational level²⁸. To assess changes in disparities 'controlling' for changes in the educational distribution, we estimated the Relative Index of Inequality (RII), regressing mortality on the mid-point of the cumulative distribution of education, thereby taking into account the size of each educational group^{28 29}. Values higher than 1 indicate educational inequalities favouring the higher educated.

We evaluated if RRs significantly changed over time and within periods on continuous scales, which was not the case (results not shown). We calculated RII for three 5-year periods (1998-2002, 2003-2007, and 2008-2012) and for three age groups (20-34, 35-49, and 50-64) in order to identify differences in inequalities along age (Figure 2).

To compare the contribution of each cause of death to mortality disparities for non-communicable diseases (classified by their ICD-10 codes, see Appendix-table 1) we calculated the Slope Index of Inequality (SII) (Figure 4). The SII is a measure of absolute disparities that represents the difference in mortality between the population at the top

and the bottom of the educational distribution ²⁸. Further details on the RII and SII are available elsewhere ²⁸.

For peer review only

Results

Table 1 shows counts of cancer deaths and population at ages 20-64 years between 1998 and 2012 in Colombia. After excluding non-imputed cases (1.5%), 106,199 cancer deaths occurred from 1998 to 2012 among women while 81,892 occurred among men, with age-standardized cancer mortality rates (ASMR, per 100,000 person-years) of 58 among women and 49 among men. Men and women with lower levels of education had higher cancer mortality rates than their higher-educated counterparts. In addition, ASMR were larger among women than among men in each educational level.

The risk of dying was significantly and consistently higher among the lower educated. The RRs show clearly and statistically significantly increased risks of dying among the lower educated compared to the highest educated, this tendency was stronger in women ($RR_{primary}=1.49$; $RR_{secondary}=1.22$, both $p<0.0001$) than in men ($RR_{primary}=1.35$; $RR_{secondary}=1.11$ both $p<0.0001$). In order to formally test this higher female risk, we calculated Rate Ratios between sexes by educational levels, using men as reference category. For all educational levels, women had significant larger rate ratios (results not shown). We also found a consistent and slight increase in rate ratios from first, to second, and then to the third period (appendix-table 2) among both men and women and for primary and secondary education compared to higher educated, but the confidence intervals overlap, indicating the differences do not reach statistical significance.

During the 15 years of observation, the general tendency was for age-standardized cancer mortality rates to decline, with recent stabilizations, reaching an average ASMR in 2012 of 47.3 per 100,000 males and 55.2 per 100,000 females. Average EAPC for males in the periods defined by joinpoint were: 1998-2000: 2.4% (95% CI -2.0; 7.0), 2000-2012: -0.92% (95% CI -1.2; -0.7); for females 1998-2003: 0.7 (95% CI -0.7; 2.1), 2003-2006: -2.8% (95% CI -8.2; 2.9), 2006-2012 -0.6 (95% CI -1.5; 0.3). The general tendency for a declining trend in mortality was not reflected equally by educational level and sex (figure 2). Women had higher mortality rates than men. No significant increases or declines over time were observed in the lowest educated groups, whereas in the middle and high educated groups there was at least one period with significantly declining cancer mortality rates. In these groups, tendencies were for mortality to decrease substantially in the period after 2002/2003, with the exception of the highest educated males who showed recent increases. The strongest declines in cancer mortality over time were observed in the group of women with secondary or tertiary education. The most consistent declines in cancer mortality rates for all educational levels in both sexes were observed in the period 2003-2007, the period with rapid increases in HIC, particularly among the poor.

RII initially decreased from 1998-2002 to 2003-2007 (reaching significance among women), but then in the period 2008-2012 they increased back to its previous levels

(Figure 3, left panel). Trends in the joinpoint for RII are in line with this finding: no significant change was found for educational inequalities among men, whereas there was an initial significant decline (-7.6%) among women up to 2004 followed by a significant increase (+3.8%) in the period 2004-2012 (Appendix-Figure 2). RII also showed that inequalities were significantly higher for the younger age categories compared to older ones and tended to be higher for women (Figure 3, right panel).

Figure 4 shows the contribution to absolute differences in ASMR by education as measured by the SII for non-communicable diseases (appendix-table 1). Cancer was the second largest cause, after cardiovascular disease: explaining 20% of male differences (SII=20.5 deaths per 100,000 population) and 24% of female differences (SII=28.5). Absolute differences in cancer mortality were 39% larger for women than for men (28.5 vs. 20.5), while this difference was only 14% for all non-communicable diseases (SII= 103.2 and 120.3 respectively).

Discussion

We observed strong and persistent educational differences in cancer mortality rates in Colombia. These differences appeared to be diminishing in the period 2002-2007³ when healthcare insurance coverage largely grew, especially for the most disadvantaged part of the population. But in recent years, despite the achievement of almost universal HIC, inequalities were widening, particularly among women, with rates in the lowest educational groups being stable or increasing and those in the higher educated groups gradually declining. This means that the lowest educated part of the population is increasingly lagging behind in the progress against cancer, despite the increase in insurance coverage.

Explanation of results

Educational level was found to be previously as a good proxy variable for Socioeconomic Status (SES)¹³. Lower SES is associated with a higher prevalence of cancer risk factors (such as smoking, alcohol, obesity, occupational risk factors, housing circumstances), and less healthcare utilization (because of lower resources but also lower awareness and poorer access due to longer distances or high impact of family income when taking time off to visit a doctor)³⁰⁻³².

Not only absolute cancer mortality rates, but also inequalities in mortality were largely higher for women than for men. In a previous paper we showed this to be almost entirely attributable to cervical cancer mortality, which accounted for 51% of inequalities in total female cancer mortality in 1998-2007 at 25-64³. The consistently higher RR for women than men are unique to cancer in Colombia; studies evaluating other causes of death consistently show a higher mortality risk for men¹², which also coincides with the usually lower life expectancy of men. This illustrates the high burden of certain cancers which are strongly SES related, probably mainly cervical and breast cancer, which, if diagnosed late, have a very poor prognosis even though early detection possibilities are ample. Several Colombian studies have identified that both characteristics of the health system such as access and delays in diagnosis and treatment, and characteristics of the populations of lower socioeconomic classes contribute to the higher burden of cervical cancer mortality^{33 34}. A previous study showed that lower access to mammography screening for early detection of breast cancer was associated with lower education, affiliation to the subsidized regime or being unaffiliated³⁵. In general, financial access to healthcare has been more limited to women^{36 37} which potentially limits access to expensive cancer therapies, particularly for most deprived women.

We also observed clearly increased educational differences in cancer mortality for the youngest age groups, in which mortality from cancer is a relatively rare event. Among males, leukaemias, stomach cancers and Non-Hodgkin lymphomas are important cancer causes of deaths in the age groups 15-39 years (representing 44% of all cancer deaths in this age group); for females these include cervical cancer and leukaemias as well as non-Hodgkin lymphomas (representing 38%)³⁸. Previous studies have shown that mortality or survival of these cancer types is very much dependent on socioeconomic status^{3 14 39}. As these tumours explain an important proportion of cancer deaths in these ages, this probably explains the high RII in the younger age groups as compared to the higher age categories. Nevertheless we cannot discard that, at least partially, the strong inequalities in younger groups are due to the fact that, when rates are low, relative inequalities tend to show an increasing pattern⁴⁰.

In order to test if RII is a good indicator of changes in inequalities⁴⁰, we compared trends in relative and absolute measures of inequalities (RII and SII), which were almost identical (results not shown). Mackenbach hypothesized that, in the case of declining mortality rates, RIIs are exaggerating the differences⁴⁰, but this was clearly not the case in our study, probably because reductions in premature mortality rates of cancer were relatively smooth, and not very strong and not very divergent between educational levels, with the exception of tertiary educated women. Among women we only found a clear reduction in inequalities during the period 1998-2004, owing to a clearly divergent pattern between educational levels: A rapid significant decrease of ASMRs in the higher socioeconomic group (tertiary educated) and an opposed rapid significant increase of ASMRs among those with secondary education. We have illustrated in previous work that these trends are most likely due to large changes in Cervical cancer mortality³.

Our results clearly illustrate that an almost complete coverage does not necessarily reduce inequalities in health^{41 42}. Particularly, having health insurance may be universal, but depending on income the type of health insurance is different (subsidized, contributory and special or exceptional regimes), with the wealthy population often buying additional private health assurance to ensure rapid and more broad access⁴³. The quality of care provided by the insurance (translating into early and timely diagnosis and adequate treatment) is not guaranteed with complete coverage, as seems to be the case: To warrant rights to get access to expensive treatments and medication, as usual in cancer, exceptional legal mechanisms are frequently launched in Colombia: Technical-scientific committees of health assurers, and an action for protection so-called "tutela"^{43 44}. A study shows that those affiliated to the contributory regime are more likely to warrant additional rights more efficiently⁴⁴, potentially increasing inequalities between regimes. Also, clear differences in gastric cancer survival by type of health insurance affiliation have been documented in a population-based Colombian study, clearly illustrating that, even

though in theory there is access to care for all, this does not translate to equal outcomes⁴⁵.

In parallel with the increases in HIC, other changes in the Colombian health care system occurred, including increases in investments in care^{5 46}. Therefore, our results reflect the impact of the entire reform rather than only the increased coverage in health insurance. However, based on our analyses, it is safe to conclude that all these reforms have not resulted in reducing inequalities in cancer mortality. This is not unique to Colombia; in several countries, as diverse as Taiwan, Thailand and European countries, inequalities in health increased upon reaching almost universal coverage⁴⁷⁻⁵².

Strengths and weaknesses

This is one of few studies investigating educational differences in mortality using population-based mortality and population-based educational level. Despite several strengths, some limitations should be considered in our study: We did not correct for misclassification within the cancer groups, but since we studied the group of cancer deaths overall, regardless of the subtype of cancer, most of these errors are cancelled out. In general “cancer” as cause of death is correctly coded⁵³, and most likely particularly in setting of ages under 65.

Data on mortality were obtained from mortality registries, while data on the population distribution by education were obtained from censuses and demographic projections. This may have led to the so-called numerator/denominator bias, which generally results in an overestimation of disparities¹⁷. Furthermore, for some years, data on population size were obtained from demographic projections combined with distributions of education from surveys. To assess the impact of this potential bias, we experimented with different education distributions from multiple data sources⁵⁴⁻⁵⁶, showing that the overall trends in our study were robust to different assumptions on the distribution of education³.

Potentially, there has been substantial under-registration of deaths in some regions. Previous studies comparing national mortality rates with indirect estimates from census⁵⁷⁵⁸ suggest that underregistration is particularly important in the poorest regions, implying that estimates of disparities in mortality by educational level are likely to be an underestimation, because those with lower education are more likely to live in areas with higher underregistration rates. This may also have led to underestimation of the extent to which inequalities have increased, because underregistration decreased over the study period^{57 58}. Our results, therefore, are indicative of potentially larger inequalities in mortality by education, which continue to increase.

Information on education was missing for 16.5% of cancer deaths, potentially leading to an underestimation of disparities, as missing values are usually more common in the least educated⁵⁹. However, we imputed values for educational level for individuals with missing educational information based on a information on age, sex marital status, region and urban/rural residence, thereby limiting the potential impact of this source of bias.

Unanswered questions and future research

Socioeconomic inequalities in mortality of non-communicable diseases have been associated with the unequal distribution of behavioural risk factors, such as smoking, alcohol consumption, an unhealthy diet, sedentary lifestyle higher risk of injuries^{2 60}. This implies underlying inequalities in exposure to risk factors and incidence of these diseases. However, data on the incidence of disease and prevalence distribution of all those behavioural and poverty risk factors by socioeconomic status are largely lacking in Colombia and most developing countries. Studies on the mechanisms underlying the way these risk factors contribute to inequalities in mortality are scarce, even in developed countries. We previously described not only large inequalities in mortality of cancers with a primary infectious aetiology such as gastric cancer³, but also documented inequalities in survival of gastric cancer measured by both type of health insurance and socioeconomic stratum¹⁴. These findings indicate that both access and quality of care (health insurance) as other factors related to socioeconomic status independently affect survival for the lower socioeconomic classes negatively. Detailed studies are needed on both types of factors: which barriers exist for timely diagnosis and adequate treatment in situations with universal health coverage and how can they be tackled? And on which other factors related to socioeconomic stratum can interventions be designed? The answers to these questions will be various and, depending on the specific disease, will be quite different in character. It is of high importance that data on socioeconomic indicators are collected in a standard way, so that for example cancer registries can calculate incidence rates by socioeconomic stratum, which, at this moment, is not possible in Colombia⁶¹.

Socioeconomic differences were substantially larger for women than for men, in our previous study this difference was almost entirely attributable to inequalities in cervical cancer mortality³. In this more recent study we see that the gender inequalities grew despite the virtually universal HIC since 2008. Gender inequalities were larger for cancer than for other non-communicable diseases according SII; more detailed analyses are needed to assess to which extent the growing inequalities are due to an increase in cervical cancer mortality amongst less favoured.

Conclusions

The recent negative curbs in mortality among the lowest educated groups, resulting in increasingly differences between educational groups, are a call for attention by the Colombian health authorities to take measures in both primary prevention, access to early detection and timely and adequate treatment. Underlying causes of the observed differences are multiple, from differences in housing, lifestyle, health-related beliefs, to differences in access to diagnostic facilities and high-quality treatment, which all should be of concern to those in charge ⁶². Insurance coverage is clearly not enough to counterbalance the deleterious effects of decreases in spending on primary prevention and other social and economic conditions which determine the growing levels of educational inequalities in cancer.

Acknowledgements

We are grateful to Mauricio Avendaño and Alex Burdorf for their contributions at initial stages of the analyses and for their continued support in this project.

References

1. Arroyave I, Cardona D, Burdorf A, Avendaño M. The Impact of Increasing Health Insurance Coverage on Disparities in Mortality: Health Care Reform in Colombia, 1998–2007. *American Journal of Public Health* 2013:e1-e7.
2. Arroyave I, Cardona D, Burdorf A, Avendaño M. Socioeconomic inequalities in premature mortality in Colombia, 1998-2007: The double burden of non-communicable diseases and injuries. *Preventive Medicine* 2014(64C):41-47.
3. de Vries E, Arroyave I, Pardo EC, Wiesner C, Murillo R, Forman D, et al. Trends in inequalities in premature cancer mortality by educational level in Colombia, 1998-2007. *Journal of Epidemiology & Community Health* 2015;69(5):408-15.
4. Agudelo-Calderón CA, Cardona-Botero J, Ortega-Bolaños J, Robledo-Martínez R. Sistema de salud en Colombia: 20 años de logros y problemas. *Ciência & Saúde Coletiva* 2011;16:2817-28.
5. Giedion U, Uribe MV. Colombia's Universal Health Insurance System. *Health Affairs* 2009;28(3):853-63.
6. Céspedes-Londoño JE, Jaramillo-Pérez I, Castaño-Yepes RA. Impacto de la reforma del sistema de seguridad social sobre la equidad en los servicios de salud en Colombia. *Cadernos de Saúde Pública* 2002;18(4):1003-24.
7. Nieto-Parra S, Olivera M. Making Reform Happen in Colombia: The Process of Regional Transfer Reform. *OECD Development Centre Working Papers: Organisation for Economic Co-operation and Development (OECD)*, 2012:39.
8. Ministerio de la Protección Social. Informe de Actividades 2009-2010 - Informe Cuatrienio 2006 - 2010, al Honorable Congreso de la República. In: Dirección General de Planeación y Análisis de Política, Grupo de Estudios y Planeación Sectorial, editors. Bogotá D.C., 2010.
9. Ruiz F, Amaya L, Venegas S. Progressive segmented health insurance: Colombian health reform and access to health services. *Health Economics* 2007;16(1):3-18.
10. Miller G, Pinto DM, Vera-Hernández M. Risk Protection, Service Use, and Health Outcomes Under Colombia's Health Insurance Program for the Poor. *American Economic Journal: Applied Economics* 2013;5(4):61-91.
11. Abadia CE, Oviedo DG. Bureaucratic Itineraries in Colombia. A theoretical and methodological tool to assess managed-care health care systems. *Social Science & Medicine* 2009;68(6):1153-60.
12. De Groote T, De Paepe P, Unger J-P. Colombia: in vivo test of health sector privatization in the developing world. *International Journal of Health Services* 2005;35(1):125 - 41.
13. Oakes JM, Rossi PH. The measurement of SES in health research: current practice and steps toward a new approach. *Social Science & Medicine* 2003;56(4):769-84.
14. de Vries E, Uribe C, Pardo C, Lemmens V, Van de Poel E, Forman D. Gastric cancer survival and affiliation to health insurance in a middle-income setting. *Cancer Epidemiology* 2014.
15. Smith JK, Ng SC, Zhou Z, Carroll JE, McDade TP, Shah SA, et al. Does increasing insurance improve outcomes for US cancer patients? *Journal of Surgical Research* 2013;185(1):15-20.
16. Kim S, Kwon S. Has the National Health Insurance improved the inequality in the use of tertiary-care hospitals in Korea? *Health Policy* 2014;118(3):377-85.
17. Kunst AE, Groenhouf F, Borgan J, Costa G, Desplanques G, Faggiano F, et al. Socio-economic inequalities in mortality. Methodological problems illustrated with three examples from Europe. *Revue d'épidémiologie et de sante publique* 1998;46(6):467-79.
18. World Health Organization. Closing the gap in a generation: Health equity through action on the social determinants of health. Geneva, Switzerland: World Health Organization, 2008.

19. Smith PC. *Performance measurement for health system improvement: experiences, challenges and prospects*: Cambridge University Press, 2010.
20. Raghunathan T, Lepkowski J, Van Hoewyk J, Solenberger P. A Multivariate Technique for Multiply Imputing Missing Values Using a Sequence of Regression Models. *Survey Methodology* 2001;27(1):85-95.
21. Profamilia. Encuesta Nacional de Demografía y Salud del 2010 (ENDS 2010), 2012.
22. Departamento Administrativo Nacional de Estadística (DANE). Colombia: Tablas de mortalidad nacionales y departamentales 1985-2020. Sep. 2007 ed. Bogotá D.C., 2007.
23. Population Analysis System (PASEX) [program]. 2.04g version. Washington DC: International Programs Center (IPC) - United States Census Bureau, 2011.
24. SAS Institute I. The MIANALYZE Procedure. In: Inc. SI, editor. *User's Guide SAS/STAT® 9.2*. Version 8 ed. Cary, NC: SAS Institute Inc., 2008:201-33.
25. Ahmad OB, Boschi-Pinto C, Lopez AD, Murray CJ, Lozano R, Inoue M. Age standardization of rates: A new WHO standard. *GPE Discussion Paper Series*. Geneva: World Health Organization, 2001.
26. Joinpoint Regression Program [program]. 4.0.4 version, 2013.
27. Kim HJ, Fay MP, Feuer EJ, Midthune DN. Permutation tests for joinpoint regression with applications to cancer rates. *Statistics in Medicine* 2000;19(3):335-51.
28. Mackenbach JP, Kunst AE. Measuring the magnitude of socio-economic inequalities in health: An overview of available measures illustrated with two examples from Europe. *Social Science & Medicine* 1997;44(6):757-71.
29. Harper S, Lynch J, Meersman SC, Breen N, Davis WW, Reichman ME. An overview of methods for monitoring social disparities in cancer with an example using trends in lung cancer incidence by area-socioeconomic position and race-ethnicity, 1992-2004. *Am J Epidemiol* 2008;167(8):889-99.
30. Arias SA. Inequidad y cáncer: una revisión conceptual. *Revista de la Facultad Nacional de Salud Pública* 2009;27(3):341-48.
31. Woods L, Rachet B, Coleman M. Origins of socio-economic inequalities in cancer survival: a review. *Annals of Oncology* 2006;17(1):5-19.
32. Auvinen A, Karjalainen S. Possible explanations for social class differences in cancer patient survival. In: Kogevinas M, Pearce N, Susser M, Boffetta P, editors. *Social Inequalities and Cancer*. Lyon, 1997:377-97.
33. Piñeros M, Cendales R, Murillo R, Wiesner C, Tovar S. Cobertura de la Citología de Cuello Uterino y Factores Relacionados en Colombia, 2005. *Revista de Salud Pública* 2007;9(3):327-41.
34. Wiesner-Ceballos C, Cendales-Duarte R, Murillo R, Piñeros M, Tovar-Murillo SL. Seguimiento de mujeres con anormalidad citológica de cuello uterino, en Colombia. *Revista de Salud Pública* 2010;12(1):1-13.
35. Charry LCd, Carrasquilla G, Roca S. Equidad en la Detección del Cáncer de Seno en Colombia. *Revista de Salud Pública* 2008;10(4):571-82.
36. Ewig C, Bello AH. Gender equity and health sector reform in Colombia: Mixed state-market model yields mixed results. *Social Science & Medicine* 2009;68(6):1145-52.
37. Guarnizo-Herreño CC, Agudelo C. Equidad de Género en el Acceso a los Servicios de Salud en Colombia. *Revista de Salud Pública* 2008;10(1):44-57.
38. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, et al. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. Lyon, France: International Agency for Research on Cancer, 2013.

39. Petridou ET, Sargentanis T, Perlepe C, Papathoma P, Tsilimidos G, Kontogeorgi E, et al. Socio-economic disparities in survival from childhood leukemia in the USA and globally: a meta-analysis. *Annals of Oncology* 2014;26(3):589-97.

40. Mackenbach JP. Should we aim to reduce relative or absolute inequalities in mortality? *The European Journal of Public Health* 2015;25(2):185-85.

41. Franks P, Clancy CM, Gold MR. Health insurance and mortality: Evidence from a national cohort. *JAMA: The Journal of the American Medical Association* 1993;270(6):737-41.

42. Katz SJ, Hofer TP. Socioeconomic disparities in preventive care persist despite universal coverage. Breast and cervical cancer screening in Ontario and the United States. *JAMA* 1994;272(7):530-4.

43. Guerrero R, Gallego AI, Becerril-Montekio V, Vásquez J. Sistema de salud de Colombia. *Salud Pública de México* 2011;53:s144-s55.

44. Uprimny R, Durán J. Equidad y protección judicial del derecho a la salud en Colombia. *Políticas Sociales*. Santiago de Chile: Comisión Económica para América Latina y el Caribe (CEPAL), 2014:70.

45. de Vries E, Uribe C, Pardo C, Lemmens V, Van de Poel E, Forman D. Gastric cancer survival and affiliation to health insurance in a middle-income setting. *Cancer epidemiology* 2015;39(1):91-96.

46. Restrepo Zea JH. El seguro de salud en Colombia: ¿cobertura universal? *Revista Gerencia y Políticas de Salud* 2002(Nº 2):25-40.

47. Shou-Hsia C, Tung-Liang C. The Effect of Universal Health Insurance on Health Care Utilization in Taiwan. *JAMA: The Journal of the American Medical Association* 1997;278(2):89-93.

48. Alvarez L, Salmon JW, Swartzman D. The Colombian health insurance system and its effect on access to health care. *Int J Health Serv* 2011;41(2):355-70.

49. Suraratdech C, Saithanu S, Tangcharoensathien V. Is universal coverage a solution for disparities in health care?: Findings from three low-income provinces of Thailand. *Health Policy* 2005;73(3):272-84.

50. Mackenbach JP, Stirbu I, Roskam A-JR, Schaap MM, Menvielle G, Leinsalu M, et al. Socioeconomic Inequalities in Health in 22 European Countries. *New England Journal of Medicine* 2008;358(23):2468-81.

51. Stirbu I, Kunst AE, Bopp M, Leinsalu M, Regidor E, Esnaola S, et al. Educational inequalities in avoidable mortality in Europe. *Journal of Epidemiology and Community Health* 2010;64(10):913-20.

52. Avendaño M, Kunst AE, Huisman M, van Lenthe F, Bopp M, Regidor E, et al. Socioeconomic status and ischaemic heart disease mortality in 10 western European populations during the 1990s. *Heart* 2006;92(4):461-67.

53. Harteloh P, de Bruin K, Kardaun J. The reliability of cause-of-death coding in The Netherlands. *European journal of epidemiology* 2010;25(8):531-38.

54. International Institute of Applied System Analysis (IIASA), Vienna Institute of Demography of the Austrian Academy of Sciences (VID). IVEP-1.0 (IIASA/VID education database). Viena, 2010.

55. World Bank. World Development Indicators (WDI). *World Bank Data (WBD)*: World Bank 2011.

56. Barro RJ, Lee J-W. A new data set of educational attainment in the world, 1950–2010. *NBER Working Paper Series*. Cambridge, MA: National Bureau of Economic Research, 2010.

57. Patiño-Villada FA, Arango-Vélez EF, Quintero-Velásquez MA, Arenas-Sosa MM. Factores de riesgo cardiovascular en una población urbana de Colombia. *Revista de Salud Pública* 2011;13:433-45.

- 1
2
3 58. Storr CL, Cheng H, Posada-Villa J, Aguilar-Gaxiola S, Anthony JC. Adult smokers in Colombia:
4 who isn't giving it up? *Addict Behav* 2008;33(3):412-21.
5
6 59. Rodríguez-García J. Desigualdades socioeconómicas entre departamentos y su asociación con
7 indicadores de mortalidad en Colombia en 2000. *Revista Panamericana de Salud Pública*
8 2007;21:111-24.
9
10 60. Adler NE, Newman K. Socioeconomic Disparities In Health: Pathways And Policies. *Health*
11 *Affairs* 2002;21(2):60-76.
12
13 61. De Vries E, Murillo R. Regarding articles about Cali Cancer Registry 2. *Salud Pública de México*
14 2015;57(3).
15
16 62. Ottersen OP, Dasgupta J, Blouin C, Buss P, Chongsuvivatwong V, Frenk J, et al. The political
17 origins of health inequity: prospects for change. *The Lancet* 2014;383(9917):630-67.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURES AND TABLES

Figure 1: Trends in national healthcare insurance coverage (HIC), Colombia, 1998-2012

Other schemes include primarily members of the military and teacher and oil workers syndicate members. Based on Annual reports of the Ministry of Health and Social Protection

Source: Annual reports of the Ministry of Health and Social Protection to the Congress of the Republic of Colombia

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 1: Descriptives of the study: deaths, population and rates separately by sex and educational level, Colombia, 1998-2012

Sex	Educational level ^a	Cancer deaths ^b	Population size	ASMR ^c (95% Confidence Intervals)	
Men	Primary	50,126	67,815,336	54.0	(53.5, 54.5)
	Secondary	22,273	67,742,806	45.1	(44.4, 45.8)
	Tertiary	9,493	30,715,631	41.4	(40.5, 42.3)
	Total	81,892	166,273,773	48.9	(48.8, 49.1)
Women	Primary	65,336	68,637,666	64.7	(64.2, 65.2)
	Secondary	30,414	72,387,214	54.2	(53.5, 54.9)
	Tertiary	10,449	35,149,442	47.2	(46.2, 48.3)
	Total	106,199	176,174,322	57.8	(57.6, 58.0)

^a: Educational attainment values registered after final imputation. Primary = up to elementary or primary school; secondary = any level of high-school; tertiary = any level of postsecondary education after high school including college and university.

^b: Cancer deaths after imputation.

^c: ASMR: Age standardized mortality rates per 100,000 population; Estimates (WHO standard population 1997) for educational level combine results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Figure 2: Age-standardized cancer mortality trends, including APC based on joinpoint models, by sex and educational level

Markers: Observed age-standardized cancer mortality rates. Lines: modelled age-standardized cancer mortality trends. Estimates are results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values. The points represent ASMR, lines represent the trendlines between joinpoints. Blue, diamonds: maximum primary education; Red, triangles: secondary education; Green, circles: tertiary education. Numbers adjacent to the lines represent estimates annual percent change (EAPC) during the corresponding periods, based on joinpoint modelling; a star indicates statistical significance at α 0.05.

Figure 3: Sex-specific Relative Index of Inequality by period and age group

Estimates from combined results from 5 databases resulting from multiple imputations, appropriately reflecting uncertainty attributable to missing values.

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 4. Slope Index of Inequality (SII) for non-communicable diseases

See Appendix-Table 1 for International Classification of Diseases (ICD-10) codes for non-communicable diseases

For peer review only




Figure 1: Trends in national healthcare insurance coverage (HIC), Colombia, 1998-2012

Other schemes include primarily members of the military and teacher and oil workers syndicate members.
Based on Annual reports of the Ministry of Health and Social Protection
Source: Annual reports of the Ministry of Health and Social Protection to the Congress of the Republic of Colombia

202x100mm (150 x 150 DPI)




Figure 2: Age-standardized cancer mortality trends, including APC based on joinpoint models, by sex and educational level

Markers: Observed age-standardized cancer mortality rates. Lines: modelled age-standardized cancer mortality trends. Estimates are results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values. The points represent ASMR, lines represent the trendlines between joinpoints. Blue, diamonds: maximum primary education; Red, triangles: secondary education; Green, circles: tertiary education. Numbers adjacent to the lines represent estimates annual percent change (EAPC) during the corresponding periods, based on joinpoint modelling; a star indicates statistical significance at α 0.05.

227x115mm (150 x 150 DPI)




Figure 3: Sex-specific Relative Index of Inequality by period and age group

Estimates from combined results from 5 databases resulting from multiple imputations, appropriately reflecting uncertainty attributable to missing values.

163x71mm (150 x 150 DPI)

review only




Figure 4: Slope Index of Inequality (SII) for non-communicable diseases

See Appendix-Table 1 for International Classification of Diseases (ICD-10) codes for non-communicable diseases

112x66mm (150 x 150 DPI)

APPENDIX

Appendix-Figure 1. Educational level trends over time, Colombia, 1998-2012

The thicker markers are those from the year with real data available (source: Colombian Demography Health Surveys in 1995, 2000, 2005 and 2010). The points in between were interpolated (see methods)

Appendix-Table 1. Classification of Causes of death, International Classification of Diseases (ICD-10) codes for non-communicable diseases

Non-Communicable Diseases		ICD-10 codes
Cardiovascular diseases		I00-I99
Malignant neoplasm		C00-D48
Diabetes mellitus		E10-E14
Chronic lower respiratory diseases (CLRD)		J40-J47
Other Non-Communicable Diseases	Mental and behavioural disorders	F00-F99
	Diseases of the nervous system	G00-G99
	Diseases of the digestive system	K00-K93
	Diseases of the genitourinary system	N00-N99
	Congenital malformations, deformations and chromosomal abnormalities	Q00-Q99


Appendix-Figure 2: Sex-specific Relative Index of Inequality (RII) trends including APC based on joinpoint models

Markers: Annual observed cancer mortality RII. Lines: modelled RII trends. Estimates are results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values.

For peer review only

Appendix-Table 2. Rate ratio in premature cancer mortality by educational level by 5-years periods and separately for men and women at 20–64 years, Colombia 1998-2012


		Period 1 (1998-2002): Relatively low coverage	Period 2 (2003-2007): Rapidly increasing coverage	Period 3 (2008-2012): Almost universal coverage
MEN	Tertiary (Ref)	1.00		
	Secondary	1.08 (0.89 , 1.30)	1.19 (1.14 , 1.24)	1.31 (1.14 , 1.52)
	Primary	1.32 (1.11 , 1.56)	1.37 (1.31 , 1.43)	1.42 (1.23 , 1.63)
	Year	1.01 (0.98 , 1.04)	0.99 (0.96 , 1.02)	1.02 (1.00 , 1.04)
	Secondary*Year	1.00 (0.96 , 1.03)	1.00 (0.97 , 1.04)	0.96 (0.94 , 0.99)
	Primary*Year	1.00 (0.96 , 1.03)	1.00 (0.97 , 1.03)	0.99 (0.96 , 1.02)
WOMEN	Tertiary (Ref)	1.00		
	Secondary	1.10 (0.91 , 1.33)	1.27 (1.22 , 1.32)	1.29 (1.13 , 1.47)
	Primary	1.25 (1.04 , 1.51)	1.44 (1.39 , 1.50)	1.45 (1.28 , 1.65)
	Year	1.03 (1.00 , 1.07)	0.98 (0.96 , 1.01)	1.00 (0.98 , 1.02)
	Secondary*Year	0.99 (0.96 , 1.03)	1.01 (0.98 , 1.04)	0.99 (0.97 , 1.02)
	Primary*Year	0.97 (0.94 , 1.01)	1.00 (0.97 , 1.02)	1.01 (0.98 , 1.03)



Appendix-Figure 1: Educational level trends over time, Colombia, 1998-2012

The thicker markers are those from the year with real data available (source: Colombian Demography Health Surveys in 1995, 2000, 2005 and 2010). The points in between were interpolated (see methods)

154x51mm (150 x 150 DPI)



Appendix- Figure 2: Sex-specific Relative Index of Inequality (RII) trends including APC based on joinpoint models

Markers: Annual observed cancer mortality RII. Lines: modelled RII trends. Estimates are results from 5 databases generated by multiple imputations, appropriately reflecting uncertainty attributable to missing values.

155x105mm (150 x 150 DPI)