

Online supplementary file 9

Table of weighted population prevalence's for calculations of sensitivity and specificity for any anxiety disorder (including PTSD and OCD) (missing PTSD data considered as PTSD cases)

	GAD-2 (<3) negative	GAD-2 (≥3) positive
No anxiety	7268 85% (80 – 89%)	723.1 62% (47 – 74%)
SCID any anxiety	1307 15% (11 – 20%)	451.3 38% (26 – 53%)
	GAD-2 (Yes/No) negative	GAD-2 (Yes/No) positive
No anxiety	5110 91% (85 – 94%)	2881 70% (62 – 77%)
SCID any anxiety	523 9% (6 – 15%)	1235 30% (23 – 38%)

Table of weighted population prevalence's for calculations of sensitivity and specificity for any anxiety disorder (including PTSD and OCD) (missing PTSD data considered as not PTSD cases)

	GAD-2 (<3) negative	GAD-2 (≥3) positive
No anxiety	7450 87% (82 – 90%)	770.8 66% (51 – 78%)
SCID any anxiety	1125 13% (10 – 18%)	403.6 34% (22 – 49%)
	GAD-2 (Yes/No) negative	GAD-2 (Yes/No) positive
No anxiety	5181 92%(87 – 95%)	3040 74% (66 – 80%)
SCID any anxiety	452.8 8% (5 – 13%)	1076 26 (20 – 34%)