

Supplemental file 2

Association between BBQ scores at baseline and LBP intensity and disability at 2, 13 and 52 weeks stratified by LBP history

| LBP intensity (NRS) | | | | | | | | | | | | | | | | | | | | | | | | |
|----------------------------------|-------------------|--------|------------------|------------------------|--------|------------------|-------------------|--------|-----------------|--------------------|-------|-----------------|--|--|--|--|--|--|--|--|--|--|--|--|
| | Acute new (n=209) | | | Acute episodic (n=932) | | | Sub acute (n=615) | | | Persistent (n=473) | | | | | | | | | | | | | | |
| Time | Coefficient | p | 95%CI | Coefficient | p | 95%CI | Coefficient | p | 95%CI | Coefficient | p | 95%CI | | | | | | | | | | | | |
| 2 weeks | -3.82 | <0.001 | -5.75 -1.89 | -2.76 | <0.001 | -3.76 -1.76 | -2.26 | 0.001 | -3.55 -0.97 | -1.004 | 0.104 | -2.35 0.34 | | | | | | | | | | | | |
| 13 weeks | -4.58 | <0.001 | -6.65 -2.52 | -3.59 | <0.001 | -4.62 -2.57 | -3.64 | <0.001 | -4.96 -2.32 | -1.80 | 0.009 | -3.17 -0.44 | | | | | | | | | | | | |
| 52 weeks | -6.73 | <0.001 | -8.8 -4.67 | -3.48 | <0.001 | -4.53 -2.43 | -3.12 | <0.001 | -4.45 -1.8 | -1.73 | 0.024 | -3.22 -0.23 | | | | | | | | | | | | |
| BBQ | -0.02 | 0.407 | -0.06 0.02 | -0.02 | 0.153 | -0.04 0.01 | -0.02 | 0.082 | -0.05 0.003 | -0.04 | 0.009 | -0.07 -0.01 | | | | | | | | | | | | |
| Interaction between BBQ and time | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 weeks | -0.01 | 0.686 | -0.07 0.05 | -0.03 | 0.087 | -0.06 0.004 | 0.002 | 0.919 | -0.04 0.04 | -0.02 | 0.330 | -0.06 0.02 | | | | | | | | | | | | |
| 13 weeks | -0.04 | 0.223 | -0.1 0.02 | -0.05 | 0.002 | -0.08 -0.02 | 0.01 | 0.753 | -0.03 0.05 | -0.02 | 0.342 | -0.06 0.02 | | | | | | | | | | | | |
| 52 weeks | 0.03 | 0.334 | -0.03 0.09 | -0.05 | 0.001 | -0.09 -0.02 | -0.01 | 0.580 | -0.05 0.03 | -0.02 | 0.329 | -0.07 0.02 | | | | | | | | | | | | |
| Interaction term | 0.2262 | | | 0.0025 | | | 0.8756 | | | 0.6618 | | | | | | | | | | | | | | |
| DISABILITY (RMDQ) | | | | | | | | | | | | | | | | | | | | | | | | |
| Time | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 weeks | -38.65 | <0.001 | -58.85 -18.44 | -34.47 | <0.001 | -44.33 -24.60 | -18.83 | 0.002 | -30.52 -7.13 | -2.08 | 0.722 | -13.51 9.36 | | | | | | | | | | | | |
| 13 weeks | -49.42 | <0.001 | -71.46 -27.38 | -44.11 | <0.001 | -54.47 -33.76 | -34.83 | <0.001 | -47.1 -22.56 | -16.21 | 0.009 | -28.41 -4.0 | | | | | | | | | | | | |
| 52 weeks | -47.51 | <0.001 | -70.78 -24.24 | -46.19 | <0.001 | -57.07 -35.31 | -40.62 | <0.001 | -53.65 -27.6 | -3.88 | 0.590 | -17.97 10.22 | | | | | | | | | | | | |
| BBQ | -0.35 | 0.127 | -0.81 0.10 | -0.56 | <0.001 | -0.79 -0.34 | -0.41 | 0.001 | -0.67 -0.16 | -0.4 | 0.002 | -0.65 -0.15 | | | | | | | | | | | | |
| Interaction between BBQ and time | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 weeks | 0.05 | 0.875 | -0.55 0.65 | 0.07 | 0.635 | -0.23 0.37 | 0.07 | 0.696 | -0.28 0.42 | -0.29 | 0.110 | -0.64 0.07 | | | | | | | | | | | | |
| 13 weeks | -0.07 | 0.838 | -0.71 0.58 | 0.02 | 0.912 | -0.3 0.33 | 0.22 | 0.235 | -0.14 0.59 | 0.02 | 0.912 | -0.36 0.4 | | | | | | | | | | | | |
| 52 weeks | -0.01 | 0.983 | -0.69 0.67 | 0.1 | 0.554 | -0.23 0.43 | 0.43 | 0.031 | 0.04 0.82 | -0.35 | 0.114 | -0.78 0.08 | | | | | | | | | | | | |
| Interaction term | 0.9895 | | | 0.9258 | | | 0.1571 | | | 0.1792 | | | | | | | | | | | | | | |

BBQ: Back Belief Questionnaire, CI: Confidence Interval, LBP: Low Back Pain, RMDQ: Roland Morris Disability Questionnaire

Analyses were controlled for age, sex, baseline LBP intensity, baseline RMDQ score and previous treatment.