




**Supplemental Figure 1.** Telehealth Modality Type among US Adults with Diabetes between Metro and Non-Metro Areas



**Supplemental Figure 2.** Reasons for Using or Not Using Telehealth Services among US Adults with Diabetes between Metro and Non-Metro Areas



**Supplemental Figure 3.** Patient-Reported Quality of Care Outcomes by Telehealth Use among US Adults with Diabetes between Metro and Non-Metro Areas



**Supplemental Table 1.** Patient-Reported Quality of Care Outcomes by Telehealth Modality

	By Video Only	By Phone Call Only	By Both	
Outcome	N=202	N=225	N=116	P-value
Overall Quality of Care				0.765
Excellent	21.9%	24.7%	25.0%	
Very good	38.7%	46.9%	35.6%	
Good	29.9%	16.9%	33.7%	
Fair	7.8%	9.7%	5.1%	
Poor	1.7%	1.7%	0.7%	
Trust in Health System				0.477
Very	39.7%	35.6%	51.4%	
Somewhat	45.3%	54.1%	41.7%	
A little/ not at all	15.0%	10.3%	6.9%	
Patient-Centered Communication Scale				0.144
1st Quartile (Poor)	23.1%	25.7%	20.7%	
2nd Quartile	27.8%	23.5%	8.4%	
3rd Quartile	20.2%	23.2%	41.7%	
4th Quartile (Best)	28.9%	27.5%	29.1%	